

---

---

**Mariana Islands  
Training and Testing Activities  
Draft Supplemental Environmental Impact  
Statement/Overseas Environmental Impact  
Statement**



**Volume 1**

**January 2019**

MITT SEIS/OEIS Project Manager  
Naval Facilities Engineering Command, Pacific  
258 Makalapa Dr., Suite 100  
Pearl Harbor, HI 96860-3134



# **DRAFT SUPPLEMENTAL ENVIRONMENTAL IMPACT STATEMENT/ OVERSEAS ENVIRONMENTAL IMPACT STATEMENT FOR MARIANA ISLANDS TRAINING AND TESTING**

**Lead Agency:** United States Department of the Navy  
**Cooperating Agency:** National Marine Fisheries Service  
**Title of the Proposed Action:** Mariana Islands Training and Testing  
Draft Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement

## **Abstract**

The United States Department of the Navy (Navy) prepared this Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement (SEIS/OEIS) in compliance with the National Environmental Policy Act (NEPA) of 1969 (42 United States Code section 4321 et seq.); the Council on Environmental Quality Regulations for Implementing the Procedural Provisions of NEPA (Title 40 Code of Federal Regulations sections 1500 et seq.); Navy Procedures for Implementing NEPA (32 Code of Federal Regulations section 775); and Executive Order 12114, *Environmental Effects Abroad of Major Federal Actions*. This SEIS/OEIS evaluates the potential environmental impacts of conducting training and testing activities after August 2020 in the Mariana Islands Training and Testing Study Area (Study Area). The Study Area is the same as was analyzed in the 2015 MITT Final EIS/OEIS and is composed of three components: (1) the Mariana Islands Range Complex, (2) additional areas on the high seas, and (3) a transit corridor between the Mariana Islands Range Complex and the Hawaii Range Complex. Three alternatives were analyzed in this SEIS/OEIS:

- The No Action Alternative represents no Navy military readiness activities at sea or on Farallon de Medinilla associated with the Proposed Action within the Study Area. Other military activities not associated with this Proposed Action would continue to occur.
- Alternative 1 reflects a representative year of military readiness activities to account for the typical fluctuation of training and testing cycles and deployment schedules that limit the maximum level of training and testing from occurring for the reasonably foreseeable future. These training and testing activities include new activities at sea, as well as activities that are currently ongoing and have historically occurred in the Study Area.
- Alternative 2 represents the same requirements as Alternative 1. It includes the same type and tempo of military readiness activities as Alternative 1 but also considers additional military readiness activities (e.g., Valiant Shield-type event) and surface warfare events every year that may be needed to respond to unforeseen world events.

Resources evaluated include sediments and water quality, air quality, marine habitats, marine mammals, sea turtles, marine birds, marine vegetation, marine invertebrates, fishes, terrestrial species and habitats, cultural resources, socioeconomic resources and environmental justice, and public health and safety.

**Prepared by:** United States Department of the Navy  
**Comments due by:** Monday, March 18, 2019  
**Point of Contact:** MITT SEIS/OEIS Project Manager  
Naval Facilities Engineering Command, Pacific/EV21  
258 Makalapa Dr., Suite 100  
Pearl Harbor, HI 96860-3134  
(808) 472-1385



---

---

# Executive Summary



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|           |   |             |
|-----------|---|-------------|
| <b>ES</b> | <b>EXECUTIVE SUMMARY .....</b>  | <b>ES-1</b> |
| ES.1      | Introduction .....  | ES-1        |
| ES.2      | Purpose of and Need for Proposed Training and Testing Activities .....  | ES-1        |
| ES.3      | Scope and Content of the Supplemental Environmental Impact Statement/Overseas<br>Environmental Impact Statement.....                    | ES-1        |
| ES.4      | Public Involvement.....   | ES-2        |
|           | ES.4.1 Draft Supplemental Environmental Impact Statement/Overseas Environmental<br>Impact Statement .....                               | ES-2        |
| ES.5      | Proposed Action and Alternatives .....  | ES-2        |
|           | ES.5.1 No Action Alternative .....  | ES-3        |
|           | ES.5.2 Alternative 1 .....  | ES-5        |
|           | ES.5.3 Alternative 2 .....  | ES-5        |
| ES.6      | Summary of Environmental Effects .....  | ES-5        |
|           | ES.6.1 Cumulative Impacts .....   | ES-22       |
| ES.7      | Standard Operating Procedures, Mitigation, and Monitoring.....  | ES-22       |
|           | ES.7.1 Standard Operating Procedures .....  | ES-23       |
|           | ES.7.2 Mitigation .....   | ES-23       |
|           | ES.7.3 Mitigation Measures Considered but Eliminated.....   | ES-23       |
|           | ES.7.4 Monitoring.....  | ES-23       |
|           | ES.7.5 Reporting .....  | ES-24       |
|           | ES.7.6 Other Considerations.....  | ES-24       |
|           | ES.7.6.1 Consistency with Other Federal, State, and Local Plans, Policies and<br>Regulations .....                                      | ES-24       |
|           | ES.7.6.2 Relationship Between Short-Term Use of the Human Environment and<br>Maintenance and Enhancement of Long-Term Productivity..... | ES-24       |
|           | ES.7.6.3 Irreversible or Irrecoverable Commitment of Resources.....   | ES-25       |
|           | ES.7.6.4 Energy Requirements and Conservation Potential of Alternatives ....  | ES-25       |

## List of Figures

Figure ES.5-1: Mariana Islands Training and Testing Study Area..... ES-4

## List of Tables

Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and  
Alternative 2..... ES-7

## ES Executive Summary

### ES.1 Introduction

The United States (U.S.) Department of the Navy (Navy) has prepared this supplement to the May 2015 Mariana Islands Training and Testing (MITT) Environmental Impact Statement/Overseas EIS (EIS/OEIS) (U.S. Department of the Navy, 2015) pursuant to Council on Environmental Quality Regulations. This Supplemental EIS (SEIS)/OEIS considers ongoing and future activities conducted at sea and on Farallon de Medinilla (FDM), updated training and testing requirements, incorporates new information from an updated acoustic effects model, updates marine mammal density data, and incorporates evolving and emergent best available science. Also, it supports the issuance of federal regulatory permits and authorizations under the Marine Mammal Protection Act (MMPA) and the Endangered Species Act (ESA) using the most current and best available science and analytical methods to reassess potential environmental impacts on the species applicable to those regulations. The Navy will consult with the National Marine Fisheries Service (NMFS) to renew these authorizations. While the Study Area remains unchanged from the 2015 MITT Final EIS/OEIS, this SEIS/OEIS focuses on the at-sea and FDM portion of that area.

The 2015 MITT Final EIS/OEIS also analyzed training and testing activities conducted at existing Mariana Islands Range Complex (MIRC) land-based training areas located on Guam, Saipan, Tinian, and Rota. As the Navy is not proposing any changes to those land based activities on Guam, Saipan, Tinian, and Rota, the Navy will continue to rely on the analysis in the 2015 MITT Final EIS/OEIS and the 2015 U.S. Fish and Wildlife Service (USFWS) consultation.

### ES.2 Purpose of and Need for Proposed Training and Testing Activities

The Navy and NMFS (as a cooperating agency) have coordinated from the outset and developed this document to meet each agency's distinct National Environmental Policy Act (NEPA) obligations and support the decision making of both agencies. The Navy's purpose of the Proposed Action is to conduct training and testing activities to ensure that the Navy and other Services meet their respective missions, which, for the Navy under Title 10 United States Code (U.S.C.) Section 5062, is to maintain, train, and equip combat-ready military forces capable of winning wars, deterring aggression, and maintaining freedom of the seas. The respective missions are achieved in part by conducting training and testing within the Study Area in accordance with established Navy military readiness requirements. NMFS's purpose is to evaluate the Navy's Proposed Action pursuant to NMFS's authority under the MMPA, and to make a determination whether to issue incidental take regulations and Letters of Authorization, including any conditions needed to meet the statutory mandates of the MMPA.

### ES.3 Scope and Content of the Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement

In this SEIS/OEIS, the Navy reanalyzed at-sea and FDM training and testing activities that could potentially impact the human environment and natural resources, especially marine mammals, sea turtles, and other marine resources. Since the completion of the 2015 MITT Final EIS/OEIS, new information has become available and is incorporated in this analysis, in addition to proposed changes in training and testing requirements. The range of alternatives in this SEIS/OEIS includes the No Action Alternative and two action alternatives. In this SEIS/OEIS, the Navy analyzes direct, indirect, cumulative, short-term, and long-term impacts, and the irreversible and ir retrievable commitment of resources that may result from the Proposed Action. The Navy is the lead agency for the Proposed Action and is

responsible for the scope and content of this SEIS/OEIS. The document is being prepared in coordination with the U.S. Air Force and U.S. Coast Guard, as their at-sea and FDM training activities in the Study Area are included in the Proposed Action.

The National Oceanic Atmospheric Administration's NMFS is serving as a cooperating agency because the scope of the Proposed Action and alternatives involve activities that have the potential to impact protected resources under their jurisdiction by law and special expertise, including marine mammals, threatened and endangered species, and Essential Fish Habitat. The National Oceanic Atmospheric Administration's authorities and special expertise is based on their statutory responsibilities under the MMPA of 1972, as amended (16 U.S.C. 1361 et seq.), the ESA of 1973 (16 U.S.C. 1531 et seq.), and the Magnuson-Stevens Fishery Conservation and Management Act (MSA). In addition, NMFS, in accordance with 40 Code of Federal Regulations (CFR) 1506.3 and 1505.2, may adopt this SEIS/OEIS and issue a separate Record of Decision associated with its decision to grant or deny the Navy's request for an incidental take authorization pursuant to Section 101(a)(5)(A) of the MMPA.

In accordance with the Council on Environmental Quality Regulations, 40 CFR 1505.2, the Navy will issue a Record of Decision that provides the rationale for choosing one of the alternatives.

#### **ES.4 Public Involvement**

In accordance with the Council on Environmental Quality regulations for implementing the requirements of NEPA, scoping is not required for a supplement to a draft or final EIS (40 CFR 1502.9(c)(4)); however, in an effort to maximize public participation and ensure the public's input is considered, the Navy chose to conduct scoping for this SEIS/OEIS.

Public scoping comments were accepted during the 45-day scoping period from August 1, 2017 to September 15, 2017. In total, the Navy received 36 comment submissions from individuals, groups, agencies, and elected officials. The Navy considered all scoping comments in preparing this SEIS/OEIS.

##### **ES.4.1 Draft Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement**

This Draft SEIS/OEIS was prepared to update the Navy's assessment of potential impacts of the Proposed Action on the environment. The Proposed Action in this SEIS/OEIS reflects an adjustment to the Proposed Action presented in the 2015 MITT Final EIS/OEIS, for which a Record of Decision was issued to support training and testing activities. The proposed training and testing activities are generally consistent with those at-sea and FDM activities analyzed in the 2015 MITT Final EIS/OEIS and are representative of activities the military has been conducting in the MITT Study Area for decades. This SEIS/OEIS assessed potential impacts of all the alternatives (Alternative 1, Alternative 2, and the No Action Alternative). On February 1, 2019, a Notice of Availability was published in the *Federal Register*, and notices were placed in local and regional newspapers announcing the availability of the Draft SEIS/OEIS. The Draft SEIS/OEIS is available for review and comment, and two public meetings are scheduled (February 26, 2019 in Guam and February 27, 2019 in Saipan, Commonwealth of the Northern Mariana Islands [CNMI]).

#### **ES.5 Proposed Action and Alternatives**

The Navy proposes to continue conducting military readiness training and testing activities throughout the Study Area (Figure ES.5-1), primarily in the existing Mariana Islands Range Complex. The proposed training and testing activities associated with the Proposed Action are to be conducted at sea (including the transit corridor between the Mariana Islands Range Complex and the Hawaii Range Complex, and

select Navy pierside and harbor locations) and on FDM. These proposed activities are generally consistent with those at-sea and FDM activities analyzed in the 2015 MITT Final EIS/OEIS. In order to achieve and maintain Fleet readiness through this SEIS/OEIS, the Navy:

- analyzes at-sea and FDM activities necessary to meet readiness requirements beyond 2020 and into the reasonably foreseeable future, including any changes to those activities previously analyzed, and reflects the most up-to-date compilation of training and testing activities deemed necessary to accomplish military readiness requirements;
- adjusts types and tempo (increases or decreases) of training and testing events from the 2015 MITT Final EIS/OEIS to the level needed to meet readiness requirements beyond 2020 and into the reasonably foreseeable future;
- presents the results of the evaluation of relevant new information, which has been incorporated into revised analyses where appropriate (each resource area analyzed within the 2015 MITT Final EIS/OEIS has been evaluated to determine the need for reanalysis within this SEIS/OEIS);
- updates the environmental impact analyses in the previous documents to account for changes to tempo of activity, renaming or combining related types of activities, acknowledging discontinuation of some activities assessed in 2015, and assessing new activities, such as those involving high energy lasers, to enable the Navy to adopt new technology and new capabilities;
- updates environmental analyses with the best available science and most current acoustic analysis methods to evaluate the potential effects of training and testing on the marine environment; and
- supports reauthorization of incidental takes of marine mammals under the MMPA and incidental takes of threatened and endangered marine species under the ESA.

#### **ES.5.1 No Action Alternative**

Under the No Action Alternative, the Navy would not conduct the proposed training and testing activities in the MITT Study Area. Other military activities not associated with this Proposed Action would continue to occur. For FDM, the lease agreement between the U.S. government and the CNMI would remain in place, and the island would continue to be maintained as a Navy range, although strike warfare would no longer continue on the island. For NMFS, denial of an application for an incidental take authorization constitutes the NMFS No Action Alternative, which is consistent with NMFS' statutory obligation under the MMPA to grant or deny requests for take incidental to specified activities. The resulting environmental effects from taking no action will be compared with the effects of the Proposed Action.

Cessation of proposed Navy at-sea training and testing activities would mean that the Navy would not meet its statutory requirements and would be unable to properly defend itself and the United States from enemy forces, unable to successfully detect enemy submarines, and unable to safely and effectively use its weapons systems or defensive countermeasures due to a lack of training of forces and testing of systems that replicate the conditions to which Naval forces must operate while executing the range of military operations required to further national security objectives. Navy personnel would essentially not obtain the unique skills or be prepared to safely and effectively use sensors, weapons, and technologies in realistic scenarios required to accomplish the overall mission. Consequently, the No Action Alternative of not conducting the proposed live, at-sea training and testing activities in the Study Area is inherently unreasonable because it does not meet the purpose and need of the Proposed Action.




Figure ES.5-1: Mariana Islands Training and Testing Study Area

### ES.5.2 Alternative 1

This Alternative consists of an adjustment from the level of training and testing activities analyzed in the 2015 MITT Final EIS/OEIS, accounting for changes in the types and tempo (increases or decreases) of activities necessary to meet current and future military readiness requirements beyond 2020.

- **Adjustments to Tempo of Training and Testing Activities.** This alternative includes changes to training and testing requirements necessary to accommodate current and future training and testing requirements at sea and on FDM, including new at-sea activities as well as activities subject to previous analysis that are currently ongoing and have historically occurred in the Study Area.

Alternative 1 reflects a level of training and testing activities to be conducted at sea and on FDM, with adjustments from the 2015 MITT Final EIS/OEIS that account for changes in the types and tempo of activities necessary to meet current and future military readiness requirements beyond 2020.

### ES.5.3 Alternative 2

Alternative 2 includes the same type of training and testing activities that would occur under Alternative 1. Alternative 2 also considers an increase in tempo of some training and testing activities, including additional Fleet exercises and associated unit-level activities, should unanticipated emergent world events require increased readiness levels. Alternative 2 includes additional electronic warfare activities for Naval Air Systems Command and additional electronic warfare, anti-submarine warfare, and surface warfare activities for Naval Sea Systems Command.

## ES.6 Summary of Environmental Effects

Environmental effects which might result from implementing the Navy's Proposed Action have been analyzed in this SEIS/OEIS. Physical resources that were considered for re-evaluation in this SEIS/OEIS are the same as those that were analyzed in the 2015 MITT Final EIS/OEIS and include sediments and water quality and air quality. Biological resources considered include marine habitats, marine mammals, sea turtles, marine birds, marine vegetation, marine invertebrates, fishes, and terrestrial species and habitats. Human resources considered in this SEIS/OEIS include cultural resources, socioeconomic resources and environmental justice, public health and safety, and cumulative impacts.

As stated previously, this SEIS/OEIS is an update to the 2015 MITT Final EIS/OEIS. New information specifically addressed in this SEIS/OEIS includes updates to military readiness requirements, an updated acoustic effects model, updated marine mammal density data, and evolving and emergent best available science<sup>1</sup>. As the science regarding the potential impacts of acoustics (sonar and explosives) on marine species has evolved since the 2015 MITT Final EIS/OEIS (new research available, updated criteria and thresholds), the acoustic analysis contained in this supplement is a complete update and does not rely

---

<sup>1</sup> The 2015 MITT Final EIS/OEIS used a new modeling system known as the Navy Acoustics Effects Model and marine mammal density information, developed by the Navy in cooperation with the National Marine Fisheries Service, that was the best available information at the time. The Navy Acoustics Effects Model has been refined, marine mammal density estimates have been updated, NMFS has published new criteria and criteria used in the acoustic model have been revised.

on the 2015 MITT Final EIS/OEIS analysis. Analysis associated with activities that result in non-acoustic impacts is updated as necessary in this SEIS/OEIS to reflect new science and refers back to the 2015 MITT Final EIS/OEIS analysis when appropriate.

Table ES.6-1 provides a listing of the potential environmental impacts of the Proposed Action. All sections of the 2015 MITT Final EIS/OEIS were reviewed to determine if there was relevant best available science that needed to be updated/incorporated into this SEIS/OEIS. To the extent there was, it is reflected in each of the sections in Chapter 3.0 (Affected Environment and Environmental Consequences). There was also a re-assessment of effects determinations for marine species. Predicted acoustic exposures are reduced 8 percent under Alternative 1 and would decrease 3 percent under Alternative 2, when compared to the impacts predicted in the 2015 MITT Final EIS/OEIS.

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2**

| Resource Category   | Summary of Impacts  |
|---|---|
| <p><b>Section 3.1</b><br/>Sediments and Water Quality</p> | <p>The Navy considered all stressors that could potentially impact sediments and water quality as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing under the No Action Alternative would result in fewer explosives and explosives byproducts, metals, chemicals, and other materials within the marine environment where training and testing activities have historically been conducted. Discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on sediments and water quality from training and testing activities.</li> </ul> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Explosives and explosives byproducts, metals, chemicals, and other materials expended during training and testing described in this SEIS/OEIS would not exceed regulatory thresholds and guidelines established for measuring impacts on sediment and water quality. Qualitative observations of nearshore waters of FDM during multi-year dive surveys included observations of generally good water quality. There was little evidence of military impacts on benthic sediments and substrates observed during the dive surveys, and, where noted, impacts were localized and shown to recover during subsequent dive surveys.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.2</b><br/>Air Quality</p>                 | <p>The Navy considered all stressors that could potentially impact air quality as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing military readiness activities under the No Action Alternative would improve the ambient air quality as the amount of pollutants being emitted would decrease.</li> </ul>  |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                             | Summary of Impacts   |
|---|--|
|   | <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Pollutant emissions expended during training and testing as described in the SEIS/OEIS would not result in an increase in emissions that would exceed the Prevention of Significant Deterioration threshold of 250 tons per year for any of the criteria pollutants.. Therefore, Alternative 1 would not have a significant impact on ambient air quality, and no applicable thresholds would be exceeded.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul>   |
| <p><b>Section 3.3</b><br/>Marine Habitats</p> | <p>The Navy considered all stressors that could potentially impact marine habitats as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Under the No Action Alternative, discontinuing the training and testing activities would result in fewer explosive and physical disturbance and strike stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for explosive or physical disturbance and strike stressor impacts on marine habitat, but would not measurably improve the overall distribution or abundance of marine habitat.</li> </ul> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Most of the explosive military expended materials would detonate at or near the water surface. Training activities that include bottom-laid in-water explosions under Alternative 1 would affect marine habitat structure in the Study Area, but these activities would occur in areas that have been previously disturbed, and impacts would be localized. Mitigation measures will help the Navy avoid or reduce impacts on seafloor resources (including shallow-water coral reefs, live hard bottom, artificial reefs, and submerged cultural resources) from explosives during applicable activities.</li> <li>• Bottom substrates could be disturbed by vessel and in-water device strikes, military expended materials, seafloor devices used for military readiness activities, and from walking, standing, or swimming in the nearshore waters</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                            | Summary of Impacts  |
|--|---|
|  | <p>during amphibious activities such as raids and assaults. The impact of vessels and in-water devices on marine habitats would remain inconsequential because (1) vessel and in-water activities that could come into contact with marine substrates would be located in previously disturbed areas (i.e., nearshore shallow waters), (2) military expended materials could be colonized by benthic organisms, and (3) seafloor devices would be used in previously disturbed areas and therefore would not be expected to affect marine substrates.</p> <p><b>Alternative 2:</b></p> <ul style="list-style-type: none"> <li>The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul>  |
| <p><b>Section 3.4</b><br/>Marine Mammals</p> | <p>The Navy considered all stressors that could potentially impact marine mammals as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b>No Action Alternative:</b></p> <ul style="list-style-type: none"> <li>Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors that potentially affect marine mammals within the marine environment. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on marine mammals, but would not measurably improve the overall habitat, distribution or abundance of marine mammals.</li> </ul> <p><b>Alternative 1:</b></p> <ul style="list-style-type: none"> <li>The use of sonar and other transducers would have the potential to expose marine mammals to sound-producing activities which would present risks to marine mammals that could range from a temporary or permanent threshold shift, auditory masking, physiological stress, or behavioral responses (only Kogia whales are predicted to have permanent threshold shift exposures). Individual animals would typically only experience a small number of behavioral responses or temporary threshold shifts per year due to exposure to acoustic stressors and are unlikely to incur substantive costs to that individual. Population-level effects are unlikely.</li> <li>The use of munitions in the water or near the water's surface present a risk to marine mammals located in close proximity to the explosion, because the resulting shock waves can cause injury or result in the death of an animal; however, there are no instances of non-auditory injury or death predicted by the acoustic modeling. If a marine mammal is farther from an explosion, the impulsive, broadband sounds introduced into the marine environment</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category | Summary of Impacts   |
|-------------------|--|
|                   | <p>may cause a temporary or permanent threshold shift, auditory masking, physiological stress, or behavioral responses. Because most estimated impacts from explosions are behavioral responses or temporary threshold shifts and because the numbers of marine mammals potentially impacted by explosives are small as compared to each species respective abundance, population-level effects are unlikely.</p> <ul style="list-style-type: none"> <li data-bbox="558 488 1852 789">• The use of in-water electromagnetic devices and high-energy lasers would have the potential to result in impacts to marine mammals. The likelihood and magnitude of energy impacts depends on the proximity of marine mammals to the activity. Based on the relatively weak strength of the electromagnetic field created by some Navy activities, a marine mammal would have to be in close proximity for there to be any effect and impacts on marine mammal migrating behaviors and navigational patterns are not anticipated. Statistical probability analyses with conservative assumptions tending to overestimation of exposures demonstrate with a high level of certainty that a marine mammal would not be struck by a high energy laser. These activities are temporary and localized in nature, and may result in short-term and minor impacts on individual marine mammals, but would not result in long-term impacts on marine mammal populations.</li> <li data-bbox="558 813 1852 1383">• The use of vessels, in-water devices, military expended materials, and seafloor devices would have the potential to result in impacts to marine mammals. The potential for impacts relies heavily on the probability that marine mammals would be in close proximity to an activity (e.g., a vessel or an expended non-explosive munition). Historical data indicates no occurrence of vessel strikes with marine mammals in the MITT over the last ten years during any training and testing activities. Since the Navy does not anticipate a substantive change in the level of vessel use compared to the last decade and there have been no strikes in that timeframe, the potential for striking a marine mammal is therefore discountable. Physical disturbance due to vessel movement and in-water devices of individual marine mammals may also occur, but any stress response of avoidance behavior would not be severe enough to have long-term fitness consequences for individual marine mammals. The use of in-water devices during Navy activities involves multiple types of vehicles or towed devices traveling on the water surface, through the water column, or along the seafloor, all of which have the potential to physically disturb or strike marine mammals. No recorded or reported instances of marine mammal strikes have resulted from in-water devices; therefore, impacts on individuals or long-term consequences to marine mammal populations are not anticipated. Potential impacts from military expended materials and seafloor devices are determined through statistical probability analyses. Analyses suggest a very low potential for marine mammals to be struck by any of these items. Long-term consequences to marine mammal populations from vessels, in-water devices, military expended materials, and seafloor devices associated with training and testing activities are not anticipated.</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category | Summary of Impacts   |
|-------------------|--|
|                   | <ul style="list-style-type: none"> <li data-bbox="558 354 1852 686">• The use of cables, wires, and decelerators/parachutes would have the potential to result in impacts to marine mammals. The potential for impacts is dependent on the probability that a marine mammal would encounter an expended material, as well as the physical properties of the expended materials and the likelihood that a marine mammal could become entangled in the item. Physical characteristics of cables, wires, and decelerators/parachutes suggest that it is not likely a marine mammal would become entangled in these items. While it may be possible for a marine mammal to become entangled in cables or wires, the sparse distribution of these items throughout the Study Area indicates a very low potential for encounter. Furthermore, fiber optic cables used during mine warfare activities are easily abraded and have a low breaking strength, which reduces the risk of entanglement should a cable be encountered. Short-term impacts on individual marine mammals and long-term impacts on marine mammal populations from entanglement associated with training and testing activities are not anticipated.</li> <li data-bbox="558 711 1852 1109">• Use of military expended materials would have the potential to result in impacts to marine mammals. The potential for impacts relies heavily on feeding behaviors of marine mammals that occur in the Study Area, the physical properties of the expended items, the feasibility that a marine mammal could ingest the items, and the likelihood that a marine mammal would encounter an item. Marine mammals that forage along the water surface or within the water column are less likely to encounter ingestion stressors as they sink through the water column to the seafloor. Most expended materials that would remain floating or suspended within the water column are typically too small to pose a risk of intestinal blockage to any marine mammal that encounters it. Bottom-feeding marine mammals would be more likely to encounter expended materials that have already sunk to the floor. In the unlikely event that a marine mammal encounters and ingests expended material, the individual might be negatively affected if the material becomes lodged in the digestive tract. The likelihood that a marine mammal would ingest a military expended item associated with training and testing activities is considered low. Long-term consequences to marine mammal populations from expended materials associated with training and testing activities are not anticipated.</li> <li data-bbox="558 1133 1852 1393">• Marine mammals would be exposed to multiple secondary causes of impact associated with training and testing activities in the Study Area. In-water explosions have the potential to injure or kill prey species that marine mammals feed on; however, impacts would not substantially impact prey availability for marine mammals. Explosion byproducts are not considered as indirect stressors to marine mammals while mixed in marine sediments or water. Marine mammals may encounter unexploded ordnance underwater or within sediments, but ingestion is very unlikely. Explosion byproducts and unexploded munitions would have no lasting or meaningful effect on water quality and would therefore not constitute a secondary indirect stressor for marine mammals. Metals are introduced into the water and sediments from targets, munitions, and other expended materials. Evidence from a</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                         | Summary of Impacts  |
|---|---|
|   | <p>number of studies indicate metal contamination is localized and ephemeral and that bioaccumulation resulting from munitions was not observed in the studies specifically designed to look for bioaccumulation. Therefore, it is unlikely that impacts on marine mammal prey availability would occur. Several training and testing activities introduce explosive byproducts into the marine environment that are potentially harmful in concentration; however, rapid dilution would occur and toxic concentrations are unlikely to be encountered. Furthermore, there is no evidence of acute toxicity or chronic accumulation in tissues of chemicals introduced by Navy activities that would alter water quality to an extent that would result in overall habitat degradation for marine mammals. Transmission of marine mammal diseases and parasites are not considered likely from the Navy’s trained marine mammals because strict protocols are in place to prevent such impacts on wild populations. Secondary stressors from training and testing activities in the Study Area are not expected to have short-term impacts on individual marine mammals or long-term impacts on marine mammal populations.</p> <p><b>Alternative 2:</b></p> <ul style="list-style-type: none"> <li>The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.5</b><br/>Sea Turtles</p> | <p>The Navy considered all stressors that could potentially impact sea turtles as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b>No Action Alternative:</b></p> <ul style="list-style-type: none"> <li>Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing the training and testing activities would result in fewer stressors that potentially affect sea turtles within the marine environment where training and testing have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on sea turtles, but would not measurably improve the status of sea turtle populations.</li> </ul> <p><b>Alternative 1:</b></p> <ul style="list-style-type: none"> <li>The 2015 MITT Final EIS/OEIS analyzed potential impacts of at-sea training and testing activities, as well as amphibious landings on beaches on Guam, Rota, and Tinian, which may support sea turtle nesting. Activities on Guam, Rota, and Tinian do not change; therefore, this SEIS only addresses potential stressors on sea turtles for training and testing activities at sea.</li> </ul>  |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category | Summary of Impacts   |
|-------------------|--|
|                   | <ul style="list-style-type: none"> <li>• The use of sonar and other transducers, explosives, in-water electromagnetic devices, vessels and in-water devices, weapons, military expended materials, seafloor devices, and military expended materials of ingestible size associated with training and testing activities may affect sea turtles present within the Study Area; however, they would not result in significant adverse impacts or jeopardize the continued existence of any sea turtle species. These findings are consistent with the analysis in the 2015 MITT Final EIS/OEIS and biological opinions provided to the Navy by NMFS and the USFWS.</li> <li>• The use of sonar and other transducers, explosives, aircraft, vessels and weapons have the potential for limited impacts to sea turtles because sea turtles have limited hearing abilities. If a sea turtle is close enough to a source using a frequency within a sea turtle’s hearing range, the sea turtle may exhibit short-term behavioral reactions or may exhibit no reaction at all. No long-term consequences to sea turtle populations would be expected.</li> <li>• In-water electromagnetic devices are not expected to result in population-level impacts for sea turtles due to the low intensity, localized potential impact area, and short duration of use. The use of high-energy lasers associated with testing activities is not expected to impact sea turtles as a result of the very low probability of a direct strike by a high-energy laser.</li> <li>• Use of vessels and in-water devices, military expended materials, and seafloor devices may cause short-term disturbance to an individual turtle within the Study Area. However, due to the low numbers of sea turtles potentially impacted by these activities that may cause physical disturbance and strike, population-level effects are unlikely.</li> <li>• The use of cables and wires, and decelerators/parachutes may cause short-term or long-term disturbance to an individual sea turtle. However, due to the physical characteristics and low number of cables, wires, and decelerators/parachutes, combined with the behavior of the species, population-level impacts are not expected.</li> <li>• The use of military expended materials and munitions may cause short-term or long-term disturbance to an individual sea turtle due to ingestion of munitions used in training activities. However, the potential impacts of exposure to munitions are not expected to result in population-level impacts.</li> </ul> <p><b>Alternative 2:</b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                          | Summary of Impacts  |
|--|---|
| <p><b>Section 3.6</b><br/>Marine Birds</p> | <p>The Navy considered all stressors that could potentially impact marine birds as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors that potentially affect marine birds within the marine environment. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on marine birds, but would not measurably improve the status of marine bird populations.</li> </ul> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• The use of sonar and other transducers, explosives, in-air electromagnetic devices, aircraft, aerial targets, vessels and in-water devices, military expended materials, and military expended materials of ingestible size associated with training and testing activities would have no effect on ESA-listed marine birds within the Study Area. In addition, the use of high-energy lasers associated with testing activities would have no effect on ESA-listed marine birds within the Study Area. This conclusion is based on the consideration that ESA-listed marine birds generally occur outside of the Study Area, with little to no overlap with at-sea training and testing activities.</li> <li>• Periodic helicopter-based surveys of FDM have occurred since 1998 (monthly up to 2009, and quarterly thereafter through September 2016) for marine birds nesting on the island. Because of a lack of commercial helicopter transit services, surveys have not been conducted since 2016. Under Alternative 1, training activities on FDM would not significantly impact populations of marine birds on the island. This conclusion is based on statistical analysis of periodic population counts of masked, brown, and red-footed boobies undertaken by the Navy from 1998 through 2016, and the relatively small increases in the number of events, munitions, and Net Explosive Weight expended on FDM proposed under Alternative 1 compared to what was analyzed in the 2015 MITT Final EIS/OEIS.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                                  | Summary of Impacts  |
|--|---|
| <p><b>Section 3.7</b><br/>Marine Vegetation</p>    | <p>The Navy considered all stressors that could potentially impact marine vegetation as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors that potentially affect marine vegetation within the marine environment. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on marine vegetation, but would not measurably improve the status of marine vegetation populations or subpopulations.</li> </ul> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Physical disturbance and strike and the use of in-water explosives could affect marine vegetation by destroying individual plants or damaging parts of plants, but are not expected to result in detectable changes in survival or propagation, and are not expected to result in population-level impacts on marine plant species. Changes in sediment and water quality are not likely to be measureable, thus no detectable changes are expected in marine vegetation growth, survival, propagation, or population-level impacts.</li> <li>• The use of explosives, military expended materials, and seafloor devices during military readiness activities under Alternative 1 could affect marine vegetation by destroying individual plants or damaging parts of plants, but are not expected to result in detectable changes in survival or propagation, and are not expected to result in population-level impacts on marine plant species.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.8</b><br/>Marine Invertebrates</p> | <p>The Navy considered all stressors that could potentially impact marine invertebrates as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors</li> </ul>   |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                    | Summary of Impacts   |
|--------------------------------------|--|
|                                      | <p>that potentially affect marine invertebrates within the marine environment. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on marine invertebrates, but would not measurably improve the status of invertebrate populations or subpopulations.</p> <p><b>Alternative 1:</b></p> <ul style="list-style-type: none"> <li>• The use of sonar and other transducers, in-water electromagnetic devices, cables, wires, decelerators/parachutes, and military expended materials of ingestible size associated with training and testing activities would have a negligible impact on marine invertebrate species. In addition, the use of high-energy lasers associated with testing activities would have a negligible impact on marine invertebrate species.</li> <li>• Use of explosives, vessels and in-water devices, military expended materials and seafloor devices, associated with training and testing activities may impact individual marine invertebrates and groups of marine invertebrates. However, these activities are unlikely to impact populations or subpopulations of marine invertebrates.</li> <li>• The use of sonar and other transducers; in-water electromagnetic devices; cables, wires, and decelerators/parachutes; metal, chemical, and other material byproducts; and secondary physical disturbances would have no adverse effect on sedentary invertebrate beds or reefs. The use of in-water explosives, vessels and in-water devices, military expended materials, and seafloor devices, explosive byproducts, and unexploded ordnance during military readiness activities may have an adverse effect on sedentary invertebrate beds or reefs.</li> </ul> <p><b>Alternative 2:</b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.9</b><br/>Fishes</p> | <p>The Navy considered all stressors that could potentially impacts fishes could potentially as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b>No Action Alternative:</b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors that potentially affect fishes within the marine environment. Therefore, discontinuing training and testing activities</li> </ul>   |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category   | Summary of Impacts  |
|---|---|
|   | <p>under the No Action Alternative would lessen the potential for stressor impacts on fishes,, but would not measurably improve the status of fish populations or subpopulations.</p> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• The use of sonar and other transducers, explosives, and in-water electromagnetic devices, may affect fishes. Impacts however are expected to be temporary and infrequent as most activities would be temporary, localized, and infrequent. More severe impacts such as mortality or injury could lead to permanent or long-term consequences for individuals, but, overall, long-term consequences for fish populations are not expected.</li> <li>• The use of vessels and in-water devices, aircraft, weapons, military expended materials, seafloor devices, cables, wires, decelerators/parachutes, and military expended materials of ingestible size associated with training and testing activities may affect fishes. However, because the number of fishes potentially impacted by these activities is low, population-level impacts are unlikely.</li> <li>• The use of sonar and other transducers, in-water explosives, in-water electromagnetic devices, vessels and in-water devices, cables, wires, decelerators/parachutes, and military expended materials associated with training and testing activities may affect fishes within the Study Area. In addition, the use of high-energy lasers associated with testing activities may affect fishes within the Study Area.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.10</b><br/>Terrestrial Species and Habitats</p> | <p>The Navy considered all stressors that could potentially impact terrestrial species and habitats as a result of the Proposed Action. This SEIS/OEIS addresses potential impacts on terrestrial species and habitats on FDM. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. For FDM, the lease agreement between the U.S. government and the Commonwealth of the Northern Mariana Islands would remain in place, and the island would continue to be maintained as a Navy range.</li> </ul> <p><b><u>Alternative 1:</u></b></p>   |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                                 | Summary of Impacts   |
|---|--|
|   | <ul style="list-style-type: none"> <li>• Under Alternative 1, more aircraft would fly over and land on FDM and more ordnance would be used on FDM. The total increase, in terms of net explosive weight (NEW) under Alternative 1 would be less than 1 percent compared to ordnance use on FDM described in the 2015 MITT Final EIS/OEIS. All of the ordnance would be used within existing impact zones, with the same avoidance and minimization measures in place as described in the 2015 MITT Final EIS/OEIS and in the 2015 USFWS Biological Opinion and 2016 USFWS concurrence letter. In accordance with 50 CFR 402.16, the 2015 and 2016 consultations remain valid as none of the factors necessary to trigger reinitiating consultation have been met. The 2015 USFWS Biological Opinion and 2016 USFWS concurrence letter would still apply to ESA-listed species occurring on FDM (the Mariana fruit bat and Micronesian megapode). The continued use of FDM as described in this SEIS/OEIS would not significantly impact populations of birds protected under the MBTA.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul>  |
| <p><b>Section 3.11</b><br/>Cultural Resources</p> | <p>The Navy considered all stressors that could potentially impact cultural resources as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Discontinuing training and testing activities under the No Action Alternative would result in fewer stressors that potentially affect submerged cultural resources. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for stressor impacts on submerged cultural resources.</li> </ul> <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Under Alternative 1, measures previously implemented to avoid and protect submerged historic properties would continue to be implemented according to the conservation measures and procedures identified and described in the 2009 MIRC Programmatic Agreement. Given the Navy avoids areas with known submerged obstructions, including submerged objects listed or eligible for listing on the National Register of Historic Places, submerged historic properties within U.S. territorial waters and National Register of Historic Places-eligible resources on Guam and the Commonwealth of the Northern Mariana Islands would not be affected by training and testing activities. In</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category  | Summary of Impacts   |
|--|--|
|  | <p>accordance with Section 402 of National Historic Preservation Act, no known World Heritage Sites would be affected.</p> <p><b>Alternative 2:</b></p> <ul style="list-style-type: none"> <li>The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul>  |
| <p><b>Section 3.12</b><br/>Socioeconomic Resources and Environmental Justice</p> | <p>The Navy considered all stressors that could potentially impact socioeconomic resources and environmental justice as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b>No Action Alternative:</b></p> <ul style="list-style-type: none"> <li>Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Limits on accessibility to the ocean and airspace associated with the proposed training and testing activities would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.</li> <li>Discontinuing training and testing activities would result in fewer stressors on socioeconomic resources within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on socioeconomic resources, such as commercial and recreational fishing, commercial transportation and shipping, tourism, and traditional fishing practices in the Study Area.</li> <li>The Navy and Navy personnel are an important and often stabilizing contributor to the local and regional economies. Therefore, not conducting the proposed at-sea training and testing activities may have negative impacts on the socioeconomic resources of Guam and the CNMI. The number of jobs and types of jobs available on Guam and to a lesser extent on the CNMI may decline. For example, vessels and associated equipment used specifically for military readiness activities would no longer be needed if training and testing activities ceased. Consequently, the civilian and Navy personnel supporting those activities may be relocated, reassigned, or have to find other employment. The secondary effects from reducing the number of personnel who support at-sea military training and testing activities could include a decline in revenue for local businesses frequented by Navy personnel and their families, such as businesses in the food services, retail, and housing sectors.</li> </ul> |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category                                       | Summary of Impacts  |
|---|---|
|   | <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>• Alternative 1 may result in impacts on commercial and recreational fishing, traditional fishing practices, or tourism when areas of co-use are temporarily inaccessible to ensure public safety during training and testing activities. No impacts on commercial transportation and shipping are anticipated, because training and testing activities are scheduled and located to avoid potential conflicts with commercial vessels and air traffic. The military will continue to collaborate with local communities to enhance existing means of communication with the public that are intended to reduce the potential effects of limiting accessibility to areas designated for use by the military.</li> <li>• Alternative 1 is not expected to result in impacts from physical disturbance and strike or airborne acoustic disturbances on commercial and recreational fishing, traditional fishing practices, other recreational activities or tourism, because the vast majority of military training and testing activities would occur in areas of the Study Area far from the locations typically used by the public for fishing and recreation activities. Furthermore, the large size of the Study Area over which the proposed military training and testing activities would be distributed, and adherence to the Navy’s standard operating procedures, would further reduce any potential for impacts.</li> <li>• Traditional fishers in Guam and the CNMI would not be disproportionately impacted by limits on accessibility, airborne acoustic disturbances, or the possibility of physical disturbance and strike, because traditional fishers typically use the same general areas as recreational fishers, specifically areas closer to shore and far from the majority of training and testing activities.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>• The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |
| <p><b>Section 3.13</b><br/>Public Health and Safety</p> | <p>The Navy considered all stressors that could potentially impact public health and safety as a result of the Proposed Action. The following conclusions have been reached for the project alternatives:</p> <p><b><u>No Action Alternative:</u></b></p> <ul style="list-style-type: none"> <li>• Under the No Action Alternative, the proposed training and testing activities would not occur in the MITT Study Area. Not conducting the proposed at-sea training and testing activities may lessen the potential for interactions between the Navy and civilians and improve public health and safety.</li> </ul>   |

**Table ES.6-1: Summary of Environmental Impacts for the No Action Alternative, Alternative 1, and Alternative 2 (continued)**

| Resource Category | Summary of Impacts  |
|-------------------|---|
|                   | <p><b><u>Alternative 1:</u></b></p> <ul style="list-style-type: none"> <li>The use of sonar, in-water explosives, radar, lasers, aircraft, vessels, in-water devices/targets, munitions, and seafloor devices would not adversely affect public health and safety because standard operating procedures are in place to ensure that there is no overlap between military and non-military activities. In addition, training and testing activities would not appreciably change the water quality in the region.</li> </ul> <p><b><u>Alternative 2:</u></b></p> <ul style="list-style-type: none"> <li>The number of training and testing activities under Alternative 2 would increase over what is proposed for Alternative 1. However, this increase would be a slight change and would not appreciably change the potential for impacts over what is analyzed for Alternative 1.</li> </ul> |

Notes: SEIS/OEIS = Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement, ESA = Endangered Species Act, FDM = Farallon de Medinilla, MBTA = Migratory Bird Treaty Act, MITT = Mariana Islands Training and Testing, MMPA = Marine Mammal Protection Act, Navy = United States Department of the Navy, U.S. = United States, USFWS = U.S. Fish and Wildlife Service, NMFS = National Marine Fisheries Service, MIRC = Mariana Islands Range Complex, CNMI = Commonwealth of the Northern Mariana Islands

### ES.6.1 Cumulative Impacts

Marine mammals, marine invertebrates, sea turtles, and socioeconomics are the primary resources of concern for cumulative impacts analysis:

- Past human activities have impacted these resources to the extent that several marine mammals, sea turtles, marine invertebrates species, and some terrestrial species occurring in the Study Area are ESA-listed. Several marine mammal species have stocks that are classified as strategic stocks under the MMPA.
- The use of sonar and other non-impulsive sound sources under Alternative 1 and Alternative 2 has the potential to disturb or injure marine mammals and sea turtles.
- Explosive detonations, and vessel strikes under Alternative 1 and Alternative 2 have the potential to disturb, injure, or kill marine mammals and sea turtles.
- Under Alternative 1 and Alternative 2, danger zones could potentially restrict access to fishing and recreational areas when ranges are in use.

The aggregate impacts of past, present, and other reasonably foreseeable future actions would continue to have significant impacts on some individual marine mammal and all sea turtle species in the Study Area. Alternative 1 or Alternative 2 would contribute to cumulative impacts; however, marine mammal and sea turtle mortality and injury from non-Navy actions associated with commercial fisheries, commercial vessel strikes, and entanglement in marine debris are leading causes of direct mortality to marine mammals and sea turtles (Carretta et al., 2017; Helker et al., 2017; Lent & Squires, 2017; National Marine Fisheries Service, 2016; National Oceanic and Atmospheric Administration Marine Debris Program, 2014; Read et al., 2006). In summary, based on the analysis presented in Sections 3.4 (Marine Mammals), 3.5 (Sea Turtles), 3.8 (Marine Invertebrates), and 3.12 (Socioeconomic Resources), the current aggregate impacts of past, present, and other reasonably foreseeable future actions are not significantly different than the assessment in the 2015 MITT Final EIS/OEIS. For marine mammals, sea turtles, and marine invertebrates Alternatives 1 or 2 would contribute to and increase cumulative impacts, but the relative contribution would be negligible compared to other non-Navy actions. Cumulative effects on socioeconomic resources may have short-term impacts on accessibility to public services, fishing sites, and tourism resources, but they are not expected to have long-term negative impacts on these resources or the economy of Guam and the Commonwealth of the Northern Mariana Islands. No new information or circumstances are significant enough to warrant further cumulative impact review.

The analysis presented in Chapter 3 (Affected Environment and Environmental Consequences) and Chapter 4 (Cumulative Impacts) indicate that the incremental contribution of Alternative 1 or Alternative 2 to cumulative impacts on sediments and water quality, air quality, marine habitats, marine birds, marine vegetation, fishes, cultural resources, and public health and safety would be negligible.

### ES.7 Standard Operating Procedures, Mitigation, and Monitoring

Within the Study Area, the Navy implements standard operating procedures, mitigation measures, and marine species monitoring and reporting. There are benefits to environmental and cultural resources resulting from the standard operating procedures discussed in this SEIS/OEIS. Mitigation measures are designed to help reduce or avoid potential impacts on marine and terrestrial resources. Marine species monitoring efforts are designed to track compliance with take authorizations, evaluate the effectiveness

of mitigation measures, and improve understanding of the effects training and testing activities on marine resources.

### **ES.7.1 Standard Operating Procedures**

For training and testing to be effective, units must be able to safely use their sensors and weapon systems as they are intended to be used in military missions and combat operations and to their optimum capabilities. Standard operating procedures applicable to training and testing have been developed through years of experience and their primary purpose is to provide for safety (including public health and safety) and mission success. Because they are essential to safety and mission success, standard operating procedures are part of the Proposed Action and are considered in the Chapter 3 (Affected Environment and Environmental Consequences) environmental analysis for applicable resources.

### **ES.7.2 Mitigation**

The Navy recognizes that the Proposed Action has the potential to impact the environment. Standard operating procedures differ from mitigation measures because mitigation is designed specifically for the purpose of avoiding or reducing environmental impacts, whereas standard operating procedures are designed to provide for safety and mission success. The Navy is coordinating with NMFS on these measures through the consultation and permitting processes. The Navy and NMFS Records of Decision, MMPA Regulations and Letter of Authorization, and ESA Biological Opinion will document all mitigation that the military will implement under the Proposed Action.

Mitigation measures that the military will implement under the Proposed Action are organized into three categories: procedural mitigation measures for at-sea activities, at-sea mitigation areas, and terrestrial mitigation measures for activities on FDM. Procedural mitigation is mitigation that will be implemented whenever and wherever an applicable military readiness activity takes place within the Study Area. Mitigation areas are geographic locations within the Study Area where the military will implement additional mitigation during all or part of the year. Terrestrial mitigation measures are measures that the Navy will implement during applicable military readiness activities that take place on land at FDM.

### **ES.7.3 Mitigation Measures Considered but Eliminated**

A number of possible additional mitigation measures have been suggested during the public scoping period of this SEIS/OEIS and comment periods of previous Navy environmental documents. Through the evaluation process, some measures were deemed to either be ineffective, have an unacceptable impact on the proposed training and testing activities, or both, and will not be carried forward for further consideration (refer to Section 5.6, Measures Considered But Eliminated).

### **ES.7.4 Monitoring**

The Navy is committed to demonstrating environmental stewardship while executing its national security mission, complying with the suite of federal environmental laws and regulations, and providing required and relevant reports to appropriate regulatory agencies. Since 2006 across all Navy range complexes (in the Marianas, Pacific, Atlantic, Gulf of Mexico, and Gulf of Alaska), the Navy has produced various reports (Major Exercise Reports, Annual Exercise Reports, and Monitoring Reports) submitted to National Marine Fisheries Service to further research goals aimed at understanding the Navy's impact on the environment as it carries out testing and training to accomplish its mission. As a complement to the Navy's commitment to avoiding and reducing impacts of the Proposed Action through mitigation, the

Navy will undertake monitoring efforts to track compliance with take authorizations, help investigate the effectiveness of implemented mitigation measures, and better understand the impacts of the Proposed Action on marine resources. Taken together, mitigation and monitoring comprise the Navy's integrated approach for reducing environmental impacts from the Proposed Action. The Navy's overall monitoring approach will seek to leverage and build on existing research efforts whenever possible.

Consistent with the cooperating agency agreement with NMFS, mitigation and monitoring measures presented in this SEIS/OEIS focus on the requirements for protection and management of marine resources. Since monitoring will be required for compliance with the Final Rule issued for the Proposed Action under the MMPA, details of the monitoring program are being developed in coordination with NMFS through the regulatory process.

The Navy developed the Integrated Comprehensive Monitoring Program to serve as the overarching framework for coordinating its marine species monitoring efforts and as a planning tool to focus its monitoring priorities pursuant to ESA and MMPA requirements (U.S. Department of the Navy, 2010). The purpose of the Integrated Comprehensive Monitoring Program is to coordinate monitoring efforts across all regions and to allocate the most appropriate level and type of monitoring effort for each range complex based on a set of standardized objectives, regional expertise, and resource availability. Additional information about the U.S. Navy Marine Species Monitoring Program, including an introduction to adaptive management and strategic planning, is provided in Section 5.1.2.2.1 (Marine Species Research and Monitoring Programs).

#### **ES.7.5 Reporting**

The Navy is committed to documenting and reporting relevant aspects of training and testing activities in order to reduce environmental impact, and improve future environmental assessments. Initiatives include training and testing activity reporting, and incident reporting.

#### **ES.7.6 Other Considerations**

##### **ES.7.6.1 Consistency with Other Federal, State, and Local Plans, Policies and Regulations**

Based on an evaluation of consistency with statutory obligations, the Navy and other Service's proposed training and testing activities would not conflict with the objectives or requirements of federal, state, regional, or local plans, policies, or legal requirements. The Navy and other Services are consulting and will continue to consult with regulatory agencies as appropriate during the NEPA process and prior to implementation of the Proposed Action to ensure all legal requirements are met.

##### **ES.7.6.2 Relationship Between Short-Term Use of the Human Environment and Maintenance and Enhancement of Long-Term Productivity**

In accordance with NEPA, this SEIS/OEIS provides an analysis of the relationship between a project's short-term impacts on the environment and the effects that these impacts may have on the maintenance and enhancement of the long-term productivity of the affected environment. The Proposed Action may result in both short- and long-term environmental effects. However, the Proposed Action would not be expected to result in any impacts that would reduce environmental productivity, permanently narrow the range of beneficial uses of the environment, or pose long-term risks to health, safety, or the general welfare of the public.

**ES.7.6.3 Irreversible or Irretrievable Commitment of Resources**

For the alternatives including the Proposed Action, most resource commitments are neither irreversible nor irretrievable. Most impacts are short-term and temporary or, if long lasting, are negligible. No habitat associated with threatened or endangered species would be lost as result of implementation of the Proposed Action. Since there would be no building or facility construction, the consumption of materials typically associated with such construction (e.g., concrete, metal, sand, fuel) would not occur. Energy typically associated with construction activities would not be expended and irreversibly lost.

Implementation of the Proposed Action would require fuels used by aircraft, ships, and ground-based vehicles. Since fixed- and rotary-wing flight and ship activities could increase, relative total fuel use could increase. Therefore, if total fuel consumption increased, this nonrenewable resource would be considered irretrievably lost.

**ES.7.6.4 Energy Requirements and Conservation Potential of Alternatives**

Resources that will be permanently and be continually consumed by project implementation include water, electricity, natural gas, and fossil fuels; however, the amount and rate of consumption of these resources would not result in significant environmental impacts or the unnecessary, inefficient, or wasteful use of resources.

Sustainable range management practices are in place that protect and conserve natural and cultural resources and preserve access to training areas for current and future training requirements while addressing potential encroachments that threaten to impact range and training area capabilities.

## **REFERENCES**

- Carretta, J. V., M. M. Muto, J. Greenman, K. Wilkinson, D. Lawson, J. Viezbicke, and J. Jannot. (2017). *Sources of Human-Related Injury and Mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2011–2015* (NOAA Technical Memorandum NMFS-SWFSC-579). La Jolla, CA: Southwest Fisheries Science Center.
- Helker, V. T., M. M. Muto, K. Savage, S. Teerlink, L. A. Jemison, K. Wilkinson, and J. Jannot. (2017). *Human-Caused Mortality and Injury of NMFS-Managed Alaska Marine Mammal Stocks, 2011–2015* (National Oceanic and Atmospheric Administration Technical Memorandum NMFS-AFSC-354). Seattle, WA: Alaska Fisheries Science Center.
- Lent, R., and D. Squires. (2017). Reducing marine mammal bycatch in global fisheries: An economics approach. *Deep-Sea Research II*, 140, 268–277.
- National Marine Fisheries Service. (2016). *U.S. National Bycatch Report First Edition Update 2*. Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service. Retrieved from <http://www.st.nmfs.noaa.gov/observer-home/first-edition-update-2>.
- National Oceanic and Atmospheric Administration Marine Debris Program. (2014). *Report on the Entanglement of Marine Species in Marine Debris with an Emphasis on Species in the United States*. Silver Spring, MD: National Oceanic and Atmospheric Administration.
- Read, A. J., P. Drinker, and S. Northridge. (2006). Bycatch of marine mammals in U.S. and global fisheries. *Conservation Biology*, 20(1), 163–169.
- U.S. Department of the Navy. (2010). *Navy Integrated Comprehensive Monitoring Plan*. Washington, DC: U.S. Department of the Navy.
- U.S. Department of the Navy. (2015). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.

**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|          |  |            |
|----------|--|------------|
| <b>1</b> | <b>PURPOSE AND NEED.....</b>   | <b>1-1</b> |
| 1.1      | Introduction .....   | 1-1        |
| 1.2      | The Navy’s Environmental Compliance and At-Sea Policy .....  | 1-3        |
| 1.3      | Proposed Action.....   | 1-4        |
| 1.4      | Purpose and Need.....  | 1-4        |
|          | 1.4.1 Why the Navy Trains.....   | 1-5        |
|          | 1.4.2 Why the Navy Tests .....   | 1-6        |
| 1.5      | Overview and Strategic Importance of Existing Range Complex.....   | 1-6        |
| 1.6      | The Environmental Planning Process.....  | 1-7        |
|          | 1.6.1 National Environmental Policy Act Requirements .....   | 1-8        |
|          | 1.6.2 Executive Order 12114 .....  | 1-8        |
|          | 1.6.3 Other Environmental Requirements Considered .....  | 1-8        |
| 1.7      | Scope and Content.....   | 1-9        |
| 1.8      | Organization of this Supplemental Environmental Impact Statement/Overseas<br>Environmental Impact Statement..... | 1-9        |
| <b>2</b> | <b>DESCRIPTION OF PROPOSED ACTION AND ALTERNATIVES.....</b>  | <b>2-1</b> |
| 2.1      | Description of the Mariana Islands Training and Testing Study Area.....  | 2-2        |
|          | 2.1.1 Description of the Mariana Islands Range Complex.....  | 2-3        |
|          | 2.1.1.1 Special Use Airspace and Air Traffic Controlled Assigned Airspace .....                                  | 2-3        |
|          | 2.1.1.2 Sea and Undersea Space.....  | 2-3        |
|          | 2.1.2 Description of the Ocean Operating Areas Outside the Bounds of the Mariana<br>Islands Range Complex .....  | 2-3        |
|          | 2.1.3 Description of Pierside Locations and Apra Harbor.....   | 2-7        |
| 2.2      | Primary Mission Areas .....  | 2-7        |
|          | 2.2.1 Air Warfare .....  | 2-7        |
|          | 2.2.2 Amphibious Warfare.....  | 2-9        |
|          | 2.2.3 Anti-Submarine Warfare.....  | 2-9        |
|          | 2.2.4 Electronic Warfare .....   | 2-9        |
|          | 2.2.5 Expeditionary Warfare.....   | 2-10       |

---

|         |   |      |
|---------|---|------|
| 2.2.6   | Mine Warfare.....   | 2-10 |
| 2.2.7   | Strike Warfare.....   | 2-11 |
| 2.2.8   | Surface Warfare.....  | 2-11 |
| 2.3     | Proposed Activities .....   | 2-11 |
| 2.3.1   | Changes to Proposed Activities .....  | 2-11 |
| 2.3.1.1 | New Technologies and Capabilities .....   | 2-12 |
| 2.3.1.2 | Renamed and Reorganized Testing Activities.....                                   | 2-12 |
| 2.3.2   | Proposed Training and Testing Activities.....                                     | 2-12 |
| 2.3.3   | Standard Operating Procedures .....   | 2-13 |
| 2.3.3.1 | High-Energy Laser Safety .....  | 2-13 |
| 2.3.3.2 | Sea Space and Airspace Deconfliction .....  | 2-14 |
| 2.3.3.3 | Target Deployment and Retrieval Safety.....                                       | 2-14 |
| 2.3.3.4 | Pierside Testing Safety.....  | 2-14 |
| 2.3.3.5 | Underwater Detonation Safety .....  | 2-15 |
| 2.3.3.6 | Sonic Booms.....  | 2-15 |
| 2.3.3.7 | Unmanned Surface Vehicle Safety .....   | 2-15 |
| 2.3.3.8 | Sinking Exercise Safety.....  | 2-15 |
| 2.3.4   | Mitigation Measures.....  | 2-15 |
| 2.4     | Action Alternatives Development.....  | 2-17 |
| 2.4.1   | Alternatives Eliminated from Further Consideration .....                          | 2-17 |
| 2.4.1.1 | Alternative Training and Testing Locations .....                                  | 2-17 |
| 2.4.1.2 | Reduced Training and Testing .....  | 2-18 |
| 2.4.1.3 | Alternatives Including Geographic Mitigation Measures within the Study Area ..... | 2-18 |
| 2.4.1.4 | Simulated Training and Testing Only .....   | 2-18 |
| 2.4.1.5 | Training and Testing Without the Use of Active Sonar .....                        | 2-19 |
| 2.4.2   | Alternatives Carried Forward.....   | 2-19 |
| 2.4.2.1 | No Action Alternative .....   | 2-19 |
| 2.4.2.2 | Alternative 1 .....   | 2-21 |
| 2.4.2.3 | Alternative 2 .....   | 2-22 |
| 2.4.3   | Comparison of Proposed Sonar and Explosive Use in the Action Alternatives .....   | 2-22 |
| 2.4.3.1 | Sonar Use .....   | 2-22 |
| 2.4.3.2 | Explosives Use.....   | 2-23 |
| 2.5     | Comparison of Alternatives .....  | 2-26 |

|          |  |            |
|----------|--|------------|
| <b>3</b> | <b>AFFECTED ENVIRONMENT AND ENVIRONMENTAL CONSEQUENCES .....</b>                         | <b>3-1</b> |
| 3.0      | Introduction .....   | 3-1        |
| 3.0.1    | Overall Approach to Analysis.....  | 3-1        |
| 3.0.1.1  | Navy Compiled and Generated Data .....   | 3-2        |
| 3.0.1.2  | Navy’s Quantitative Analysis to Determine Impacts on Sea Turtles and Marine Mammals..... | 3-3        |
| 3.0.2    | Regulatory Framework .....   | 3-7        |
| 3.0.3    | Resources and Issues Not Carried Forward for More Detailed Discussion .....              | 3-8        |
| 3.0.4    | Identification of Stressors for Analysis .....   | 3-8        |
| 3.0.4.1  | Acoustic Stressors.....  | 3-9        |
| 3.0.4.2  | Explosive Stressors.....   | 3-21       |
| 3.0.4.3  | Energy Stressors .....   | 3-24       |
| 3.0.4.4  | Physical Disturbance and Strike Stressors .....  | 3-25       |
| 3.0.4.5  | Entanglement Stressors.....  | 3-30       |
| 3.0.4.6  | Ingestion Stressors.....   | 3-34       |
| 3.0.4.7  | Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities .....  | 3-35       |
| 3.1      | Sediments and Water Quality.....   | 3.1-1      |
| 3.1.1    | Affected Environment.....  | 3.1-1      |
| 3.1.1.1  | Existing Conditions.....   | 3.1-1      |
| 3.1.2    | Environmental Consequences .....   | 3.1-5      |
| 3.1.2.1  | Explosives and Explosives Byproducts.....  | 3.1-6      |
| 3.1.2.2  | Metals .....   | 3.1-7      |
| 3.1.2.3  | Chemicals Other Than Explosives .....  | 3.1-8      |
| 3.1.2.4  | Other Materials .....  | 3.1-11     |
| 3.1.3    | Public Scoping Comments .....  | 3.1-12     |
| 3.2      | Air Quality .....  | 3.2-1      |
| 3.2.1    | Affected Environment.....  | 3.2-1      |
| 3.2.1.1  | Climate of the Study Area.....   | 3.2-1      |
| 3.2.1.2  | Regional Emissions .....   | 3.2-2      |
| 3.2.1.3  | Existing Air Quality.....  | 3.2-2      |
| 3.2.2    | Environmental Consequences .....   | 3.2-2      |
| 3.2.2.1  | Criteria Pollutants .....  | 3.2-2      |

---

|          |   |        |
|----------|---|--------|
| 3.2.2.2  | Hazardous Pollutants.....   | 3.2-4  |
| 3.2.2.3  | Greenhouse Gases.....   | 3.2-5  |
| 3.2.3    | Public Scoping Comments.....                                      | 3.2-5  |
| 3.3      | Marine Habitats.....  | 3.3-1  |
| 3.3.1    | Affected Environment.....   | 3.3-1  |
| 3.3.1.1  | Existing Conditions.....  | 3.3-1  |
| 3.3.2    | Environmental Consequences.....                                   | 3.3-3  |
| 3.3.2.1  | Explosive Stressors.....  | 3.3-3  |
| 3.3.2.2  | Physical Disturbance and Strike Stressors.....                    | 3.3-4  |
| 3.3.3    | Public Scoping Comments.....                                      | 3.3-6  |
| 3.4      | Marine Mammals.....   | 3.4-1  |
| 3.4.1    | Affected Environment.....   | 3.4-1  |
| 3.4.1.1  | General Background.....   | 3.4-1  |
| 3.4.1.2  | Species Unlikely to Be Present in the Study Area.....             | 3.4-3  |
| 3.4.1.3  | Group Size.....   | 3.4-3  |
| 3.4.1.4  | Habitat Use.....  | 3.4-4  |
| 3.4.1.5  | Dive Behavior.....  | 3.4-4  |
| 3.4.1.6  | Hearing and Vocalization.....                                     | 3.4-4  |
| 3.4.1.7  | General Threats.....  | 3.4-8  |
| 3.4.1.8  | Blue Whale ( <i>Balaenoptera musculus</i> ).....                  | 3.4-15 |
| 3.4.1.9  | Bryde’s Whale ( <i>Balaenoptera edeni</i> ).....                  | 3.4-16 |
| 3.4.1.10 | Fin Whale ( <i>Balaenoptera physalus</i> ).....                   | 3.4-18 |
| 3.4.1.11 | Humpback Whale ( <i>Megaptera novaeangliae</i> ).....             | 3.4-18 |
| 3.4.1.12 | Minke Whale ( <i>Balaenoptera acutorostrata</i> ).....            | 3.4-22 |
| 3.4.1.13 | Omura’s Whale ( <i>Balaenoptera omurai</i> ).....                 | 3.4-23 |
| 3.4.1.14 | Sei Whale ( <i>Balaenoptera borealis</i> ).....                   | 3.4-23 |
| 3.4.1.15 | Blainville’s Beaked Whale ( <i>Mesoplodon densirostris</i> )..... | 3.4-24 |
| 3.4.1.16 | Common Bottlenose Dolphin ( <i>Tursiops truncatus</i> ).....      | 3.4-25 |
| 3.4.1.17 | Cuvier’s Beaked Whale ( <i>Ziphius cavirostris</i> ).....         | 3.4-27 |
| 3.4.1.18 | Dwarf Sperm Whale ( <i>Kogia sima</i> ).....                      | 3.4-28 |
| 3.4.1.19 | False Killer Whale ( <i>Pseudorca crassidens</i> ).....           | 3.4-29 |
| 3.4.1.20 | Fraser’s Dolphin ( <i>Lagenodelphis hosei</i> ).....              | 3.4-30 |

|          |   |         |
|----------|---|---------|
| 3.4.1.21 | Ginkgo-Toothed Beaked Whale ( <i>Mesoplodon ginkgodens</i> ) .....                | 3.4-31  |
| 3.4.1.22 | Killer Whale ( <i>Orcinus orca</i> ) .....  | 3.4-32  |
| 3.4.1.23 | Longman’s Beaked Whale ( <i>Indopacetus pacificus</i> ) .....                     | 3.4-34  |
| 3.4.1.24 | Melon-Headed Whale ( <i>Peponocephala electra</i> ) .....                         | 3.4-35  |
| 3.4.1.25 | Pantropical Spotted Dolphin ( <i>Stenella attenuata</i> ) .....                   | 3.4-36  |
| 3.4.1.26 | Pygmy Killer Whale ( <i>Feresa attenuata</i> ) .....                              | 3.4-37  |
| 3.4.1.27 | Pygmy Sperm Whale ( <i>Kogia breviceps</i> ) .....                                | 3.4-38  |
| 3.4.1.28 | Risso’s Dolphin ( <i>Grampus griseus</i> ) .....                                  | 3.4-39  |
| 3.4.1.29 | Rough-Toothed Dolphin ( <i>Steno bredanensis</i> ).....                           | 3.4-40  |
| 3.4.1.30 | Short-Finned Pilot Whale ( <i>Globicephala macrorhynchus</i> ) .....              | 3.4-41  |
| 3.4.1.31 | Sperm Whale ( <i>Physeter macrocephalus</i> ).....                                | 3.4-42  |
| 3.4.1.32 | Spinner Dolphin ( <i>Stenella longirostris</i> ) .....                            | 3.4-44  |
| 3.4.1.33 | Striped Dolphin ( <i>Stenella coeruleoalba</i> ).....                             | 3.4-45  |
| 3.4.2    | Environmental Consequences .....  | 3.4-46  |
| 3.4.2.1  | Acoustic Stressors .....  | 3.4-50  |
| 3.4.2.2  | Explosive Stressors.....  | 3.4-175 |
| 3.4.2.3  | Energy Stressors .....  | 3.4-245 |
| 3.4.2.4  | Physical Disturbance and Strike Stressors .....                                   | 3.4-247 |
| 3.4.2.5  | Entanglement Stressors.....   | 3.4-249 |
| 3.4.2.6  | Ingestion Stressors.....  | 3.4-250 |
| 3.4.2.7  | Secondary Stressors.....  | 3.4-252 |
| 3.4.3    | Summary of Potential Impacts on Marine Mammals.....                               | 3.4-253 |
| 3.4.3.1  | Combined Impacts of All Stressors Under Alternative 1 .....                       | 3.4-255 |
| 3.4.3.2  | Combined Impacts of All Stressors Under Alternative 2 .....                       | 3.4-255 |
| 3.4.3.3  | Combined Impacts of All Stressors Under the No Action Alternative<br>.....        | 3.4-255 |
| 3.4.3.4  | Summary of Monitoring and Observations During Navy Activities Since<br>2015 ..... | 3.4-255 |
| 3.4.4    | Marine Mammal Protection Act Determinations.....                                  | 3.4-258 |
| 3.4.5    | Endangered Species Act Determinations .....                                       | 3.4-259 |
| 3.4.6    | Public Scoping Comments .....   | 3.4-259 |
| 3.5      | Sea Turtles.....  | 3.5-1   |
| 3.5.1    | Affected Environment.....   | 3.5-1   |

---

|          |  |        |
|----------|--|--------|
| 3.5.1.1  | Group Size.....  | 3.5-2  |
| 3.5.1.2  | Habitat Use .....  | 3.5-2  |
| 3.5.1.3  | Dive Behavior.....   | 3.5-3  |
| 3.5.1.4  | Hearing and Vocalization .....   | 3.5-4  |
| 3.5.1.5  | General Threats .....  | 3.5-5  |
| 3.5.1.6  | Green Sea Turtle ( <i>Chelonia mydas</i> ) .....                                 | 3.5-7  |
| 3.5.1.7  | Hawksbill Sea Turtle ( <i>Eretmochelys imbricata</i> ) .....                     | 3.5-8  |
| 3.5.1.8  | Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) .....                           | 3.5-9  |
| 3.5.1.9  | Olive Ridley Sea Turtle ( <i>Lepidochelys olivacea</i> ) .....                   | 3.5-9  |
| 3.5.1.10 | Leatherback Sea Turtle ( <i>Dermochelys coriacea</i> ) .....                     | 3.5-9  |
| 3.5.2    | Environmental Consequences .....   | 3.5-9  |
| 3.5.2.1  | Acoustic Stressors.....  | 3.5-11 |
| 3.5.2.2  | Explosive Stressors.....   | 3.5-23 |
| 3.5.2.3  | Energy Stressors .....   | 3.5-45 |
| 3.5.2.4  | Physical Disturbance and Strike Stressors .....                                  | 3.5-47 |
| 3.5.2.5  | Entanglement Stressors.....  | 3.5-49 |
| 3.5.2.6  | Ingestion Stressors.....   | 3.5-50 |
| 3.5.2.7  | Secondary Stressors.....   | 3.5-51 |
| 3.5.3    | Public Scoping Comments .....  | 3.5-51 |
| 3.6      | Marine Birds.....  | 3.6-1  |
| 3.6.1    | Affected Environment.....  | 3.6-1  |
| 3.6.1.1  | Group Size.....  | 3.6-2  |
| 3.6.1.2  | Diving Behavior .....  | 3.6-2  |
| 3.6.1.3  | Flight Altitudes.....  | 3.6-2  |
| 3.6.1.4  | Distance from Shore .....  | 3.6-3  |
| 3.6.1.5  | Hearing and Vocalization .....   | 3.6-3  |
| 3.6.1.6  | General Threats .....  | 3.6-5  |
| 3.6.1.7  | Rookery Locations and Breeding Activities on FDM.....                            | 3.6-5  |
| 3.6.2    | Environmental Consequences .....   | 3.6-6  |
| 3.6.2.1  | Acoustic Stressors.....  | 3.6-11 |
| 3.6.2.2  | Explosives Stressors (explosive shock wave and sound, explosive fragments) ..... | 3.6-19 |

---

|         |   |        |
|---------|---|--------|
| 3.6.2.3 | Energy Stressors .....  | 3.6-22 |
| 3.6.2.4 | Physical Disturbance and Strike Stressors .....                                 | 3.6-25 |
| 3.6.2.5 | Ingestion Stressors.....  | 3.6-28 |
| 3.6.2.6 | Secondary Stressors (Impacts on Habitat; Impacts on Prey Availability)<br>..... | 3.6-29 |
| 3.6.2.7 | Endangered Species Act Determinations .....                                     | 3.6-30 |
| 3.6.2.8 | Migratory Bird Treaty Act .....   | 3.6-30 |
| 3.6.3   | Public Scoping Comments .....   | 3.6-30 |
| 3.7     | Marine Vegetation .....   | 3.7-1  |
| 3.7.1   | Affected Environment.....   | 3.7-1  |
| 3.7.1.1 | General Threats .....   | 3.7-1  |
| 3.7.1.2 | Marine Vegetation Groups .....  | 3.7-1  |
| 3.7.1.3 | Seagrasses.....   | 3.7-1  |
| 3.7.1.4 | Mangroves .....   | 3.7-1  |
| 3.7.2   | Environmental Consequences .....  | 3.7-1  |
| 3.7.2.1 | Explosive Stressors.....  | 3.7-2  |
| 3.7.2.2 | Physical Disturbance and Strike Stressors .....                                 | 3.7-3  |
| 3.7.2.3 | Secondary Stressors.....  | 3.7-5  |
| 3.7.3   | Public Scoping Comments .....   | 3.7-5  |
| 3.8     | Marine Invertebrates .....  | 3.8-1  |
| 3.8.1   | Affected Environment.....   | 3.8-1  |
| 3.8.1.1 | Sound Sensing and Production .....  | 3.8-2  |
| 3.8.1.2 | General Threats .....   | 3.8-2  |
| 3.8.1.3 | Endangered Species Act-Listed Species.....                                      | 3.8-2  |
| 3.8.1.4 | Taxonomic Groups.....   | 3.8-3  |
| 3.8.2   | Environmental Consequences .....  | 3.8-3  |
| 3.8.2.1 | Acoustic Stressors.....   | 3.8-4  |
| 3.8.2.2 | Explosive Stressors.....  | 3.8-6  |
| 3.8.2.3 | Energy Stressors .....  | 3.8-8  |
| 3.8.2.4 | Physical Disturbance and Strike Stressors .....                                 | 3.8-10 |
| 3.8.2.5 | Entanglement Stressors.....   | 3.8-12 |
| 3.8.2.6 | Ingestion Stressors.....  | 3.8-13 |

---

|          |  |         |
|----------|--|---------|
| 3.8.2.7  | Secondary Stressors.....                                     | 3.8-15  |
| 3.8.3    | Public Scoping Comments .....                                | 3.8-16  |
| 3.9      | Fishes.....  | 3.9-1   |
| 3.9.1    | Affected Environment.....                                    | 3.9-1   |
| 3.9.1.1  | Hearing and Vocalization .....                               | 3.9-1   |
| 3.9.1.2  | General Threats .....  | 3.9-3   |
| 3.9.1.3  | Endangered Species Act Species.....                          | 3.9-3   |
| 3.9.1.4  | Federally Managed Species .....                              | 3.9-7   |
| 3.9.1.5  | Taxonomic Group Descriptions .....                           | 3.9-12  |
| 3.9.2    | Environmental Consequences .....                             | 3.9-12  |
| 3.9.2.1  | Acoustic Stressors.....                                      | 3.9-13  |
| 3.9.2.2  | Explosive Stressors.....                                     | 3.9-39  |
| 3.9.2.3  | Energy Stressors .....                                       | 3.9-52  |
| 3.9.2.4  | Physical Disturbance and Strike Stressors .....              | 3.9-54  |
| 3.9.2.5  | Entanglement Stressors.....                                  | 3.9-56  |
| 3.9.2.6  | Ingestion Stressors.....                                     | 3.9-57  |
| 3.9.2.7  | Secondary Stressors.....                                     | 3.9-59  |
| 3.9.3    | Public Scoping Comments .....                                | 3.9-61  |
| 3.10     | Terrestrial Species and Habitats .....                       | 3.10-1  |
| 3.10.1   | Affected Environment.....                                    | 3.10-1  |
| 3.10.1.1 | Vegetation Communities on Farallon de Medinilla .....        | 3.10-2  |
| 3.10.1.2 | Wildlife Communities on Farallon de Medinilla .....          | 3.10-3  |
| 3.10.1.3 | Endangered Species Act Listed Species .....                  | 3.10-3  |
| 3.10.1.4 | Major Terrestrial Species Taxonomic Group Descriptions ..... | 3.10-4  |
| 3.10.2   | Environmental Consequences .....                             | 3.10-5  |
| 3.10.2.1 | Acoustic Stressors.....                                      | 3.10-5  |
| 3.10.2.2 | Explosives Stressors.....                                    | 3.10-11 |
| 3.10.2.3 | Physical Disturbance and Strike Stressors .....              | 3.10-13 |
| 3.10.2.4 | Secondary Stressors.....                                     | 3.10-15 |
| 3.10.3   | Public Scoping Comments .....                                | 3.10-16 |
| 3.11     | Cultural Resources .....                                     | 3.11-1  |
| 3.11.1   | Affected Environment.....                                    | 3.11-1  |

---

|          |  |         |
|----------|--|---------|
| 3.11.1.1 | Guam .....   | 3.11-1  |
| 3.11.1.2 | Commonwealth of the Northern Mariana Islands .....             | 3.11-1  |
| 3.11.1.3 | Mariana Islands Training and Testing Transit Corridor.....     | 3.11-2  |
| 3.11.1.4 | Current Requirements, Practices, and Protective Measures ..... | 3.11-2  |
| 3.11.2   | Environmental Consequences .....                               | 3.11-2  |
| 3.11.2.1 | Explosive Stressors.....                                       | 3.11-3  |
| 3.11.2.2 | Physical Disturbance and Strike.....                           | 3.11-4  |
| 3.11.3   | Public Scoping Comments .....                                  | 3.11-6  |
| 3.12     | Socioeconomic Resources and Environmental Justice .....        | 3.12-1  |
| 3.12.1   | Affected Environment.....                                      | 3.12-1  |
| 3.12.1.1 | Commercial Transportation and Shipping .....                   | 3.12-3  |
| 3.12.1.2 | Commercial and Recreational Fishing .....                      | 3.12-7  |
| 3.12.1.3 | Tourism .....  | 3.12-11 |
| 3.12.1.4 | Environmental Justice.....                                     | 3.12-12 |
| 3.12.2   | Environmental Consequences .....                               | 3.12-15 |
| 3.12.2.1 | Accessibility (to the Ocean and Airspace) .....                | 3.12-15 |
| 3.12.2.2 | Airborne Acoustics.....  | 3.12-22 |
| 3.12.2.3 | Physical Disturbance and Strike Stressors .....                | 3.12-25 |
| 3.12.2.4 | Secondary Stressors.....                                       | 3.12-29 |
| 3.12.3   | Public Scoping Comments .....                                  | 3.12-30 |
| 3.13     | Public Health and Safety .....                                 | 3.13-1  |
| 3.13.1   | Affected Environment.....                                      | 3.13-1  |
| 3.13.1.1 | Existing Conditions.....                                       | 3.13-1  |
| 3.13.1.2 | Airspace .....   | 3.13-3  |
| 3.13.1.3 | Safety and Inspection Procedures .....                         | 3.13-3  |
| 3.13.2   | Environmental Consequences .....                               | 3.13-5  |
| 3.13.2.1 | Underwater Energy.....   | 3.13-6  |
| 3.13.2.2 | In-Air Energy .....  | 3.13-7  |
| 3.13.2.3 | Physical Interactions.....                                     | 3.13-8  |
| 3.13.2.4 | Secondary Stressors.....                                       | 3.13-9  |
| 3.13.3   | Public Scoping Comments .....                                  | 3.13-9  |

---

|          |   |            |
|----------|---|------------|
| <b>4</b> | <b>CUMULATIVE IMPACTS.....</b>  | <b>4-1</b> |
| 4.1      | Principles of Cumulative Impacts Analysis.....                                    | 4-1        |
| 4.1.1    | Determination of Significance .....   | 4-1        |
| 4.1.2    | Identifying Region of Influence for Cumulative Impacts Analysis .....             | 4-1        |
| 4.2      | Projects and Other Activities Analyzed for Cumulative Impacts .....               | 4-2        |
| 4.3      | Cumulative Impacts on Environmental Resources .....                               | 4-22       |
| 4.4      | Resource-Specific Cumulative Impacts .....  | 4-22       |
| 4.4.1    | Sediments and Water Quality.....  | 4-22       |
| 4.4.2    | Air Quality .....   | 4-23       |
| 4.4.3    | Marine Habitats .....   | 4-24       |
| 4.4.4    | Marine Mammals.....   | 4-25       |
| 4.4.4.1  | Region of Influence.....  | 4-25       |
| 4.4.4.2  | Resource Trends .....   | 4-25       |
| 4.4.4.3  | Impacts of Other Actions .....  | 4-26       |
| 4.4.4.4  | Impacts of the Proposed Action that May Contribute to Cumulative<br>Impacts ..... | 4-29       |
| 4.4.4.5  | Cumulative Impacts on Marine Mammals .....  | 4-30       |
| 4.4.5    | Sea Turtles .....   | 4-31       |
| 4.4.5.1  | Region of Influence.....  | 4-31       |
| 4.4.5.2  | Resource Trends .....   | 4-32       |
| 4.4.5.3  | Impacts of Other Actions .....  | 4-32       |
| 4.4.5.4  | Impacts of the Proposed Action That May Contribute to Cumulative<br>Impacts ..... | 4-35       |
| 4.4.5.5  | Cumulative Impacts on Sea Turtles .....   | 4-36       |
| 4.4.6    | Marine Birds .....  | 4-37       |
| 4.4.7    | Marine Vegetation.....  | 4-38       |
| 4.4.8    | Marine Invertebrates.....   | 4-39       |
| 4.4.8.1  | Region of Influence.....  | 4-39       |
| 4.4.8.2  | Resource Trends .....   | 4-39       |
| 4.4.8.3  | Impacts of Other Actions .....  | 4-39       |
| 4.4.8.4  | Impacts of the Proposed Action That May Contribute to Cumulative<br>Impacts ..... | 4-40       |
| 4.4.8.5  | Cumulative Impacts on Marine Invertebrates.....                                   | 4-41       |
| 4.4.9    | Marine Fishes.....  | 4-42       |
| 4.4.10   | Terrestrial Species and Habitats .....  | 4-43       |

---

---

|          |  |            |
|----------|--|------------|
| 4.4.11   | Cultural Resources .....                                     | 4-43       |
| 4.4.12   | Socioeconomic Resources .....                                | 4-44       |
| 4.4.12.1 | Resource Trends .....  | 4-45       |
| 4.4.12.2 | Impacts of Other Actions .....                               | 4-45       |
| 4.4.12.3 | Cumulative Impacts on Socioeconomic Resources .....          | 4-45       |
| 4.4.13   | Public Health and Safety .....                               | 4-45       |
| 4.5      | Summary of Cumulative Impacts .....                          | 4-46       |
| 4.6      | Public Scoping Comments .....                                | 4-46       |
| <b>5</b> | <b>MITIGATION .....</b>                                      | <b>5-1</b> |
| 5.1      | Introduction .....   | 5-1        |
| 5.1.1    | Benefits of Mitigation .....                                 | 5-2        |
| 5.1.2    | Compliance Initiatives.....                                  | 5-3        |
| 5.1.2.1  | Protective Measures Assessment Protocol .....                | 5-3        |
| 5.1.2.2  | Monitoring, Research, and Reporting Initiatives.....         | 5-3        |
| 5.2      | Mitigation Development Process.....                          | 5-8        |
| 5.2.1    | At-Sea Procedural Mitigation Development.....                | 5-9        |
| 5.2.1.1  | Lookouts .....   | 5-10       |
| 5.2.1.2  | Mitigation Zones .....                                       | 5-11       |
| 5.2.1.3  | Procedural Mitigation Implementation .....                   | 5-12       |
| 5.2.2    | At-Sea Mitigation Area Development.....                      | 5-12       |
| 5.2.3    | Terrestrial Mitigation Measure Development.....              | 5-13       |
| 5.2.4    | Practicality of Implementation .....                         | 5-13       |
| 5.2.4.1  | Assessment Criteria .....                                    | 5-14       |
| 5.2.4.2  | Factors Affecting Practicality .....                         | 5-17       |
| 5.3      | At-Sea Procedural Mitigation to be Implemented.....          | 5-19       |
| 5.3.1    | Environmental Awareness and Education.....                   | 5-19       |
| 5.3.2    | Acoustic Stressors.....                                      | 5-20       |
| 5.3.2.1  | Active Sonar .....   | 5-20       |
| 5.3.2.2  | Weapons Firing Noise .....                                   | 5-24       |
| 5.3.3    | Explosive Stressors.....                                     | 5-25       |
| 5.3.3.1  | Explosive Sonobuoys .....                                    | 5-25       |
| 5.3.3.2  | Explosive Torpedoes .....                                    | 5-28       |
| 5.3.3.3  | Explosive Medium-Caliber and Large-Caliber Projectiles ..... | 5-30       |
| 5.3.3.4  | Explosive Missiles and Rockets .....                         | 5-33       |

|          |   |            |
|----------|---|------------|
| 5.3.3.5  | Explosive Bombs .....   | 5-37       |
| 5.3.3.6  | Sinking Exercises .....   | 5-39       |
| 5.3.3.7  | Explosive Mine Countermeasure and Neutralization Activities .....                                 | 5-42       |
| 5.3.3.8  | Explosive Mine Neutralization Activities Involving Navy Divers.....                               | 5-45       |
| 5.3.3.9  | Maritime Security Operations – Anti-Swimmer Grenades .....  | 5-48       |
| 5.3.4    | Physical Disturbance and Strike Stressors .....   | 5-50       |
| 5.3.4.1  | Vessel Movement .....   | 5-50       |
| 5.3.4.2  | Towed In-Water Devices.....   | 5-53       |
| 5.3.4.3  | Small-, Medium-, and Large-Caliber Non-Explosive Practice Munitions                               | 5-54       |
| 5.3.4.4  | Non-Explosive Missiles and Rockets .....  | 5-55       |
| 5.3.4.5  | Non-Explosive Bombs and Mine Shapes .....   | 5-55       |
| 5.4      | At-Sea Mitigation Areas to be Implemented .....   | 5-56       |
| 5.4.1    | Mitigation Areas for Seafloor Resources .....   | 5-56       |
| 5.4.1.1  | Resource Description.....   | 5-60       |
| 5.4.1.2  | Mitigation Area Assessment.....   | 5-60       |
| 5.5      | Terrestrial Mitigation Measures to be Implemented .....   | 5-61       |
| 5.5.1    | Farallon De Medinilla.....  | 5-61       |
| 5.6      | Measures Considered but Eliminated.....   | 5-63       |
| 5.6.1    | Active Sonar .....  | 5-64       |
| 5.6.2    | Explosives.....   | 5-66       |
| 5.6.3    | Active and Passive Acoustic Monitoring Devices .....  | 5-67       |
| 5.6.4    | Thermal Detection Systems.....  | 5-69       |
| 5.6.5    | Third-Party Observers.....  | 5-71       |
| 5.6.6    | Foreign Navy Mitigation .....   | 5-71       |
| 5.6.7    | Reporting Requirements.....   | 5-72       |
| 5.7      | Mitigation Summary .....  | 5-73       |
| <b>6</b> | <b>ADDITIONAL REGULATORY CONSIDERATIONS .....</b>   | <b>6-1</b> |
| 6.1      | Consistency with Other Applicable Federal, State, and Local Plans, Policies, and Regulations..... | 6-1        |
| 6.1.1    | Coastal Zone Management Act Compliance.....   | 6-6        |
| 6.1.1.1  | Guam Coastal Management Program .....   | 6-6        |
| 6.1.1.2  | Commonwealth of the Northern Mariana Islands Coastal Zone Management Program.....                 | 6-6        |
| 6.1.2    | Marine Protected Areas.....   | 6-6        |
| 6.1.3    | Magnuson-Stevens Fishery Conservation and Management Act.....                                     | 6-7        |

|          |   |            |
|----------|---|------------|
| 6.2      | Relationship Between Short-Term Use of the Environment and Maintenance and Enhancement of Long-Term Productivity .....              | 6-7        |
| 6.3      | Irreversible or Irrecoverable Commitment of Resources.....  | 6-8        |
| 6.4      | Energy Requirements and Conservation Potential of Alternatives .....  | 6-8        |
| <b>7</b> | <b>LIST OF PREPARERS .....</b>  | <b>7-1</b> |
| <b>8</b> | <b>PUBLIC INVOLVEMENT AND DISTRIBUTION .....</b>  | <b>8-1</b> |
| 8.1      | Project Website.....  | 8-1        |
| 8.2      | Scoping Period .....  | 8-1        |
| 8.2.1    | Public Scoping Notification.....  | 8-1        |
| 8.2.1.1  | Notification Letters .....  | 8-1        |
| 8.2.1.2  | Postcard Mailers .....  | 8-11       |
| 8.2.1.3  | Newspaper Advertisements .....  | 8-14       |
| 8.2.1.4  | Press Release .....   | 8-15       |
| 8.2.2    | Public Scoping Comments .....   | 8-18       |
| 8.2.2.1  | Description of the Proposed Action and Alternatives .....   | 8-18       |
| 8.2.2.2  | National Environmental Policy Act and Public Involvement .....  | 8-18       |
| 8.2.2.3  | Location of Activities .....  | 8-18       |
| 8.2.2.4  | Indirect and Cumulative Impacts.....  | 8-18       |
| 8.2.2.5  | Sediments and Water Quality Impacts.....  | 8-18       |
| 8.2.2.6  | Socioeconomic Resources .....   | 8-19       |
| 8.2.2.7  | Terrestrial Species and Habitats/Marine Birds.....  | 8-19       |
| 8.2.2.8  | Marine Resources .....  | 8-19       |
| 8.2.2.9  | Marine Mammal Impacts/Sea Turtles .....   | 8-19       |
| 8.2.2.10 | Fish/Marine Habitat.....  | 8-19       |
| 8.2.2.11 | Cultural Resources .....  | 8-19       |
| 8.2.2.12 | Public Health and Safety.....   | 8-19       |
| 8.2.2.13 | Mitigation Measures.....  | 8-20       |
| 8.2.2.14 | Other.....  | 8-20       |
| 8.3      | Notification of Availability of the Draft Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement ..... | 8-20       |
| 8.3.1    | Notification of Draft Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement and Public Meetings ..... | 8-20       |
| 8.3.1.1  | Notification Letters .....  | 8-20       |

|         |   |      |
|---------|---|------|
| 8.3.1.2 | Postcards .....   | 8-20 |
| 8.3.1.3 | Press Releases.....   | 8-21 |
| 8.3.1.4 | Newspaper Advertisements .....  | 8-21 |
| 8.3.2   | Public Meetings .....   | 8-21 |
| 8.4     | Distribution of the Draft Supplemental Environmental Impact Statement/Overseas<br>Environmental Impact Statement..... | 8-21 |
| 8.4.1   | Federal Agencies .....  | 8-21 |
| 8.4.2   | Information Repositories .....  | 8-21 |

## List of Figures

### 1 Purpose and Need

|   |     |
|---|-----|
| Figure 1.1-1: Mariana Islands Training and Testing Study Area .....                       | 1-3 |
| Figure 1.5-1: National Environmental Policy Act Process Conducted for this SEIS/OEIS..... | 1-7 |

### 2 Description of Proposed Action and Alternatives

|   |      |
|---|------|
| Figure 2.1-1: Mariana Islands Training and Testing Study Area .....   | 2-2  |
| Figure 2.1-2: Mariana Islands Range Complex Airspace.....   | 2-4  |
| Figure 2.1-3: Warning Area 517 .....  | 2-5  |
| Figure 2.1-4: Farallon de Medinilla Restricted Area 7201, 7201A, and Danger Zone .....  | 2-6  |
| Figure 2.2-1: Apra Harbor Naval Complex (Main Base) and Main Base/Polaris Point .....   | 2-8  |
| Figure 2.4-1: Proposed Annual Total Hull-Mounted Mid-Frequency Sonar Hour Use Compared to the<br>Number Authorized in the 2015–2020 Marine Mammal Protection Act Permit ..... | 2-23 |
| Figure 2.4-2: Proposed Annual Explosives Use (Bins E1–E5) Compared to the 2015–2020 Marine<br>Mammal Protection Act Permit .....  | 2-24 |
| Figure 2.4-3: Proposed Annual Explosives Use (Bins E6–E12) Compared to the 2015–2020 Marine<br>Mammal Protection Act Permit .....   | 2-24 |

### 3 Affected Environment and Environmental Consequences

|  |      |
|--|------|
| Figure 3.0-1: Sonobuoy Launch Depicting the Relative Size of a Small Decelerator/Parachute ..... | 3-33 |
| Figure 3.0-2: Aerial Target (Drone) with Parachute Deployed .....                                | 3-33 |
| Figure 3.0-3: Flow Chart of the Evaluation Process of Sound-Producing Activities.....            | 3-37 |
| Figure 3.0-4: Two Hypothetical Threshold Shifts.....   | 3-41 |

#### 3.1 Sediments and Water Quality

There are no figures in this section.

#### 3.2 Air Quality

There are no figures in this section.

**3.3 Marine Habitats**

Figure 3.3-1: Nearshore Marine Habitats around Saipan ..... 3.3-2

**3.4 Marine Mammals**

Figure 3.4-1: Composite Audiograms for Hearing Groups Likely Found in the Study Area ..... 3.4-7

Figure 3.4-2: Two Hypothetical Threshold Shifts ..... 3.4-55

Figure 3.4-3: Odontocete Critical Ratios ..... 3.4-62

Figure 3.4-4: Critical Ratios for Different Noise Types ..... 3.4-63

Figure 3.4-5: Navy Auditory Weighting Functions for all Species Groups ..... 3.4-93

Figure 3.4-6: TTS and PTS Exposure Functions for Sonar and Other Transducers ..... 3.4-94

Figure 3.4-7: Behavioral Response Function for Odontocetes ..... 3.4-96

Figure 3.4-8: Behavioral Response Function for Mysticetes ..... 3.4-96

Figure 3.4-9: Behavioral Response Function for Beaked Whales ..... 3.4-97

Figure 3.4-10: Relative Likelihood of a Response Being Significant Based on the Duration and Severity of Behavioral Reactions ..... 3.4-98

Figure 3.4-11: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-115

Figure 3.4-12: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-116

Figure 3.4-13: Bryde’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-117

Figure 3.4-14: Bryde’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-118

Figure 3.4-15: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-119

Figure 3.4-16: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-120

Figure 3.4-17: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-121

Figure 3.4-18: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-122

Figure 3.4-19: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-123

Figure 3.4-20: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-124

Figure 3.4-21: Omura’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-125

Figure 3.4-22: Omura’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-126

Figure 3.4-23: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-127

Figure 3.4-24: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-129

Figure 3.4-25: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-134

Figure 3.4-26: Cuvier’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-135

Figure 3.4-27: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1..... 3.4-136

Figure 3.4-28: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-137

Figure 3.4-29: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-138

Figure 3.4-30: Cuvier’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-139

Figure 3.4-31: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2..... 3.4-140

Figure 3.4-32: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-141

Figure 3.4-33: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-142

Figure 3.4-34: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-143

Figure 3.4-35: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-144

Figure 3.4-36: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-145

Figure 3.4-37: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-146

Figure 3.4-38: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-147

Figure 3.4-39: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-148

Figure 3.4-40: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-149

Figure 3.4-41: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-150

Figure 3.4-42: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-151

Figure 3.4-43: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-153

Figure 3.4-44: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-154

Figure 3.4-45: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-155

Figure 3.4-46: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-156

Figure 3.4-47: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1..... 3.4-157

Figure 3.4-48: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2..... 3.4-158

Figure 3.4-49: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-159

Figure 3.4-50: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-160

Figure 3.4-51: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-161

Figure 3.4-52: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-162

Figure 3.4-53: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-163

Figure 3.4-54: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-164

Figure 3.4-55: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-165

Figure 3.4-56: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-166

Figure 3.4-57: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-167

Figure 3.4-58: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-168

Figure 3.4-59: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-169

Figure 3.4-60: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-170

Figure 3.4-61: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... 3.4-171

Figure 3.4-62: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... 3.4-172

Figure 3.4-63: Navy Phase III Weighting Functions for All Species Groups .....3.4-188

Figure 3.4-64: Navy Phase III Behavioral, TTS and PTS Exposure Functions for Explosives.....3.4-189

Figure 3.4-65: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 1 .....3.4-205

Figure 3.4-66: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 2 .....3.4-206

Figure 3.4-67: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 1 .....3.4-208

Figure 3.4-68: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 2 .....3.4-209

Figure 3.4-69: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 1 .....3.4-210

Figure 3.4-70: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 2 .....3.4-211

Figure 3.4-71: Omura’s Whale Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 2 .....3.4-212

Figure 3.4-72: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During  
Training and Testing Under Alternative 1 .....3.4-214

Figure 3.4-73: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During  
Training and Testing Under Alternative 2 .....3.4-215

Figure 3.4-74: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number  
of Explosions During Training and Testing Under Alternative 1 .....3.4-218

Figure 3.4-75: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 1 .....3.4-219

Figure 3.4-76: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number  
of Explosions During Training and Testing Under Alternative 2 .....3.4-220

Figure 3.4-77: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 2 .....3.4-221

Figure 3.4-78: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 1 .....3.4-223

Figure 3.4-79: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 1 .....3.4-224

Figure 3.4-80: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 2 .....3.4-225

Figure 3.4-81: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of  
Explosions During Training and Testing Under Alternative 2 .....3.4-226

Figure 3.4-82: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 1 .....3.4-228

Figure 3.4-83: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-229

Figure 3.4-84: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.4-231

Figure 3.4-85: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-232

Figure 3.4-86: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.4-233

Figure 3.4-87: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-234

Figure 3.4-88: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.4-236

Figure 3.4-89: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-237

Figure 3.4-90: Short-Finned Pilot Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-239

Figure 3.4-91: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.4-241

Figure 3.4-92: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-242

Figure 3.4-93: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.4-243

Figure 3.4-94: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 ..... 3.4-244

**3.5 Sea Turtles**

Figure 3.5-1: Dive Depth and Duration Summaries for Sea Turtle Species ..... 3.5-3

Figure 3.5-2: Generalized Dive Profiles and Activities Described for Sea Turtles..... 3.5-4

Figure 3.5-3: Composite Underwater Audiogram for Sea Turtles ..... 3.5-5

Figure 3.5-4: Auditory Weighting Function for Sea Turtles ..... 3.5-17

Figure 3.5-5: TTS and PTS Exposure Functions for Sonar and Other Transducers..... 3.5-18

Figure 3.5-6: Auditory Weighting Functions for Sea Turtles..... 3.5-32

Figure 3.5-7: TTS and PTS Exposure Functions for Impulsive Sounds..... 3.5-33

Figure 3.5-8: Green Sea Turtle Estimated Impacts per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 ..... 3.5-41

Figure 3.5-9: Green Sea Turtle Impacts Estimated per Year from Explosions During Training and Testing Under Alternative 2..... 3.5-44

**3.6 Marine Birds**

There are no figures in this section.

**3.7 Marine Vegetation**

There are no figures in this section.

**3.8 Marine Invertebrates**

There are no figures in this section.

**3.9 Fishes**

Figure 3.9-1: Fish Hearing Group and Navy Sonar Bin Frequency Ranges.....3.9-30

**3.10 Terrestrial Species and Habitats**

Figure 3.10-1: Farallon de Medinilla Impact Zones and Micronesian Megapode Occurrences .....3.10-7

**3.11 Cultural Resources**

There are no figures in this section.

**3.12 Socioeconomic Resources and Environmental Justice**

Figure 3.12-1: Commercial Fisheries Landings in Guam from the Years 2010 through 2015 .....3.12-8

Figure 3.12-2: Commercial Fisheries Landings in the CNMI from the Years 2010 through 2015.....3.12-10

Figure 3.12-3: Number of NOTMARs Issued for FDM and W-517 from the Years 2010 through 2017  
.....3.12-17

Figure 3.12-4: Number of Days per Year Affected by Military Activities at FDM and W-517 .....3.12-18

Figure 3.12-5: Joint Region Marianas Facebook Post Announcing Military Training Activities at FDM and  
W-517 .....3.12-20

**3.13 Public Health and Safety**

There are no figures in this section.

**4 Cumulative Impacts**

There are no figures in this chapter.

**5 Mitigation**

Figure 5.4-1: Seafloor Resource Mitigation Areas off Guam .....5-58

Figure 5.4-2: Seafloor Resource Mitigation Areas off Tinian and Saipan .....5-59

**6 Additional Regulatory Considerations**

There are no figures in this chapter.

**7 List of Preparers**

There are no figures in this chapter.

**8 Distribution List**

Figure 8-1: Notice of Intent..... 8-2  
Figure 8-2: Stakeholder Scoping Notification Letter..... 8-8  
Figure 8-3: Postcard Mailer for Scoping (Front) ..... 8-12  
Figure 8-4: Newspaper Announcement of Scoping ..... 8-14  
Figure 8-5: Commander, Joint Region Marianas Press Release..... 8-16

**List of Tables**

**1 Purpose and Need**

There are no tables in this chapter.

**2 Description of Proposed Action and Alternatives**

Table 2.3-1: Overview of Mitigation Categories ..... 2-16  
Table 2.5-1: Current and Proposed Training Activities ..... 2-26  
Table 2.5-2: Current and Proposed Testing Activities ..... 2-40

**3 Affected Environment and Environmental Consequences**

Table 3.0-1: Comparison of Stressors Analyzed..... 3-8  
Table 3.0-2: Sonar and Transducer Sources Quantitatively Analyzed ..... 3-13  
Table 3.0-3: Sonar and Transducers Qualitatively Analyzed..... 3-15  
Table 3.0-4: Representative Aircraft Sound Characteristics ..... 3-17  
Table 3.0-5: Sonic Boom Underwater Sound Levels Modeled for F/A-18 Hornet Supersonic Flight ..... 3-18  
Table 3.0-6: Example Weapons Noise ..... 3-19  
Table 3.0-7: Explosive Sources Quantitatively Analyzed that Could Be Used Underwater or at the Water  
Surface ..... 3-22  
Table 3.0-8: Typical Air Explosive Munitions During Navy Activities ..... 3-23  
Table 3.0-9: Annual Number of Events in the Study Area Including In-Water Electromagnetic Devices 3-24  
Table 3.0-10: Annual Number of Events in the Study Area Including High-Energy Lasers ..... 3-25  
Table 3.0-11: Annual Number of Events in the Study Area Including Aircraft Movement..... 3-26  
Table 3.0-12: Annual Number of Events in the Study Area Including Vessel Movement..... 3-26  
Table 3.0-13: Annual Number of Events in the Study Area Including In-Water Devices..... 3-26  
Table 3.0-14: Annual Number of Non-Explosive Practice Munitions Expended At Sea in the Study Area  
..... 3-27  
Table 3.0-15: Annual Number of Other Military Expended Materials Used At Sea in the Study Area.... 3-28

Table 3.0-16: Annual Number of Explosive Munitions Expended At Sea in the Study Area ..... 3-29

Table 3.0-17: Annual Number of Targets Expended At Sea in the Study Area ..... 3-29

Table 3.0-18: Annual Number of Events in the Study Area Including Seafloor Devices ..... 3-30

Table 3.0-19: Annual Number of Munitions Used on Farallon de Medinilla ..... 3-30

Table 3.0-20: Annual Number of Wires and Cables Expended in the Study Area ..... 3-32

Table 3.0-21: Size Categories for Decelerators/Parachutes Expended During Training and Testing  
Activities ..... 3-32

Table 3.0-22: Annual Number of Decelerators/Parachutes Expended in the Study Area ..... 3-34

Table 3.0-23: Annual Number of Chaff Cartridges Expended in the Study Area ..... 3-35

Table 3.0-24: Annual Number of Flares Expended in the Study Area ..... 3-35

**3.1 Sediments and Water Quality**

Table 3.1-1: Rates of Failure and Low-Order Detonations ..... 3.1-3

**3.2 Air Quality**

Table 3.2-1: Baseline Pollutant Emissions for At-Sea and FDM Training and Testing Activities that Occur  
Within 3 Nautical Miles of the Coast from Aircraft, Vessels, and Ordnance (tpy) ..... 3.2-3

Table 3.2-2: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3  
Nautical Miles of the Coast Under Alternative 1 from Aircraft, Vessels, and Ordnance (tpy)  
..... 3.2-4

Table 3.2-3: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3  
Nautical Miles of the Coast Under Alternative 2 from Aircraft, Vessels, and Ordnance (tpy)  
..... 3.2-4

Table 3.2-4: Annual Greenhouse Gas Emissions Under All Three Alternatives ..... 3.2-5

**3.3 Marine Habitats**

There are no tables in this section.

**3.4 Marine Mammals**

Table 3.4-1: Marine Mammal Occurrence within the Study Area ..... 3.4-2

Table 3.4-2: Species within Marine Mammal Hearing Groups Likely Found in the Study Area ..... 3.4-6

Table 3.4-3: Cutoff Distances for Moderate Source Level, Single Platform Training and Testing Events and  
for All Other Events with Multiple Platforms or Sonar with Source Levels at or Exceeding  
215 dB re 1  $\mu$ Pa at 1 m ..... 3.4-97

Table 3.4-4: Range to Permanent Threshold Shift for Five Representative Sonar Systems ..... 3.4-101

Table 3.4-5: Ranges to Temporary Threshold Shift for Sonar Bin HF4 over a Representative Range of  
Environments within the Study Area ..... 3.4-102

Table 3.4-6: Ranges to Temporary Threshold Shift for Sonar Bin LF4 over a Representative Range of  
Environments within the Study Area ..... 3.4-102

Table 3.4-7: Ranges to Temporary Threshold Shift for Sonar Bin MF1 over a Representative Range of Environments within the Study Area ..... 3.4-103

Table 3.4-8: Ranges to Temporary Threshold Shift for Sonar Bin MF4 over a Representative Range of Environments within the Study Area ..... 3.4-103

Table 3.4-9: Ranges to Temporary Threshold Shift for Sonar Bin MF5 over a Representative Range of Environments within the Study Area ..... 3.4-104

Table 3.4-10: Ranges to a Potentially Significant Behavioral Response for Sonar Bin HF4 over a Representative Range of Environments within the Study Area..... 3.4-104

Table 3.4-11: Ranges to a Potentially Significant Behavioral Response for Sonar Bin LF4 over a Representative Range of Environments within the Study Area..... 3.4-105

Table 3.4-12: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF1 over a Representative Range of Environments within the Study Area..... 3.4-106

Table 3.4-13: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF4 over a Representative Range of Environments within the Study Area..... 3.4-107

Table 3.4-14: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF5 over a Representative Range of Environments within the Study Area..... 3.4-108

Table 3.4-15: Criteria to Quantitatively Assess Non-Auditory Injury Due to Underwater Explosions 3.4-187

Table 3.4-16: Navy Phase III Weighted Sound Exposure Thresholds for Underwater Explosive Sounds ..... 3.4-190

Table 3.4-17: Ranges to Non-Auditory Injury (in meters) for All Marine Mammal Hearing Groups.. 3.4-192

Table 3.4-18: Ranges to Mortality (in meters) for All Marine Mammal Hearing Groups as a Function of Animal Mass ..... 3.4-193

Table 3.4-19: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for High-Frequency Cetaceans ..... 3.4-194

Table 3.4-20: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for High-Frequency Cetaceans ..... 3.4-195

Table 3.4-21: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Low-Frequency Cetaceans ..... 3.4-197

Table 3.4-22: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Low-Frequency Cetaceans ..... 3.4-198

Table 3.4-23: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Mid-Frequency Cetaceans ..... 3.4-200

Table 3.4-24: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Mid-Frequency Cetaceans ..... 3.4-201

**3.5 Sea Turtles**

Table 3.5-1: Endangered Species Act Status and Presence of Endangered Species Act Listed Sea Turtles in the Mariana Islands Training and Testing Study Area ..... 3.5-2

Table 3.5-2: Criteria to Quantitatively Assess Non-Auditory Injury due to Underwater Explosions.... 3.5-31

Table 3.5-3: TTS and PTS Peak Pressure Thresholds Derived for Sea Turtles Exposed to Impulsive Sounds ..... 3.5-33

Table 3.5-4: Ranges to Non-Auditory Injury (in meters) for Sea Turtles Exposed to Explosives as a Function of Animal Mass..... 3.5-35

Table 3.5-5: Ranges to Mortality for Sea Turtles Exposed to Explosives as a Function of Animal Mass ..... 3.5-36

Table 3.5-6: Peak Pressure Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives ..... 3.5-37

Table 3.5-7: SEL Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives..... 3.5-38

**3.6 Marine Birds**

Table 3.6-1: Number of Events by Activity Type on Farallon de Medinilla..... 3.6-8

Table 3.6-2: Number of Munitions by Activity Type on Farallon de Medinilla..... 3.6-9

Table 3.6-3: Munitions Use on Farallon de Medinilla, Net Explosive Weight Comparisons..... 3.6-9

**3.7 Marine Vegetation**

There are no tables in this section.

**3.8 Marine Invertebrates**

Table 3.8-1: Status of Endangered Species Act-Listed Species Within the Study Area ..... 3.8-3

**3.9 Fishes**

Table 3.9-1: Endangered Species Act Listed and Special Status Fish Species in the Mariana Islands Study Area ..... 3.9-4

Table 3.9-2: Federally Managed Fish Species within the Mariana Islands Study Area for Each Fishery Management Unit in the FEP ..... 3.9-8

Table 3.9-3: Sound Exposure Criteria for TTS from Sonar ..... 3.9-31

Table 3.9-4: Ranges to Temporary Threshold Shift from Four Representative Sonar Bins ..... 3.9-32

Table 3.9-5: Range to 10 Percent Mortality from In-water Explosions for Fishes with a Swim Bladder ..... 3.9-42

Table 3.9-6: Sound Exposure Criteria for Mortality and Injury from Explosives ..... 3.9-47

Table 3.9-7: Sound Exposure Criteria for Hearing Loss from Explosives ..... 3.9-48

Table 3.9-8: Range to Mortality and Injury for All Fishes from Explosives ..... 3.9-49

Table 3.9-9: Range to TTS for Fishes with a Swim Bladder from Explosives..... 3.9-50

**3.10 Terrestrial Species and Habitats**

Table 3.10-1: Endangered Species Act Listed Species on Farallon de Medinilla ..... 3.10-2

**3.11 Cultural Resources**

There are no tables in this section.

**3.12 Socioeconomic Resources and Environmental Justice**

There are no tables in this section.

**3.13 Public Health and Safety**

Table 3.13-1: Marine Protected Areas within the Study Area ..... 3.13-2

**4 Cumulative Impacts**

Table 4.2-1: Past, Present, and Reasonably Foreseeable Actions ..... 4-3

**5 Mitigation**

Table 5.3-1: Environmental Awareness and Education ..... 5-19

Table 5.3-2: Procedural Mitigation for Active Sonar ..... 5-21

Table 5.3-3: Procedural Mitigation for Weapons Firing Noise ..... 5-24

Table 5.3-4: Procedural Mitigation for Explosive Sonobuoys ..... 5-26

Table 5.3-5: Procedural Mitigation for Explosive Torpedoes ..... 5-28

Table 5.3-6: Procedural Mitigation for Explosive Medium-Caliber and Large-Caliber Projectiles ..... 5-31

Table 5.3-7: Procedural Mitigation for Explosive Missiles and Rockets ..... 5-34

Table 5.3-8: Procedural Mitigation for Explosive Bombs ..... 5-37

Table 5.3-9: Procedural Mitigation for Sinking Exercises ..... 5-40

Table 5.3-10: Procedural Mitigation for Explosive Mine Countermeasure and Neutralization Activities  
..... 5-43

Table 5.3-11: Procedural Mitigation for Explosive Mine Neutralization Activities Involving Navy Divers  
..... 5-46

Table 5.3-12: Procedural Mitigation for Maritime Security Operations – Anti-Swimmer Grenades ..... 5-49

Table 5.3-13: Procedural Mitigation for Vessel Movement ..... 5-51

Table 5.3-14: Procedural Mitigation for Towed In-Water Devices ..... 5-53

Table 5.3-15: Procedural Mitigation for Small-, Medium-, and Large-Caliber Non-Explosive Practice  
Munitions ..... 5-54

Table 5.3-16: Procedural Mitigation for Non-Explosive Missiles and Rockets ..... 5-55

Table 5.3-17: Procedural Mitigation for Non-Explosive Bombs and Mine Shapes ..... 5-56

Table 5.4-1: Mitigation Areas for Seafloor Resources ..... 5-57

Table 5.5-1: Farallon de Medinilla Mitigation Measures ..... 5-62

Table 5.7-1: Summary of At-Sea Procedural Mitigation ..... 5-73

Table 5.7-2: Summary of Mitigation Areas ..... 5-74

Table 5.7-3: Summary of Terrestrial Mitigation ..... 5-74

**6 Additional Regulatory Considerations**

Table 6.1-1: Summary of Environmental Compliance for the Proposed Action ..... 6-2

**7 List of Preparers**

There are no tables in this chapter.

**8 Distribution List**

Table 8-1: Federal and Local Entities that Received the Scoping Notification Letter.....8-4  
Table 8-2: Newspaper Publications ..... 8-14  
Table 8-3: Newspaper Announcements of Draft Supplemental EIS/OEIS and Public Meetings .....8-21  
Table 8-4: Information Repositories ..... 8-22

## Appendices

**Appendix A** Training and Testing Activities Descriptions  
**Appendix B** Federal Register Notices  
**Appendix C** Agency Correspondence  
**Appendix D** Air Quality Emissions Calculations  
**Appendix E** Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities  
**Appendix F** Training and Testing Activities Matrices  
**Appendix G** Conceptual Framework for Assessing Effects on Biological Resources  
**Appendix H** Acoustic and Explosive Concepts  
**Appendix I** Geographic Mitigation Assessment  
**Appendix J** Statistical Probability Analysis for Estimating Direct Strike Impact and Number of Potential Exposures from Military Expended Materials

## Acronyms and Abbreviations

| Acronym                     | Definition   | Acronym | Definition                                    |
|-----------------------------|--|---------|---|
| μPa                         | Micropascal(s)   | EO      | Executive Order                               |
| ac.                         | Acre(s)  | EPA     | Environmental Protection Agency               |
| AGL                         | Above Ground Level   | ESA     | Endangered Species Act                        |
| AMRAAM                      | Advanced Medium-Range Air-to-Air Missile                             | FAA     | Federal Aviation Administration               |
| ANSI                        | American National Standards Institute                                | FDM     | Farallon de Medinilla                         |
| BO                          | Biological Opinion   | FEP     | Fishery Ecosystem Plan                        |
| BOBO                        | 2nd Lt. John P. Bobo class vessel                                    | FR      | Federal Register                              |
| CD                          | Consistency Determination  | ft.     | Foot/feet                                     |
| CEQ                         | Council on Environmental Quality                                     | FY      | Fiscal Year                                   |
| CFR                         | Code of Federal Regulations  | GDP     | Gross Domestic Product                        |
| CJMT                        | Commonwealth of the Northern Mariana Islands Joint Military Training | GUNEX   | Gunnery Exercise                              |
| CNMI                        | Commonwealth of the Northern Mariana Islands                         | ha      | Hectare                                       |
| CO                          | Carbon Monoxide  | HAPC    | Habitat Area of Particular Concern            |
| CV                          | Coefficient of Variation   | HARM    | High-Speed Anti-Radiation Missile             |
| dB                          | Decibel(s)   | HE-ET   | High Explosive-Electronic Time                |
| dB re 1 μPa                 | Decibels referenced to 1 micropascal                                 | HF      | High Frequency                                |
| dB re 1 μPa <sup>2</sup> s  | Decibels referenced to 1 micropascal squared seconds                 | HMX     | High Melting Explosive                        |
| dBA                         | A-weighted decibel   | hr.     | Hour(s)                                       |
| dBA re 1 μPa                | A-weighted db referenced to 1 micropascal                            | HRC     | Hawaii Range Complex                          |
| dBA re 1 μPa <sup>2</sup> s | A-weighted db referenced to 1 micropascal squared seconds            | HUMMA   | Hawaii Undersea Military Munitions Assessment |
| DDT                         | Dichlorodiphenyltrichloroethane                                      | Hz      | Hertz   |
| DNT                         | Dinitrotoluene   | in.     | Inch(es)                                      |
| DoD                         | Department of Defense  | JRM     | Joint Region Marianas                         |
| DPS                         | Distinct Population Segment  | kHz     | Kilohertz                                     |
| EEZ                         | Exclusive Economic Zone  | km      | Kilometers                                    |
| EFH                         | Essential Fish Habitat   | lb.     | Pound(s)                                      |
| EFHA                        | Essential Fish Habitat Assessment                                    | LF      | Low Frequency                                 |
| EIS                         | Environmental Impact Statement                                       | LMSR    | Large, Medium Speed, Roll On/Roll Off         |
|                             |  | LOA     | Letter of Authorization                       |
|                             |  | m       | Meter(s)                                      |
|                             |  | MBTA    | Migratory Bird Treaty Act                     |
|                             |  | MF      | Mid-Frequency                                 |
|                             |  | MHz     | Megahertz                                     |
|                             |  | mi.     | Mile(s)                                       |

| Acronym         | Definition   | Acronym           | Definition                                  |
|-----------------|--|-------------------|---|
| MIRC            | Mariana Islands Range Complex                            | PM <sub>10</sub>  | Particulate matter less than 10 microns     |
| MISSILEX        | Missile Exercise   | PM <sub>2.5</sub> | Particulate matter less than 2.5 microns    |
| MITT            | Mariana Islands Training and Testing                     | PSD               | Prevention of Significant Deterioration     |
| mm              | millimeter(s)  | psi               | Pounds per square inch                      |
| MMPA            | Marine Mammal Protection Act                             | Psi-ms            | Pounds per square inch per millisecond      |
| MPA             | Marine Protected Area                                    | PTS               | Permanent Threshold Shift                   |
| MSA             | Magnuson-Stevens Fishery Conservation and Management Act | R                 | Restricted Area                             |
| MSC             | Military Sealift Command                                 | RDX               | Royal Demolition Explosive                  |
| Navy            | U.S. Department of the Navy                              | RL                | Received SPL                                |
| NEPA            | National Environmental Policy Act                        | rms               | Root-Mean-Square                            |
| NEPM            | Non-Explosive Practice Munition                          | ROD               | Record of Decision                          |
| NEW             | Net explosive weight                                     | SAR               | Stock Assessment Report                     |
| NM              | Nautical Mile(s)   | SEIS              | Supplemental Environmental Impact Statement |
| NM <sup>2</sup> | Square Nautical Miles                                    | SEL               | Sound Exposure Level                        |
| NMFS            | National Marine Fisheries Service                        | SL                | Source Level                                |
| NOAA            | National Oceanic and Atmospheric Administration          | SO <sub>x</sub>   | Sulfur Oxide                                |
| NOTAM           | Notice to Airmen   | SPL               | Sound Pressure Level                        |
| NOTMAR          | Notice to Mariners                                       | tpy               | Tons per year                               |
| NO <sub>x</sub> | Nitrogen oxide   | TS                | Threshold Shift                             |
| OEIS            | Overseas Environmental Impact Statement                  | TTS               | Temporary Threshold Shift                   |
| Pa              | Pascal   | U.S.              | United States                               |
| Pa-s            | Pascal-second(s)   | U.S.C.            | United States Code                          |
| PCB             | Polychlorinated Biphenyl                                 | UNDET             | Underwater Detonation                       |
|                 |  | USFWS             | U.S. Fish and Wildlife Service              |
|                 |  | VOC               | Volatile Organic Compound                   |
|                 |  | W                 | Warning Area                                |

---

---

# 1 Purpose and Need



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

**1**            **PURPOSE AND NEED .....1-1**

1.1      Introduction ..... 1-1

1.2      The Navy’s Environmental Compliance and At-Sea Policy ..... 1-3

1.3      Proposed Action..... 1-4

1.4      Purpose and Need..... 1-4

            1.4.1 Why the Navy Trains..... 1-5

            1.4.2 Why the Navy Tests ..... 1-6

1.5      Overview and Strategic Importance of Existing Range Complex..... 1-6

1.6      The Environmental Planning Process..... 1-7

            1.6.1 National Environmental Policy Act Requirements ..... 1-8

            1.6.2 Executive Order 12114 ..... 1-8

            1.6.3 Other Environmental Requirements Considered ..... 1-8

1.7      Scope and Content..... 1-9

1.8      Organization of this Supplemental Environmental Impact Statement/Overseas  
            Environmental Impact Statement..... 1-9

**List of Figures**

Figure 1.1-1: Mariana Islands Training and Testing Study Area ..... 1-3

Figure 1.5-1: National Environmental Policy Act Process Conducted for this SEIS/OEIS..... 1-7

**List of Tables**

There are no tables in this chapter.

This page intentionally left blank.

# 1 Purpose and Need

## 1.1 Introduction

The United States (U.S.) Department of the Navy (Navy) has prepared this supplement to the May 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) (U.S. Department of the Navy, 2015) pursuant to Council on Environmental Quality Regulations. The Navy proposes to conduct training activities (referred to as “training”), and research, development, testing and evaluation (referred to as “testing”) activities in the MITT Study Area, primarily within the existing Mariana Islands Range Complex (MIRC), as represented in Figure 1.1-1. Training and testing activities, collectively referred to as “military readiness activities,” that prepare the Navy to fulfill its mission to protect and defend the United States and its allies, have the potential to impact the environment. The Navy prepared this Supplemental EIS (SEIS)/OEIS to comply with the National Environmental Policy Act (NEPA) and Executive Order 12114, *Environmental Effects Abroad of Major Federal Actions*, by reassessing the potential environmental impacts associated with the proposed military readiness activities to be conducted within the Study Area.

This SEIS/OEIS was prepared to update the Navy’s assessment of the potential environmental impacts associated with proposed training and testing to be conducted at sea and on Farallon de Medinilla (FDM). These proposed activities are generally consistent with those at-sea and FDM activities analyzed in the May 2015 *Final Mariana Islands Training and Testing Activities Environmental Impact Statement/Overseas Environmental Impact Statement* (U.S. Department of the Navy, 2015), referred to as the 2015 MITT Final EIS/OEIS, and are representative of activities the military has been conducting in the Study Area for decades. These military readiness activities include the use of active sonar and explosives at sea off the coasts of Guam and the Commonwealth of the Northern Mariana Islands (CNMI), throughout the in-water areas around the MIRC, the transit corridor between the MIRC and the Hawaii Range Complex, and at select Navy pierside and harbor locations.

The 2015 MITT Final EIS/OEIS also analyzed training and testing activities conducted at existing MIRC land-based training areas located on Guam, Saipan, Tinian, and Rota. The Navy consulted with the U.S. Fish and Wildlife Service regarding effects of the land-based training activities on terrestrial species listed under the Endangered Species Act (ESA) and received a Biological Opinion (U.S. Fish and Wildlife Service, 2015) and concurrence letter (U.S. Fish and Wildlife Service, 2016). As the Navy is not proposing any changes to those land-based activities on Guam, Saipan, Tinian, and Rota, the Navy will continue to rely on the 2015 MITT Final EIS/OEIS because there is no new information that would affect the EIS analysis. In addition, in accordance with 50 Code of Federal Regulations (CFR) Section 402.16, the 2015 and 2016 consultations remain valid as none of the factors necessary to trigger reinitiating consultation have been met.

New information specifically addressed in this SEIS/OEIS includes updates to training and testing requirements, an updated acoustic effects model, updated marine mammal density data, and evolving

and emergent best available science.<sup>1</sup> Using the updated information, the Navy will seek the reissuance of federal regulatory permits and authorizations under the Marine Mammal Protection Act (MMPA) and ESA to support training and testing requirements within the Study Area beyond the 2020 expiration of current authorizations and consultation. The Navy will consult with the National Marine Fisheries Service (NMFS) to renew these authorizations. While the 2015 MITT Final EIS/OEIS Study Area remains unchanged, this SEIS/OEIS focuses on the at-sea and FDM portion of that area. The Study Area consists of three primary components: (1) the MIRC, (2) additional areas on the high seas, and (3) a transit corridor between the MIRC and the Hawaii Range Complex. Collectively, these areas continue to be referred to as the MITT Study Area (Figure 1.1-1).

The United States is facing increased global disorder, characterized by decline in the long-standing rules-based international order—creating a more complex and volatile security environment. Major conflicts, terrorism, outlaw actions, and natural disasters all have the potential to threaten national security of the United States. The security, prosperity, and vital interests of the United States are increasingly tied to other nations because of the close relationships between the United States and other national economies. The Navy operates on the world’s oceans, seas, and coastal areas—the international maritime domain—on which 90 percent of the world’s trade and two-thirds of its oil are transported. The majority of the world’s population also lives within a few hundred miles of an ocean. The U.S. Navy carries out training and testing activities to be able to protect the United States against its potential adversaries, to protect and defend the rights and interests of the United States and its allies to move freely on the oceans, and to provide humanitarian assistance.

Department of Defense realignment/reassignment efforts in the Western Pacific have previously been, or are currently, the subject of various environmental planning processes, including the EIS and Supplemental EIS, which studied the realignment of Marine Corps forces to Guam, ongoing EIS efforts to address joint training and land-based training requirements in the CNMI, and EIS efforts to discuss Air Force divert landing and training requirements in the CNMI. The training and testing activities covered by the 2015 MITT Final EIS/OEIS, as well as in this supplement, are separate and distinct and have independent utility from the actions proposed by Marine Corps forces and those of the U.S. Air Force within the CNMI. This SEIS/OEIS only addresses ongoing and future at-sea and FDM training and testing activities that are independent and do not rely on any realignment efforts. Further, the training and testing activities covered by the 2015 MITT Final EIS/OEIS and this SEIS/OEIS have been occurring in the Study Area for decades and would continue regardless of whether any of the other Department of Defense efforts in the Western Pacific come to fruition.

---

<sup>1</sup> The 2015 MITT Final EIS/OEIS used a new modeling system known as the Navy Acoustics Effects Model and marine mammal density information, developed by the Navy in cooperation with the National Marine Fisheries Service, that was the best available information at the time. The Navy Acoustics Effects Model has been refined, marine mammal density estimates have been updated, NMFS has published new criteria, and criteria used in the acoustic model have been revised.




Figure 1.1-1: Mariana Islands Training and Testing Study Area

## 1.2 The Navy’s Environmental Compliance and At-Sea Policy

In 2000, the Navy completed a review of its environmental compliance requirements for exercises and training at sea. The Navy then instituted the “At-Sea Policy” (U.S. Department of the Navy, 2000) to ensure compliance with applicable environmental regulations and policies, and preserve the flexibility necessary for the Navy to train and test at sea. This policy directed, in part, that Fleet Commanders develop a programmatic approach to environmental compliance at sea for ranges and operational areas within their respective geographic areas of responsibility (U.S. Department of the Navy, 2000). Those ranges affected by the “At-Sea Policy” are designated water areas, sometimes containing instrumentation, which are managed and used to conduct training and testing activities. Some ranges are further broken down into operational areas, to better manage and deconflict military readiness activities.

In 2005, the Navy and the National Oceanic and Atmospheric Administration reached an agreement on a coordinated programmatic strategy for assessing certain environmental effects of military readiness activities at sea.

The Navy is currently in the third phase of implementing this programmatic approach, which covers similar types of military readiness training and testing activities in the same MITT Study Area analyzed in Phase II. As was done in Phase I and Phase II, the Navy will use the Phase III analysis to support regulatory consultations and a request for a letter of authorization under the MMPA and incidental take statements under the ESA. Given that the training and testing activities and many areas of environmental analysis remain similar to those addressed in Phase II, and the same Study Area is used for the proposed activities, the Navy determined an SEIS/OEIS to be appropriate for Phase III of the Navy's environmental compliance planning in the MITT Study Area. For further discussion of the first two phases, please see Section 1.2 (The Navy's Environmental Compliance and At-Sea Policy) of the 2015 MITT Final EIS/OEIS.

### 1.3 Proposed Action

The Navy's Proposed Action, described in detail in Chapter 2 (Description of Proposed Action and Alternatives), is to conduct military readiness training and testing activities in the Study Area (Figure 1.1-1).

### 1.4 Purpose and Need

The Navy and NMFS (as a cooperating agency) have coordinated from the outset and developed this document to meet each agency's distinct NEPA obligations and support the decision making of both agencies. The Navy's purpose of the Proposed Action is to conduct training and testing activities to ensure that the Navy and other Services meet their respective missions, which, for the Navy, is to maintain, train, and equip combat-ready military forces capable of winning wars, deterring aggression, and maintaining freedom of the seas. The respective missions are achieved in part by conducting training and testing within the Study Area in accordance with established Navy military readiness requirements. The sections that follow provide a description of the need for military readiness activities. Appendix A (Training and Testing Activities Descriptions) provides detailed Navy and other Services' activities descriptions.

Title 10 section 5062 of the U.S. Code provides: "The Navy shall be organized, trained, and equipped primarily for prompt and sustained combat incident to operations at sea. It is responsible for the preparation of naval forces necessary for the effective prosecution of war except as otherwise assigned and, in accordance with integrated joint mobilization plans, for the expansion of the peacetime components of the Navy to meet the needs of war."

The Navy has requested authorization to take marine mammals incidental to conducting their training and testing activities in the Study Area by Level A and B harassment, serious injury, and/or mortality. Take under the MMPA is defined as "to harass, hunt, capture, or kill, or attempt to harass, hunt, capture, or kill any marine mammal." For military readiness activities, harassment is defined as "(i) any act that injures or has the significant potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment] or (ii) any act that disturbs or is likely to disturb a marine mammal or marine mammal stock in the wild by causing disruption of natural behavioral patterns, including, but not limited to, migration, surfacing, nursing, breeding,

feeding, or sheltering, to a point where such behavioral patterns are abandoned or significantly altered [Level B harassment].”

NMFS has issued proposed regulations and is considering issuance of a subsequent Letter of Authorization (LOA) under section 101(a)(5)(A) of the MMPA of 1972, as amended (16 United States Code [U.S.C.] 1361 et seq.) that would govern the taking of marine mammals incidental to the Navy training and testing activities within the Study Area. The issuance of regulations and associated LOA to the Navy is a major federal action requiring NMFS to analyze the effects of their issuance on the human environment pursuant to NEPA requirements and National Oceanic and Atmospheric Administration policies.

The purpose of issuing an incidental take authorization is to provide an exception to the take prohibition in the MMPA and to ensure that the action complies with the MMPA and implementing regulations. Incidental take authorizations may be issued as either (1) regulations and associated LOA under section 101(a)(5)(A) of the MMPA or (2) Incidental Harassment Authorization under section 101(a)(5)(D) of the MMPA. An Incidental Harassment Authorization can be issued only when there is no potential for serious injury or mortality or where any such potential can be negated through required mitigation measures. Because some of the activities under the Proposed Action may create a potential for lethal takes or takes that may result in serious injury that could lead to mortality, the Navy is requesting rulemaking and the issuance of an LOA for this action.

NMFS’s purpose is to evaluate the Navy’s Proposed Action pursuant to NMFS’s authority under the MMPA, and to make a determination whether to issue incidental take regulations and an LOA, including any conditions needed to meet the statutory mandates of the MMPA. To authorize the incidental take of marine mammals, NMFS evaluates the best available scientific information to determine whether the take would have a negligible impact on the affected marine mammal species or stocks and an unmitigable impact on their availability for taking for subsistence uses (not relevant here for Navy’s proposed action). NMFS must also prescribe permissible methods of taking, other “means of effecting the least practicable adverse impact” on the affected species or stocks and their habitat, and monitoring and reporting requirements. NMFS cannot issue an incidental take authorization unless it can make the required findings. The need for NMFS’s action is to consider the impacts of the Navy’s activities on marine mammals and meet NMFS’ obligations under the MMPA. This SEIS/OEIS analyzes the environmental impacts associated with issuance of the requested authorization of the take of marine mammals incidental to the training and testing activities within the Study Area, to include a variety of mitigation measures that were considered during the MMPA authorization process. The analysis of mitigation measures considers benefits to species or stocks and their habitat, and analyzes the practicability and efficacy of each measure. This analysis of mitigation measures was used to support requirements pertaining to mitigation, monitoring, and reporting that would be specified in final MMPA regulations and subsequent LOA.

#### **1.4.1 Why the Navy Trains**

As described above, the Navy is statutorily mandated to protect U.S. national security by being ready, at all times, to effectively prosecute war and defend the nation by conducting operations at sea. The Navy is essential to protecting U.S. national interests, considering that 70 percent of the earth is covered in water, 80 percent of the planet’s population lives within close proximity to coastal areas, and 90 percent of global commerce is conducted by sea. Naval forces must be ready for a variety of military operations to address the dynamic, social, political, economic, and environmental issues that occur in today’s rapidly evolving world. Through its continuous presence on the world’s oceans, the Navy can respond to

a wide range of situations because, on any given day, over one-third of its ships, submarines, and aircraft are deployed overseas. Units must be able to respond promptly and effectively while forward deployed. This presence helps to dissuade aggression, which prevents conflict escalation, and provides the President with options to promptly address global contingencies. Before deploying, naval forces must train to develop a broad range of capabilities to respond to threats, from full-scale armed conflict in a variety of different geographic areas and environmental conditions to humanitarian assistance and disaster relief efforts. Training prepares Navy personnel to be proficient in safely operating and maintaining the equipment, weapons, and systems they will use to conduct their assigned missions. The training process provides personnel with an in-depth understanding of their individual limits and capabilities; the training process also helps the testing community improve new weapon systems' capabilities and effectiveness. Refer to Chapter 1, Section 1.4.1 (Why the Navy Trains) in the 2015 MITT Final EIS/OEIS for additional information on Navy training.

#### **1.4.2 Why the Navy Tests**

The Navy's research and acquisition community, including research-funding organizations, laboratory facilities, and systems commands, has a mission to provide weapons, systems, and platforms for the Navy to support its missions and ensure a technological edge over the United States' potential adversaries. This community is at the forefront of researching, developing, testing, evaluating, acquiring, and delivering modern platforms, systems, and related equipment to meet Fleet capability and readiness requirements. The Navy's research funding organizations and laboratories concentrate primarily on the development of new science and technology, and the initial testing of concepts that are relevant to the Navy of the future. As a result, systems commands develop ship, aircraft, and weapons products that support all Navy platforms throughout their lifecycles from systems acquisition through sustainment to end of life. Refer to Chapter 1, Section 1.5.1 (Why the Navy Tests) in the 2015 MITT Final EIS/OEIS for additional information on Navy testing. The Navy's research and acquisition community operating in the Study Area includes the following:

- The Naval Sea Systems Command, which develops, acquires, delivers, and maintains surface ships, submarines, unmanned vehicles, and weapon system platforms.
- The Naval Air Systems Command, which develops, tests, acquires, delivers, and sustains naval aviation aircraft, unmanned aerial systems, weapons, and systems.
- The Office of Naval Research, which plans, fosters, and encourages scientific research that promotes future naval sea power and enhances national security.

#### **1.5 Overview and Strategic Importance of Existing Range Complex**

The Navy has historically used areas in the Study Area for training and testing. The Navy has designated a portion of the Study Area as a "range complex." A range complex provides a controlled environment where military ship, submarine, and aircraft crews can train in realistic conditions while safely deconflicting with non-military activities, such as civilian shipping and aircraft. Sufficient sea and airspace in proximity to land training ranges, airfields, nearshore amphibious landing sites, and special use airspace is critical to realistic training and testing. Diverse and realistic training is critical to ensuring U.S. Forces, when needed, are both ready and able to effectively conduct operations in myriad environments.

Systems commands also require access to a realistic environment to conduct testing. The systems commands frequently conduct tests on Fleet range complexes and use Fleet assets to support the testing. The MIRC, which is primarily used by the systems commands, must provide the flexibility to

meet diverse testing requirements, given the wide range of various advanced platforms and systems and capabilities that the fleets and systems commands must demonstrate before certification for deployment for the Fleet. This is important because testing in conditions that reflect (or are similar to) those in which the technology could be employed enhances combat readiness.

The MIRC is characterized by a unique combination of attributes that make it a strategically important range complex, including:


- Location within and adjacent to a U.S. territory
- Ranges and operating areas on the islands of Guam, Rota, Saipan, Tinian, and FDM
- Expansive airspace, surface sea space, and underwater sea space
- Authorized use of multiple types of explosive and non-explosive ordnance on FDM
- Support for all Navy warfare areas and numerous other service roles, missions, and tactical tasks
- Support for service units based at military installations on Guam
- Training support for deployed forces
- Ability to conduct joint and combined force exercises, including those in which foreign partners and allies participate
- Rehearsal area for Western Pacific contingencies

### 1.6 The Environmental Planning Process

NEPA and Executive Order 12114 requires federal agencies to examine the environmental impacts of their proposed actions within the United States and its territories. An EIS is a detailed public document that assesses the potential effects that a major federal action might have on the human environment. The Navy undertakes environmental planning for major Navy actions in accordance with applicable laws, regulations, and Executive Orders.

Pursuant to 40 CFR section 1502.9(c), a supplemental EIS is prepared when the agency makes substantial changes in the proposed action that are relevant to environmental concerns (40 CFR section 1502.9(c)(1)(i)); or there are significant new circumstances or information relevant to environmental concerns and bearing on the proposed action or its impacts (40 CFR section 1502.9(c)(1)(ii)). An agency may also supplement a final EIS when the agency determines that the purpose of NEPA will be furthered by doing so (40 CFR section 1502(c)(2)).

Pursuant to Council on Environmental Quality Regulations, the Navy has prepared this supplement to the 2015 MITT Final EIS/OEIS to consider future activities conducted at sea and on FDM, and updated training and testing requirements; incorporate new information from an updated acoustic effects model and updated marine mammal density data; and incorporate evolving and emergent best available science. It will also support any reissuance of federal regulatory permits and authorizations under the MMPA and the ESA using the best available science and analytical methods to assess potential environmental impacts.



**Figure 1.5-1: National Environmental Policy Act Process Conducted for this SEIS/OEIS**

### **1.6.1 National Environmental Policy Act Requirements**

When developing a supplement to an existing EIS/OEIS, the first step in the NEPA process (Figure 1.5-1) is to prepare a Notice of Intent. The Notice of Intent is published in the Federal Register and in local newspapers, and provides an overview of the proposed action and the scope of this SEIS/OEIS (see Appendix B, Federal Register Notices). The Notice of Intent is also the first step in engaging the public, initiating the scoping process.

Scoping is an early and open process for developing the “scope” of issues to be addressed in an EIS and for identifying significant issues related to a proposed action. In accordance with the Council on Environmental Quality regulations for implementing the requirements of NEPA, scoping is not required for a supplement to a draft or final EIS; however, in an effort to maximize public participation and ensure the public’s input was considered, the Navy chose to conduct a scoping period for this SEIS/OEIS.

After the scoping process, a Draft SEIS/OEIS is prepared to assess potential impacts of the proposed action and alternatives on the environment. When completed, a Notice of Availability is published in the Federal Register and notices are placed in local or regional newspapers announcing the availability of the Draft SEIS/OEIS. The Draft SEIS/OEIS is circulated for public review and comment. Public meetings may also be scheduled to further inform the public and solicit their comments.

The Final SEIS/OEIS addresses all public comments received on the Draft SEIS/OEIS. Responses to public comments may include factual corrections, supplements or modifications to analysis, and inclusion of new information. Additionally, responses may explain why the comments do not warrant further agency response.

Finally, the decision-maker will issue a Record of Decision no earlier than 30 days after the Final SEIS/OEIS is made available to the public.

For a description of how the Navy complies with each of these requirements during the development of this SEIS/OEIS, please see Chapter 8 (Public Involvement and Distribution).

### **1.6.2 Executive Order 12114**

Executive Order 12114 of 1979, *Environmental Impacts Abroad of Major Federal Actions*, furthers the purpose of NEPA by directing federal agencies to provide for informed environmental decision-making for major federal actions outside the United States and its territories. Presidential Proclamation 5928, issued December 27, 1988, extended the exercise of U.S. sovereignty and jurisdiction under international law to 12 nautical miles (NM) from the shoreline; however, the proclamation expressly provides that it does not extend or otherwise alter existing federal law or any associated jurisdiction, rights, legal interests, or obligations. Thus, as a matter of policy, the Navy analyzes environmental effects and actions within 12 NM under NEPA (an EIS) and those effects occurring beyond 12 NM under the provisions of Executive Order 12114 (an OEIS).

### **1.6.3 Other Environmental Requirements Considered**

The Navy must comply with all applicable federal environmental laws, regulations, and executive orders as discussed in the 2015 MITT Final EIS/OEIS. Further information can be found in Chapter 6 (Additional Regulatory Considerations). Since the publication of the 2015 MITT Final EIS/OEIS, Executive Order 13840, *Ocean Policy to Advance the Economic, Security, and Environmental Interests of the United States*

revoked and replaced Executive Order 13547, *Stewardship of the Ocean, Our Coasts, and the Great Lakes*.

**1.6.3.1.1 Executive Order 13840, *Ocean Policy to Advance the Economic, Security, and Environmental Interests of the United States***

On June 19, 2018, President Trump signed Executive Order 13840. The Executive Order is intended to advance the economic, security, and environmental interests of the United States through improved public access to marine data and information; efficient federal agency coordination on ocean-related matters; and engagement with marine industries, the science and technology community, and other ocean stakeholders, including Regional Ocean Partnerships. The Executive Order continues to require federal agencies to coordinate activities regarding ocean-related matters for effective management of the ocean as well as promote lawful use of the ocean by agencies, including the Armed Forces. The Navy continues to engage with regional and state ocean planning entities. This Executive Order revokes and replaces Executive Order 13547, *Stewardship of the Ocean, Our Coasts, and the Great Lakes*.

**1.7 Scope and Content**

In this SEIS/OEIS, the Navy reanalyzed at-sea and FDM military readiness activities that could potentially impact the human environment and natural resources. Since the completion of the 2015 MITT Final EIS/OEIS, new information has become available and is incorporated in this analysis, in addition to proposed changes in training and testing requirements. The range of alternatives in this SEIS/OEIS includes the No Action Alternative and two action alternatives. In this SEIS/OEIS, the Navy analyzed direct, indirect, and cumulative impacts that may result from the Proposed Action. The Navy is the lead agency for the Proposed Action and is responsible for the scope and content of this SEIS/OEIS. The document is being prepared in coordination with the U.S. Air Force, U.S. Coast Guard, as their at-sea and FDM training activities are included in the Proposed Action.

The National Oceanic Atmospheric Administration's NMFS is serving as a cooperating agency because the scope of the Proposed Action and alternatives involves activities that have the potential to impact protected resources under their jurisdiction by law and special expertise, including marine mammals, threatened and endangered species, and Essential Fish Habitat. The National Oceanic Atmospheric Administration's authorities and special expertise is based on their statutory responsibilities under the MMPA of 1972, as amended 16 U.S.C. 1361 et seq.), the Endangered Species Act of 1973 (16 U.S.C. 1531 et seq.), and the Magnuson-Stevens Fishery Conservation and Management Act. In addition, NMFS, in accordance with 40 CFR 1506.3 and 1505.2, may adopt this SEIS/OEIS and issue a separate Record of Decision associated with its decision to grant or deny the Navy's request for an incidental take authorization pursuant to Section 101(a)(5)(A) of the MMPA.

**1.8 Organization of this Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement**

This SEIS/OEIS is organized as follows:

- Chapter 1 (Purpose and Need) describes the purpose of and need for the Proposed Action.
- Chapter 2 (Description of Proposed Action and Alternatives) describes the Proposed Action and proposed changes to the 2015 MITT Final EIS/OEIS implemented actions projected to take place starting in 2020, and alternatives to be carried forward for analysis.
- Chapter 3 (Affected Environment and Environmental Consequences) describes the existing conditions of the affected environment and potential environmental consequences on those

resources requiring additional discussion or analysis beyond what was analyzed in the 2015 MITT Final EIS/OEIS.

- Chapter 4 (Cumulative Impacts) describes the analysis of cumulative impacts, which are the impacts of the Proposed Action when added to past, present, and reasonably foreseeable future actions.
- Chapter 5 (Mitigation) describes the measures the Navy evaluated that could mitigate impacts to the environment.
- Chapter 6 (Additional Regulatory Considerations) describes considerations required by NEPA and describes how the Navy complies with other federal, state, and local plans, policies, and regulations.
- Chapter 7 (List of Preparers) includes a list of preparers of this SEIS/OEIS.
- Chapter 8 (Public Involvement and Distribution) describes the public participation process.
- References are provided at the end of each section.
- Appendices provide technical information that support the SEIS/OEIS analyses and its conclusions.

## **REFERENCES**

- U.S. Department of the Navy. (2000). Compliance with Environmental Requirements in the Conduct of Naval Exercises or Training at Sea. Washington, DC: The Under Secretary of the Navy.
- U.S. Department of the Navy. (2015). Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Fish and Wildlife Service. (2015). Biological Opinion for the Mariana Islands Training and Testing Program. Honolulu, HI: U.S. Fish and Wildlife Service Pacific Islands Fish and Wildlife Office.
- U.S. Fish and Wildlife Service. (2016). Informal Consultation on Mariana Islands Training and Testing Program Affects to Eighteen Newly-Listed Species, Guam and Tinian. Honolulu, HI: Fish and Wildlife Service, Pacific Islands Fish and Wildlife Office.

This page intentionally left blank.

---

---

## 2 Description of Proposed Action and Alternatives



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

**2 DESCRIPTION OF PROPOSED ACTION AND ALTERNATIVES.....2-1**

2.1 Description of the Mariana Islands Training and Testing Study Area..... 2-2

2.1.1 Description of the Mariana Islands Range Complex..... 2-3

2.1.1.1 Special Use Airspace and Air Traffic Controlled Assigned Airspace ..... 2-3

2.1.1.2 Sea and Undersea Space..... 2-3

2.1.2 Description of the Ocean Operating Areas Outside the Bounds of the Mariana Islands Range Complex ..... 2-3

2.1.3 Description of Pierside Locations and Apra Harbor ..... 2-7

2.2 Primary Mission Areas ..... 2-7

2.2.1 Air Warfare ..... 2-7

2.2.2 Amphibious Warfare..... 2-9

2.2.3 Anti-Submarine Warfare..... 2-9

2.2.4 Electronic Warfare ..... 2-9

2.2.5 Expeditionary Warfare..... 2-10

2.2.6 Mine Warfare..... 2-10

2.2.7 Strike Warfare..... 2-11

2.2.8 Surface Warfare..... 2-11

2.3 Proposed Activities ..... 2-11

2.3.1 Changes to Proposed Activities ..... 2-11

2.3.1.1 New Technologies and Capabilities ..... 2-12

2.3.1.2 Renamed and Reorganized Testing Activities..... 2-12

2.3.2 Proposed Training and Testing Activities..... 2-12

2.3.3 Standard Operating Procedures ..... 2-13

2.3.3.1 High-Energy Laser Safety ..... 2-13

2.3.3.2 Sea Space and Airspace Deconfliction ..... 2-14

2.3.3.3 Target Deployment and Retrieval Safety..... 2-14

2.3.3.4 Pierside Testing Safety..... 2-14

2.3.3.5 Underwater Detonation Safety ..... 2-15

2.3.3.6 Sonic Booms..... 2-15

2.3.3.7 Unmanned Surface Vehicle Safety ..... 2-15

|         |   |      |
|---------|---|------|
| 2.3.3.8 | Sinking Exercise Safety.....  | 2-15 |
| 2.3.4   | Mitigation Measures.....  | 2-15 |
| 2.4     | Action Alternatives Development.....  | 2-17 |
| 2.4.1   | Alternatives Eliminated from Further Consideration .....                          | 2-17 |
| 2.4.1.1 | Alternative Training and Testing Locations .....                                  | 2-17 |
| 2.4.1.2 | Reduced Training and Testing .....  | 2-17 |
| 2.4.1.3 | Alternatives Including Geographic Mitigation Measures within the Study Area ..... | 2-18 |
| 2.4.1.4 | Simulated Training and Testing Only .....   | 2-18 |
| 2.4.1.5 | Training and Testing Without the Use of Active Sonar .....                        | 2-18 |
| 2.4.2   | Alternatives Carried Forward .....  | 2-18 |
| 2.4.2.1 | No Action Alternative .....   | 2-19 |
| 2.4.2.2 | Alternative 1 .....   | 2-20 |
| 2.4.2.3 | Alternative 2 .....   | 2-21 |
| 2.4.3   | Comparison of Proposed Sonar and Explosive Use in the Action Alternatives .....   | 2-22 |
| 2.4.3.1 | Sonar Use.....  | 2-22 |
| 2.4.3.2 | Explosives Use.....   | 2-23 |
| 2.5     | Comparison of Alternatives .....  | 2-25 |

## List of Figures

|  |      |
|--|------|
| Figure 2.1-1: Mariana Islands Training and Testing Study Area .....  | 2-2  |
| Figure 2.1-2: Mariana Islands Range Complex Airspace .....   | 2-4  |
| Figure 2.1-3: Warning Area 517 .....   | 2-5  |
| Figure 2.1-4: Farallon de Medinilla Restricted Area 7201, 7201A, and Danger Zone .....   | 2-6  |
| Figure 2.2-1: Apra Harbor Naval Complex (Main Base) and Main Base/Polaris Point .....  | 2-8  |
| Figure 2.4-1: Proposed Annual Total Hull-Mounted Mid-Frequency Sonar Hour Use Compared to the Number Authorized in the 2015–2020 Marine Mammal Protection Act Permit ..... | 2-23 |
| Figure 2.4-2: Proposed Annual Explosives Use (Bins E1–E5) Compared to the 2015–2020 Marine Mammal Protection Act Permit .....  | 2-24 |
| Figure 2.4-3: Proposed Annual Explosives Use (Bins E6–E12) Compared to the 2015–2020 Marine Mammal Protection Act Permit .....   | 2-24 |

---

## List of Tables

Table 2.3-1: Overview of Mitigation Categories .....2-16  
Table 2.5-1: Current and Proposed Training Activities .....2-26  
Table 2.5-2: Current and Proposed Testing Activities.....2-40

This page intentionally left blank.

## 2 Description of Proposed Action and Alternatives

The United States (U.S.) Department of the Navy (Navy) proposes to conduct military readiness activities which includes training activities (referred to as “training”), and research, development, testing, and evaluation (referred to as “testing”) activities in the Mariana Islands Training and Testing (MITT) Study Area, primarily within the existing Mariana Islands Range Complex (MIRC). This Supplemental Environmental Impact Statement/Overseas Environmental Impact Statement (SEIS/OEIS) is being prepared to assess the potential environmental impacts associated with proposed training and testing activities to be conducted at sea and on Farallon de Medinilla (FDM). These proposed activities are generally consistent with those at-sea and FDM activities analyzed in the 2015 MITT Final EIS/OEIS and are representative of activities the military has been conducting in the Study Area since the 1940s. These training and testing activities include the use of active sonar and explosives at sea off the coasts of Guam and the Commonwealth of the Northern Mariana Islands (CNMI), throughout the in-water areas around the MIRC, the transit corridor between the MIRC and the Hawaii Range Complex (HRC), and at select Navy pierside and harbor locations.

In this chapter, the military builds upon the purpose and need to train and test (as described in Chapter 1) by describing the Study Area and identifying the primary mission areas for which these training and testing activities are conducted. Each warfare community (e.g., aviation, surface, submarine, and expeditionary) conducts training and testing activities that contribute to the success of these primary mission areas. Each primary mission area requires unique skills, sensors, weapons, and technologies to accomplish the overall mission. For example, under the anti-submarine warfare primary mission area, surface, submarine, and aviation warfare communities each utilize different skills, sensors, and weapons to locate, track, and eliminate submarine threats. The testing community contributes to the success of anti-submarine warfare by anticipating and identifying technologies and systems that respond to the needs of the warfare communities. See Section 2.2 (Primary Mission Areas) and Section 2.3 (Descriptions of Sonar, Ordnance/Munitions, Targets and Other Systems Employed in MITT Events) for additional information.

This chapter describes the activities that comprise the Proposed Action for this SEIS/OEIS necessary to meet training and testing requirements beyond 2020 and into the reasonably foreseeable future. These at-sea and FDM activities are then analyzed for their potential effects on the environment in the resource-specific chapters of this SEIS/OEIS. For further details regarding specific training and testing activities, please see Appendix A (Training and Testing Activities Descriptions). The Navy intends to request from the National Marine Fisheries Service (NMFS) an incidental take authorization under the Marine Mammal Protection Act (MMPA), and an incidental take statement under the Endangered Species Act (ESA) for marine species. Relative to compliance with the National Environmental Policy Act, NMFS’ Proposed Action will be a direct outcome of responding to the Navy’s request for an incidental take authorization pursuant to the MMPA.

The 2015 MITT Final EIS/OEIS also analyzed training and testing activities conducted at existing MIRC land-based training areas located on Guam, Saipan, Tinian, and Rota. The Navy consulted with the U.S. Fish and Wildlife Service regarding effects of the land-based training activities on terrestrial species listed under the ESA and received a Biological Opinion (U.S. Fish and Wildlife Service, 2015) and concurrence letter (U.S. Fish and Wildlife Service, 2016). As the Navy is not proposing any changes to those land-based activities on Guam, Saipan, Tinian, and Rota, the Navy will continue to rely on the 2015 MITT Final EIS/OEIS because there is no new information that would affect the EIS analysis. In addition,

in accordance with 50 Code of Federal Regulations (CFR) section 402.16, the 2015 and 2016 consultations remain valid as none of the factors necessary to trigger reinitiating consultation have been met.

## 2.1 Description of the Mariana Islands Training and Testing Study Area

The Study Area (Figure 2.1-1) for this SEIS/OEIS is the same used for the analysis in the 2015 MITT Final EIS/OEIS (Section 2.1, Description of the Mariana Islands Training and Testing Study Area), and is composed of three components: (1) the MIRC, (2) additional areas on the high seas outside of the MIRC, and (3) a transit corridor between the MIRC and the HRC. Collectively, for the purposes of this SEIS/OEIS, these areas continue to be referred to as the MITT Study Area (Figure 2.1-1). The transit corridor is outside the geographic boundaries of the MIRC and is a direct route across the high seas for Navy ships transiting between the MIRC and the HRC.

Section 2.1.1 (Description of the Mariana Islands Range Complex) and the 2015 MITT Final EIS/OEIS (Section 2.1, Description of the Mariana Islands Training and Testing Study Area) provide complete descriptions of range components that comprise the MIRC. For more information on the areas outside the boundaries of the MIRC but within the Study Area, see Section 2.1.2 (Description of the Ocean Operating Areas Outside the Bounds of the Mariana Islands Range Complex) and Section 2.1.3 (Description of Pierside Locations and Apra Harbor) below and in the 2015 MITT Final EIS/OEIS.




Figure 2.1-1: Mariana Islands Training and Testing Study Area

### **2.1.1 Description of the Mariana Islands Range Complex**

The MIRC includes land training areas, ocean surface and subsurface areas, and special use airspace. These areas extend from the waters south of Guam to north of Pagan (Commonwealth of the Northern Mariana Islands), and from the Pacific Ocean east of the Mariana Islands to the Philippine Sea to the west, encompassing 501,873 square nautical miles (NM<sup>2</sup>) of open ocean (Figure 2.1-1). The Department of Defense leases FDM for use as a live and inert gunnery, missile, and bombing range.

#### **2.1.1.1 Special Use Airspace and Air Traffic Controlled Assigned Airspace**

The MIRC includes approximately 40,000 NM<sup>2</sup> of special use airspace. Special use airspace is airspace of defined dimensions where activities must be confined because of their nature or where limitations may be imposed upon aircraft operations that are not part of those activities (Federal Aviation Administration, 2013). As described in the 2015 MITT Final EIS/OEIS, Section 2.1 (Description of the Mariana Islands Training and Testing Study Area), special use airspace includes restricted areas, military operations areas, and warning areas. As depicted in Figure 2.1-2 and Figure 2.1-3, most of this airspace is almost entirely over the ocean and includes warning areas, and restricted areas:

Warning Areas (W): W-517 and W-12 include approximately 11,800 NM<sup>2</sup> of special use airspace (Figure 2.1-2 and Figure 2.1-3); W-11 (A/B) is approximately 10,500 NM<sup>2</sup> of special use airspace, and W-13 (A/B/C) is approximately 18,000 NM<sup>2</sup> of special use airspace.

Restricted Area Airspace (R): Over or near land areas within the MIRC includes approximately 2,463 NM<sup>2</sup> of special use airspace and includes restricted areas R-7201 and R-7201A, which extends in a 12 nautical mile radius around FDM (Figure 2.1-2 and Figure 2.1-4).

#### **2.1.1.2 Sea and Undersea Space**

The MIRC includes the sea and undersea space from the ocean surface to the ocean floor. The MIRC also consists of designated sea and undersea space training areas, which include designated drop zones, underwater demolition and floating mine exclusion zones, danger zones associated with live fire ranges, and training areas associated with military controlled beaches, harbors, and littoral areas.

W-517, W-12, W-11 and, W-13 (Figure 2.1-2) are designated as special use airspace where the sea space underneath may be restricted from public access during hazardous training events. Portions of the Marianas Trench Marine National Monument, established in January 2009 by Presidential Proclamation under the authority of the Antiquities Act (16 U.S. Code sections 431–433), lie within the MIRC and under all MIRC Warning Areas. However, the prohibitions required by the Proclamation do not apply to activities and exercises of the Armed Forces (including those carried out by the U.S. Coast Guard).

### **2.1.2 Description of the Ocean Operating Areas Outside the Bounds of the Mariana Islands Range Complex**

In addition to the MIRC, the Study Area includes the area to the north of the MIRC that is within the U.S. Exclusive Economic Zone of the CNMI and areas to the west of the MIRC, as depicted in Figure 2.1-1.

The transit corridor between MIRC and HRC, although not part of any defined range complex, is important to the Navy in that it provides available air, sea, and undersea space where vessels and aircraft conduct training and testing while in transit. The transit corridor is the shortest distance between the MIRC and the HRC.




Figure 2.1-2: Mariana Islands Range Complex Airspace





Figure 2.1-3: Warning Area 517



**Figure 2.1-4: Farallon de Medinilla Restricted Area 7201, 7201A, and Danger Zone**

### 2.1.3 Description of Pierside Locations and Apra Harbor

The Study Area includes pierside locations in Apra Harbor. For purposes of this SEIS/OEIS, pierside locations include channels and routes to and from the Navy port in the Apra Harbor Naval Complex, and associated wharves and facilities within the Navy port (Figure 2.2-1).

## 2.2 Primary Mission Areas

The Navy categorizes its at-sea activities into functional warfare areas called primary mission areas. Training and testing activities generally fall into the following eight primary mission areas:

- air warfare
- amphibious warfare
- anti-submarine warfare
- electronic warfare
- expeditionary warfare
- mine warfare
- strike warfare
- surface warfare

Most activities addressed in this SEIS/OEIS are categorized under one of these primary mission areas; activities that do not fall within one of these areas are listed as “other activities” (e.g., precision anchoring, search and rescue at sea). Each warfare community (e.g., surface, subsurface, aviation, and expeditionary warfare) may train in some or all of these primary mission areas. The research and acquisition community also categorizes most, but not all, of its testing activities under these primary mission areas. A description of the sonar, munitions, targets, systems, and other material used during training and testing activities within these primary mission areas is provided in Appendix A (Training and Testing Activities Descriptions).

### 2.2.1 Air Warfare

The mission of air warfare (referred to as anti-air warfare in the 2015 MITT Final EIS/OEIS) is to destroy or reduce enemy air and missile threats (including unmanned airborne threats) and serves two purposes: to protect U.S. forces from attacks from the air and to gain air superiority. Air warfare provides U.S. forces with adequate attack warnings, while denying hostile forces the ability to gather intelligence about U.S. forces.

Aircraft conduct air warfare through radar search, detection, identification, and engagement of airborne threats. Surface ships conduct air warfare through an array of modern anti-aircraft weapon systems such as aircraft-detecting radar, naval guns linked to radar-directed fire-control systems, surface-to-air missile systems, and radar-controlled guns for close-in point defense.

Testing of air warfare systems is required to ensure the equipment is fully functional under the conditions in which it will be used. Tests may be conducted on radar and other early-warning detection and tracking systems, new guns or gun rounds, and missiles. Testing of these systems may be conducted on new ships and aircraft, and on existing ships and aircraft following maintenance, repair, or modification. For some systems, tests are conducted periodically to assess operability. Additionally, tests may be conducted in support of scientific research to assess new and emerging technologies.




Figure 2.2-1: Apra Harbor Naval Complex (Main Base) and Main Base/Polaris Point

### **2.2.2 Amphibious Warfare**

The mission of amphibious warfare is to project military power from the sea to the shore (i.e., attack a threat on land by a military force embarked on ships) through the use of naval firepower and expeditionary landing forces. Amphibious warfare operations include small unit reconnaissance or raid missions to large-scale amphibious exercises involving multiple ships and aircraft combined into a strike group.

Amphibious warfare training ranges from individual, crew, and small unit events to large task force exercises. Individual and crew training include amphibious vehicles and naval gunfire support training. Such training includes shore assaults, boat raids, airfield or port seizures, and reconnaissance. Large-scale amphibious exercises involve ship-to-shore maneuver, naval fire support, such as shore bombardment, and air strike and attacks on targets that are in close proximity to friendly forces.

### **2.2.3 Anti-Submarine Warfare**

The mission of anti-submarine warfare is to locate, neutralize, and defeat hostile submarine forces that threaten Navy surface forces. Anti-submarine warfare is based on the principle that surveillance and attack aircraft, ships, and submarines all search for hostile submarines. These forces operate together or independently to gain early warning and detection, and to localize, track, target, and attack submarine threats.

Anti-submarine warfare training addresses basic skills such as detection and classification of submarines, as well as evaluating sounds to distinguish between enemy submarines and friendly submarines, ships, and marine life. More advanced training integrates the full spectrum of anti-submarine warfare from detecting and tracking a submarine to attacking a target using either exercise torpedoes (i.e., torpedoes that do not contain a warhead) or simulated weapons. These integrated anti-submarine warfare training exercises are conducted in coordinated, at-sea training events involving submarines, ships, and aircraft.

Testing of anti-submarine warfare systems is conducted to develop new technologies and assess weapon performance and operability with new systems and platforms, such as unmanned systems. Testing uses ships, submarines, and aircraft to demonstrate capabilities of torpedoes, missiles, countermeasure systems, and underwater surveillance and communications systems. Tests may be conducted as part of a large-scale Fleet training event involving submarines, ships, fixed-wing aircraft, and helicopters. These integrated training events offer opportunities to conduct research and acquisition activities and to train aircrew in the use of new or newly enhanced systems during a large-scale, complex exercise.

### **2.2.4 Electronic Warfare**

The mission of electronic warfare is to degrade the enemy's ability to use electronic systems, such as communication systems and radar, and to confuse or deny them the ability to defend their forces and assets. Electronic warfare is also used to detect enemy threats and counter their attempts to degrade the electronic capabilities of the Navy.

Typical electronic warfare activities include threat avoidance training, signals analysis for intelligence purposes, and use of airborne and surface electronic jamming devices (that block or interfere with other devices) to defeat tracking, navigation, and communications systems.

Testing of electronic warfare systems is conducted to improve the capabilities of systems and ensure compatibility with new systems. Testing involves the use of aircraft, surface ships, and submarine crews

to evaluate the effectiveness of electronic systems. Similar to training activities, typical electronic warfare testing activities include the use of airborne and surface electronic jamming devices (including testing chaff and flares; see Appendix A (Training and Testing Activities Descriptions) for a description of these devices) to defeat tracking and communications systems. Chaff tests evaluate newly developed or enhanced chaff, chaff dispensing equipment, or modified aircraft systems' use against chaff deployment. Flare tests evaluate deployment performance and crew competency with newly developed or enhanced flares, flare dispensing equipment, or modified aircraft systems' use against flare deployment.

### **2.2.5 Expeditionary Warfare**

The mission of expeditionary warfare is to provide security and surveillance in the littoral (at the shoreline), riparian (along a river), or coastal environments. Expeditionary warfare is wide ranging and includes defense of harbors, operation of remotely operated vehicles, defense against swimmers, and boarding/seizure operations.

### **2.2.6 Mine Warfare**

The mission of mine warfare is to detect, classify, and avoid or neutralize (disable) mines to protect Navy ships and submarines and to maintain free access to ports and shipping lanes. Mine warfare also includes offensive mine laying to gain control of or deny the enemy access to sea space. Naval mines can be laid by ships, submarines, or aircraft.

Mine warfare neutralization training includes exercises in which ships, aircraft, submarines, underwater vehicles, unmanned vehicles, or marine mammal detection systems search for mine shapes. Personnel train to destroy or disable mines by attaching underwater explosives to or near the mine or using remotely operated vehicles to destroy the mine.

Testing and development of mine warfare systems is conducted to improve sonar, laser, and magnetic detectors intended to hunt, locate, and record the positions of mines for avoidance or subsequent neutralization. Mine warfare testing and development falls into two primary categories: mine detection and classification, and mine countermeasure and neutralization. Mine detection and classification testing involves the use of air, surface, and subsurface vessels and uses sonar, including towed and side-scan sonar, and unmanned vehicles to locate and identify objects underwater. Mine detection and classification systems are sometimes used in conjunction with a mine neutralization system. Mine countermeasure and neutralization testing includes the use of air, surface, and subsurface units to evaluate the effectiveness of tracking devices, countermeasure and neutralization systems, and general purpose bombs to neutralize mine threats. Most neutralization tests use mine shapes, or non-explosive practice mines, to evaluate a new or enhanced capability. For example, during a mine neutralization test, a previously located mine is destroyed or rendered nonfunctional using a helicopter or manned/unmanned surface vehicle-based system that may involve the deployment of a towed neutralization system.

A small percentage of mine warfare tests require the use of high-explosive mines to evaluate and confirm the ability of the system to neutralize a high-explosive mine under operational conditions. The majority of mine warfare systems are deployed by ships, helicopters, and unmanned vehicles. Tests may also be conducted in support of scientific research to support these new technologies.

### **2.2.7 Strike Warfare**

The mission of strike warfare is to conduct offensive attacks on land-based targets, such as refineries, power plants, bridges, major roadways, and ground forces to reduce the enemy's ability to wage war. Strike warfare employs weapons by manned and unmanned air, surface, submarine, and Navy special warfare assets in support of extending dominance over enemy territory (power projection).

Strike warfare includes training fixed-wing attack aircraft pilots and aircrews in the delivery of precision-guided munitions, non-guided munitions, rockets, and other ordnance against land-based targets. Not all strike mission training activities involve dropping ordnance and instead the activity is simulated with video footage obtained by onboard sensors.

### **2.2.8 Surface Warfare**

The mission of surface warfare (referred to as anti-surface warfare in the 2015 MITT Final EIS/OEIS) is to obtain control of sea space from which naval forces may operate, and entails offensive action against other surface, subsurface, and air targets while also defending against enemy forces. In surface warfare, aircraft use guns, air-launched cruise missiles, or other precision-guided munitions; ships employ torpedoes, naval guns, and surface-to-surface missiles; and submarines attack surface ships using torpedoes or submarine-launched, anti-ship cruise missiles.

Surface warfare training includes surface-to-surface gunnery and missile exercises, air-to-surface gunnery and missile exercises, and submarine missile or torpedo launch activities, and other munitions against surface targets.

Testing of weapons used in surface warfare is conducted to develop new technologies and to assess weapon performance and operability with new systems and platforms, such as unmanned systems. Tests include various air-to-surface guns and missiles, surface-to-surface guns and missiles, and bombing tests. Testing activities may be integrated into training activities to test aircraft or aircraft systems in the delivery of munitions on a surface target. In most cases the tested systems are used in the same manner in which they are used for Fleet training activities.

## **2.3 Proposed Activities**

The Navy has been conducting training and testing activities in the Study Area for decades. The tempo and types of training and testing activities have fluctuated because of the introduction of new technologies, the evolving nature of international events, advances in warfighting doctrine and procedures, and changes in force structure (organization of ships, submarines, aircraft, weapons, and Sailors). Such developments influence the frequency, duration, intensity, and location of required training and testing activities. The activities analyzed in this SEIS/OEIS are largely a continuation of activities that have been ongoing and were analyzed previously in the 2015 MITT Final EIS/OEIS. This SEIS/OEIS includes the analysis of those at sea and FDM activities necessary to meet readiness requirements beyond 2020 and into the reasonably foreseeable future, includes any changes to those activities previously analyzed, and reflects the most up-to-date compilation of training and testing activities deemed necessary to accomplish military readiness requirements.

### **2.3.1 Changes to Proposed Activities**

The majority of proposed modifications to the activities included in the Proposed Action are changes to tempo of activity, and renaming or combining related types of activities for greater clarity in this document and for consistency across all Navy at-sea planning documents. A few activities assessed in

2015 have been discontinued from analysis, and a few new activities have been added to the proposed activities to enable the Navy to adopt new technology and new capabilities. The new and renamed/combined training and testing activities are listed in Table 2.5-1 and Table 2.5-2 and discussed in greater detail below.

### **2.3.1.1 New Technologies and Capabilities**

As described above, new technologies and capabilities are introduced to be evaluated in testing. Some systems have been used and tested by the Navy in other locations, but not the MITT Study Area. Those systems that are new to the Study Area will be analyzed for environmental impacts in this SEIS/OEIS.

The Navy is proposing the testing of new systems and technologies for Naval Sea Systems Command. Radar and Other Systems Testing may include the use of military or commercial radar, communication systems or simulators, or high-energy lasers. Testing may occur aboard a ship against drones, small boats, rockets, missiles, or other targets. Simulant Testing involves the testing of simulated chemical-biological agents and simulants that are deployed against surface ships. Naval Air Systems Command and the Office of Naval Research are not proposing any new testing capabilities in this SEIS/OEIS. Information on all testing activities is provided at the end of this Chapter in Table 2.5-1.

### **2.3.1.2 Renamed and Reorganized Testing Activities**

Some Naval Sea Systems Command testing activities have been renamed. Following is a list of testing activities that have been renamed since the 2015 MITT Final EIS/OEIS:

- Undersea Warfare Testing (previously named Torpedo Testing)
- Mine Countermeasure and Neutralization Testing (previously named Mine Countermeasure Mission Package Testing)
- Anti-Submarine Warfare Mission Package Testing (previously named Anti-Submarine Warfare)

In addition, some Naval Sea Systems Command testing activities have been reorganized under a different primary mission area. Following is a list of testing activities that have been reorganized since the 2015 MITT Final EIS/OEIS:

- Kinetic Energy Weapons Testing (now analyzed under Surface Warfare)
- At-Sea Sonar Testing (now analyzed under Anti-Submarine Warfare)
- Torpedo (Explosive) Testing (now analyzed under Anti-Submarine Warfare)
- Torpedo (Non-explosive) Testing (now analyzed under Anti-Submarine Warfare)
- Undersea Warfare Testing (now analyzed under Vessel Evaluation)
- Anti-Submarine Warfare Mission Package Testing (now analyzed under Anti-Submarine Warfare)
- Mine Countermeasure and Neutralization Testing (now analyzed under Mine Warfare)

### **2.3.2 Proposed Training and Testing Activities**

Table 2.5-1 and Table 2.5-2 at the end of this chapter provides additional information on all training and testing activities, respectively, such as location, number of events per year, and ordnance used, if any. More information about each training and testing activity can be found in Appendix A (Training and Testing Activities Descriptions) and Appendix F (Training and Testing Activities Matrices). Except for the new activities described in Table 2.5-2, the activities proposed by the Navy in this SEIS/OEIS were described in the 2015 MITT Final EIS/OEIS in Table 2.4-2 and Table 2.4-3.

As described in the 2015 MITT Final EIS/OEIS, the Navy's research and acquisition community engages in a broad spectrum of testing activities in support of the Fleet. These activities include, but are not limited

to, basic and applied scientific research and technology development; testing, evaluation, and maintenance of systems (missiles, radar, and sonar) and platforms (surface ships, submarines, and aircraft); and acquisition of systems and platforms. The individual commands within the research and acquisition community included in this SEIS/OEIS are Naval Air Systems Command, Naval Sea Systems Command, and Office of Naval Research.

### **2.3.3 Standard Operating Procedures**

For training and testing to be effective, units must be able to safely use their sensors and weapon systems as they are intended to be used in military missions and combat operations and to their optimum capabilities. Standard operating procedures applicable to training and testing have been developed through years of experience, and their primary purpose is to provide for safety (including public health and safety) and mission success. Because they are essential to safety and mission success, standard operating procedures are part of the Proposed Action and are considered in the Chapter 3 (Affected Environment and Environmental Consequences) environmental analysis for applicable resources.

In many cases, there are benefits to environmental and cultural resources (some of which have high socioeconomic value in the Study Area) resulting from standard operating procedures. Those standard operating procedures that are recognized as providing a benefit to the resources analyzed in this Draft SEIS/OEIS are included in Appendix A (Training and Testing Activities Descriptions), as applicable. The following standard operating procedure categories apply to the Proposed Action and are generally consistent with those included in the specified sections in Chapter 5 (Standard Operating Procedures, Mitigation, and Monitoring) of the 2015 MITT Final EIS/OEIS:

- Section 5.1.1 (Vessel Safety)
- Section 5.1.2 (Aircraft Safety)
- Section 5.1.3 (Laser Procedures)
- Section 5.1.4 (Weapons Firing Procedures), except Section 5.1.4.3 (Target Deployment Safety), which has been updated in Section 2.3.3.3 (Target Deployment and Retrieval Safety) of this Draft SEIS/OEIS
- Section 5.1.6 (Unmanned Aerial and Underwater Vehicle Procedures)
- Section 5.1.7 (Towed In-Water Device Procedures)
- Section 5.1.8 (Amphibious Assault and Amphibious Raid Procedures)
- Section 5.7.3 (Farallon de Medinilla Access Restrictions)

Standard operating procedures that apply to the Proposed Action and were not included in, or require a clarification from, the 2015 MITT Final EIS/OEIS, are discussed in the following sections.

#### **2.3.3.1 High-Energy Laser Safety**

The Navy operates laser systems approved for fielding by the Laser Safety Review Board or service equivalent. Only properly trained and authorized personnel operate high-energy lasers within designated areas. Designated areas where lasers are used are required to have a Laser Range Safety Certification Report that is updated every three years. Prior to commencing activities involving high-energy lasers, the operator performs a search of the intended impact location to ensure that the area is clear of unauthorized persons. These standard operating procedures benefit public health and safety by reducing the potential for interaction with high-energy lasers.

### **2.3.3.2 Sea Space and Airspace Deconfliction**

The Navy schedules training and testing activities to minimize conflicts with the use of sea space and airspace within ranges and throughout the Study Area to ensure the safety of military personnel, the public, commercial aircraft, commercial and recreational vessels, and military assets. The Navy deconflicts its own use of sea space and airspace to allow for the necessary separation of multiple military units to prevent interference with equipment sensors and to avoid interaction with established commercial air traffic routes and commercial shipping lanes. The Navy also minimizes conflicts within areas used for commercial and recreational fishing, subsistence use, and tourism. For example, during applicable seasons around the islands of Guam and the CNMI, the Navy works collaboratively with local communities to deconflict sea space used for fishing to the maximum extent practicable, such as avoiding known fishery infrastructures (e.g., fish aggregating devices) and high-use fishing areas. To help civilian mariners better plan fishing and boating activities that involve accessing the waters around FDM, the Navy notifies them through various means, such as U.S. Coast Guard-issued Notices to Mariners and social media of the time periods when FDM will not be in use for several consecutive days. Announcing in advance when FDM will be in use (and when it will not be in use for an extended period of time) facilitates use of waters around FDM by the public during time periods that will not conflict with training and testing activities. These standard operating procedures benefit public health and safety (including persons participating in activities that have subsistence benefits and socioeconomic value, such as recreational or commercial fishing) by reducing potential interactions with training and testing activities.

### **2.3.3.3 Target Deployment and Retrieval Safety**

The standard operating procedures for target deployment and retrieval safety are consistent with the procedures described in Section 5.1.4.3 (Target Deployment Safety) of the 2015 MITT Final EIS/OEIS, except for the description of which activities will implement them. Under the Proposed Action, the standard operating procedure for target deployment and retrieval safety applies to weapons firing activities that involve small boats deploying or retrieving targets. These activities are typically conducted in daylight hours in Beaufort sea state number 4 conditions or better to ensure safe operating conditions during target deployment and recovery. These standard operating procedures benefit public health and safety, and marine mammals and sea turtles by increasing the effectiveness of visual observations for mitigation, thereby reducing the potential for interactions with the weapons firing activities associated with the use of applicable deployed targets.

During activities that involve recoverable targets (e.g., aerial drones), the military recovers the target and any associated decelerators/parachutes to the maximum extent practicable consistent with personnel and equipment safety. Recovery of these items helps minimize the amount of materials that remain, which could potentially alert enemy forces to the presence of military assets during military missions and combat operations. This standard operating procedure benefits biological resources (e.g., marine mammals, sea turtles, fish, marine birds) by reducing the potential for physical disturbance and strike, entanglement, or ingestion of applicable targets and any associated decelerators/parachutes.

### **2.3.3.4 Pierside Testing Safety**

The *U.S. Navy Dive Manual* (U.S. Department of the Navy, 2011) prescribes safe distances for divers from active sonar sources and in-water explosions. Safety distances for the use of electromagnetic energy are specified in Department of Defense Instruction 6055.11 (U.S. Department of Defense, 2009) and Military Standard 464A (U.S. Department of Defense, 2002). These distances are used as the standard safety buffers for in-water energy to protect military divers. If an unauthorized person is

detected within the exercise area, the activity will be temporarily halted until the area is again cleared and secured. These standard operating procedures benefit public health and safety (including persons participating in activities that have socioeconomic value, such as commercial or recreational diving) by reducing the potential for interaction with pierside testing activities.

#### **2.3.3.5 Underwater Detonation Safety**

Underwater detonation training takes place in designated areas that are located away from popular recreational dive sites, primarily for human safety. Recreational dive sites often include shallow-water coral reefs, artificial reefs, and wrecks. If an unauthorized person (e.g., a recreational diver) or vessel is detected within the exercise area, the activity will be temporarily halted until the area is cleared and secured. Notices to Mariners are issued when the events are scheduled to alert the public to stay clear of the area. These standard operating procedures benefit public health and safety, environmental resources (e.g., shallow-water coral reefs, artificial reefs, and the biological resources that inhabit, shelter in, or feed among them), and cultural resources by reducing the potential for interaction with underwater detonation activities.

#### **2.3.3.6 Sonic Booms**

As a general policy, aircraft do not intentionally generate sonic booms below 30,000 feet of altitude unless over water and more than 30 miles from inhabited land areas or islands. The military may authorize deviations from this policy for tactical missions, phases of formal training syllabus flights, or research, test, and operational suitability test flights. The standard operating procedures for sonic booms benefit public health and safety by reducing the potential for exposure to sonic booms.

#### **2.3.3.7 Unmanned Surface Vehicle Safety**

For activities involving unmanned surface vehicles, the Navy evaluates the need to publish a Notice to Airmen or Notice to Mariners based on the scale, location, and timing of the activity. When necessary, Notices to Airmen and Notices to Mariners are issued to alert the public to stay clear of the area. These standard operating procedures benefit public health and safety by reducing the potential for interaction with unmanned surface vehicles.

#### **2.3.3.8 Sinking Exercise Safety**

The Navy is required to conduct sinking exercises greater than 50 nautical miles from land and in waters at least 6,000 feet deep (40 CFR section 229.2). The Navy selects sinking exercise areas to avoid established commercial air traffic routes, commercial vessel shipping lanes, and areas used for recreational activities, and to allow for the necessary separation of Navy units to ensure safety for Navy personnel, the public, commercial aircraft and vessels, and Navy assets. These standard operating procedures benefit public health and safety (including persons participating in activities that have socioeconomic value, such as recreational or commercial fishing) by reducing the potential for interaction with sinking exercises.

### **2.3.4 Mitigation Measures**

The military will implement mitigation measures to avoid or reduce potential impacts from the Proposed Action on environmental and cultural resources. Mitigation measures that the Navy will implement under the Proposed Action are organized into three categories: at-sea procedural mitigation measures, at-sea mitigation areas, and terrestrial mitigation measures. The Navy will implement procedural mitigation measures whenever and wherever applicable training or testing activities take place within the Study Area. Mitigation areas are geographic locations within the Study Area where the military will

implement additional mitigation during all or part of the year. Terrestrial mitigation measures will be implemented during activities conducted on FDM.

A list of the activity categories, stressors, and geographic locations that have mitigation measures is provided in Table 2.3-1. Chapter 5 (Mitigation) of this Draft SEIS/OEIS provides a full description of each mitigation measure that will be implemented under Alternative 1 and Alternative 2 of the Proposed Action. It also presents a discussion of how the Navy developed and assessed each measure and includes maps of the mitigation area locations. Mitigation developed for the Proposed Action is generally in line with the type and level of mitigation included in Chapter 5 (Standard Operating Procedures, Mitigation, and Monitoring) of the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015). The Navy has updated Chapter 5 (Mitigation) of this Draft SEIS/OEIS in its entirety based on its ongoing analysis of the best available science and practicality of implementing potential mitigation measures. A full analysis of the mitigation areas that the Navy has considered for marine mammals and sea turtles in the Study Area is provided in Appendix I (Geographic Mitigation Assessment). Relevant mitigation details are also provided throughout Appendix A (Training and Testing Activities Descriptions). The Navy and NMFS Records of Decision, MMPA Regulations and Letter of Authorization, and ESA Biological Opinion will document all mitigation measures that the military will implement under the Proposed Action.

**Table 2.3-1: Overview of Mitigation Categories**

| <i>Mitigation Category</i> | <i>Draft SEIS/OEIS Section</i>                                  | <i>Applicable Activity Category, Stressor, Mitigation Area, or Location</i>   |
|----------------------------|---|---|
| Procedural Mitigation      | Section 5.3.2 (Acoustic Stressors)                              | Active Sonar<br>Weapons Firing Noise  |
|                            | Section 5.3.3 (Explosive Stressors)                             | Explosive Sonobuoys<br>Explosive Torpedoes<br>Explosive Medium-Caliber and Large-Caliber Projectiles<br>Explosive Missiles and Rockets<br>Explosive Bombs<br>Sinking Exercises<br>Explosive Mine Countermeasure and Neutralization Activities<br>Explosive Mine Neutralization Activities Involving Navy Divers<br>Maritime Security Operations – Anti-Swimmer Grenades |
|                            | Section 5.3.4 (Physical Disturbance and Strike Stressors)       | Vessel Movement<br>Towed In-Water Devices<br>Small-, Medium-, and Large-Caliber Non-Explosive Practice Munitions<br>Non-Explosive Missiles and Rockets<br>Non-Explosive Bombs and Mine Shapes   |
| Mitigation Areas           | Section 5.4 (At-Sea Mitigation Areas to be Implemented)         | Areas with seafloor resources (shallow-water coral reefs, live hard bottom, artificial reefs, and shipwrecks)   |
|                            | Appendix I (Geographic Mitigation Assessment)                   | Marine mammal and/or sea turtle mitigation areas (Marpi Reef, Chalan Kanoa Reef, Agat Bay Nearshore)  |
| Terrestrial Mitigation     | Section 5.5 (Terrestrial Mitigation Measures to be Implemented) | Farallon de Medinilla   |

## **2.4 Action Alternatives Development**

The identification, consideration, and analysis of alternatives are critical components of the National Environmental Policy Act process and contribute to the goal of objective decision-making. The Council on Environmental Quality (CEQ) developed regulations to implement National Environmental Policy Act and these regulations require the decision maker to consider the environmental effects of the proposed action and a range of alternatives (including the No Action Alternative) to the proposed action (40 Code of Federal Regulations section 1502.14). CEQ guidance further provides that an EIS must rigorously and objectively explore all reasonable alternatives for implementing the proposed action and, for alternatives eliminated from detailed study, briefly discuss the reasons for their having been eliminated. To be reasonable, an alternative, except for the no action alternative, must meet the stated purpose of and need for the proposed action.

The action alternatives, and in particular the mitigation measures that are incorporated in the action alternatives, were developed to meet both the Navy's purpose and need to train and test, and NMFS' independent purpose and need to evaluate the potential impacts of the Navy's activities, determine whether incidental take resulting from the Navy's activities will have a negligible impact on affected marine mammal species and stocks, and to prescribe measures to effect the least practicable adverse impact on species or stocks and their habitat, as well as monitoring and reporting requirements.

The Navy developed the alternatives considered in this SEIS/OEIS after careful assessment by subject matter experts, including military commands that utilize the ranges, military range management professionals, and Navy environmental managers and scientists.

### **2.4.1 Alternatives Eliminated from Further Consideration**

This SEIS/OEIS serves as an update to the 2015 MITT Final EIS/OEIS; therefore, alternatives eliminated from consideration in the 2015 MITT Final EIS/OEIS were evaluated to determine if they should be reconsidered for this SEIS/OEIS. In response to the comments received during the public scoping period, the Navy also considered developing an alternative that included geographic mitigation. Alternatives eliminated from further consideration are described in the subsections below. The Navy determined that these alternatives did not meet the purpose of and need for the Proposed Action after a thorough consideration of each.

#### **2.4.1.1 Alternative Training and Testing Locations**

As described in Section 2.5.1.1 (Alternative Training and Testing Locations) in the 2015 MITT Final EIS/OEIS, the diverse and multi-dimensional environment provided within the Study Area allows the military to develop and maintain high levels of readiness and interoperability with foreign partners in the Western Pacific. There are no other proximate alternative locations that provide for this capability. As a result, this alternative is neither reasonable or practicable and does not meet the purpose of and need for the Proposed Action and has been eliminated from detailed study.

#### **2.4.1.2 Reduced Training and Testing**

As described in Section 2.5.1.2 (Reduced Training and Testing) in the 2015 MITT Final EIS/OEIS, a reduction or cessation of training and testing would prevent the armed forces from meeting its statutory requirements and adequately preparing forces for operations ranging from disaster relief to armed conflict. Therefore, this alternative does not meet the purpose of and need for the Proposed Action and has been eliminated from detailed study.

### **2.4.1.3 Alternatives Including Geographic Mitigation Measures within the Study Area**

The Navy considered developing an alternative based on geographic mitigation which would impose time/area restrictions on an expanded list of specific areas of the MITT Study Area associated with the presence of specific species. However, such an alternative would present a patchwork of areas and time periods in which the Navy could conduct required training and testing, preventing the Navy from conducting the full scope of activities necessary to fulfill its Title 10 responsibilities and running counter to the Purpose and Need of the Proposed Action. Thus, such an alternative would not be reasonable. Further, regulations governing the National Environmental Policy Act allow agencies to, "Include appropriate mitigation measures not already included in the proposed action or alternatives." (40 CFR 1502.14(f)). Under both alternatives, the Navy would implement limited geographic mitigation areas that are biologically supported and are practicable to implement. Such areas are more fully described in Appendix I (Geographic Mitigation Assessment). Therefore, appropriate mitigation protective of impacted species would be implemented regardless of alternative selected..

### **2.4.1.4 Simulated Training and Testing Only**

As described in Section 2.5.1.4 (Simulated Training and Testing) in the 2015 MITT Final EIS/OEIS, the Navy continues to use computer simulation for training and testing activities whenever possible; however, there are limits to the realism that current simulation technology can presently provide, and its use cannot substitute for live training or testing. Training and testing through simulated means cannot replicate the conditions in which Navy personnel and platforms are required to conduct military operations. While beneficial as a complementing medium to train and test personnel and platforms, simulation alone cannot accurately replicate both the conditions and the stresses that must be placed on personnel and platforms during training. These conditions and stresses are absolutely vital to adequately preparing Naval forces to conduct the broad spectrum of military operations required of them by operational Commanders. Therefore, simulation as an alternative that completely replaces training and testing in the field does not meet the purpose of and need for the Proposed Action and has been eliminated from detailed study.

### **2.4.1.5 Training and Testing Without the Use of Active Sonar**

As explained in Section 2.4.1.4 (Simulated Training and Testing Only), in order to detect and counter submerged mines and potentially hostile submarines, the Navy uses both passive and active sonar. Sonar proficiency is a complex and perishable skill that requires regular, hands-on training in realistic and diverse conditions. Training and testing with active sonar is needed to find and counter newer-generation submarines around the world, which are growing in number, as are torpedoes and underwater mines, which are true threats to global commerce, national security, and the safety of military personnel. As a result, defense against enemy submarines is a top priority for the Navy. The detection and countering of submarines is paramount to national security. Naval forces cannot counter this threat without the use of active sonar. Because the Navy is statutorily responsible to provide combat ready forces to operational Commanders, it must train in a manner in which it will be utilized in military operations. Accordingly, training and testing without active sonar is not a reasonable alternative and will not be carried forward.

## **2.4.2 Alternatives Carried Forward**

The military's anticipated level of training and testing activity evolves over time based on numerous factors. Over the past several years, the Navy's ongoing sonar reporting program has gathered classified data regarding the number of sonar hours used to meet anti-submarine warfare requirements. These

data allow for a more accurate projection of the number of active sonar hours required to meet anti-submarine warfare training requirements into the reasonably foreseeable future. Alternatives carried forward for analysis in this SEIS/OEIS are discussed in the following subsections and presented in Table 2.5-1 and Table 2.5-2 at the end of this Chapter. As previously discussed, in addition to meeting the Navy's purpose and need to train and test, the action alternatives, and in particular the mitigation measures that are incorporated in the action alternatives, were developed to meet NMFS' independent purpose and need to evaluate the potential impacts of the Navy's activities, determine whether incidental take resulting from the Navy's activities would have a negligible impact on affected marine mammal species and stocks, and prescribe measures to effect the least practicable adverse impact on species or stocks and their habitat, as well as monitoring and reporting requirements.

#### **2.4.2.1 No Action Alternative**

As mentioned above in Section 2.4 (Action Alternatives Development), the Council on Environmental Quality implementing regulations require inclusion of a No Action Alternative and analysis of all reasonable alternatives to provide a clear basis for choice among options by the decision maker and the public (40 CFR section 1502.14). Council on Environmental Quality guidance identifies two approaches in developing the No Action Alternative (46 *Federal Register* 18026). One approach for activities that have been ongoing for long periods of time is for the No Action Alternative to be thought of in terms of continuing the present course of action, or current management direction or intensity, such as the continuation of Navy training and testing at sea in the MITT Study Area at current levels, even if separate legal authorizations under the MMPA and ESA are required. Under this approach, which was used in the 2015 MITT Final EIS/OEIS, the analysis compares the effects of continuing current activity levels (i.e., the "status quo") with the effects of the Proposed Action. The second approach depicts a scenario where no authorizations or permits are issued, the Navy's training and testing activities do not take place, and the resulting environmental effects from taking no action are compared with the effects of the Proposed Action. The Navy applied the second approach in this SEIS/OEIS as it further supports NMFS' regulatory process by presenting the scenario where no authorization will be issued. Additionally, the second approach responds to comments submitted at various stages regarding the 2015 MITT Final EIS/OEIS and during the scoping process of this SEIS/OEIS.

Under the No Action Alternative analyzed in this SEIS/OEIS, the Navy would not conduct the proposed training and testing activities in the MITT Study Area. Other military activities not associated with this Proposed Action would continue to occur. For FDM, the lease agreement between the U.S. government and the Commonwealth of the Northern Mariana Islands would remain in place, and the island would continue to be maintained as a Navy range, although strike warfare would no longer continue on the island. Consequently, the No Action Alternative of not conducting the proposed at-sea training and testing activities in the Study Area is inherently unreasonable in that it does not meet the purpose and need (see Section 1.4, Purpose and Need for Proposed Training and Testing Activities) for the reasons stated below. However, the analysis associated with the No Action Alternative is carried forward in order to compare the degree of the potential environmental effects of the Proposed Action with the conditions that would occur if the Proposed Action did not occur (see Section 3.0.1, Overall Approach to Analysis).

From NMFS' perspective, pursuant to its obligation to grant or deny permit applications under the MMPA, the No Action Alternative involves NMFS denying Navy's application for an incidental take authorization under Section 101(a)(5)(A) of the MMPA. If NMFS were to deny the Navy's application, the

Navy would not be authorized to incidentally take marine mammals and the Navy would not conduct the at-sea proposed training and testing activities in the MITT Study Area.

Cessation of proposed Navy at-sea training and testing activities would mean that the Navy would not meet its statutory requirements and would be unable to properly defend itself and the United States from enemy forces, unable to successfully detect enemy submarines, and unable to effectively use its weapons systems or defensive countermeasures due to a lack of training of forces and testing of systems that replicate the conditions to which Naval forces must operate while executing the range of military operations required to further national security objectives. Navy personnel would essentially not obtain the unique skills or be prepared to safely and effectively use sensors, weapons, and technologies in realistic scenarios required to accomplish the overall mission. For example, sonar proficiency, which is a complex and perishable skill, requires regular, hands-on training in realistic and diverse conditions. In order to detect and counter potentially hostile submarines, the Navy uses both passive and active sonar. Inability to train with active sonar would result in no or greatly diminished anti-submarine warfare capability.

Additionally, without proper training, individual Sailors and Marines serving onboard Navy vessels would not be taught how to properly operate complex equipment in inherently dynamic and dangerous environments. Even with high levels of training, injuries and sometimes even death occur. Therefore, without proper training, it is likely that there would be an increase in the number of mishaps, potentially resulting in the death or serious injury of Sailors and Marines. Failing to allow our Sailors and Marines to achieve and maintain the skills necessary to defend the United States and its interests would result in an unacceptable increase in the danger they willingly face.

Finally, the lack of live training and testing would require a higher reliance on simulated training and testing. While the Navy continues to research new ways to provide realistic training through simulation, there are limits to the realism that technology provides. While simulators are used for the basic training of sonar technicians, they are of limited utility beyond basic training. A simulator cannot match the dynamic nature of the environment, such as bathymetry and sound propagation properties, or the training activities involving several units with multiple crews interacting in a variety of acoustic environments. Sole reliance on simulation would deny service members the ability to develop battle-ready required proficiency in the employment of active sonar during military operations (Section 2.4.1.4, Simulated Training and Testing Only).

#### **2.4.2.2 Alternative 1**

Alternative 1 reflects a representative year of training and testing to account for the typical fluctuation of training cycles, testing programs, and deployment schedules that generally limit the maximum level of training and testing from occurring for the reasonably foreseeable future.

##### **2.4.2.2.1 Training**

Under this alternative, the Navy proposes to conduct training activities into the reasonably foreseeable future, as necessary to meet current and future readiness requirements. These include training activities subject to previous analysis that are currently ongoing and have historically occurred in the Study Area. The requirements for the types of activities to be conducted, as well as the intensity at which they need to occur, have been validated by senior leadership. Specifically, training activities are based on changing world events, advances in technology, and U.S. tactical and strategic priorities. These activities account for force structure changes and include training with new aircraft, vessels, unmanned/autonomous

systems, and weapon systems that will be introduced to the Fleets after August 2020. The numbers and locations of all proposed training activities are provided in Table 2.5-1.

#### **2.4.2.2.2 Testing**

Alternative 1 reflects a level of testing activities to be conducted into the reasonably foreseeable future, with adjustments from the 2015 MITT Final EIS/OEIS that account for changes in the types and tempo (increases or decreases) of testing activities to meet current and future military readiness requirements. The majority of testing activities that would be conducted under this alternative are the same as or similar as those conducted currently or in the past. This alternative includes the testing of new systems using new technologies and takes into account inherent uncertainties in this type of testing. The numbers and locations of all proposed testing activities are listed in Table 2.5-2.

#### **2.4.2.2.3 Mitigation Measures**

The Navy's entire suite of mitigation measures was applied to Alternative 1 to ensure that: (1) the benefit of mitigation measures to environmental and cultural resources was considered during the applicable environmental analyses, and (2) Navy Senior Leadership approved each mitigation measure included in this Draft SEIS/OEIS under Alternative 1. Navy Senior Leadership reviewed relevant supporting information to make a fully informed decision, including the benefit of mitigation measures to environmental and cultural resources, and the impacts that implementing mitigation will have on training and testing activities under Alternative 1. As discussed in Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment), the final suite of mitigation measures that will be included in the Final SEIS/OEIS will represent the maximum level of mitigation that is practicable for the Navy to implement when balanced against impacts to safety, sustainability, and the ability to continue meeting its mission requirements.

#### **2.4.2.3 Alternative 2**

##### **2.4.2.3.1 Training**

Alternative 2 includes the same types of training activities as Alternative 1 but also considers additional Fleet exercises and associated unit-level activities should unanticipated emergent world events require increased readiness levels. For example, Alternative 2 contemplates Joint Multi-Strike Group Exercises (i.e., Valiant Shield) occurring annually as compared to every other year under Alternative 1 (see Table 2.5-1). Additionally, Alternative 2 contemplates three (vice two) Small Joint Coordinated Anti-Submarine Warfare exercises (Multi-Sail/Guam Exercises) per year with a 50 percent increase in associated unit-level events (e.g., Missile Exercise [Surface-to-Air]). The numbers and locations of all proposed training activities are provided in Section 2.3 (Proposed Activities) and listed in Table 2.5-1.

Alternative 2 reflects the maximum number of training activities that could occur within a given year, and assumes that the maximum number of Fleet exercises would occur annually. This allows for the greatest flexibility for the Navy to maintain readiness when considering potential changes in the national security environment, fluctuations in training and deployment schedules, and anticipated in-theater demands.

##### **2.4.2.3.2 Testing**

Alternative 2 entails a level of testing activities to be conducted into the reasonably foreseeable future. Under Alternative 2, types and tempo of testing activities would increase compared to Alternative 1 (see Table 2.5-2). This alternative includes the contingency for augmenting some weapon systems tests in response to potential increased world conflicts and changing Navy leadership priorities as the result of a

direct challenge from a naval opponent that possesses near peer capabilities. The numbers and locations of all proposed testing activities are listed in Table 2.5-2.

#### **2.4.2.3.3 Mitigation Measures**

The Navy's entire suite of mitigation measures was applied to Alternative 2 to ensure that: (1) the benefit of mitigation measures to environmental and cultural resources was considered during the applicable environmental analyses, and (2) Navy Senior Leadership approved each mitigation measure included in this Draft SEIS/OEIS under Alternative 2. Navy Senior Leadership reviewed relevant supporting information to make a fully informed decision, including the benefit of mitigation measures to environmental and cultural resources, and the impacts that implementing mitigation will have on training and testing activities under Alternative 2. As discussed in Chapter 5 (Mitigation), the final suite of mitigation measures that will be included in the Final SEIS/OEIS will represent the maximum level of mitigation that is practical for the Navy to implement when balanced against impacts to safety, sustainability, and the ability to continue meeting its mission requirements.


### **2.4.3 Comparison of Proposed Sonar and Explosive Use in the Action Alternatives**

#### **2.4.3.1 Sonar Use**

As part of the 2015 MITT Final EIS/OEIS and 2015 MMPA Letter of Authorization, NMFS authorized the Navy to use non-impulsive sound sources including sonars and other transducers. Sonars and other transducers were grouped into classes that share one or more attributes, such as frequency range or purpose of use. The classes were further sorted into sound source bins. These bins are defined and quantified in Section 3.0.5.1 (Acoustic Stressors).

In the 2015 analysis, the Navy identified the type of sonar source that resulted in the highest number of exposures to marine mammals, which was hull-mounted mid-frequency active sonar in bin MF1. The Navy was authorized 1,872 hours of MF1 annually in the 2015 MITT Final EIS/OEIS and by NMFS under the MMPA permit and ESA Biological Opinion.

In this SEIS/OEIS, the Navy is evaluating the potential impacts associated with 1,729 hours of MF1 annually under Alternative 1, a reduction of approximately 8 percent from the currently authorized total (Figure 2.4-1). Under Alternative 2, the Navy is evaluating the potential impacts associated with 1,818 hours of MF1 annually, which is a decrease of approximately 3 percent over currently permitted levels.




**Figure 2.4-1: Proposed Annual Total Hull-Mounted Mid-Frequency Sonar Hour Use Compared to the Number Authorized in the 2015–2020 Marine Mammal Protection Act Permit**


**2.4.3.2 Explosives Use**

As part of the 2015 MITT Final EIS/OEIS and 2015 MMPA Letter of Authorization, NMFS authorized the Navy to use impulsive sources (i.e., explosives). Similar to non-impulsive sources, the Navy sorted explosive sources into bins based on the net explosive weight of the explosive. After analyzing the level of explosive activities conducted during Phase II, the Navy identified that some explosive sources were incorrectly classed into bins with greater net explosive weights than actually is present in the munition. For example, 20 millimeter rounds were considered in bin E1 (defined as 0.1–0.25 pounds net explosive weight) during Phase II, but have less than 0.1 pound of net explosive weight (defined as bin E0) and are, therefore, analyzed qualitatively instead of quantitatively for Phase III. Additionally, in Phase II, munitions within the same category were all analyzed with the highest net explosive weight for all munitions in that category. For example, most bombs were analyzed as bin E12 (to account for the largest potential for environmental impact), whereas many fall within bins E9 and E10. For Phase III, munitions were divided into more appropriate bins based on current and anticipated weapon inventory. Bins used to sort explosive munitions are further defined and quantified in Section 3.0.4.2 (Explosive Stressors).

See Figure 2.4-2 and Figure 2.4-3 for a comparison between explosives authorized for training and testing in the 2015 MITT Final EIS/OEIS and proposed in this SEIS/OEIS. The number of impulsive sources in bins E2, E5, E8, E9, and E10 would increase in this SEIS/OEIS compared with the totals analyzed in the 2015 MITT Final EIS/OEIS. The number of impulsive sources that would decrease under this SEIS/OEIS are in bins E1, E3, E4, E6, E11, and E12.



**Figure 2.4-2: Proposed Annual Explosives Use (Bins E1–E5) Compared to the 2015–2020 Marine Mammal Protection Act Permit**



**Figure 2.4-3: Proposed Annual Explosives Use (Bins E6–E12) Compared to the 2015–2020 Marine Mammal Protection Act Permit**

## 2.5 Comparison of Alternatives

The following tables compare the proposed SEIS/OEIS action alternatives (Alternative 1 and Alternative 2) with the ongoing training and testing activities (Table 2.5-1, Table 2.5-2). Each table describes the activities in terms of the activity name and where in the Study Area the Navy proposes to conduct it (first two columns). The next two columns show the annual occurrence and ordnance or other expended items (if any) involved in the activity as is currently ongoing (under the heading “2015 MITT EIS/OEIS Ongoing Activities”). The final two pairs of columns present the same information (annual occurrence and ordnance/items) as the activities are analyzed in this SEIS/OEIS for Alternative 1 and Alternative 2, respectively. Table 2.5-1 is the table of training activities, Table 2.5-2 is the table of Naval Sea Systems Command testing activities and Naval Air Systems Command testing activities.

**Table 2.5-1: Current and Proposed Training Activities**

| Range Activity  | Location  | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |   | Supplemental EIS/OEIS<br>(Alternative 1) |   | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|---|--|---|--|---|--|----------------------------------|
|   |   | No. of events<br>(per year)  | Ordnance (Number<br>per year)                                   | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)                                | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| <b>Legend:</b>  |   |  | = Decrease in number of events<br>from 2015 Final MITT EIS/OEIS |  | = Increase in number of events<br>from 2015 Final MITT EIS/OEIS |  |                                  |
| <b>Major Training Events</b>  |   |  |   |  |   |  |                                  |
| Joint Expeditionary Exercise  | Study Area; MIRC  | 1  | Note 1  | 1  | Note 1  | 1  | Note 1                           |
| Joint Multi-Strike Group<br>Exercise  | Study Area; MIRC  | 1  | Note 1  | 1 every other<br>year                    | Note 1  | 1  | Note 1                           |
| Marine Air Ground Task<br>Force Exercise (Amphibious)<br>– Battalion                    | Study Area to<br>nearshore; MIRC;<br>Tinian; Guam;<br>Rota; Saipan; FDM | 4  | Note 1  | 4  | Note 1  | 4  | Note 1                           |
| <b>Air Warfare (AW) (previously named Anti-Air Warfare in 2015 MITT Final EIS/OEIS)</b> |   |  |   |  |   |  |                                  |
| Air Combat Maneuver   | Study Area ><br>12 NM from land:<br>SUA                                 | 4,800  | None  | 3,800                                    | None  | 3,800                                    | None                             |
| Air Defense Exercise<br>(ADEX)  | Study Area ><br>12 NM from land:<br>SUA                                 | 100  | None  | 100                                      | None  | 100                                      | None                             |
| Air Intercept Control<br>(AIC)  | Study Area ><br>12 NM from land:<br>SUA                                 | 4,800  | None  | 5,300                                    | None  | 5,300                                    | None                             |
| Gunnery Exercise (GUNEX)<br>(Air-to-Air [A-A]) – Medium-<br>caliber                     | Study Area SUA ><br>12 NM from land                                     | 36   | 9,000 rounds  | 36                                       | 9,000<br>rounds   | 36                                       | 9,000<br>rounds                  |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity  | Location                            | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|-------------------------------------|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|   |                                     | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| Missile Exercise (Missilex)<br>(A-A)  | Study Area SUA ><br>12 NM from land | 18   | 36 explosive missiles         | 18                                       | 36 explosive<br>missiles         | 18                                       | 36 explosive<br>missiles         |
| GUNEX (Surface-to-Air [S-A])<br>– Large-caliber                               | Study Area SUA ><br>12 NM from land | 5  | 40 rounds                     | 6  | 60 rounds                        | 9  | 90 rounds                        |
| GUNEX [S-A] – Medium-<br>caliber  | Study Area SUA ><br>12 NM from land | 12   | 24,000 rounds                 | 13                                       | 26,000<br>rounds                 | 19                                       | 38,000<br>rounds                 |
| MISSILEX [S-A]  | Study Area SUA ><br>12 NM from land | 15   | 15 explosive missiles         | 18                                       | 18 explosive<br>missiles         | 27                                       | 27 explosive<br>missiles         |
| <b>Amphibious Warfare (AMW)</b>   |                                     |  |                               |  |                                  |  |                                  |
| Naval Surface Fire Support<br>Exercise (FIREX) – Land-<br>based target (Land) | FDM                                 | 10   | 1,800 NEPM rounds             | 10                                       | 2,800<br>explosive<br>rounds     | 15                                       | 4,200<br>explosive<br>rounds     |
|   |                                     |  | 1,000 explosive<br>rounds     |  |                                  |  |                                  |
| Amphibious Rehearsal, No<br>Landing   | Study Area and<br>Nearshore         | 12   | None                          | 12                                       | None                             | 12                                       | None                             |
| Amphibious Assault  | MIRC; Tinian;<br>Guam               | 6  | Blanks; Simunitions           | 6  | Blanks;<br>Simunitions           | 6  | Blanks;<br>Simunitions           |
| Amphibious Raid   | MIRC; Tinian;<br>Guam; Rota         | 6  | Blanks; Simunitions           | 6  | Blanks;<br>Simunitions           | 6  | Blanks;<br>Simunitions           |
| Noncombatant Evacuation<br>Operation  | MIRC; Guam;<br>Tinian; Rota         | 5  | Blanks; Simunitions           | 5  | Blanks;<br>Simunitions           | 5  | Blanks;<br>Simunitions           |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity   | Location  | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|--|---|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|  |   | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| Humanitarian Assistance/Disaster Relief Operations                       | MIRC; Guam; Tinian; Rota                                  | 5  | Blanks; Simunitions           | 5  | Blanks; Simunitions              | 5  | Blanks; Simunitions              |
| Unmanned Aerial Vehicle – Intelligence, Surveillance, and Reconnaissance | MIRC; SUA   | 100  | None                          | 100                                      | None                             | 100                                      | None                             |
| Special Purpose Marine Air Ground Task Force Exercise                    | Study Area to nearshore; MIRC; Tinian; Guam; Rota; Saipan | 2  | Note 1                        | 2  | Note 1                           | 2  | Note 1                           |
| <b>Anti-Submarine Warfare (ASW)</b>                                      |   |  |                               |  |                                  |  |                                  |
| Tracking Exercise (TRACKEX) –Helicopter (Helo)                           | Study Area > 3 NM from land; Transit Corridor             | 62   | None/ REXTORP                 | 10                                       | None                             | 10                                       | None/ REXTORP                    |
| Torpedo Exercise (TORPEX)–Helo   | Study Area > 3 NM from land                               | 4  | 4 EXTORP                      | 4  | 4 EXTORP                         | 6  | 6 EXTORP                         |
| TRACKEX – Maritime Patrol (Extended Echo Ranging Sonobuoys)              | Study Area > 3 NM from land                               | 11   | None                          | 0  | 0                                | 0  | 0                                |
| TRACKEX – Maritime Patrol Aircraft                                       | Study Area > 3 NM from land                               | 34   | None/ REXTORP                 | 36                                       | None                             | 36                                       | None/ REXTORP                    |
| TORPEX – Maritime Patrol Aircraft  | Study Area > 3 NM from land                               | 4  | 4 EXTORP                      | 4  | 4 EXTORP                         | 6  | 6 EXTORP                         |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity  | Location  | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative)   |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|---|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|   |   | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| TRACKEX – Surface   | Study Area > 3 NM<br>from land                      | CG/DDG-92<br>FFG-30<br>LCS-10  | None/<br>REXTORP              | 91                                       | None/<br>REXTORP                 | 91                                       | None/<br>REXTORP                 |
| TORPEX – Surface  | Study Area > 3 NM<br>from land                      | 3  | 3 EXTORP                      | 4  | 4 EXTORP                         | 6  | 6 EXTORP                         |
| TRACKEX – Submarine (Sub)   | Study Area > 3 NM<br>from land; Transit<br>Corridor | 12   | None                          | 4  | None                             | 4  | None                             |
| TORPEX – Sub  | Study Area > 3 NM<br>from land                      | 10   | 40 MK-48 EXTORP               | 6  | 24 MK-48<br>EXTORP               | 9  | 36 MK-48<br>EXTORP               |
| Small Joint Coordinated<br>ASW exercise (e.g., Multi-<br>Sail/GUAMEX/SWATT) (see<br>Note 2) | Study Area > 3 NM<br>from land                      | Not called<br>out in<br>previous<br>document,<br>but<br>components<br>were covered<br>under several<br>unit-level<br>exercises | None                          | 2  | None                             | 3  | None                             |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                                       | Location   | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|--|--|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|  |  | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| <b>Electronic Warfare (EW)</b>                       |  |  |                               |  |                                  |  |                                  |
| Electronic Warfare Operations (EW Ops)               | Study Area   | 480  | None                          | 522                                      | None                             | 522                                      | None                             |
| Counter Targeting Flare Exercise (FLAREX) – Aircraft | Study Area > 12 NM from land                         | 3,200  | 25,600 rounds                 | 2,200                                    | 17,600 rounds                    | 2,200                                    | 17,600 rounds                    |
| Counter Targeting Chaff Exercise (CHAFFEX) – Ship    | Study Area > 12 NM from land                         | 40   | 240 rounds                    | 41                                       | 244 rounds                       | 60                                       | 360 rounds                       |
| CHAFFEX –Aircraft                                    | Study Area > 12 NM from land                         | 3,200  | 25,600 rounds                 | 2,200                                    | 17,600 rounds                    | 2,200                                    | 17,600 rounds                    |
| <b>Expeditionary Warfare</b>                         |  |  |                               |  |                                  |  |                                  |
| Personnel Insertion/ Extraction                      | MIRC; Guam; Tinian; Rota                             | 240  | None                          | 365                                      | None                             | 365                                      | None                             |
| Parachute Insertion                                  | MIRC parachute drop zones; Guam; Tinian; Rota        | 20   | None                          | 64                                       | None                             | 64                                       | None                             |
| <b>Mine Warfare (MIW)</b>                            |  |  |                               |  |                                  |  |                                  |
| Civilian Port Defense                                | Mariana littorals; MIRC; Inner and Outer Apra Harbor | 1  | None                          | 1  | None                             | 1  | None                             |
| Mine Laying  | MIRC Warning Areas                                   | 4  | 480 mine shapes               | 4  | 480 mine shapes                  | 4  | 480 mine shapes                  |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity  | Location   | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|--|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|   |  | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| Mine Neutralization –<br>Explosive Ordnance Disposal<br>(EOD)                             | Agat Bay<br>underwater<br>detonation site<br><br>Piti and Outer<br>Apra Harbor<br>underwater<br>detonation sites | 20   | 20 explosive charges          | 20                                       | 20 explosive<br>charges          | 20                                       | 20 explosive<br>charges          |
| Limpet Mine Neutralization<br>System  | Mariana littorals;<br>Inner and Outer<br>Apra Harbor   | 40   | 40 charges                    | 60                                       | 60<br>charges                    | 60                                       | 60 charges                       |
| Airborne Mine<br>Countermeasure – Towed<br>Mine Detection                                 | Study Area;<br>nearshore   | 4  | None                          | 4  | None                             | 4  | None                             |
| Mine Countermeasure<br>Exercise – Towed Sonar<br>(AQS-20, LCS)                            | Study Area   | 4  | None                          | 4  | None                             | 4  | None                             |
| Mine Countermeasure<br>Exercise – Surface Ship<br>Sonar (SQQ-32, MCM)                     | Study Area   | 4  | None                          | 4  | None                             | 4  | None                             |
| Mine Neutralization –<br>Remotely Operated Vehicle<br>Sonar (ASQ-235 [AQS-20],<br>SLQ-48) | Study Area   | 4  | 4 explosive<br>neutralizers   | 4  | 4 explosive<br>neutralizers      | 4  | 4 explosive<br>neutralizers      |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity   | Location   | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|--|--|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|  |  | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| Mine Countermeasure –<br>Towed Mine Neutralization       | Study Area   | 4  | None                          | 4  | None                             | 4  | None                             |
| Underwater Demolition<br>Qualification/<br>Certification | Agat Bay<br>underwater<br>detonation site<br>Piti and Outer<br>Apra Harbor<br>underwater<br>detonation sites | 30   | 30 explosive charges          | 45                                       | 45 explosive<br>charges          | 45                                       | 45 explosive<br>charges          |
| Submarine Mine Exercise                                  | Mariana Littorals,<br>Inner/Outer Apra<br>Harbor   | 16   | None                          | 1  | None                             | 1  | None                             |
| Surface Ship Object<br>Detection                         | Study Area   | Not<br>previously<br>analyzed  | Not previously<br>analyzed    | 6  | None                             | 6  | None                             |
| <b>Strike Warfare (STW)</b>                              |  |  |                               |  |                                  |  |                                  |
| Bombing Exercise (BOMBEX)<br>(Air-to-Ground [A-G])       | FDM  | 2,300  | 2,670 NEPM                    | 2,300                                    | 2,670 NEPM                       | 2,300                                    | 2,670 NEPM                       |
|  |  |  | 6,242 explosive<br>rounds     |  | 6,242<br>explosive<br>rounds     |  | 6,242<br>explosive<br>rounds     |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                                  | Location                            | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                                      | Supplemental EIS/OEIS<br>(Alternative 1) |                                     | Supplemental EIS/OEIS<br>(Alternative 2) |                                     |
|---|-------------------------------------|--|--------------------------------------|--|-------------------------------------|--|-------------------------------------|
|   |                                     | No. of events<br>(per year)  | Ordnance (Number<br>per year)        | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)    | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)    |
| GUNEX (A-G)                                     | FDM                                 | 96   | 24,000 small-caliber rounds          | 96                                       | 24,000 small-caliber rounds         | 96                                       | 24,000 small-caliber rounds         |
|   |                                     |  | 94,150 medium-caliber rounds         |  | 94,650 medium-caliber rounds        |  | 94,650 medium-caliber rounds        |
|   |                                     |  | 17,350 explosive med.-caliber rounds |  | 17,500 explosive med-caliber rounds |  | 17,500 explosive med-caliber rounds |
|   |                                     |  | 200 explosive large-caliber rounds   |  | 200 explosive large-caliber rounds  |  | 200 explosive large-caliber rounds  |
| MISSILEX  | FDM                                 | 85   | 2,000 explosive rockets              | 115                                      | 2,000 explosive rockets             | 115                                      | 2,000 explosive rockets             |
|   |                                     |  | 85 explosive missiles                |  | 115 explosive missiles              |  | 115 explosive missiles              |
| <b>Surface Warfare (SUW)</b>                    |                                     |  |                                      |  |                                     |  |                                     |
| GUNEX (Air-to-Surface [A-S])<br>– Small-caliber | Study Area SUA ><br>12 NM from land | 242  | 48,040 rounds                        | 321                                      | 128,400 rounds                      | 321                                      | 128,400 rounds                      |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                          | Location   | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|--|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|   |  | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| GUNEX (A-S) – Medium-caliber            | Study Area SUA > 12 NM from land; Transit Corridor | 295  | 29,500 non-explosive rounds   | 120                                      | 3,600 explosive rounds           | 120                                      | 3,600 explosive rounds           |
|   |  |  | 7,150 explosive rounds        |  |                                  |  |                                  |
| MISSILEX (A-S) – Rocket)                | Study Area SUA > 12 NM from land                   | 3  | 114 rockets (114 explosive)   | 111                                      | 2,109                            | 111                                      | 2,109                            |
| MISSILEX (A-S)                          | Study Area SUA > 12 NM from land                   | 20   | 20 explosive missiles         | 10                                       | 18 explosive missiles            | 10                                       | 18 explosive missiles            |
| Laser Targeting (at sea)                | Study Area SUA > 12 NM from land                   | 600  | None                          | 600                                      | None                             | 600                                      | None                             |
| BOMBEX (A-S)                            | Study Area > 50 NM from land                       | 37   | 368 NEPM                      | 37                                       | 368 NEPM                         | 37                                       | 368 NEPM                         |
|   |  |  | 184 explosive rounds          |  | 184 explosive rounds             |  |                                  |
| Torpedo Exercise (Submarine to Surface) | Study Area > 3 NM from land                        | 5  | 10 EXTORP                     | 0  | None                             | 0  | None                             |
| MISSILEX (Surface-to-Surface [S-S])     | Study Area > 50 NM from land                       | 12   | 12 explosive missiles         | 19                                       | 19 explosive missiles            | 28                                       | 28 explosive missiles            |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                                  | Location   | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                                | Supplemental EIS/OEIS<br>(Alternative 1) |  | Supplemental EIS/OEIS<br>(Alternative 2) |  |
|---|--|--|--------------------------------|--|--|--|--|
|   |  | No. of events<br>(per year)  | Ordnance (Number<br>per year)  | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)       | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)       |
| GUNEX (S-S) Ship – Large-<br>caliber            | Study Area SUA ><br>12 NM from land;<br>Transit Corridor | 140  | 5,198 non-explosive<br>rounds  | 170                                      | 16,320 non-<br>explosive<br>rounds     | 255                                      | 24,480 non-<br>explosive<br>rounds     |
|   |  |  | 500 explosive rounds           |  | 510<br>explosive<br>rounds             |  | 765<br>explosive<br>rounds             |
| GUNEX (S-S) Ship – Small-<br>and Medium-caliber | Study Area SUA ><br>12 NM from land;<br>Transit Corridor | 100  | 21,000 non-explosive<br>rounds | 162                                      | 172,010<br>non-<br>explosive<br>rounds | 234                                      | 250,800<br>non-<br>explosive<br>rounds |
|   |  |  | 900 explosive rounds           |  | 480<br>explosive<br>rounds             |  | 720<br>explosive<br>rounds             |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity  |                    | Location  | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |  | Supplemental EIS/OEIS<br>(Alternative 1) |  | Supplemental EIS/OEIS<br>(Alternative 2) |  |
|---|--------------------|---|--|--|--|--|--|--|
|   |                    |   | No. of events<br>(per year)  | Ordnance (Number<br>per year)  | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)   | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)   |
| Sinking Exercise (SINKEX)<br>Representative ordnance.<br>Actual ordnance used will<br>vary. |                    | Study Area ><br>50 NM from land<br>and > 1,000<br>fathoms depth | 2  | 28 explosive Bombs<br>42 explosive Missiles<br>800 explosive Large-<br>caliber rounds<br>2 MK-48 explosive<br>4 explosive<br>Demolitions | 1  | 28 explosive<br>Bombs<br>42 explosive<br>Missiles<br>800<br>explosive<br>Large-<br>caliber<br>rounds<br>2 MK-48<br>explosive<br>4 explosive<br>Demolitions | 1  | 28 explosive<br>Bombs<br>42 explosive<br>Missiles<br>800<br>explosive<br>Large-<br>caliber<br>rounds<br>2 MK-48<br>explosive<br>4 explosive<br>Demolitions |
| GUNEX [S-S]<br>Boat – Small<br>and Medium-<br>caliber                                       | Medium-<br>caliber | Study Area SUA ><br>12 NM from land;<br>Transit Corridor        | 10   | 2,000 non-explosive<br>rounds  | 20                                       | 4,000 non-<br>explosive<br>rounds  | 20                                       | 4,000 non-<br>explosive<br>rounds  |
|   |                    |   |  | 100 explosive rounds   |  | 200<br>explosive<br>rounds   |  | 200<br>explosive<br>rounds   |
|   | Small-<br>caliber  | Study Area > 3 NM<br>from land; Transit<br>Corridor             | 40   | 36,000 rounds  | 43                                       | 36,600<br>rounds   | 43                                       | 36,600<br>rounds   |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                             | Location                                | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                                 | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|--|---|--|---------------------------------|--|----------------------------------|--|----------------------------------|
|  |   | No. of events<br>(per year)  | Ordnance (Number<br>per year)   | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| Maritime Security Operations (MSO)         | Study Area; MIRC                        | 40   | 200 G911 anti-swimmer grenades  | 40                                       | 200 G911 anti-swimmer grenades   | 40                                       | 200 G911 anti-swimmer grenades   |
| <b>Other</b>                               |   |  |                                 |  |                                  |  |                                  |
| Direct Action (Tactical Air Control Party) | FDM                                     | 18   | 18,000 small-caliber rounds     | 18                                       | 30,000 small-caliber rounds      | 18                                       | 30,000 small caliber rounds      |
|  |   |  | 600 explosives (grenade/mortar) |  | 1,000 med-caliber explosive      |  | 1,000 med-caliber explosive      |
|  |   |  |                                 |  | 1,000 explosive (grenade mortar) |  | 1,000 explosive (grenade mortar) |
| Intelligence, Surveillance, Reconnaissance | MIRC; Guam; Tinian; Rota; Saipan        | 16   | None                            | 44                                       | None                             | 44                                       | None                             |
| Precision Anchoring                        | Apra Harbor; Mariana Islands anchorages | 18   | None                            | 18                                       | None                             | 18                                       | None                             |
| Search and Rescue At Sea                   | Study Area                              | 40   | None                            | 45                                       | None                             | 45                                       | None                             |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity                                       | Location  | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                   | Supplemental EIS/OEIS<br>(Alternative 2) |                                   |
|--|---|--|-------------------------------|--|-----------------------------------|--|-----------------------------------|
|  |   | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)  | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)  |
| Small Boat Attack                                    | Study Area > 3 NM<br>from land  | 6  | 2,100 small-caliber<br>rounds | 18                                       | 2,100 small-<br>caliber<br>rounds | 27                                       | 3,150 small-<br>caliber<br>rounds |
|  | Study Area  | 12   | 4,000 blank rounds            |  | 4,000 blank<br>rounds             |  | 6,000 blank<br>rounds             |
| Submarine Navigation                                 | Apra Harbor and<br>Mariana littorals                                      | 8  | None                          | 8  | None                              | 8  | None                              |
| Submarine Sonar<br>Maintenance                       | Study Area > 3 NM<br>from land; Inner<br>Apra Harbor;<br>Transit Corridor | 48   | None                          | 86                                       | None                              | 86                                       | None                              |
| Surface Ship Sonar<br>Maintenance                    | Study Area > 3 NM<br>from land; Inner<br>Apra Harbor;<br>Transit Corridor | 42   | None                          | 44                                       | None                              | 44                                       | None                              |
| Underwater Survey                                    | Mariana littorals   | 16   | None                          | 32                                       | None                              | 32                                       | None                              |
| Unmanned Aerial Training<br>and Certification        | Study Area; MIRC<br>airfields; <sup>1</sup> MIRC<br>SUA                   | 1,000  | None                          | 951                                      | None                              | 951                                      | None                              |
| Unmanned Underwater<br>Vehicle <sup>2</sup> Training | MIRC, Warning<br>Areas  | N/A  | N/A                           | 64                                       | None                              | 64                                       | None                              |

**Table 2.5-1: Current and Proposed Training Activities (continued)**

| Range Activity | Location | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |                               | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|----------------|----------|--|-------------------------------|--|----------------------------------|--|----------------------------------|
|                |          | No. of events<br>(per year)  | Ordnance (Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |

Legend:  = Decrease in number of events from 2015 Final MITT EIS/OEIS

= Increase in number of events from 2015 Final MITT EIS/OEIS

Note 1: All ordnance use during the conduct of these exercises is analyzed within the Primary Mission events listed in this table.

Note 2: Small Joint Coordinated ASW exercise was not called out in previous document, but the components of the exercise were covered under several unit level activities.

Notes: MITT = Mariana Islands Training and Testing, ROD = Record of Decision, EIS = Environmental Impact Statement, OEIS = Overseas Environmental Impact Statement, MIRC = Mariana Islands Range Complex, FDM = Farallon de Medinilla, N/A = Not Applicable, No. = Number, SUA = Special Use Airspace, NM = Nautical Mile(s), NEPM = Non-Explosive Practice Munitions, EXTORP = Exercise Torpedo, REXTORP = Recoverable Exercise Torpedo

**Table 2.5-2: Current and Proposed Testing Activities**

| Range Activity   | Location                        | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |   | Supplemental EIS/OEIS<br>(Alternative 1) |   | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|--|---------------------------------|--|---|--|---|--|----------------------------------|
|  |                                 | No. of events<br>(per year)  | Ordnance<br>(Number per<br>year)                                | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)                                | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| <b>Legend:</b>   |                                 |  | = Decrease in number of events<br>from 2015 Final MITT EIS/OEIS |  | = Increase in number of events<br>from 2015 Final MITT EIS/OEIS |  |                                  |
| <b>NAVAL AIR SYSTEMS COMMAND PROPOSED TESTING ACTIVITIES</b>   |                                 |  |   |  |   |  |                                  |
| <b>Surface Warfare (SUW)</b>   |                                 |  |   |  |   |  |                                  |
| Air-to-Surface Missile Test  | Study Area > 50 NM<br>from land | 8  | 8 Harpoon<br>Missiles   | 4  | 4 Harpoon<br>Missiles   | 4  | 4 Harpoon<br>Missiles            |
|  |                                 |  | (up to 4<br>explosive)  |  | (up to 4<br>explosive)  |  | (up to 4<br>explosive)           |
| <b>Anti-Submarine Warfare (ASW)</b>  |                                 |  |   |  |   |  |                                  |
| Anti-Submarine Warfare<br>Tracking Test – Maritime<br>Patrol Aircraft (Sonobuoys)  | Study Area > 3 NM from<br>land  | 188  | 240 IEER<br>553 SUS   | 26                                       | 392 SUS   | 26                                       | 392 SUS                          |
| Anti-Submarine Warfare<br>Torpedo Test   | Study Area > 3 NM from<br>land  | 40   | 40 EXTORP   | 20                                       | 20<br>REXTORPs  | 20                                       | 20<br>REXTORPs                   |
| <b>Electronic Warfare (EW)</b>   |                                 |  |   |  |   |  |                                  |
| Intelligence, Surveillance,<br>Reconnaissance /Electronic<br>Warfare Testing (previously<br>named Broad Area Maritime<br>Surveillance Testing – MQ-<br>4C) | Study Area > 3 NM from<br>land  | 10   | None  | 20                                       | None  | 20                                       | None                             |

**Table 2.5-2: Current and Proposed Testing Activities (continued)**

| Range Activity  | Location                        | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |  | Supplemental EIS/OEIS<br>(Alternative 1) |                                      | Supplemental EIS/OEIS<br>(Alternative 2) |                                       |
|---|---------------------------------|--|--|--|--------------------------------------|--|---------------------------------------|
|   |                                 | No. of events<br>(per year)  | Ordnance<br>(Number per<br>year)             | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)     | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)      |
| <b>NAVAL SEA SYSTEMS COMMAND TESTING ACTIVITIES</b>   |                                 |  |  |  |                                      |  |                                       |
| <b>Anti-Submarine Warfare</b>   |                                 |  |  |  |                                      |  |                                       |
| Anti-Submarine Warfare<br>Mission Package Testing   | Mariana Island Range<br>Complex | 33   | None   | 100                                      | 8 torpedoes<br>(non-<br>explosive)   | 100                                      | 8 torpedoes<br>(non-<br>explosive)    |
| At-Sea Sonar Testing  | Study Area                      | 20   | None   | 3  | None                                 | 7  | None                                  |
| Countermeasure Testing  | Study Area                      | 2  | 56 torpedoes                                 | 0  | None                                 | 0  | None                                  |
| Torpedo (Explosive) Testing   | Mariana Island Range<br>Complex | 2*   | 20 torpedoes<br>(up to 8 non-<br>explosive*) | 2  | 4 explosive<br>(8 non-<br>explosive) | 3  | 6 explosive<br>(12 non-<br>explosive) |
| Torpedo (Non-explosive)<br>Testing  | Mariana Island Range<br>Complex |  |  | 6  | 28 non-<br>explosive                 | 7  | 37 non-<br>explosive                  |
| <b>Mine Warfare</b>   |                                 |  |  |  |                                      |  |                                       |
| Mine Countermeasure and<br>Neutralization Testing<br>(Previously covered under<br>Mine Countermeasure<br>Mission Package Testing) | Mariana Island Range<br>Complex | 32   | 48 neutralizers<br>(up to 24<br>explosive)   | 3  | 40<br>neutralizers                   | 3  | 40<br>neutralizers                    |
| <b>Electronic Warfare (EW)</b>  |                                 |  |  |  |                                      |  |                                       |
| Radar and Other System<br>Testing   | Study Area                      | Not<br>Previously<br>Analyzed  | Not Previously<br>Analyzed                   | 54                                       | None                                 | 60                                       | None                                  |

**Table 2.5-2: Current and Proposed Testing Activities (continued)**

| Range Activity  | Location                     | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |  | Supplemental EIS/OEIS<br>(Alternative 1) |   | Supplemental EIS/OEIS<br>(Alternative 2) |  |
|---|------------------------------|--|--|--|---|--|--|
|   |                              | No. of events<br>(per year)  | Ordnance<br>(Number per<br>year)                   | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)                          | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year)                           |
| <b>Shipboard Protection Systems and Swimmer Defense Testing</b> |                              |  |  |  |   |  |  |
| Pierside Integrated Swimmer Defense                             | Inner Apra Harbor            | 11   | None   | 0  | None  | 0  | None   |
| <b>Surface Warfare</b>  |                              |  |  |  |   |  |  |
| Gun Testing – Large Caliber                                     | Mariana Island Range Complex | 4  | 5,600 rounds<br>(Up to 3,290 in-air explosives)    | 0  | None  | 0  | None   |
| Gun Testing – Medium Caliber                                    | Mariana Island Range Complex | 4  | 4,080 rounds<br>(up to 2,040 explosives)           | 0  | None  | 0  | None   |
| Gun Testing – Small Caliber                                     | Study Area                   | 4  | 2,000 rounds                                       | 0  | None  | 0  | None   |
| Missile and Rocket Testing                                      | Mariana Island Range Complex | 4  | 32<br>missiles/rockets<br>(up to 16<br>explosives) | 0  | None  | 0  | None   |
| Kinetic Energy Weapon Testing                                   | Study Area                   | 50   | 2,000<br>projectiles                               | 4  | 80<br>projectiles<br>160 non-<br>explosive<br>projectiles | 9  | 180<br>projectiles<br>360 non-<br>explosive<br>projectiles |
|   |                              | 1 time-only<br>event   | 5,000<br>projectiles                               |  |   |  |  |

**Table 2.5-2: Current and Proposed Testing Activities (continued)**

| Range Activity  | Location                        | 2015 MITT FINAL EIS/OEIS<br>Ongoing Activities (MITT ROD<br>Alternative) |   | Supplemental EIS/OEIS<br>(Alternative 1) |                                  | Supplemental EIS/OEIS<br>(Alternative 2) |                                  |
|---|---------------------------------|--|---|--|----------------------------------|--|----------------------------------|
|   |                                 | No. of events<br>(per year)  | Ordnance<br>(Number per<br>year)        | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) | No. of events<br>(per year)              | Ordnance<br>(Number<br>per year) |
| <b>Vessel Evaluation (previously named Life Cycle Activities)</b>   |                                 |  |   |  |                                  |  |                                  |
| Ship Signature Testing  | MITT Study Area                 | 17   | None                                    | 0  | None                             | 0  | None                             |
| Undersea Warfare Testing<br>(previously covered under<br>torpedo testing)   | Mariana Island Range<br>Complex | 2*   | 20 torpedoes<br>(up to 8<br>explosive)* | 1  | 8 non-<br>explosive<br>torpedoes | 1  | 8 non-<br>explosive<br>torpedoes |
| <b>Other Testing Activities</b>   |                                 |  |   |  |                                  |  |                                  |
| Simulant Testing  | Study Area                      | Not<br>Previously<br>Analyzed  | Not Previously<br>Analyzed              | 100                                      | None                             | 100                                      | None                             |
| <b>OFFICE OF NAVAL RESEARCH</b>   |                                 |  |   |  |                                  |  |                                  |
| Acoustic and Oceanographic<br>Research (previously named<br>North Pacific Acoustic Lab<br>Philippine Sea 2018–19<br>Experiment, Deep Water) | Study Area                      | 1  | None                                    | 1  | None                             | 1  | None                             |

Legend:  = Decrease in number of events from 2015 Final MITT EIS/OEIS  
 = Increase in number of events from 2015 Final MITT EIS/OEIS

\*Torpedo (Explosive) Testing, Torpedo (Non-explosive) Testing, and Undersea Warfare Testing were previously covered under torpedo testing in the 2015 MITT EIS/OEIS.

Notes: MITT = Mariana Islands Training and Testing, ROD = Record of Decision, EIS = Environmental Impact Statement, OEIS = Overseas Environmental Impact Statement, MIRC = Mariana Islands Range Complex, NM = Nautical Mile(s), No. = Number, EXTORP = Exercise Torpedo (non-explosive), REXTORP = Recoverable Exercise Torpedo (non-explosive), IEER = Improved Extended Echo Ranging, SUS = Signal Underwater Sound

## **REFERENCES**

- U.S. Department of Defense. (2002). *Electromagnetic Environmental Effects: Requirements for Systems*. (MIL-STD-464A). Wright-Patterson Air Force Base, OH: U.S. Air Force/Aeronautical Systems Center.
- U.S. Department of Defense. (2009). *Protecting Personnel from Electromagnetic Fields*. (DoD Instruction 6055.11). Washington, DC: Under Secretary of Defense for Acquisition, Technology, and Logistics.
- U.S. Department of the Navy. (2011). *U.S. Navy Dive Manual*. Washington, DC: Commander, Naval Sea Systems Command.
- U.S. Department of the Navy. (2015). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Fish and Wildlife Service. (2015). *Biological Opinion for the Mariana Islands Training and Testing Program*. Honolulu, HI: U.S. Fish and Wildlife Service Pacific Islands Fish and Wildlife Office.
- U.S. Fish and Wildlife Service. (2016). *Informal Consultation on Mariana Islands Training and Testing Program Affects to Eighteen Newly-Listed Species, Guam and Tinian*. Honolulu, HI: Fish and Wildlife Service, Pacific Islands Fish and Wildlife Office.

---

---

## 3 Affected Environment and Environmental Consequences



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

**3**            **AFFECTED ENVIRONMENT AND ENVIRONMENTAL CONSEQUENCES .....3-1**

    3.0      Introduction ..... 3-1

        3.0.1    Overall Approach to Analysis..... 3-1

            3.0.1.1    Navy Compiled and Generated Data .....3-2

            3.0.1.2    Navy’s Quantitative Analysis to Determine Impacts on Sea Turtles and  
                    Marine Mammals..... 3-3

        3.0.2    Regulatory Framework ..... 3-7

        3.0.3    Resources and Issues Not Carried Forward for More Detailed Discussion ..... 3-8

        3.0.4    Identification of Stressors for Analysis ..... 3-8

            3.0.4.1    Acoustic Stressors ..... 3-9

            3.0.4.2    Explosive Stressors..... 3-20

            3.0.4.3    Energy Stressors ..... 3-23

            3.0.4.4    Physical Disturbance and Strike Stressors ..... 3-25

            3.0.4.5    Entanglement Stressors..... 3-29

            3.0.4.6    Ingestion Stressors..... 3-33

            3.0.4.7    Conceptual Framework for Assessing Effects from Acoustic and Explosive  
                    Activities ..... 3-33

**List of Figures**

Figure 3.0-1: Sonobuoy Launch Depicting the Relative Size of a Small Decelerator/Parachute ..... 3-32

Figure 3.0-2: Aerial Target (Drone) with Parachute Deployed ..... 3-32

Figure 3.0-3: Flow Chart of the Evaluation Process of Sound-Producing Activities..... 3-35

Figure 3.0-4: Two Hypothetical Threshold Shifts ..... 3-39

**List of Tables**

Table 3.0-1: Comparison of Stressors Analyzed..... 3-8

Table 3.0-2: Sonar and Transducer Sources Quantitatively Analyzed ..... 3-13

Table 3.0-3: Sonar and Transducers Qualitatively Analyzed..... 3-15

Table 3.0-4: Representative Aircraft Sound Characteristics ..... 3-17

---

|   |      |
|---|------|
| Table 3.0-5: Sonic Boom Underwater Sound Levels Modeled for F/A-18 Hornet Supersonic Flight .....                     | 3-18 |
| Table 3.0-6: Example Weapons Noise .....  | 3-19 |
| Table 3.0-7: Explosive Sources Quantitatively Analyzed that Could Be Used Underwater or at the Water<br>Surface ..... | 3-22 |
| Table 3.0-8: Typical Air Explosive Munitions During Navy Activities .....   | 3-23 |
| Table 3.0-9: Annual Number of Events in the Study Area Including In-Water Electromagnetic Devices                     | 3-24 |
| Table 3.0-10: Annual Number of Events in the Study Area Including High-Energy Lasers .....                            | 3-25 |
| Table 3.0-11: Annual Number of Events in the Study Area Including Aircraft Movement.....                              | 3-26 |
| Table 3.0-12: Annual Number of Events in the Study Area Including Vessel Movement.....                                | 3-26 |
| Table 3.0-13: Annual Number of Events in the Study Area Including In-Water Devices.....                               | 3-26 |
| Table 3.0-14: Annual Number of Non-Explosive Practice Munitions Expended At Sea in the Study Area<br>.....            | 3-26 |
| Table 3.0-15: Annual Number of Other Military Expended Materials Used At Sea in the Study Area....                    | 3-27 |
| Table 3.0-16: Annual Number of Explosive Munitions Expended At Sea in the Study Area .....                            | 3-28 |
| Table 3.0-17: Annual Number of Targets Expended At Sea in the Study Area .....  | 3-28 |
| Table 3.0-18: Annual Number of Events in the Study Area Including Seafloor Devices .....                              | 3-29 |
| Table 3.0-19: Annual Number of Munitions Used on Farallon de Medinilla .....  | 3-29 |
| Table 3.0-20: Annual Number of Wires and Cables Expended in the Study Area .....                                      | 3-31 |
| Table 3.0-21: Size Categories for Decelerators/Parachutes Expended During Training and Testing<br>Activities.....     | 3-31 |
| Table 3.0-22: Annual Number of Decelerators/Parachutes Expended in the Study Area.....                                | 3-33 |
| Table 3.0-23: Annual Number of Chaff Cartridges Expended in the Study Area .....                                      | 3-33 |
| Table 3.0-24: Annual Number of Flares Expended in the Study Area .....  | 3-33 |

## 3 Affected Environment and Environmental Consequences

This chapter describes the United States (U.S.) Department of the Navy's (Navy's) approach to analysis, existing environmental conditions in the Mariana Islands Training and Testing (MITT) Study Area, as well as the analysis of resources potentially impacted by the Proposed Action described in Chapter 2 (Description of Proposed Action and Alternatives). The Study Area is described in Section 2.1 (Description of the Mariana Islands Training and Testing Study Area) and depicted in Figure 2.1-1.

### 3.0 Introduction

In May 2015, the Navy released the MITT Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) (U.S. Department of the Navy, 2015a), hereafter referred to as the 2015 MITT Final EIS/OEIS, for which a Record of Decision was released (U.S. Department of the Navy, 2015b). The Navy applied the Navy Acoustics Effects Model for the 2015 MITT Final EIS/OEIS to quantitatively analyze potential acoustic effects from Navy training and testing activities. For this Supplemental EIS (SEIS)/OEIS, the Navy refined the Navy Acoustics Effects Model (U.S. Department of the Navy, 2018b) and updated marine mammal density estimates (U.S. Department of the Navy, 2018a), as well as the acoustic criteria and activity data inputs used in the acoustic model (U.S. Department of the Navy, 2017c).

The following subsections are included in Section 3.0:

- Section 3.0.1 (Overall Approach to Analysis) identifies the methodology used in this SEIS/OEIS to assess resource impacts associated with the Proposed Action.
- Section 3.0.2 (Regulatory Framework) presents the regulatory framework on which this SEIS/OEIS is based. It identifies applicable laws, regulations, executive orders (EOs), and directives used to develop the analyses.
- Section 3.0.3 (Resources and Issues Not Carried Forward for More Detailed Discussion) identifies the resources that were eliminated from further consideration in this SEIS/OEIS.
- Section 3.0.4 (Identification of Stressors for Analysis) discusses the stressors used in the analysis of impacts on resources

#### 3.0.1 Overall Approach to Analysis

The methods used in this SEIS/OEIS to assess resource impacts associated with the Proposed Action include the procedural steps outlined below:

- Review the existing 2015 MITT Final EIS/OEIS and Record of Decision.
- Determine if the affected environment has changed.
- Identify new activities and proposed changes to existing activities.
- Identify the stressors associated with the updated list of activities.
- Review existing and identify new federal and state regulations and standards relevant to resource-specific management or protection and determine if there has been any change since the 2015 MITT Final EIS/OEIS.
- Review and apply new literature, including science, surveys, and information on how resources could be affected by stressors.
- Determine if there is a new method of analysis for those activities.
- Review and consider comments received from members of the public and other stakeholders during the scoping period.

- Identify past, present, and reasonably foreseeable future actions to analyze the cumulative impacts.
- Consider mitigation measures to reduce identified potential impacts.

### **3.0.1.1 Navy Compiled and Generated Data**

While preparing this document, the Navy used the best available data, science, and information accepted by the relevant and appropriate regulatory and scientific communities to establish a baseline in the environmental analyses for all resources in accordance with the National Environmental Policy Act (NEPA), the Administrative Procedure Act (5 United States Code sections 551–596), and EO 12114.

In support of the environmental baseline and environmental consequences sections for this and other environmental documents, the Navy has sponsored and supported both internal and independent research and monitoring efforts. The Navy’s research and monitoring programs, as described below, are largely focused on filling data gaps and obtaining the most up-to-date science.

#### **3.0.1.1.1 Marine Species Monitoring and Research Programs**

The Navy has been conducting marine species monitoring for compliance with the Marine Mammal Protection Act (MMPA) and Endangered Species Act (ESA) since 2006, both in association with training and testing events and independently. In addition to monitoring activities associated with regulatory compliance, two other U.S. Navy research programs provide extensive investments in basic and applied research: the Office of Naval Research Marine Mammals & Biology program, and the Living Marine Resources program. In fact, the U.S. Navy is one of the largest sources of funding for marine mammal research in the world. A survey of federally funded marine mammal research and conservation conducted by the Marine Mammal Commission found that the Navy was the second-largest source of funding for marine mammal activities (direct project expenditures, as well as associated indirect or support costs) in the United States in 2014, second only to National Oceanic and Atmospheric Administration Fisheries (Purdy, 2016).

The monitoring program has historically focused on collecting baseline data that supports analysis of marine mammal occurrence, distribution, abundance, and habitat use preferences in and around ocean areas in the Atlantic and Pacific where the Navy conducts training and testing. More recently, the priority has begun to shift towards assessing the potential response of individual species to training and testing activities. Data collected through the monitoring program serves to inform the analysis of impacts on marine mammals with respect to species distribution, habitat use, and potential responses to training and testing activities. Monitoring is performed using various methods, including visual surveys from surface vessels and aircraft, passive acoustics, and tagging. Additional information on the program is available on the U.S. Navy Marine Species Monitoring Program website, which serves as a public online portal for information on the background, history, and progress of the program and also provides access to reports, documentation, data, and updates on current monitoring projects and initiatives.

The two other Navy programs previously mentioned invest in research on the potential effects of sound on marine species and develop scientific information and analytic tools that support preparation of environmental impact statements and associated regulatory processes under the MMPA and ESA, as well as support development of improved monitoring and detection technology and advance overall knowledge about marine species. These programs support coordinated science, technology, research, and development focused on understanding the effects of sound on marine mammals and other marine

species, including physiological, behavioral, ecological, and population-level effects. Additional information on these programs and other ocean resources-oriented initiatives can be found at the Department of the Navy – Energy, Environment and Climate Change website.

### **3.0.1.2 Navy’s Quantitative Analysis to Determine Impacts on Sea Turtles and Marine Mammals**

If proposed Navy activities introduce sound or explosive energy into the marine environment, an analysis of potential impacts on marine species is conducted. Data on the density of animals (number of animals per unit area) of each species and stock is needed, along with criteria and thresholds defining the levels of sound and energy that may cause certain types of impacts. The Navy’s acoustic effects model takes the density and the criteria and thresholds as inputs and analyzes Navy training and testing activities. Finally, mitigation and animal avoidance behaviors are considered to determine the number of impacts that could occur. The inputs and process are described below. A detailed explanation of this analysis is provided in the technical report titled *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018b).

#### **3.0.1.2.1 Marine Species Density Database**

A quantitative analysis of impacts on a species requires data on their abundance and distribution in the potentially impacted area. The most appropriate metric for this type of analysis is density, which is the number of animals present per unit area. Estimating marine species density requires substantial surveys and effort to collect and analyze data to produce a usable estimate. The National Marine Fisheries Service (NMFS) is the primary agency responsible for estimating marine mammal and sea turtle density within the U.S. Exclusive Economic Zone. Other agencies and independent researchers often publish density data for species in specific areas of interest, including areas outside the U.S. Exclusive Economic Zone. In areas where surveys have not produced adequate data to allow robust density estimates, methods such as model extrapolation from surveyed areas, Relative Environmental Suitability models, or expert opinion are used to estimate occurrence. These density estimation methods rely on information such as animal sightings from adjacent locations, amount of survey effort, and the associated environmental variables (e.g., depth, sea surface temperature).

There is no single source of density data for every area of the world, species, and season because of the fiscal limitations, resources, effort involved in providing survey coverage to sufficiently estimate density, and practical limitations. Therefore, to characterize marine species density for large areas, such as the MITT Study Area, the Navy compiled data from multiple sources and developed a protocol to select the best available density estimates based on species, area, and time (i.e., season). When multiple data sources were available, the Navy ranked density estimates based on a hierarchical approach to ensure that the most accurate estimates were selected. The highest tier included peer-reviewed published studies of density estimates from spatial models, since these provide spatially explicit density estimates with relatively low uncertainty. Other preferred sources included peer-reviewed published studies of density estimates derived from systematic line-transect survey data, the method typically used for the NMFS marine mammal stock assessment reports. In the absence of survey data, information on species occurrence and known or inferred habitat associations have been used to predict densities using model-based approaches, including Relative Environmental Suitability models. Because these estimates inherently include a high degree of uncertainty, they were considered the least preferred data source. In cases where a preferred data source was not available, density estimates were selected based on expert opinion from scientists.

The resulting Geographic Information System database includes seasonal density values for every marine mammal and sea turtle species present within the Study Area. This database is described in the technical report titled *U.S. Navy Marine Species Density Database Phase III for the Mariana Islands Training and Testing Study Area* (U.S. Department of the Navy, 2018a), hereafter referred to as the Density Technical Report. These data were used as an input into the Navy Acoustic Effects Model.

The Density Technical Report describes the models that were utilized in detail and provides detailed explanations of the models applied to each species density estimate. The below list describes models in order of preference.

1. Spatial density models are preferred and used when available because they provide an estimate with the least amount of uncertainty by deriving estimates for divided segments of the sampling area. These models (see Becker et al., 2016; Forney et al., 2015) predict spatial variability of animal presence as a function of habitat variables (e.g., sea surface temperature, seafloor depth). This model is developed for areas, species, and, when available, specific timeframes (months or seasons) with sufficient survey data.
2. Stratified design-based density estimates use line-transect survey data with the sampling area divided (stratified) into sub-regions, and a density is predicted for each sub-region (see Barlow, 2016; Becker et al., 2016; Bradford et al., 2017; Campbell et al., 2015; Jefferson et al., 2014). While geographically stratified density estimates provide a good indication of a species' distribution within the Study Area, the uncertainty is typically high because each sub-region estimate is based on a smaller stratified segment of the overall survey effort.
3. Design-based density estimations use line-transect survey data from land and aerial surveys designed to cover a specific geographic area (see Carretta et al., 2015). These estimates use the same survey data as stratified design-based estimates, but they are not segmented into sub-regions and instead provide one estimate for a large surveyed area.
4. Although relative environmental suitability models provide estimates for areas of the oceans that have not been surveyed, using information on species occurrence and inferred habitat associations, and have been used in past density databases, these models were not used in the current quantitative analysis.

When interpreting the results of the quantitative analysis, as described in the Density Technical Report, it is important to consider that "each model is limited to the variables and assumptions considered by the original data source provider. No mathematical model representation of any biological population is perfect, and with regards to marine mammal biodiversity, any single model will not completely explain the results" (U.S. Department of the Navy, 2017b). These factors and others described in the Density Technical Report should be considered when examining the estimated impact numbers in comparison to current population abundance information for any given species or stock.

#### **3.0.1.2.2 Developing Acoustic and Explosive Criteria and Thresholds**

Information about the numerical sound and energy levels that are likely to elicit certain types of physiological and behavioral reactions is needed to analyze potential impacts on marine species. Revised Phase III criteria and thresholds for quantitative modeling of impacts use the best available existing data from scientific journals, technical reports, and monitoring reports to develop thresholds and functions for estimating impacts on marine species. Working with NMFS, the Navy has developed updated criteria for marine mammals and sea turtles. Criteria for estimating impacts on marine fishes are also used in this analysis, which largely follows the *Sound Exposure Guidelines for Fishes and Sea Turtles* (Popper et al., 2014).

Since the release of the *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effect Analysis* in 2012 (U.S. Department of the Navy, 2012b), recent and emerging science has necessitated an update to these criteria and thresholds for assessing potential impacts on marine mammals and sea turtles. A detailed description of the Phase III acoustic and explosive criteria and threshold development is included in the supporting technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Impact to Marine Mammals and Sea Turtles* (U.S. Department of the Navy, 2017c), and details are provided in each resource section. A series of behavioral studies, largely funded by the U.S. Navy, has led to a new understanding of how some species of marine mammals react to military sonar. This understanding resulted in developing new behavioral response functions for estimating alterations in behavior. Additional information on auditory weighting functions has also emerged [e.g., (Mulsow et al., 2015)], leading to the development of a new methodology to predict auditory weighting functions for each hearing group along with the accompanying hearing loss thresholds. These criteria for predicting hearing loss in marine mammals were largely adopted by NMFS for species within their purview (National Marine Fisheries Service, 2016).

The Navy also uses criteria for estimating effects to fishes and the ranges to which those effects are likely to occur. A working group of experts generated a technical report that provides numerical criteria and relative likelihood of effects to fish within different hearing groups (i.e., fishes with no swim bladder versus fishes with a swim bladder involved in hearing) (Popper et al., 2014). Where applicable, thresholds and relative risk factors presented in the technical report were used to assist in the analysis of effects to fishes from Navy activities. Details on criteria used to estimate impacts on marine fishes are contained within the appropriate stressor section (e.g., sonar and other transducers, explosives) within Section 3.9 (Fishes). This panel of experts also estimated parametric criteria for the effects of sea turtle exposure to sources located at "near," "intermediate," and "far" distances, assigning "low," "medium," and "high" probability to specific categories of behavioral impacts (Popper et al., 2014).

#### **3.0.1.2.3 The Navy's Acoustic Effects Model**

The Navy's Acoustic Effects Model calculates sound energy propagation from sonar and other transducers, air guns, and explosives during naval activities and the energy or sound received by animat dosimeters. Animat dosimeters are virtual representations of marine mammals or sea turtles distributed in the area around the modeled naval activity; each animat records its individual sound "dose." The model bases the distribution of animats over the Study Area on the density values in the Navy Marine Species Density Database and distributes animats in the water column proportional to the known time that species spend at varying depths.

The model accounts for environmental variability of sound propagation in both distance and depth when computing the received sound level on the animats. The model conducts a statistical analysis based on multiple model runs to compute the estimated effects on animals. The number of animats that exceed the received threshold for an effect is tallied to provide an estimate of the number of marine mammals or sea turtles that could be affected.

Assumptions in the Navy model intentionally err on the side of overestimation when there are unknowns:

- Naval activities are modeled as though they would occur regardless of proximity to marine mammals or sea turtles (i.e., mitigation and implementation of standard operating procedures that employ protective measures are not modeled) and without any avoidance of the activity by the animal. The final step of the quantitative analysis of acoustic effects is to consider the

implementation of mitigation. For sonar and other transducers, the possibility that marine mammals or sea turtles would avoid continued or repeated sound exposures is also considered.

- Many explosions from munitions such as bombs and missiles actually occur upon impact with above-water targets and at the water's surface. However, for this analysis, sources such as these were modeled as exploding underwater. This modeling overestimates the amount of explosive and acoustic energy entering the water.

The model estimates the impacts caused by individual training and testing activities. During any individual modeled event, impacts on individual animals are considered over 24-hour periods. The animals do not represent actual animals, but rather allow for a statistical analysis of the number of instances that marine mammals or sea turtles may be exposed to sound levels resulting in an effect. Therefore, the model estimates the number of instances in which an effect threshold was exceeded over the course of a year, but it does not estimate the number of individual marine mammals or sea turtles that may be impacted over a year (i.e., some marine mammals or sea turtles could be impacted several times, while others would not experience any impact). A detailed explanation of the Navy's Acoustic Effects Model is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018b).

#### **3.0.1.2.4 Accounting for Mitigation**

##### **3.0.1.2.4.1 Sonar and Other Transducers**

The Navy implements mitigation measures (described in Section 5.3.2, Acoustic Stressors), including the power-down or shut-down (i.e., power off) of sonar when a marine mammal or sea turtle is observed in the mitigation zone, during activities that use sonar and other transducers. The mitigation zones encompass the estimated ranges to injury (including permanent threshold shift [PTS]) for a given sonar exposure. Therefore, the impact analysis quantifies the potential for mitigation to reduce the risk of PTS. Two factors are considered when quantifying the effectiveness of mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., active sonar) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018b).

In the quantitative analysis, consideration of mitigation measures means that, for activities where mitigation is feasible, some model-estimated PTS is considered mitigated to the level of temporary threshold shift (TTS). The quantitative analysis does not analyze the potential for mitigation to reduce TTS or behavioral effects, even though mitigation could also reduce the likelihood of these effects. In practice, mitigation also protects all unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

The ability to observe the range to PTS was estimated for each training or testing event. The ability of Navy Lookouts to detect marine mammals or sea turtles in or approaching the mitigation zone is dependent on the animal's presence at the surface and the characteristics of the animal that influence

its sightability (such as group size or surface active behavior). The behaviors and characteristics of some species may make them easier to detect. For example, based on small boat surveys between 2000 and 2012 in the Hawaiian Islands, pantropical spotted dolphins and striped dolphins were frequently observed leaping out of the water, and Cuvier's beaked whales (Baird, 2013) and Blainville's beaked whales (HDR, 2012) were occasionally observed breaching. These behaviors are visible from a great distance and likely increase sighting distances and detections of these species. Environmental conditions under which the training or testing activity could take place are also considered, such as the sea surface conditions, weather (e.g., fog or rain), and day versus night.

#### **3.0.1.2.4.2 Explosions**

The Navy implements mitigation measures (described in Section 5.3.3, Explosive Stressors) during explosive activities, including delaying detonations when a marine mammal or sea turtle is observed in the mitigation zone. The mitigation zones encompass the estimated ranges to mortality for a given explosive. Therefore, the impact analysis quantifies the potential for mitigation to reduce the risk of mortality due to exposure to explosives. Two factors are considered when quantifying the effectiveness of mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., gunnery exercise) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018b).

In the quantitative analysis, consideration of mitigation measures means that, for activities where mitigation is feasible, model-estimated mortality is considered mitigated to the level of injury. The impact analysis does not analyze the potential for mitigation to reduce non-auditory injury, PTS, TTS, or behavioral effects, even though mitigation would also reduce the likelihood of these effects. In practice, mitigation also protects all unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

#### **3.0.1.2.5 Marine Mammal Avoidance of Sonar and other Transducers**

Because a marine mammal is assumed to initiate avoidance behavior (tens of meters away for most species groups) after an initial startle reaction when exposed to relatively high received levels of sound, a marine mammal could reduce its cumulative sound energy exposure over a sonar event with multiple pings. This would reduce risk of both PTS and TTS, although the quantitative analysis conservatively only considers the potential to reduce instances of PTS by accounting for marine mammals swimming away to avoid repeated high-level sound exposures. All reductions in PTS impacts from likely avoidance behaviors are instead considered TTS impacts.

### **3.0.2 Regulatory Framework**

In accordance with the Council on Environmental Quality regulations that implement the requirements of the NEPA, other planning and environmental review procedures are integrated in this SEIS/OEIS to the fullest extent possible. Chapter 6 (Additional Regulatory Considerations) provides a status of compliance with the applicable environmental laws, regulations, and EOs that were considered in preparing this SEIS/OEIS (including those that may be secondary considerations in the resource evaluations).

The federal statutes and EOs considered in this SEIS/OEIS that were described in Section 3.0.1 (Regulatory Framework) of the 2015 MITT Final EIS/OEIS have not changed.

**3.0.3 Resources and Issues Not Carried Forward for More Detailed Discussion**

Considerations under EO 13045, *Protection of Children from Environmental Health Risks and Safety Risks*, were eliminated from further analysis because all of the proposed activities occur in the ocean where there are no child populations present. Therefore, the Proposed Action would not lead to disproportionate environmental health risks or safety risks to children.

**3.0.4 Identification of Stressors for Analysis**

Some of the stressors identified for consideration in this SEIS/OEIS in the analysis of resources have been refined from those considered in the 2015 MITT Final EIS/OEIS. The list of stressors analyzed in this SEIS/OEIS and changes from the 2015 MITT Final EIS/OEIS are shown in Table 3.0-1. Although the names of some stressors have changed, the analysis conducted on that stressor did not change. Where useful, an explanation of the change is provided in italics.

**Table 3.0-1: Comparison of Stressors Analyzed**

| <i>2015 MITT Final EIS/OEIS</i>   | <i>Supplemental EIS/OEIS</i>   |
|---|--|
| <b>Components and Stressors for Physical Resources</b>  |  |
| <b><i>Sediments and Water Quality Stressors</i></b>   |  |
| <ul style="list-style-type: none"> <li>• Explosives and explosive byproducts</li> <li>• Metals</li> <li>• Chemicals other than explosives</li> <li>• Other materials</li> </ul>   | <ul style="list-style-type: none"> <li>• Explosives</li> <li>• Metals</li> <li>• Chemicals</li> <li>• Other materials</li> </ul>   |
| <b><i>Air Quality Stressors</i></b>   |  |
| <ul style="list-style-type: none"> <li>• Criteria pollutants</li> <li>• Hazardous air pollutants</li> </ul>   | <ul style="list-style-type: none"> <li>• Criteria pollutants</li> <li>• Hazardous air pollutants</li> </ul>  |
| <b>Components and Stressors for Biological Resources</b>  |  |
| <b><i>Acoustic Stressors</i></b>  |  |
| <ul style="list-style-type: none"> <li>• Sonar and other active acoustic sources</li> <li>• Vessel noise</li> <li>• Aircraft noise</li> <li>• Weapons firing, launch, and impact noise</li> <li>• Underwater explosives</li> <li>• Swimmer defense airguns</li> </ul> | <ul style="list-style-type: none"> <li>• Sonar and other transducers</li> <li>• Vessel noise</li> <li>• Aircraft noise</li> <li>• Weapons noise</li> <li>• (<i>“Underwater explosives” is moved to next category of “In-water explosions”</i>)</li> <li>• (<i>Swimmer defense airguns are not proposed or analyzed in this SEIS/OEIS</i>)</li> </ul> |
| <b><i>Explosive Stressors</i></b>   |  |
| ( <i>In the 2015 MITT Final EIS/OEIS, Explosives were included under Acoustic Stressors</i> )   | <ul style="list-style-type: none"> <li>• In-water explosions</li> <li>• In-air explosions</li> </ul>   |
| <b><i>Energy Stressors</i></b>  |  |
| <ul style="list-style-type: none"> <li>• Electromagnetic devices</li> <li>• Lasers</li> </ul>   | <ul style="list-style-type: none"> <li>• In-air electromagnetic devices (<i>included under Electromagnetic Devices</i>)</li> <li>• In-water electromagnetic devices (<i>included under Electromagnetic Devices</i>)</li> <li>• Lasers</li> </ul>   |

**Table 3.0-1: Comparison of Stressors Analyzed (continued)**

| <i>2015 MITT Final EIS/OEIS</i>  | <i>Supplemental EIS/OEIS</i>  |
|--|---|
| <b>Components and Stressors for Physical Resources</b>   |   |
| <b><i>Physical Disturbance and Strike Stressors</i></b>  |   |
| <ul style="list-style-type: none"> <li>• Aircraft and aerial targets</li> <li>• Vessels</li> <li>• In-water devices</li> <li>• Military expended materials</li> <li>• Seafloor devices</li> <li>• Ground disturbance</li> <li>• Wildfires</li> </ul> | <ul style="list-style-type: none"> <li>• Aircraft and aerial targets</li> <li>• Vessels and in-water devices</li> <li>• Military expended materials</li> <li>• Seafloor devices</li> <li>• Ground disturbance (FDM only)</li> <li>• Wildfires (FDM only)</li> </ul> |
| <b><i>Entanglement Stressors</i></b>   |   |
| <ul style="list-style-type: none"> <li>• Fiber optic cables and guidance wires</li> <li>• Decelerators/parachutes</li> </ul>   | <ul style="list-style-type: none"> <li>• Wires and cables</li> <li>• Decelerators/parachutes</li> </ul>   |
| <b><i>Ingestion Stressors</i></b>  |   |
| <ul style="list-style-type: none"> <li>• Military expended materials from munitions</li> <li>• Military expended materials other than munitions</li> </ul>   | <ul style="list-style-type: none"> <li>• Military expended materials from munitions</li> <li>• Military expended materials other than munitions</li> </ul>  |
| <b><i>Secondary Stressors</i></b>  |   |
| <ul style="list-style-type: none"> <li>• Habitat</li> <li>• Prey availability</li> </ul>   | <ul style="list-style-type: none"> <li>• Impacts on habitat</li> <li>• Invasive species introductions into terrestrial habitats (FDM only)</li> <li>• Impacts on prey availability</li> </ul>   |
| <b>Components and Stressors for Human Resources</b>  |   |
| <b><i>Cultural Resources Stressors</i></b>   |   |
| <ul style="list-style-type: none"> <li>• Acoustic</li> <li>• Physical Disturbance and Strike</li> </ul>  | <ul style="list-style-type: none"> <li>• Explosives (<i>previously referred to as Acoustic</i>)</li> <li>• Physical Disturbance and Strike</li> </ul>   |
| <b><i>Socioeconomic Resources Stressors</i></b>  |   |
| <ul style="list-style-type: none"> <li>• Accessibility</li> <li>• Airborne acoustics</li> <li>• Physical disturbance and strike</li> <li>• Secondary impacts from availability of resources</li> </ul>   | <ul style="list-style-type: none"> <li>• Accessibility</li> <li>• Airborne acoustics</li> <li>• Physical disturbance and strike</li> <li>• Secondary impacts from availability of resources</li> </ul>  |
| <b><i>Public Health and Safety Stressors</i></b>   |   |
| <ul style="list-style-type: none"> <li>• Underwater energy</li> <li>• In-air energy</li> <li>• Physical interactions</li> <li>• Secondary stressors (sediments and water quality)</li> </ul>   | <ul style="list-style-type: none"> <li>• Underwater energy</li> <li>• In-air energy</li> <li>• Physical interactions</li> <li>• Secondary stressors (sediments and water quality)</li> </ul>  |

Notes: (1) *Italics* reflect changes in stressors/stressor analysis in this SEIS/OEIS as compared to 2015 MITT Final EIS/OEIS; (2) FDM = Farallon de Medinilla

**3.0.4.1 Acoustic Stressors**

This section describes the characteristics of sounds produced during naval training and testing and the relative magnitude and location of these sound-producing activities. This section provides the basis for analysis of acoustic impacts on resources in the remainder of Chapter 3 (Affected Environment and Environmental Consequences). Explanations of the terminology and metrics used when describing sound in this SEIS/OEIS are in Appendix H (Acoustic and Explosive Concepts).

Acoustic stressors include acoustic signals emitted into the water from a specific source such as sonar and other transducers (devices that convert energy from one form to another – in this case, to sound waves), as well as incidental sources of broadband sound produced as a byproduct of vessel movement, aircraft transits, and use of weapons or other deployed objects. Explosives also produce broadband sound but are characterized separately from other acoustic sources due to their unique hazardous characteristics (Section 3.0.4.2, Explosive Stressors). Characteristics of each of these sound sources are described in the following sections.

In order to better organize and facilitate the analysis of approximately 300 sources of underwater sound used for testing and training by the Navy including sonars, other transducers, and explosives, a series of source classifications, or source bins, were developed. The source classification bins do not include the broadband sounds produced incidental to vessel and aircraft transits and weapons firing.

The use of source classification bins provides the following benefits:

- Provides the ability for new sensors or munitions to be covered under existing authorizations, as long as those sources fall within the parameters of a “bin.”
- Improves efficiency of source utilization data collection and reporting requirements anticipated under the MMPA authorizations.
- Ensures a conservative approach to all impact estimates, as all sources within a given class are modeled as the most impactful source (highest source level, longest duty cycle [i.e., the proportion of time signals are emitted in a given period of time], or largest net explosive weight) within that bin.
- Allows analyses to be conducted in a more efficient manner, without any compromise of analytical results.
- Provides a framework to support the reallocation of source usage (hours/explosives) between different source bins, as long as the total numbers of takes remain within the overall analyzed and authorized limits. This flexibility is required to support evolving Navy training and testing requirements, which are linked to military missions and combat operations.

#### **3.0.4.1.1 Sonar and Other Transducers**

Active sonar and other transducers emit non-impulsive sound waves into the water to detect objects, safely navigate, and communicate. Passive sonars differ from active sound sources in that they do not emit acoustic signals; rather, they only receive acoustic information about the environment, or listen. In this SEIS/OEIS, the terms sonar and other transducers will be used to indicate active sound sources unless otherwise specified.

The Navy employs a variety of sonars and other transducers to obtain and transmit information about the undersea environment. Some examples are mid-frequency hull-mounted sonars used to find and track potential enemy submarines, high-frequency small object detection sonars used to detect mines, high-frequency underwater modems used to transfer data over short ranges, and extremely high-frequency (greater than 200 kilohertz [kHz]) Doppler sonars used for navigation, like those used on commercial and private vessels. The characteristics of these sonars and other transducers, such as source level, beam width, directivity, and frequency, depend on the purpose of the source. Higher frequencies can carry or provide more information about objects off which they reflect, but attenuate more rapidly. Lower frequencies attenuate less rapidly, so may detect objects over a longer distance, but with less detail.

Propagation of sound produced underwater is highly dependent on environmental characteristics such as bathymetry, bottom type, water depth, temperature, and salinity. The sound received at a particular location will be different than near the source due to the interaction of many factors, including propagation loss; how the sound is reflected, refracted, or scattered; the potential for reverberation; and interference due to multi-path propagation. In addition, absorption greatly affects the distance over which higher frequency sounds propagate. The effects of these factors are explained in Appendix H (Acoustic and Explosive Concepts). Because of the complexity of analyzing sound propagation in the ocean environment, the Navy relies on acoustic models in its environmental analyses that consider sound source characteristics and varying ocean conditions across the Study Area.

The sound sources and platforms typically used in naval activities analyzed in the SEIS/OEIS are described in Appendix A (Training and Testing Activities Descriptions). Sonars and other transducers used to obtain and transmit information underwater during Navy training and testing activities generally fall into several categories of use described below.

#### **3.0.4.1.1.1 Anti-Submarine Warfare**

Sonar used during anti-submarine warfare would impart the greatest amount of acoustic energy of any category of sonar and other transducers analyzed in this SEIS/OEIS. Types of sonars used to detect potential enemy vessels include hull-mounted, towed, line array, sonobuoy, helicopter dipping, and torpedo sonars. In addition, acoustic targets and decoys (countermeasures) may be deployed to emulate the sound signatures of vessels or repeat received signals.

Most anti-submarine warfare sonars are mid-frequency (1–10 kHz) because mid-frequency sound balances sufficient resolution to identify targets with distance over which threats can be identified. However, some sources may use higher or lower frequencies. Duty cycles can vary widely, from rarely used to continuously active. Anti-submarine warfare sonars can be wide-angle in a search mode or highly directional in a track mode.

Most anti-submarine warfare activities involving submarines or submarine targets would occur in waters greater than 600 feet (ft.) deep due to safety concerns about running aground at shallower depths. Sonars used for anti-submarine warfare activities would typically be used beyond 12 nautical miles (NM) from shore. Exceptions include use of dipping sonar by helicopters, maintenance of systems while in port, and system checks while transiting to or from port.

#### **3.0.4.1.1.2 Mine Warfare, Small Object Detection, and Imaging**

Sonars used to locate mines and other small objects, as well as those used in imaging (e.g., for hull inspections or imaging of the seafloor), are typically high frequency or very high-frequency. Higher frequencies allow for greater resolution and, due to their greater attenuation, are most effective over shorter distances. Mine detection sonar can be deployed (towed or vessel hull-mounted) at variable depths on moving platforms (ships, helicopters, or unmanned vehicles) to sweep a suspected mined area. Hull-mounted anti-submarine sonars can also be used in an object detection mode known as “Kingfisher” mode. Sonars used for imaging are usually used in close proximity to the area of interest, such as pointing downward near the seafloor.

Mine detection sonar use would be concentrated in areas where practice mines are deployed, typically in water depths less than 200 ft., and at established training minefields, temporary minefields close to strategic ports and harbors, or at targets of opportunity such as navigation buoys. Kingfisher mode on

vessels is most likely to be used when transiting to and from port. Sound sources used for imaging could be used throughout the Study Area.

#### **3.0.4.1.1.3 Navigation and Safety**

Similar to commercial and private vessels, Navy vessels employ navigational acoustic devices including speed logs, Doppler sonars for ship positioning, and fathometers. These may be in use at any time for safe vessel operation. These sources are typically highly directional to obtain specific navigational data.

#### **3.0.4.1.1.4 Communication**

Sound sources used to transmit data (such as underwater modems), provide location (pingers), or send a single brief release signal to bottom-mounted devices (acoustic release) may be used throughout the Study Area. These sources typically have low duty cycles and are usually only used when it is desirable to send a detectable acoustic message.

#### **3.0.4.1.1.5 Classification of Sonar and Other Transducers**

Sonars and other transducers are grouped into classes that share an attribute, such as frequency range or purpose of use. As detailed below, classes are further sorted by bins based on the frequency or bandwidth; source level; and, when warranted, the application in which the source would be used. Unless stated otherwise, a reference distance of 1 meter is used for sonar and other transducers.

- Frequency of the non-impulsive acoustic source:
  - Low-frequency sources operate below 1 kHz.
  - Mid-frequency sources operate at and above 1 kHz, up to and including 10 kHz.
  - High-frequency sources operate above 10 kHz, up to and including 100 kHz.
  - Very high-frequency sources operate above 100 kHz but below 200 kHz.
- Sound pressure level:
  - Greater than 160 decibels (dB) referenced to (re) 1 micropascal (dB re 1  $\mu$ Pa), but less than 180 dB re 1  $\mu$ Pa
  - Equal to 180 dB re 1  $\mu$ Pa at 1 m and up to 200 dB re 1  $\mu$ Pa
  - Greater than 200 dB re 1  $\mu$ Pa
- Application in which the source would be used:
  - Sources with similar functions that have similar characteristics, such as pulse length (duration of each pulse), beam pattern, and duty cycle

The bins used for classifying active sonars and transducers that are quantitatively analyzed in the Study Area are shown in Table 3.0-2. While general parameters or source characteristics are shown in the table, actual source parameters are classified.

Table 3.0-2 also shows the bin use that could occur in any year under each action alternative for training and testing activities and Phase II amounts are included for comparison.

**Table 3.0-2: Sonar and Transducer Sources Quantitatively Analyzed**

| Source Class Category   | Bin   | Unit* | Training & Testing  |               |               |
|---|-------|-------|---------------------|---------------|---------------|
|   |       |       | 2015 Final EIS/OEIS | Alternative 1 | Alternative 2 |
| <b>Low-Frequency (LF):</b> Sources that produce signals less than 1 kHz   | LF4   | H     | 123                 | 1             | 1             |
|   | LF5   | H     | 11                  | 10            | 10            |
|   | LF6   | H     | 40                  | 0             | 0             |
| <b>Mid-Frequency (MF):</b> Tactical and non-tactical sources that produce signals between 1 and 10 kHz  | MF1   | H     | 1,872               | 1,729         | 1,818         |
|   | MF1K  | H     | 0                   | 3             | 3             |
|   | MF2   | H     | 625                 | 0             | 0             |
|   | MF3   | H     | 192                 | 189           | 228           |
|   | MF4   | H     | 214                 | 172           | 185           |
|   | MF5   | C     | 2,588               | 2,024         | 2,094         |
|   | MF6   | C     | 33                  | 62            | 74            |
|   | MF8   | H     | 123                 | 0             | 0             |
|   | MF9   | H     | 47                  | 15            | 29            |
|   | MF10  | H     | 231                 | 0             | 0             |
|   | MF11  | H     | 324                 | 292           | 304           |
|   | MF12  | H     | 656                 | 608           | 616           |
| <b>High-Frequency (HF):</b> Tactical and non-tactical sources that produce signals between 10 and 100 kHz   | HF1   | H     | 113                 | 63            | 73            |
|   | HF3   | H     | 0                   | 4             | 4             |
|   | HF4   | H     | 1,060               | 1,472         | 1,472         |
|   | HF5   | H     | 336                 | 0             | 0             |
|   | HF6   | H     | 1,173               | 163           | 309           |
| <b>Anti-Submarine Warfare (ASW):</b> Tactical sources (e.g., active sonobuoys and acoustic countermeasures systems) used during ASW training and testing activities | ASW1  | H     | 144                 | 192           | 192           |
|   | ASW2  | C     | 660                 | 538           | 554           |
|   | ASW3  | H     | 3,935               | 3,024         | 3,124         |
|   | ASW4  | C     | 11                  | 268           | 332           |
|   | ASW5  | H     | 0                   | 50            | 50            |
| <b>Torpedoes (TORP):</b> Source classes associated with the active acoustic signals produced by torpedoes   | TORP1 | C     | 115                 | 62            | 71            |
|   | TORP2 | C     | 62                  | 40            | 62            |
|   | TORP3 | C     | 0                   | 6             | 6             |
| <b>Forward Looking Sonar (FLS):</b> Forward or upward looking object avoidance sonars used for ship navigation and safety   | FLS2  | H     | 0                   | 4             | 4             |
| <b>Acoustic Modems (M):</b> Systems used to transmit data through the water   | M3    | H     | 112                 | 17            | 31            |

**Table 3.0-2: Sonar and Transducer Sources Quantitatively Analyzed (continued)**

| Source Class Category   | Bin  | Unit* | Training & Testing  |               |               |
|---|------|-------|---------------------|---------------|---------------|
|   |      |       | 2015 Final EIS/OEIS | Alternative 1 | Alternative 2 |
| <b>Swimmer Detection Sonar (SD):</b> Used to detect divers and submerged swimmers   | SD1  | H     | 2,341               | 0             | 0             |
| <b>Air Guns (AG):</b> Used during swimmer defense and diver deterrent training and testing activities   | AG   | C     | 308                 | 0             | 0             |
| <b>Synthetic Aperture Sonars (SAS):</b> Sonars in which active acoustic signals are post-processed to form high-resolution images of the seafloor | SAS2 | H     | 0                   | 449           | 449           |
|   | SAS4 | H     | 0                   | 6             | 6             |

\* H = hours; C = count (e.g., number of individual pings or individual sonobuoys)

There are in-water active acoustic sources with narrow beam widths, downward directed transmissions, short pulse lengths, frequencies above known hearing ranges, low source levels, or combinations of these factors, which are not anticipated to result in takes of protected species. These sources are categorized as *de minimis* sources and are qualitatively analyzed to determine the appropriate determinations under NEPA in the appropriate resource impact analyses, as well as under the MMPA and the ESA. When used during routine training and testing activities, and in a typical environment, *de minimis* sources fall into one or more of the following categories:

- Transmit primarily above 200 kHz: Sources above 200 kHz are above the hearing range of the most sensitive marine mammals and far above the hearing range of other protected species in the Study Area.
- Source levels of 160 dB re 1 μPa or less: Low-powered sources with source levels less than 160 dB re 1 μPa are typically hand-held sonars, range pingers, transponders, and acoustic communication devices. Assuming spherical spreading for a 160 dB re 1 μPa source, the sound will attenuate to less than 140 dB re 1 μPa within 10 m and less than 120 dB re 1 μPa within 100 m of the source. Ranges would be even shorter for a source less than 160 dB re 1 μPa source level.
- Acoustic source classes listed in Table 3.0-3: Sources with operational characteristics, such as short pulse length, narrow beam width, downward-directed beam, and low energy release, or manner of system operation, which exclude the possibility of any significant impact on a protected species (actual source parameters are classified). Even if there is a possibility that some species may be exposed to and detect some of these sources, any response is expected to be short-term and inconsequential.

**Table 3.0-3: Sonar and Transducers Qualitatively Analyzed**

| <i>Source Class Category</i>   | <i>Bin</i>  | <i>Characteristics</i>  |
|--|-------------|---|
| <b>Broadband Sound Sources (BB):</b><br>Sources with wide frequency spectra  | BB3         | <ul style="list-style-type: none"> <li>• very high frequency</li> <li>• very short pulse length</li> </ul>  |
|  | BB8         | <ul style="list-style-type: none"> <li>• small imploding source (light bulb)</li> </ul>   |
| <b>Doppler Sonar/Speed Logs (DS):</b><br>High-frequency/very high-frequency navigation transducers   | DS2–DS4     | <i>Required for safe navigation</i> <ul style="list-style-type: none"> <li>• downward focused</li> <li>• narrow beam width</li> <li>• very short pulse lengths</li> </ul>   |
| <b>Fathometers (FA):</b> High-frequency sources used to determine water depth  | FA1–FA4     | <i>Required for safe navigation</i> <ul style="list-style-type: none"> <li>• downward focused directly below the vessel</li> <li>• narrow beam width (typically much less than 30°)</li> <li>• short pulse lengths (less than 10 milliseconds)</li> </ul> |
| <b>Hand-Held Sonar (HHS):</b> High-frequency sonar devices used by Navy divers for object location   | HHS1        | <ul style="list-style-type: none"> <li>• very high frequency sound at low power levels</li> <li>• narrow beam width</li> <li>• short pulse lengths</li> <li>• under control of the diver (power and direction)</li> </ul>                                 |
| <b>Imaging Sonar (IMS):</b> Sonars with high or very high frequencies used to obtain images of objects underwater  | IMS1–IMS3   | <ul style="list-style-type: none"> <li>• High-frequency or very high-frequency</li> <li>• downward directed</li> <li>• narrow beam width</li> <li>• very short pulse lengths (typically 20 milliseconds)</li> </ul>                                       |
| <b>High-Frequency Acoustic Modems (M):</b> Systems that send data underwater<br><b>Tracking Pingers (P):</b> Devices that send a ping to identify an object location | M2<br>P1–P4 | <ul style="list-style-type: none"> <li>• low duty cycles (single pings in some cases)</li> <li>• short pulse lengths (typically 20 milliseconds)</li> <li>• low source levels</li> </ul>  |
| <b>Acoustic Releases (R):</b> Systems that ping to release a bottom-mounted object from its housing in order to retrieve the device at the surface                   | R1–R3       | <ul style="list-style-type: none"> <li>• typically emit only several pings to send release order</li> </ul>   |
| <b>Side-Scan Sonars (SSS):</b> Sonars that use active acoustic signals to produce high-resolution images of the seafloor   | SSS1–SSS2   | <ul style="list-style-type: none"> <li>• downward-directed beam</li> <li>• short pulse lengths (less than 20 milliseconds)</li> </ul>   |

Notes: ° = degree(s), kHz = kilohertz, lb. = pound(s)

### 3.0.4.1.2 Vessel Noise

Vessel noise, in particular commercial shipping, is a major contributor to underwater anthropogenic noise in the ocean within the Study Area. Naval vessels (e.g., ships and small craft) and civilian vessels (e.g., commercial ships, tugs, work boats, pleasure craft) produce low-frequency, broadband underwater sound, though the exact level of noise produced varies by vessel type. Frisk (2012) reported that between 1950 and 2007 ocean noise in the 25–50 Hertz (Hz) frequency range has increased 3.3 dB per decade, resulting in a cumulative increase of approximately 19 dB over a baseline of 52 dB. The increase in noise is associated with an increase in commercial shipping, which correlates with global economic growth (Frisk, 2012). Within the Study Area, Navy vessels represent a small amount of overall vessel traffic and an even smaller amount of overall vessel traffic noise (Mintz & Filadelfo, 2011; Mintz, 2012).

The Center for Naval Analyses conducted studies to determine traffic patterns of Navy and non-Navy vessels (Mintz & Parker, 2006; Mintz & Filadelfo, 2011; Mintz, 2012; Mintz, 2016). The most recent analysis covered the period 2011–2015 (Mintz, 2016) and included U.S. Navy surface ship traffic and non-military vessels such as cargo vessels, bulk carriers, commercial fishing vessels, oil tankers, passenger vessels, tugs, and research vessels. Caveats to this analysis include that only vessels over 65 ft. in length are reported, so smaller Navy vessels and civilian craft are not included, and vessel position records are much more frequent for Navy vessels than for commercial vessels. Therefore, the Navy is likely overrepresented in the data and the reported fraction of total energy is likely the upper limit of its contribution (Mintz & Filadelfo, 2011; Mintz, 2012).

Although the aforementioned studies did not include analysis of vessel traffic and associated vessel noise in the Study Area (the geographic scope was the continental United States and Hawaii), the conclusions of the studies are relevant to vessel noise in the Study Area. Overall, the contribution of Navy vessel traffic to broadband noise levels was relatively small compared with the contribution from commercial vessel traffic.

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.5, Vessel Noise) provides detailed information regarding vessel noise characteristics and production, and timing and duration of vessel activity.

#### **3.0.4.1.3 Aircraft Noise**

Fixed-wing, tiltrotor, and rotary-wing aircraft are used for a variety of training and testing activities throughout the Study Area, contributing both airborne and underwater sound to the ocean environment. Sounds in air are often measured using A-weighting, which adjusts received sound levels based on human hearing abilities (see Appendix H, Acoustic and Explosive Concepts). Aircraft used in training and testing generally have turboprop or jet engines. Motors, propellers, and rotors produce the most noise, with some noise contributed by aerodynamic turbulence. Aircraft sounds have more energy at lower frequencies. Aircraft may transit to or from vessels at sea throughout the Study Area from established airfields on land. The majority of aircraft noise would be generated at air stations, which are outside the Study Area. Takeoffs and landings occur at established airfields as well as on vessels at sea across the Study Area. Takeoffs and landings from Navy vessels produce in-water noise at a given location for a brief period as the aircraft climbs to cruising altitude. Military activities involving aircraft generally are dispersed over large expanses of open ocean but can be highly concentrated in time and location. Table 3.0-4 provides source levels for some typical aircraft used during training and testing in the Study Area and depicts comparable airborne source levels for the F-35A, EA-18G, and F/A-18C/D during takeoff.

##### **3.0.4.1.3.1 Underwater Transmission of Aircraft Noise**

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.6, Aircraft Overflight Noise) describes underwater transmission of aircraft noise. Since information regarding underwater transmission of aircraft noise has not changed, this SEIS/OEIS will not further analyze underwater transmission of aircraft noise.

##### **3.0.4.1.3.2 Helicopters**

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.6, Aircraft Overflight Noise) describes characteristics and production of noise from helicopters. Since information regarding characteristics and production of noise from helicopters has not changed, this SEIS/OEIS will not further analyze characteristics and production of noise from helicopters.

**Table 3.0-4: Representative Aircraft Sound Characteristics**

| <i>Noise Source</i>                                | <i>Sound Pressure Level</i>  |
|--|--|
| <b><i>In-Water Noise Level</i></b>                 |  |
| F/A-18 Subsonic at 1,000 ft. (300 m) Altitude      | 152 dB re 1 $\mu$ Pa at 2 m below water surface <sup>1</sup>   |
| F/A-18 Subsonic at 10,000 ft. (3,000 m) Altitude   | 128 dB re 1 $\mu$ Pa at 2 m below water surface <sup>1</sup>   |
| H-60 Helicopter Hovering at 82 ft. (25 m) Altitude | Approximately 125 dB re 1 $\mu$ Pa at 1 m below water surface, estimate based on in-air level <sup>2</sup>         |
| <b><i>Airborne Noise Level</i></b>                 |  |
| F/A-18C/D Under Military Power                     | 143 dBA re 20 $\mu$ Pa at 13 m from source <sup>3</sup>  |
| F/A-18C/D Under Afterburner                        | 146 dBA re 20 $\mu$ Pa at 13 m from source <sup>3</sup>  |
| F35-A Under Military Power                         | 145 dBA re 20 $\mu$ Pa at 13 m from source <sup>3</sup>  |
| F-35-A Under Afterburner                           | 148 dBA re 20 $\mu$ Pa at 13 m from source <sup>3</sup>  |
| H-60 Helicopter Hovering at 82 ft. (25 m) Altitude | 113 dBA re 20 $\mu$ Pa at 25 m from source <sup>2</sup>  |
| F-35A Takeoff Through 1,000 ft. (300 m) Altitude   | 119 dBA re 20 $\mu$ Pa <sup>2</sup> s <sup>4</sup> (per second of duration), based on average sound exposure level |
| EA-18G Takeoff Through 1,622 ft. (500 m) Altitude  | 115 dBA re 20 $\mu$ Pa <sup>2</sup> s <sup>5</sup> (per second of duration), based on average sound exposure level |

Notes: dB re 1  $\mu$ Pa = decibel(s) referenced to 1 micropascal, dBA re 20  $\mu$ Pa = A-weighted decibel(s) referenced to 20 micropascals, m = meter(s), ft. = feet, dBA re 20  $\mu$ Pa<sup>2</sup>s = A-weighted decibel(s) referenced to 20 micropascals squared seconds

Sources: <sup>1</sup>Eller and Cavanagh (2000), <sup>2</sup>Bousman and Kufeld (2005), <sup>3</sup>U.S. Naval Research Advisory Committee (2009), <sup>4</sup>U.S. Department of the Air Force (2016), <sup>5</sup>U.S. Department of the Navy (2012a).

### 3.0.4.1.3.3 Sonic Booms

An intense but infrequent type of aircraft noise is the sonic boom, produced when an aircraft exceeds the speed of sound. An intense but infrequent type of aircraft noise is the sonic boom, produced when an aircraft exceeds the speed of sound. Per Navy Instruction *Naval Air Training and Operating Procedures General Flight and Operating Instructions Manual, Commander Naval Air Forces Manual-3710.7* (U.S. Department of the Navy, 2017a), it is incumbent on every pilot flying aircraft capable of generating sonic booms to reduce such disturbances and damage to the absolute minimum dictated by operational/training requirements. Supersonic flight operations shall be strictly controlled and supervised by operational commanders. Supersonic flight over land or within 30 NM offshore shall be conducted in specifically designated areas. Such areas must be chosen to ensure minimum possibility of disturbance. As a general policy, sonic booms shall not be intentionally generated below 30,000 ft. of altitude unless over water and more than 30 miles from inhabited land areas or islands. Deviations from the foregoing general policy may be authorized only under one of the following conditions:

- tactical missions that require supersonic speeds;
- phases of formal training syllabus flights requiring supersonic speeds;
- research, test, and operational suitability test flights requiring supersonic speeds; or
- when specifically authorized by the Chief of Naval Operations for flight demonstration purposes.

Several factors that influence sonic booms include weight, size, and shape of aircraft or vehicle; altitude; flight paths; and atmospheric conditions. A larger and heavier aircraft must displace more air and create more lift to sustain flight, compared with small, light aircraft. Therefore, larger aircraft create sonic

booms that are stronger than those of smaller, lighter aircraft. Consequently, the larger and heavier the aircraft, the stronger the shock waves (U.S. Department of the Navy & Department of Defense, 2007). Aircraft maneuvers that result in changes to acceleration, flight path angle, or heading can also affect the strength of a boom. In general, an increase in flight path angle (lifting the aircraft’s nose) will diffuse a boom while a decrease (lowering the aircraft’s nose) will focus it. In addition, acceleration will focus a boom while deceleration will weaken it. Any change in horizontal direction will focus a boom, causing two or more wave fronts that originated from the aircraft at different times to coincide exactly (U.S. Department of the Navy, 2001). Atmospheric conditions such as wind speed and direction, and air temperature and pressure can also influence the sound propagation of a sonic boom.

Of all the factors influencing sonic booms, increasing altitude is the most effective method of reducing sonic boom intensity. The width of the boom “carpet” or area exposed to sonic boom beneath an aircraft is about 1 mile for each 1,000 ft. of altitude. For example, an aircraft flying supersonic, straight and level at 50,000 ft. can produce a sonic boom carpet about 50 miles wide. The sonic boom, however, would not be uniform, and its intensity at the water surface would decrease with greater aircraft altitude. Maximum intensity is directly beneath the aircraft and decreases as the lateral distance from the flight path increases, until shock waves refract away from the ground or water surface and the sonic boom attenuates. The lateral spreading of the sonic boom depends only on altitude, speed, and the atmosphere and is independent of the vehicle’s shape, size, and weight. The ratio of the aircraft length to maximum cross-sectional area also influences the intensity of the sonic boom. The longer and more slender the aircraft, the weaker the shock waves. The wider and more blunt the aircraft, the stronger the shock waves can be (U.S. Department of the Navy & Department of Defense, 2007).

In air, the energy from a sonic boom is concentrated in the frequency range from 0.1 to 100 Hz. The underwater sound field due to transmitted sonic boom waveforms is primarily composed of low-frequency components (Sparrow, 2002), and frequencies greater than 20 Hz have been found to be difficult to observe at depths greater than 33 ft. (10 meters) (Sohn et al., 2000). F/A-18 Hornet supersonic flight was modeled to obtain peak sound pressure levels and energy flux density at the water surface and at depth (U.S. Department of the Air Force, 2000). Table 3.0-5 shows these results.

**Table 3.0-5: Sonic Boom Underwater Sound Levels Modeled for F/A-18 Hornet Supersonic Flight**

| Mach Number* | Aircraft Altitude (km) | Peak SPL (dB re 1 $\mu$ Pa) |            |             | Energy Flux Density (dB re 1 $\mu$ Pa <sup>2</sup> -s) <sup>1</sup> |            |             |
|--------------|------------------------|-----------------------------|------------|-------------|---|------------|-------------|
|              |                        | At surface                  | 50 m Depth | 100 m Depth | At surface  | 50 m Depth | 100 m Depth |
| 1.2          | 1                      | 176                         | 138        | 126         | 160   | 131        | 122         |
|              | 5                      | 164                         | 132        | 121         | 150   | 126        | 117         |
|              | 10                     | 158                         | 130        | 119         | 144   | 124        | 115         |
| 2            | 1                      | 178                         | 146        | 134         | 161   | 137        | 128         |
|              | 5                      | 166                         | 139        | 128         | 150   | 131        | 122         |
|              | 10                     | 159                         | 135        | 124         | 144   | 127        | 119         |

<sup>1</sup> Equivalent to SEL for a plane wave.

\* Mach number equals aircraft speed divided by the speed of sound.

Notes: SPL = sound pressure level, dB re 1  $\mu$ Pa = decibel(s) referenced to 1 micropascal, dB re 1  $\mu$ Pa<sup>2</sup>-s = decibel(s) referenced to 1 micropascal squared seconds, m = meter(s)

### 3.0.4.1.4 Weapon Noise

The Navy trains and tests using a variety of weapons, as described in Appendix A (Training and Testing Activities Descriptions). Depending on the weapon, incidental (unintentional) noise may be produced at launch or firing, while in flight, or upon impact. Other devices intentionally produce noise to serve as a non-lethal deterrent. Not all weapons utilize explosives, either by design or because they are non-explosive practice munitions. Noise produced by explosives, both in air and water, are discussed in Section 3.0.4.2 (Explosive Stressors).

Noise associated with large-caliber weapons firing and the impact of non-explosive practice munitions or kinetic weapons would occur at locations greater than 12 NM from shore in warning areas or special use airspace for safety reasons, with the exception of areas near Farallon de Medinilla (FDM). Small- and medium-caliber weapons firing could occur throughout the Study Area in identified training areas.

Table 3.0-6 shows examples of some types of weapons noise and provides examples of launch noise. Noise produced by other weapons and devices are described further below.

**Table 3.0-6: Example Weapons Noise**

| <i>Noise Source</i>                          | <i>Sound Level</i>  |
|--|---|
| <b><i>In-Water Noise Level</i></b>           |   |
| Naval Gunfire Muzzle Blast (5-inch)          | Approximately 200 dB re 1 $\mu$ Pa peak directly under gun muzzle at 1.5 m below the water surface <sup>1</sup> |
| <b><i>Airborne Noise Level</i></b>           |   |
| Naval Gunfire Muzzle Blast (5-inch)          | 178 dB re 20 $\mu$ Pa peak directly below the gun muzzle above the water surface <sup>1</sup>                   |
| Hellfire Missile Launch from Aircraft        | 149 dB re 20 $\mu$ Pa at 4.5 m <sup>2</sup>   |
| Advanced Gun System Missile (115-millimeter) | 133-143 dBA re 20 $\mu$ Pa between 12 and 22 m from the launcher on shore <sup>3</sup>                          |
| RIM 116 Surface-to-Air Missile               | 122-135 dBA re 20 $\mu$ Pa between 2 and 4 m from the launcher on shore <sup>3</sup>                            |
| Tactical Tomahawk Cruise Missile             | 92 dBA re 20 $\mu$ Pa 529 m from the launcher on shore <sup>3</sup>   |

Notes: dB re 1  $\mu$ Pa = decibel(s) referenced to 1 micropascal, dB re 20  $\mu$ Pa = decibel(s) referenced to 20 micropascals, dBA re 20  $\mu$ Pa = A-weighted decibel(s) referenced to 20 micropascals, m = meter(s)  
Sources: <sup>1</sup>Yagla and Stiegler (2003); <sup>2</sup>U.S. Department of the Army (1999); <sup>3</sup>U.S. Department of the Navy (2013)

#### 3.0.4.1.4.1 Muzzle Blast from Naval Gunfire

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.4, Weapons Firing, Launch, and Impact Noise) describes the characteristics of the 5-inch (in.) large caliber naval gun, which is the most prevalent large weapon fired. Since information regarding characteristics of muzzle blast from naval gunfire has not changed, this SEIS/OEIS will not further analyze muzzle blast from naval gunfire. Examples of noise measurements from naval gunfire muzzle blast are provided in Table 3.0-6.

#### 3.0.4.1.4.2 Supersonic Projectile Bow Shock Wave

Supersonic projectiles, such as a fired gun shell or kinetic energy weapon, create a bow shock wave along the line of fire. A bow shock wave is an impulsive sound caused by a projectile exceeding the speed of sound (for more explanation, see Appendix H, Acoustic and Explosive Concepts). The bow

shock wave itself travels at the speed of sound in air. The projectile bow shock wave created in air by a shell in flight at supersonic speeds propagates in a cone (generally about 65°) behind the projectile in the direction of fire (U.S. Department of the Navy, 1981). Exposure to the bow shock wave is very brief.

Projectiles from a 5 in./54 caliber gun would travel at approximately 2,600 ft./second, and the associated bow shock wave is subjectively described as a “crack” noise (U.S. Department of the Navy, 1981). Measurements of a 5 in. projectile shock wave ranged from 140 to 147 dB re 20 µPa SPL peak taken at the ground surface at 0.59 NM distance from the firing location and 10° off the line of fire for safety (approximately 190 meters from the shell’s trajectory) (U.S. Department of the Navy, 1981).

Hyperkinetic projectiles may travel up to and exceeding approximately six times the speed of sound in air, or about 6,500 ft./second (U.S. Department of the Navy, 2014). For a hyperkinetic projectile sized similar to the 5 in. shell, peak pressures would be expected to be several dB higher than those described for the 5 in. projectile above, following the model in U.S. Department of the Navy (1981).

Like sound from the gun muzzle blast, sound waves from a projectile in flight could only enter the water in a narrow cone beneath the sound source, with in-air sound being totally reflected from the water surface outside of the cone. The region of underwater sound influence from a single traveling shell would be relatively narrow, and the duration of sound influence would be brief at any location.

#### **3.0.4.1.4.3 Launch Noise**

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.4, Weapons Firing, Launch, and Impact Noise) describes launch noise. Since information regarding launch noise has not changed, this SEIS/OEIS will not further analyze launch noise. Table 3.0-6 provides examples of launch noise measurements.

#### **3.0.4.1.4.4 Impact Noise (Non-Explosive)**

The 2015 MITT Final EIS/OEIS (Section 3.0.5.2.1.4, Weapons Firing, Launch, and Impact Noise) describes characteristics and production of non-explosive impact noise. Since information regarding non-explosive impact noise has not changed, this SEIS/OEIS will not further analyze non-explosive impact noise.

#### **3.0.4.1.4.5 Long Range Acoustic Device**

The Long Range Acoustic Device is a communication device that can be used to warn vessels against continuing towards a high-value asset by emitting loud sounds in air. Although not a weapon, the Long Range Acoustic Device (and other hailing and deterrent devices) is considered along with in-air sounds produced by Navy sources. The system would typically be used in training activities nearshore, and use would be intermittent during these activities. Source levels at 1 meter range between 137 A-weighted decibels re 1 µPa for small portable systems and 153 A-weighted decibels re 1 µPa for large systems. Sound would be directed within a 30–60° wide zone and would be directed over open water.

#### **3.0.4.2 Explosive Stressors**

This section describes the characteristics of explosions during naval training and testing. The activities analyzed in the SEIS/OEIS that use explosives are described in Appendix A (Training and Testing Activities Descriptions). This section provides the basis for analysis of explosive impacts on resources in the remainder of Chapter 3 (Affected Environment and Environmental Consequences). Explanations of the terminology and metrics used when describing explosives in this SEIS/OEIS are in Appendix H (Acoustic and Explosive Concepts).

The near-instantaneous rise from ambient to an extremely high peak pressure is what makes an explosive shock wave potentially damaging. Farther from an explosive, the peak pressures decay and the

explosive waves propagate as an impulsive, broadband sound. Several parameters influence the effect of an explosive: the weight of the explosive warhead, the type of explosive material, the boundaries and characteristics of the propagation medium, and, in water, the detonation depth. The net explosive weight, the explosive power of a charge expressed as the equivalent weight of trinitrotoluene (TNT), accounts for the first two parameters. The effects of these factors are explained in Appendix H (Acoustic and Explosive Concepts).

#### 3.0.4.2.1.1 Explosions in Water

Explosive detonations during training and testing activities are associated with high-explosive munitions, including, but not limited to, bombs, missiles, rockets, naval gun shells, torpedoes, mines, demolition charges, and explosive sonobuoys. Explosive detonations during training and testing activities involving the use of high-explosive munitions, including bombs, missiles, and naval gun shells, could occur in the air or near the water's surface. Explosive detonations associated with torpedoes and explosive sonobuoys would occur in the water column; mines and demolition charges could be detonated in the water column or on the ocean bottom. Detonations would typically occur in waters greater than 200 ft. in depth, and greater than 3 NM from shore, with the exception of existing mine warfare areas, including Outer Apra Harbor, Piti, and Agat. Section 5.3.3 (Explosive Stressors) outlines the procedural mitigation measures for explosive stressors to reduce potential impacts on biological resources.

In order to better organize and facilitate the analysis of Navy training and testing activities using explosives that could detonate in water or at the water surface, explosive classification bins were developed. The use of explosive classification bins provides the same benefits as described for acoustic source classification bins in Section 3.0.4.1 (Acoustic Stressors).

Explosives detonated in water are binned by net explosive weight. The bins of explosives that are proposed for use in the Study Area are shown in Table 3.0-7. This table shows the number of explosive items that could be used in any year under each action alternative for training and testing activities. A range of annual bin use indicates that use of that bin is anticipated to vary annually, consistent with the variation in the number of annual activities described in Chapter 2 (Description of Proposed Action and Alternatives).

In addition to the explosives quantitatively analyzed for impacts on protected species shown in Table 3.0-7, the Navy uses some very small impulsive sources (less than 0.1 pounds net explosive weight), categorized in bin E0, that are not anticipated to result in takes of protected species. Quantitative modeling in multiple locations has validated that these sources have a very small zone of influence. These E0 charges, therefore, are categorized as *de minimis* sources and are qualitatively analyzed to determine the appropriate determinations under NEPA in the appropriate resource impact analyses, as well as under the MMPA and the ESA.

Propagation of explosive pressure waves in water is highly dependent on environmental characteristics such as bathymetry, bottom type, water depth, temperature, and salinity, which affect how the pressure waves are reflected, refracted, or scattered; the potential for reverberation; and interference due to multi-path propagation. In addition, absorption greatly affects the distance over which higher frequency components of explosive broadband noise can propagate. Appendix H (Acoustic and Explosive Concepts) explains the characteristics of explosive detonations and how the above factors affect the propagation of explosive energy in the water. Because of the complexity of analyzing sound propagation in the ocean environment, the Navy relies on acoustic models in its environmental analyses that consider sound source characteristics and varying ocean conditions across the Study Area.

**Table 3.0-7: Explosive Sources Quantitatively Analyzed that Could Be Used Underwater or at the Water Surface**

| <i>Explosives</i>            | <i>Training &amp; Testing Activities<br/>(Annual In-Water Detonations)</i> |                      |                      |
|------------------------------|--|----------------------|----------------------|
|                              | <i>2015 Final<br/>EIS/OEIS</i>   | <i>Alternative 1</i> | <i>Alternative 2</i> |
| E1<br>(0.1–0.25 lb. NEW)     | 10,140   | 512                  | 768                  |
| E2<br>(>0.25–0.5 lb. NEW)    | 106  | 400                  | 400                  |
| E3<br>(>0.5–2.5 lb. NEW)     | 932  | 683                  | 683                  |
| E4<br>(> 2.5–5 lb. NEW)      | 420  | 44                   | 44                   |
| E5<br>(> 5–10 lb. NEW)       | 684  | 965                  | 1,221                |
| E6<br>(> 10–20 lb. NEW)      | 76   | 29                   | 29                   |
| E8<br>(> 60–100 lb. NEW)     | 16   | 132–134              | 132–134              |
| E9<br>(> 100–250 lb. NEW)    | 4  | 110                  | 110                  |
| E10<br>(> 250–500 lb. NEW)   | 12   | 69                   | 78                   |
| E11<br>(> 500–650 lb. NEW)   | 6  | 1–3                  | 1–5                  |
| E12<br>(> 650–1,000 lb. NEW) | 184  | 48                   | 48                   |

Notes: lb. = pound(s), NEW = Net Explosive Weight

**3.0.4.2.1.2 Explosions in Air**

Explosions in air include detonations of projectiles and missiles during surface-to-air gunnery and air-to-air missile exercises conducted during air warfare. These explosions typically occur far above the water surface in special use airspace. Some typical types of explosive munitions that would be detonated in air during Navy activities are shown in Table 3.0-8. Various missiles, rockets, and medium- and large-caliber projectiles may be explosive or non-explosive, depending on the objective of the training or testing activity in which they are used. Quantities of explosive and non-explosive missiles, rockets, and projectiles proposed for use during Navy training and testing are provided in the tables below.

**Table 3.0-8: Typical Air Explosive Munitions During Navy Activities**

| <i>Weapon Type<sup>1</sup></i>                | <i>Net Explosive Weight (lb.)</i> | <i>Typical Altitude of Detonation (ft.)</i> |
|---|-----------------------------------|---|
| <b>Surface-to-Air Missile</b>                 |                                   |   |
| RIM-66 SM-2 Standard Missile                  | 80                                | > 15,000                                    |
| RIM-116 Rolling Airframe Missile              | 39                                | < 3,000                                     |
| RIM-7 Sea Sparrow                             | 36                                | > 15,000 (can be used on low targets)       |
| FIM-92 Stinger                                | 7                                 | < 3,000                                     |
| <b>Air-to-Air Missile</b>                     |                                   |   |
| AIM-9 Sidewinder                              | 38                                | > 15,000                                    |
| AIM-7 Sparrow                                 | 36                                | > 15,000                                    |
| AIM-120 AMRAAM                                | 17                                | > 15,000                                    |
| <b>Air-to-Surface Missile</b>                 |                                   |   |
| AGM-88 HARM                                   | 45                                | < 100                                       |
| <b>Projectile – Large-Caliber<sup>2</sup></b> |                                   |   |
| 5"/54 caliber HE-ET                           | 7                                 | < 100                                       |
| 5"/54 caliber Other                           | 8                                 | < 3,000                                     |

<sup>1</sup> Mission Design Series and popular name shown for missiles.

<sup>2</sup> Most medium and large caliber projectiles used during training and testing activities do not contain high explosives.

Notes: AMRAAM = Advanced Medium-Range Air-to-Air Missile, HARM = High-Speed Anti-Radiation Missile, HE-ET = High Explosive-Electronic Time, lb. = pound(s), ft. = foot/feet

Bombs and projectiles that detonate at or near the water surface, which are considered for underwater impacts (see Table 3.0-8), would also release some explosive energy into the air. Appendix A (Training and Testing Activities Descriptions) describes where activities with these stressors typically occur.

The explosive energy released by detonations in the air has been well-studied (see Appendix H, Acoustic and Explosive Concepts) and basic methods are available to estimate the explosive energy exposure with distance from the detonation (U.S. Department of the Navy, 1975). In air, the propagation of impulsive noise from an explosion is highly influenced by atmospheric conditions, including temperature and wind. While basic estimation methods do not consider the unique environmental conditions that may be present on a given day, they allow for approximation of explosive energy propagation under neutral atmospheric conditions. Explosions that occur during air warfare would typically be at a sufficient altitude that a large portion of the sound refracts upward due to cooling temperatures with increased altitude.

Missiles, rockets, projectiles, and other cased weapons will produce casing fragments upon detonation. These fragments may be of variable size and are ejected at supersonic speed from the detonation. The casing fragments will be ejected at velocities much greater than debris from any target due to the proximity of the casing to the explosive material. Unlike detonations on land targets, in-air detonations during Navy training and testing would not result in other propelled materials such as crater debris.

#### **3.0.4.3 Energy Stressors**

Energy stressors are discussed in the 2015 MITT Final EIS/OEIS. Changes to energy stressors analyzed in this SEIS/OEIS are described below.

**3.0.4.3.1 Electromagnetic Devices**

In the 2015 MITT Final EIS/OEIS, electromagnetic devices included those used in water. For this SEIS/OEIS, electromagnetic devices are further categorized as either in-water electromagnetic devices or in-air electromagnetic devices.

**3.0.4.3.1.1 In-Water Electromagnetic Devices**

In-water electromagnetic devices were described in Section 3.0.5.2.2.1 (Electromagnetic Devices) of the 2015 MITT Final EIS/OEIS. Table 3.0-9 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of in-water electromagnetic devices.

**Table 3.0-9: Annual Number of Events in the Study Area Including In-Water Electromagnetic Devices**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 5                             | 4                    | 4                    |

**3.0.4.3.1.2 In-Air Electromagnetic Devices**

Sources of electromagnetic energy in the air include communications transmitters, radars, and electronic countermeasures transmitters. Electromagnetic devices on Navy platforms operate across a wide range of frequencies and power. On a single ship the source frequencies may range from 2 megahertz (MHz) to 14,500 MHz, and transmitter maximum average power may range from 0.25 watts to 1,280,00 watts.

The Navy originally coined the term “radar” to refer to Radio Detection And Ranging. A radar system is an electromagnetic device that emits radio waves to detect and locate objects. In most cases, basic radar systems operate by generating pulses of radio frequency energy and transmitting these pulses via directional antennae into space (Courbis & Timmel, 2008). Some of this energy is reflected by the target back to the antenna, and the signal is processed to provide useful information to the operator.

Radars come in a variety of sizes and power, ranging from wide-band milliwatt systems to very high-power systems that are used primarily for long-range search and surveillance (Courbis & Timmel, 2008). In general, radars operate at radio frequencies that range between 300 MHz and 300 gigahertz, and are often classified according to their frequency range. Navy vessels commonly operate radar systems that include S-band and X-band electronically steered radar. S-band radar serves as the primary search and acquisition sensor capable of tracking and collecting data on a large number of objects, while X-band radar can provide high-resolution data on particular objects of interest and discrimination for weapons systems. Both systems employ a variety of waveforms and bandwidths to provide high-quality data collection and operational flexibility (Baird et al., 2016).

It is assumed that most Navy platforms associated with the Proposed Action will be transmitting from a variety of in-air electromagnetic devices at all times when underway, with very limited exceptions. Most of these transmissions (e.g., for routine surveillance, communications, and navigation) will be at low power. High-power settings are used for a small number of activities, including ballistic missile defense training, radar and other system testing, and signature analysis operations. The number of Navy vessels or aircraft in the Study Area at any given time varies and is dependent on local training or testing

requirements. Therefore, in-air electromagnetic energy as part of the Proposed Action would be widely dispersed throughout the Study Area, but more concentrated in portions of the Study Area near ports, naval installations, and range complexes. Because these stressors are operated at power levels, altitudes, and distances from people and animals to ensure that energy received is well below levels that could disrupt behavior or cause injury and because most in-air electromagnetic energy is reflected by water, in-air electromagnetic energy is not analyzed further in Section 3.4 (Marine Mammals) and Section 3.6 (Marine Birds).

**3.0.4.3.2 Lasers**

Laser devices can be organized into two categories: (1) low-energy lasers and (2) high-energy lasers.

**3.0.4.3.2.1 Low-Energy Lasers**

Low-energy lasers are proposed to be used as described in the 2015 MITT Final EIS/OEIS, where they would have an extremely low potential to impact marine biological resources (U.S. Department of the Navy, 2010). Therefore, as in the 2015 MITT Final EIS/OEIS, low-energy lasers will not be further analyzed in this SEIS/OEIS for possible impacts on biological resources.

**3.0.4.3.2.2 High-Energy Lasers**

While no high-energy lasers were proposed to be used in the Study Area previously, they are now proposed for use as part of the Proposed Action in this SEIS/OEIS. High-energy laser weapons testing involves the use of directed energy as a weapon against small surface vessels and airborne targets. High-energy lasers would be employed from surface ships and are designed to create small but critical failures in potential targets. The high-energy laser is expected to be used at short ranges. Marine life or birds at or near the ocean surface could be susceptible to injury by high-energy lasers. Table 3.0-10 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of high-energy lasers.

**Table 3.0-10: Annual Number of Events in the Study Area Including High-Energy Lasers**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 0                             | 54                   | 60                   |

**3.0.4.4 Physical Disturbance and Strike Stressors**

As described in the 2015 MITT Final EIS/OEIS, physical disturbance and strike stressors can result from the Navy’s proposed use of aircraft and aerial targets, vessels, in-water devices, military expended materials, seafloor devices, and, on the island of FDM, ground disturbance and wildfires.

**3.0.4.4.1 Aircraft and Aerial Targets**

Section 3.0.5.2.3.1 (Aircraft and Aerial Targets) in the 2015 MITT Final EIS/OEIS described aircraft and aerial targets. Table 3.0-11 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of aircraft.

**Table 3.0-11: Annual Number of Events in the Study Area Including Aircraft Movement**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 22,397                        | 20,058               | 20,094               |

**3.0.4.4.2 Vessels**

Section 3.0.5.2.3.2 (Vessels) in the 2015 MITT Final EIS/OEIS described vessels. Table 3.0-12 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of vessels.

**Table 3.0-12: Annual Number of Events in the Study Area Including Vessel Movement**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 3,968                         | 4,249                | 4,493                |

**3.0.4.4.3 In-Water Devices**

Section 3.0.5.2.3.3 (In-Water Devices) in the 2015 MITT Final EIS/OEIS described in-water devices. Table 3.0-13 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of towed in-water devices.

**Table 3.0-13: Annual Number of Events in the Study Area Including In-Water Devices**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 2,205                         | 2,289                | 2,397                |

**3.0.4.4.4 Military Expended Materials**

Section 3.0.5.2.3.4 (Military Expended Materials) in the 2015 MITT Final EIS/OEIS described military expended materials. Table 3.0-14 shows the number of non-explosive practice munitions analyzed in the 2015 MITT Final EIS/OEIS and the number proposed in this SEIS/OEIS. Other military expended materials are listed in Table 3.0-15, explosive munitions in Table 3.0-16, and targets in Table 3.0-17.

**Table 3.0-14: Annual Number of Non-Explosive Practice Munitions Expended At Sea in the Study Area**

| <i>Non-Explosive Ordnance</i>           | <i>Training &amp; Testing</i> |                      |                      |
|---|-------------------------------|----------------------|----------------------|
|   | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Mine Neutralization System Neutralizers | 24                            | 0                    | 0                    |
| Anti-Torpedo Torpedoes                  | N/A <sup>1</sup>              | 8                    | 11                   |
| Torpedoes <sup>2</sup>                  | 169                           | 104                  | 132                  |

**Table 3.0-14: Annual Number of Non-Explosive Practice Munitions Expended At Sea in the Study Area (continued)**

| <i>Non-Explosive Ordnance</i>               | <i>Training &amp; Testing</i> |                      |                      |
|---|-------------------------------|----------------------|----------------------|
|   | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Bombs                                       | 848                           | 368                  | 368                  |
| Rockets                                     | 0                             | 1,697                | 1,697                |
| Rockets (Flechette)                         | Note 1                        | 89                   | 89                   |
| Missiles                                    | 20                            | 0                    | 0                    |
| Kinetic Energy Rounds                       | Note 1                        | 80                   | 180                  |
| Large-Caliber Projectiles                   | 6,918                         | 14,772               | 22,268               |
| Large-Caliber Projectile Land-Based Casings | Note 1                        | 2,800                | 4,200                |
| Medium-Caliber Projectiles                  | 87,540                        | 223,150              | 280,750              |
| Small-Caliber Projectiles                   | 88,140                        | 308,364              | 354,318              |

Note 1: These items were not calculated in the 2015 Final EIS/OEIS.

**Table 3.0-15: Annual Number of Other Military Expended Materials Used At Sea in the Study Area**

| <i>Other Military Expended Materials</i> | <i>Training &amp; Testing</i> |                      |                      |
|--|-------------------------------|----------------------|----------------------|
|  | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Acoustic Countermeasures                 | 294                           | 387                  | 466                  |
| Anchor (Expended)                        | Note 1                        | 20                   | 28                   |
| Anti-Torpedo Torpedo Accessories         | Note 1                        | 8                    | 11                   |
| Buoy (Non-Explosive)                     | 314                           | 70                   | 82                   |
| Canister – Miscellaneous                 | Note 1                        | 1                    | 1                    |
| Compression Pad or Plastic Pistons       | Note 1                        | 17,600               | 17,600               |
| Endcap – Chaff and Flares                | Note 1                        | 35,218               | 35,218               |
| Expended Bathythermograph                | 520                           | 341                  | 364                  |
| Fiber Optic Can                          | 28                            | 44                   | 44                   |
| Flare O-ring                             | Note 1                        | 17,618               | 17,618               |
| Heavyweight Torpedo Accessories          | 54                            | 49                   | 73                   |
| Lightweight Torpedo Accessories          | 72                            | 60                   | 66                   |
| Illumination Flare                       | 18                            | 18                   | 18                   |
| JATO Bottle                              | 20                            | 20                   | 20                   |

**Table 3.0-15: Annual Number of Other Military Expended Materials Used At Sea in the Study Area (continued)**

| <i>Other Military Expended Materials</i> | <i>Training &amp; Testing</i> |                      |                      |
|--|-------------------------------|----------------------|----------------------|
|  | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Marine Marker                            | 617                           | 538                  | 538                  |
| Sonobuoys                                | 11,912                        | 5,386                | 5,876                |

Note 1: These items were not calculated in the 2015 Final EIS/OEIS.

**Table 3.0-16: Annual Number of Explosive Munitions Expended At Sea in the Study Area**

| <i>Explosive Ordnance</i>               | <i>Training &amp; Testing</i> |                      |                      |
|---|-------------------------------|----------------------|----------------------|
|   | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Mine Neutralization System Neutralizers | 28                            | 44                   | 44                   |
| Grenades                                | Note 1                        | 400                  | 400                  |
| Torpedoes                               | 10                            | 5                    | 7                    |
| Bombs                                   | 212                           | 198                  | 198                  |
| Rockets                                 | 114                           | 323                  | 323                  |
| Missiles                                | 145                           | 231                  | 249                  |
| Large-Caliber Projectiles               | 12,220                        | 1,372                | 1,658                |
| Medium-Caliber Projectiles              | 10,190                        | 22,224               | 22,480               |
| Buoys                                   | 804                           | 392                  | 392                  |

Note 1: These items were not calculated in the 2015 Final EIS/OEIS.

**Table 3.0-17: Annual Number of Targets Expended At Sea in the Study Area**

| <i>Target</i>                  | <i>Training &amp; Testing</i> |                      |                      |
|--------------------------------|-------------------------------|----------------------|----------------------|
|                                | <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| Air Targets –Decoy             | Note 1                        | 153                  | 168                  |
| Air Targets –Drone             | Note 1                        | 1                    | 1                    |
| Mine Shape (Non-Explosive)     | Note 1                        | 599                  | 599                  |
| Ship Hulk                      | 2                             | 1                    | 1                    |
| Subsurface Target (Mobile)     | Note 1                        | 254                  | 265                  |
| Subsurface Target (Stationary) | Note 1                        | 4                    | 5                    |
| Surface Target (Mobile)        | Note 1                        | 1,499                | 1,581                |
| Surface Target (Stationary)    | 786                           | 879                  | 1,107                |

Note 1: These items were not calculated in the 2015 Final EIS/OEIS.

**3.0.4.4.5 Seafloor Devices**

Seafloor devices represent items used during training or testing activities that are deployed onto the seafloor and recovered. These items include moored mine shapes, anchors, and bottom placed instruments. In certain cases, weights that anchor a device would be expended when the device is recovered (e.g., pop up buoys). Seafloor devices are either stationary or move very slowly along the bottom and do not pose a threat to highly mobile organisms. The effect of devices on the bottom will be discussed as an alteration of the bottom substrate and associated living resources (i.e., invertebrates and vegetation). Table 3.0-18 shows the number of ongoing events (from the 2015 MITT Final EIS/OEIS) and the number of events proposed in this SEIS/OEIS that include the use of seafloor devices.

**Table 3.0-18: Annual Number of Events in the Study Area Including Seafloor Devices**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 200                           | 180                  | 182                  |

**3.0.4.4.6 Ground Disturbance and Wildfires**

Section 3.0.5.2.3.6 (Ground Disturbance and Wildfires) in the 2015 MITT Final EIS/OEIS described ground disturbance and wildfires on FDM. Table 3.0-19 shows the number and type of munitions analyzed in the 2015 MITT Final EIS/OEIS and proposed in this SEIS/OEIS.

**Table 3.0-19: Annual Number of Munitions Used on Farallon de Medinilla**

| <i>Ordnance Use</i>                 | <i>2015 Final EIS/OEIS</i>       | <i>Alternative 1</i>             | <i>Alternative 2</i>             |
|-------------------------------------|----------------------------------|----------------------------------|----------------------------------|
| Small-caliber Rounds                | 42,000                           | 44,096                           | 44,096                           |
| NEPM Bombs ≤ 2,000 lb.              | 2,670                            | 2,670                            | 2,670                            |
| Explosive Bombs ≤ 2,000 lb.         | 6,242                            | 6,242                            | 6,242                            |
| Explosive Missiles and Rockets ≤ 5" | 85 missiles;<br>2,000 rockets    | 115 missiles;<br>2,000 rockets   | 115 missiles;<br>2,000 rockets   |
| Explosive Grenades and Mortars      | 600                              | 1,000                            | 1,000                            |
| Medium-caliber Projectiles          | 17,350 explosive;<br>94,150 NEPM | 18,144 explosive;<br>94,150 NEPM | 18,144 explosive;<br>94,150 NEPM |
| Large-caliber Projectiles           | 1,200 explosive;<br>1,800 NEPM   | 400 explosive                    | 400 explosive                    |

Notes: lb. = pound, NEPM = Non-Explosive Practice Munition

**3.0.4.5 Entanglement Stressors**

As described in the 2015 MITT Final EIS/OEIS, entanglement stressors can result from the Navy’s proposed use of fiber optic cables, guidance wires, and decelerators/parachutes. In addition, sonobuoy wires, not previously identified as entanglement stressors, can be entanglement stressors and are included in this SEIS/OEIS for analysis.

### **3.0.4.5.1 Wires and Cables**

#### **3.0.4.5.1.1 Fiber Optic Cables**

Although a portion may be recovered, some fiber optic cables used during Navy training and testing associated with remotely operated mine neutralization activities would be expended. The length of the expended tactical fiber would vary (up to about 3,000 meters) depending on the activity. Tactical fiber has an 8-micrometer (0.008 millimeter [mm]) silica core and acrylate coating, and looks and feels like thin monofilament fishing line. Other characteristics of tactical fiber are a 242-micrometer (0.24 mm) diameter, 12-pound tensile strength, and 3.4-mm bend radius (Corning Incorporated, 2005; Raytheon Company, 2015). Tactical fiber is relatively brittle; it readily breaks if knotted, kinked, or abraded against a sharp object. Deployed tactical fiber will break if looped beyond its bend radius (3.4 mm), or exceeds its tensile strength (12 pounds). If the fiber becomes looped around an underwater object or marine animal, it will not tighten unless it is under tension. Such an event would be unlikely based on its method of deployment and its resistance to looping after it is expended. The tactical fibers are often designed with controlled buoyancy to minimize the fiber's effect on vehicle movement. The tactical fiber would be suspended within the water column during the activity, and then be expended and sink to the seafloor (effective sink rate of 1.45 centimeters/second (Raytheon Company, 2015)) where it would be susceptible to abrasion and burial by sedimentation.

#### **3.0.4.5.1.2 Guidance Wires**

Section 3.0.5.2.4.1 (Guidance Wires) in the 2015 MITT Final EIS/OEIS described guidance wires.

#### **3.0.4.5.1.3 Sonobuoy Wires**

Sonobuoys consist of a surface antenna and float unit and a subsurface hydrophone assembly unit. The two units are attached through a thin-gauge, dual-conductor, and hard-draw copper strand wire, which is then wrapped by a hollow rubber tubing or bungee in a spiral configuration. The tensile breaking strength of the wire and rubber tubing is no more than 40 pounds. The length of the wire is housed in a plastic canister dispenser, which remains attached upon deployment. The length of wire that extends out is no more than 1,500 ft. and is dependent on the water depth and type of sonobuoy. Attached to the wire is a kite-drogue and damper disk stabilizing system made of non-woven nylon fabric. The nylon fabric is very thin and can be broken by hand. The wire runs through the stabilizing system, and leads to the hydrophone components. The hydrophone components may be covered by thin plastic netting depending on type of sonobuoy, but pose no entanglement risk. Each sonobuoy has a saltwater activated polyurethane float that inflates when the sonobuoy is submerged and keeps the sonobuoy components floating vertically in the water column below it. Sonobuoys remain suspended in the water column for no more than 30 hours, after which they sink to the seafloor.

Bathythermographs are similar to sonobuoys in that they consist of a subsurface unit (to measure temperature of the water column in the case of the bathythermograph) that is connected by wire to the float unit (for air-deployed bathythermographs) or directly to the ship (for ship-deployed bathythermographs). The bathythermograph wire is similar to the sonobuoy wire as described above.

Table 3.0-20 shows the number of wires and cables analyzed in the 2015 MITT Final EIS/OEIS and the number proposed in this SEIS/OEIS.

**Table 3.0-20: Annual Number of Wires and Cables Expended in the Study Area**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| <b>Fiber Optic Cables</b>     |                      |                      |
| 144                           | 44                   | 44                   |
| <b>Guidance Wires</b>         |                      |                      |
| 60                            | 49                   | 73                   |
| <b>Sonobuoy Wires</b>         |                      |                      |
| Note 1                        | 5,386                | 5,876                |
| <b>Bathythermograph Wires</b> |                      |                      |
| Note 1                        | 341                  | 364                  |

Note 1: These items were not calculated in the 2015 Final EIS/OEIS.

**3.0.4.5.2 Decelerators/Parachutes**

Decelerators/parachutes used during training and testing activities are classified into four different categories based on size: small, medium, large, and extra-large (Table 3.0-21). Aircraft-launched sonobuoys and lightweight torpedoes (such as the MK 46 and MK 54) use nylon decelerators/parachutes ranging in size from 18 to 48 in. in diameter (small). The majority of the decelerators/parachutes in the small size category are smaller (18 in.) cruciform shape decelerators/parachutes associated with sonobuoys (Figure 3.0-1). Illumination flares use medium decelerators/parachutes, up to approximately 19 ft. in diameter. Both small- and medium-sized decelerators/parachutes are made of cloth and nylon, many with weights attached to their short attachment lines to speed their sinking. At water impact, the decelerator/parachute assembly is expended and sinks away from the unit. The decelerator/parachute assembly may remain at the surface for 5–15 seconds before the decelerator/parachute and its housing sink to the seafloor, where it becomes flattened (Environmental Sciences Group, 2005). Once settled on the bottom the canopy may temporarily billow if bottom currents are present.

**Table 3.0-21: Size Categories for Decelerators/Parachutes Expended During Training and Testing Activities**

| <i>Size Category</i> | <i>Diameter (ft.)</i> | <i>Associated Activity</i>   |
|----------------------|-----------------------|--|
| Small                | 1.5–6                 | Air-launched sonobuoys, lightweight torpedoes, and drones (drag decelerator/parachute) |
| Medium               | 19                    | Illumination flares  |
| Large                | 30–50                 | Drones (main decelerator/parachute)  |
| Extra-large          | 82                    | Drones (main decelerator/parachute)  |



**Figure 3.0-1: Sonobuoy Launch Depicting the Relative Size of a Small Decelerator/Parachute**

Aerial targets (drones) use large (between 30 and 50 ft. in diameter) and extra-large (82 ft. in diameter) decelerators/parachutes (Figure 3.0-2). Large and extra-large decelerators/parachutes are also made of cloth and nylon, with suspension lines of varying lengths (large: 40–70 ft. in length [with up to 28 lines per decelerator/parachute]; extra-large: 82 ft. in length [with up to 64 lines per decelerator/parachute]). Some aerial targets also use a small drag parachute (6 ft. in diameter) to slow their forward momentum prior to deploying the larger primary decelerator/parachute. Unlike the small- and medium-sized decelerators/parachutes, drone decelerators/parachutes do not have weights attached and may remain at the surface or suspended in the water column for some time prior to eventual settlement on the seafloor.



**Figure 3.0-2: Aerial Target (Drone) with Parachute Deployed**

Table 3.0-22 shows the number of decelerators/parachutes analyzed in the 2015 MITT Final EIS/OEIS and the number proposed in this SEIS/OEIS.

**Table 3.0-22: Annual Number of Decelerators/Parachutes Expended in the Study Area**

| <i>Training &amp; Testing</i> |                                  |                                  |
|-------------------------------|----------------------------------|----------------------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i>             | <i>Alternative 2</i>             |
| 12,572                        | 10 Large, 18 Medium, 5,437 Small | 10 Large, 18 Medium, 5,934 Small |

**3.0.4.6 Ingestion Stressors**

As described in the 2015 MITT Final EIS/OEIS, ingestion stressors can result from the Navy’s proposed use of non-explosive practice munitions (small- and medium-caliber), fragments from explosives, fragments from targets, chaff, flare casings (including plastic end caps and pistons), and decelerator/parachutes. The annual number of non-explosive practice munitions expended is shown in Table 3.0-14, the number of explosive munitions that could fragment is shown in Table 3.0-16, the number of targets that could fragment is shown in Table 3.0-17, the number of decelerator/parachutes is shown in Table 3.0-22, the number of chaff cartridges is shown in Table 3.0-23, and the number of flares is shown in Table 3.0-24.

**Table 3.0-23: Annual Number of Chaff Cartridges Expended in the Study Area**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| <b>Chaff – Air Cartridge</b>  |                      |                      |
| 26,000                        | 17,600               | 17,600               |
| <b>Chaff – Ship Cartridge</b> |                      |                      |
| 440                           | 246                  | 360                  |

**Table 3.0-24: Annual Number of Flares Expended in the Study Area**

| <i>Training &amp; Testing</i> |                      |                      |
|-------------------------------|----------------------|----------------------|
| <i>2015 Final EIS/OEIS</i>    | <i>Alternative 1</i> | <i>Alternative 2</i> |
| 25,900                        | 17,600               | 17,600               |

**3.0.4.7 Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities**

This conceptual framework describes the potential effects from exposure to acoustic and explosive activities and the accompanying short-term costs to the animal (e.g., expended energy or missed feeding opportunity). It then outlines the conditions that may lead to long-term consequences for the individual if the animal cannot fully recover from the short-term costs and how these in turn may affect the population. Within each biological resource section (e.g., marine mammals, birds, and fishes) the detailed methods to predict effects on specific taxa are derived from this conceptual framework.

An animal is considered “exposed” to a sound if the received sound level at the animal’s location is above the background ambient noise level within a similar frequency band. A variety of effects may result from exposure to acoustic and explosive activities.

The categories of potential effects are:

- **Injury** - Injury to organs or tissues of an animal.
- **Hearing loss** - A noise-induced decrease in hearing sensitivity that can be either temporary or permanent and may be limited to a narrow frequency range of hearing.
- **Masking** - When the perception of a biologically important sound (i.e., signal) is interfered with by a second sound (i.e., noise).
- **Physiological stress** - An adaptive process that helps an animal cope with changing conditions; although, too much stress can result in physiological problems.
- **Behavioral response** - A reaction ranging from very minor and brief changes in attentional focus, changes in biologically important behaviors, and avoidance of a sound source or area, to aggression or prolonged flight.

Figure 3.0-3 is a flowchart that diagrams the process used to evaluate the potential effects to marine animals exposed to sound-producing activities. The shape and color of each box on the flowchart represents either a decision point in the analysis (green diamonds); specific processes such as responses, costs, or recovery (blue rectangles); external factors to consider (purple parallelograms); and final outcomes for the individual or population (orange ovals and rectangles). Each box is labeled for reference throughout the following sections. For simplicity, sound is used here to include not only sound waves but also blast waves generated from explosive sources. Box A1, the Sound-Producing Activity, is the source of this stimuli and therefore the starting point in the analysis.

The first step in predicting whether an activity is capable of affecting a marine animal is to define the stimuli experienced by the animal. The stimuli include the overall level of activity, the surrounding acoustical environment, and characteristics of the sound when it reaches the animal.




Figure 3.0-3: Flow Chart of the Evaluation Process of Sound-Producing Activities

This page intentionally left blank.

Sounds emitted from a sound-producing activity (Box A1) travel through the environment to create a spatially variable sound field. The received sound by the animal (Box A2) determines the range of possible effects. The received sound can be evaluated in several ways, including number of times the sound is experienced (repetitive exposures), total received energy, or highest SPL experienced. Sounds that are higher than the ambient noise level and within an animal's hearing sensitivity range (Box A3) have the potential to cause effects. There can be any number of individual sound sources in a given activity, each with its own unique characteristics. For example, a Navy training exercise may involve several ships and aircraft using several types of sonar. Environmental factors such as temperature and bottom type impact how sound spreads and attenuates through the environment. Additionally, independent of the sounds, the overall level of activity and the number and movement of sound sources are important to help predict the probable reactions.

The magnitude of the responses is based on the characteristics of the acoustic stimuli and the characteristics of the animal (species, susceptibility, life history stage, size, and past experiences). Very high exposure levels close to explosives have the potential to cause injury. High-level, long-duration, or repetitive exposures may potentially cause some hearing loss. All perceived sounds may lead to behavioral responses, physiological stress, and masking. Many sounds, including sounds that are not detectable by the animal, could have no effect (Box A4).

#### **3.0.4.7.1 Injury**

Injury (Box B1) refers to the direct injury of tissues and organs by shock or pressure waves impinging upon or traveling through an animal's body. Marine animals are well adapted to large, but relatively slow, hydrostatic pressure changes that occur with changing depth. However, injury may result from exposure to rapid pressure changes, such that the tissues do not have time to adequately adjust.

Therefore, injury is normally limited to relatively close ranges from explosions. Injury can be mild and fully recoverable or, in some cases, lead to mortality.

Injury includes both auditory and non-auditory injury. Auditory injury is the direct mechanical injury to hearing-related structures, including tympanic membrane rupture, disarticulation of the middle ear ossicles, and injury to the inner ear structures such as the organ of Corti and the associated hair cells. Auditory injury differs from auditory fatigue in that the latter involves the overstimulation of the auditory system at levels below those capable of causing direct mechanical damage. Auditory injury is always injurious but can be temporary. One of the most common consequences of auditory injury is hearing loss.

Non-auditory injury can include hemorrhaging of small blood vessels and the rupture of gas-containing tissues such as the lung, swim bladder, or gastrointestinal tract. After the ear (or other sound-sensing organs), these are usually the organs and tissues most sensitive to explosive injury. An animal's size and anatomy are important in determining its susceptibility to non-auditory injury (Box B2). Larger size indicates more tissue to protect vital organs. Therefore, larger animals should be less susceptible to injury than smaller animals. In some cases, acoustic resonance of a structure may enhance the vibrations resulting from noise exposure and result in an increased susceptibility to injury. The size, geometry, and material composition of a structure determine the frequency at which the object will resonate. Because most biological tissues are heavily damped, the increase in susceptibility from resonance is limited.

Vascular and tissue bubble formation resulting from sound exposure is a hypothesized mechanism of injury to breath-holding marine animals. Bubble formation and growth due to direct sound exposure

have been hypothesized (Crum & Mao, 1996; Crum et al., 2005); however, the experimental laboratory conditions under which these phenomena were observed would not be replicated in the wild. Certain dive behaviors by breath-holding animals are predicted to result in conditions of blood nitrogen super-saturation, potentially putting an animal at risk for decompression sickness (Fahlman et al., 2014), although this phenomena has not been observed (Houser et al., 2009). In addition, animals that spend long periods of time at great depths are predicted to have super-saturated tissues that may slowly release nitrogen if the animal then spends a long time at the surface (i.e., stranding) (Houser et al., 2009).

Injury could increase the animal's physiological stress (Box B8), which feeds into the stress response (Box B7) and also increases the likelihood or severity of a behavioral response. Injury may reduce an animal's ability to secure food by reducing its mobility or the efficiency of its sensory systems, making the injured individual less attractive to potential mates, increasing an individual's chances of contracting diseases, falling prey to a predator (Box D2), or increasing an animal's overall physiological stress level (Box D10). Severe injury can lead to the death of the individual (Box D1).

Damaged tissues from mild to moderate injury may heal over time. The predicted recovery of direct injury is based on the severity of the injury, availability of resources, and characteristics of the animal. The animal may also need to recover from any potential costs due to a decrease in resource gathering efficiency and any secondary effects from predators or disease. Severe injuries can lead to reduced survivorship (longevity), elevated stress levels, and prolonged alterations in behavior that can reduce an animal's lifetime reproductive success. An animal with decreased energy stores or a lingering injury may be less successful at mating for one or more breeding seasons, thereby decreasing the number of offspring produced over its lifetime.


#### **3.0.4.7.2 Hearing Loss**

Hearing loss, also called a noise-induced threshold shift, is possibly the most studied type of effect from sound exposures to animals. Hearing loss manifests itself as loss in hearing sensitivity across part of an animal's hearing range, which is dependent upon the specifics of the noise exposure. Hearing loss may be either PTS or TTS. If the threshold shift eventually returns to zero (the animal's hearing returns to pre-exposure value), the threshold shift is a TTS. If the threshold shift does not return to zero but leaves some finite amount of threshold shift, then that remaining threshold shift is a PTS. Figure 3.0-4 shows one hypothetical threshold shift that completely recovers, a TTS, and one that does not completely recover, leaving some PTS.

The characteristics of the received sound stimuli are used and compared to the animal's hearing sensitivity and susceptibility to noise (Box A3) to determine the potential for hearing loss. The amplitude, frequency, duration, and temporal pattern of the sound exposure are important parameters for predicting the potential for hearing loss over a specific portion of an animal's hearing range. Duration is particularly important because hearing loss increases with prolonged exposure time. Longer exposures with lower sound levels can cause more threshold shift than a shorter exposure using the same amount of energy overall. The frequency of the sound also plays an important role. Experiments show that animals are most susceptible to hearing loss (Box B3) within their most sensitive hearing range. Sounds outside of an animal's audible frequency range do not cause hearing loss.

The mechanisms responsible for hearing loss may consist of a variety of mechanical and biochemical processes in the inner ear, including physical damage or distortion of the tympanic membrane (not including tympanic membrane rupture which is considered auditory injury), physical damage or

distortion of the cochlear hair cells, hair cell death, changes in cochlear blood flow, and swelling of cochlear nerve terminals (Henderson et al., 2006; Kujawa & Liberman, 2009). Although the outer hair cells are the most prominent target for fatigue effects, severe noise exposures may also result in inner hair cell death and loss of auditory nerve fibers (Henderson et al., 2006).



**Figure 3.0-4: Two Hypothetical Threshold Shifts**

The relationship between TTS and PTS is complicated and poorly understood, even in humans and terrestrial mammals, where numerous studies failed to delineate a clear relationship between the two. Relatively small amounts of TTS (e.g., less than 40–50 decibels measured two minutes after exposure) will recover with no apparent permanent effects; however, terrestrial mammal studies revealed that larger amounts of threshold shift can result in permanent neural degeneration, despite the hearing thresholds returning to normal (Kujawa & Liberman, 2009). The amounts of threshold shift induced by Kujawa and Liberman (2009) were described as being “at the limits of reversibility.” It is unknown whether smaller amounts of threshold shift can result in similar neural degeneration, or if effects would translate to other species such as marine animals.

Hearing loss can increase an animal’s physiological stress (Box B8), which feeds into the stress response (Box B7). Hearing loss increases the likelihood or severity of a behavioral response and increase an animal’s overall physiological stress level (Box D10). Hearing loss reduces the distance over which animals can communicate and detect other biologically important sounds (Box D3). Hearing loss could also be inconsequential for an animal if the frequency range affected is not critical for that animal to hear within, or the hearing loss is of such short duration (e.g., a few minutes) that there are no costs to the individual.

Small to moderate amounts of hearing loss may recover over a period of minutes to days, depending on the amount of initial threshold shift. Severe noise-induced hearing loss may not fully recover, resulting in some amount of PTS. An animal whose hearing does not recover quickly and fully could suffer a reduction in lifetime reproductive success. An animal with PTS may be less successful at mating for one or more breeding seasons, thereby decreasing the number of offspring it can produce over its lifetime.

#### **3.0.4.7.3 Masking**

Masking occurs if the noise from an activity interferes with an animal’s ability to detect, understand, or recognize biologically relevant sounds of interest (Box B4). In this context noise refers to unwanted or

unimportant sounds that mask an animal's ability to hear sounds of interest. Sounds of interest include those from conspecifics such as offspring, mates, and competitors; echolocation clicks; sounds from predators; natural, abiotic sounds that may aid in navigation; and reverberation, which can give an animal information about its location and orientation within the ocean. The probability of masking increases as the noise and sound of interest increase in similarity and the masking noise increases in level. The frequency, received level, and duty cycle of the noise determines the potential degree of auditory masking. Masking only occurs during the sound exposure.

A behavior decision (either conscious or instinctive) is made by the animal when the animal detects increased background noise, or possibly, when the animal recognizes that biologically relevant sounds are being masked (Box C1). An animal's past experiences can be important in determining the behavioral response when dealing with masking (Box C4). For example, an animal may modify its vocalizations to reduce the effects of masking noise. Other stimuli present in the environment can influence an animal's behavior decision (Box C5) such as the presence of predators, prey, or potential mates.

An animal may exhibit a passive behavioral response when coping with masking (Box C2). It may simply not respond and keep conducting its current natural behavior. An animal may also stop calling until the background noise decreases. These passive responses do not present a direct energetic cost to the animal; however, masking will continue, depending on the acoustic stimuli.

An animal may actively compensate for masking (Box C3). An animal can vocalize more loudly to make its signal heard over the masking noise. An animal may also shift the frequency of its vocalizations away from the frequency of the masking noise. This shift can actually reduce the masking effect for the animal and other animals that are listening in the area.

If masking impairs an animal's ability to hear biologically important sounds (Box D3) it could reduce an animal's ability to communicate with conspecifics or reduce opportunities to detect or attract more distant mates, gain information about their physical environment, or navigate. An animal that modifies its vocalization in response to masking could also incur a cost (Box D4). Modifying vocalizations may cost the animal energy, interfere with the behavioral function of a call, or reduce a signaler's apparent quality as a mating partner. For example, songbirds that shift their calls up an octave to compensate for increased background noise attract fewer or less-desirable mates, and many terrestrial species advertise body size and quality with low-frequency vocalizations (Slabbekoorn & Ripmeester, 2007). Masking may also lead to no measurable costs for an animal. Masking could be of short duration or intermittent such that biologically important sounds that are continuous or repeated are received by the animal between masking noise.

Masking only occurs when the sound source is operating; therefore, direct masking effects stop immediately upon cessation of the sound-producing activity. Masking could have long-term consequences for individuals if the activity was continuous or occurred frequently enough.

#### **3.0.4.7.4 Physiological Stress**

Marine animals naturally experience physiological stress as part of their normal life histories. The physiological response to a stressor, often termed the stress response, is an adaptive process that helps an animal cope with changing external and internal environmental conditions. Sound-producing activities have the potential to cause additional stress. However, too much of a stress response can be harmful to an animal, resulting in physiological dysfunction.

If a sound is detected (i.e., heard or sensed) by an animal, a stress response can occur (Box B7). The severity of the stress response depends on the received sound level by the animal (Box A2), the details of the sound-producing activity (Box A1), and the animal's life history stage (e.g., juvenile or adult, breeding or feeding season), and past experience with the stimuli (Box B5). An animal's life history stage is an important factor to consider when predicting whether a stress response is likely (Box B5). An animal's life history stage includes its level of physical maturity (i.e., larva, infant, juvenile, sexually mature adult) and the primary activity in which it is engaged such as mating, feeding, or rearing/caring for young. Prior experience with a stressor may be of particular importance because repeated experience with a stressor may dull the stress response via acclimation (St. Aubin & Dierauf, 2001) or increase the response via sensitization. Additionally, if an animal suffers injury or hearing loss, a physiological stress response will occur (Box B8).

The generalized stress response is characterized by a release of hormones (Reeder & Kramer, 2005) and other chemicals (e.g., stress markers) such as reactive oxidative compounds associated with noise-induced hearing loss (Henderson et al., 2006). Stress hormones include norepinephrine and epinephrine (i.e., the catecholamines), which produce elevations in the heart and respiration rate, increase awareness, and increase the availability of glucose and lipid for energy. Other stress hormones are the glucocorticoid steroid hormones cortisol and aldosterone, which are classically used as an indicator of a stress response and to characterize the magnitude of the stress response (Hennessy et al., 1979).

An acute stress response is traditionally considered part of the startle response and is hormonally characterized by the release of the catecholamines. Annoyance type reactions may be characterized by the release of either or both catecholamines and glucocorticoid hormones. Regardless of the physiological changes that make up the stress response, the stress response may contribute to an animal's decision to alter its behavior.

Elevated stress levels may occur whether or not an animal exhibits a behavioral response (Box D10). Even while undergoing a stress response, competing stimuli (e.g., food or mating opportunities) may overcome any behavioral response. Regardless of whether the animal displays a behavioral response, this tolerated stress could incur a cost to the animal. Reactive oxygen compounds produced during normal physiological processes are generally counterbalanced by enzymes and antioxidants; however, excess stress can lead to damage of lipids, proteins, and nucleic acids at the cellular level (Berlett & Stadtman, 1997; Sies, 1997; Touyz, 2004).

Frequent physiological stress responses may accumulate over time increasing an animal's chronic stress level. Each component of the stress response is variable in time, and stress hormones return to baseline levels at different rates. Elevated chronic stress levels are usually a result of a prolonged or repeated disturbance. Chronic elevations in the stress levels (e.g., cortisol levels) may produce long-term health consequences that can reduce lifetime reproductive success.

#### **3.0.4.7.5 Behavioral Reactions**

Behavioral responses fall into two major categories: alterations in natural behavior patterns and avoidance. These types of reactions are not mutually exclusive, and many overall reactions may be combinations of behaviors or a sequence of behaviors. Severity of behavioral reactions can vary drastically between minor and brief reorientations of the animal to investigate the sound, to severe reactions such as aggression or prolonged flight. The type and severity of the behavioral response will determine the cost to the animal. The total number of vessels and platforms involved, the size of the

activity area, the distance between the animal and activity, and the duration of the activity are important considerations when predicting the initial behavioral responses.

A physiological stress response (Box B7) such as an annoyance or startle reaction, or cueing or alerting (Box B6) may cause an animal to make a behavior decision (Box C6). Any exposure that produces an injury or hearing loss is also assumed to produce a stress response (Box B7) and increase the severity or likelihood of a behavioral reaction. Both an animal's experience (Box C4) and competing and reinforcing stimuli (Box C5) can affect an animal's behavior decision. The decision can result in three general types of behavioral reactions: no response (Box C9), area avoidance (Box C8), or alteration of a natural behavior (Box C7).

An animal's past experiences can be important in determining what behavior decision it may make when dealing with a stress response (Box C4). Habituation is the process by which an animal learns to ignore or tolerate stimuli over some period and return to a normal behavior pattern, perhaps after being exposed to the stimuli with no negative consequences. Sensitization is when an animal becomes more sensitive to a set of stimuli over time, perhaps as a result of a past, negative experience that could result in a stronger behavioral response.

Other stimuli (Box C5) present in the environment can influence an animal's behavioral response. These stimuli may be conspecifics or predators in the area or the drive to engage in a natural behavior. Other stimuli can also reinforce the behavioral response caused by acoustic stimuli. For example, the awareness of a predator in the area coupled with the sound-producing activity may elicit a stronger reaction than the activity alone would have.

An animal may reorient, become more vigilant, or investigate if it detects a sound-producing activity (Box C7). These behaviors all require the animal to divert attention and resources, therefore slowing or stopping their presumably beneficial natural behavior. This can be a very brief diversion, or an animal may not resume its natural behaviors until after the activity has concluded. An animal may choose to leave or avoid an area where a sound-producing activity is taking place (Box C8). A more severe form of this comes in the form of flight or evasion. Avoidance of an area can help the animal avoid further effects by avoiding or reducing further exposure. An animal may also choose not to respond to a sound-producing activity (Box C9).

An animal that alters its natural behavior in response to stress or an auditory cue may slow or cease its natural behavior and instead expend energy reacting to the sound-producing activity (Box D5). Natural behaviors include feeding, breeding, sheltering, and migrating. The cost of feeding disruptions depends on the energetic requirements of individuals and the potential amount of food missed during the disruption. Alteration in breeding behavior can result in delaying reproduction. The costs of a brief interruption to migrating or sheltering are less clear.

An animal that avoids a sound-producing activity may expend additional energy moving around the area, be displaced to poorer resources, miss potential mates, or have social interactions affected (Box D6). The amount of energy expended depends on the severity of the behavioral response. Missing potential mates can result in delaying reproduction. Groups could be separated during a severe behavioral response such as flight and offspring that depend on their parents may die if they are permanently separated. Splitting up an animal group can result in a reduced group size, which can have secondary effects on individual foraging success and susceptibility to predators.

Some severe behavioral reactions can lead to stranding (Box D7) or secondary injury (Box D8). Animals that take prolonged flight, a severe avoidance reaction, may injure themselves or strand in an

environment for which they are not adapted. Some injury is likely to occur to an animal that strands (Box D8). Injury can reduce the animal's ability to secure food and mates, and increase the animal's susceptibility to predation and disease (Box D2). An animal that strands and does not return to a hospitable environment may die (Box D9).

#### **3.0.4.7.6 Long-Term Consequences**

The potential long-term consequences from behavioral responses are difficult to discern. Animals displaced from their normal habitat due to an avoidance reaction may return over time and resume their natural behaviors. This is likely to depend upon the severity of the reaction and how often the activity is repeated in the area. In areas of repeated and frequent acoustic disturbance, some animals may habituate to the new baseline; conversely, species that are more sensitive may not return, or return but not resume use of the habitat in the same manner. For example, an animal may return to an area to feed but no longer rest in that area. Long-term abandonment or a change in the utilization of an area by enough individuals can change the distribution of the population. Frequent disruptions to natural behavior patterns may not allow an animal to recover between exposures, which increase the probability of causing long-term consequences to individuals.

The magnitude and type of effect and the speed and completeness of recovery (i.e., return to baseline conditions) must be considered in predicting long-term consequences to the individual animal (Box E4). The predicted recovery of the animal (Box E1) is based on the cost to the animal from any reactions, behavioral or physiological. Available resources fluctuate by season, location, and year and can play a major role in an animal's rate of recovery (Box E2). Recovery can occur more quickly if plentiful food resources, many potential mates, or refuge or shelter is available. An animal's health, energy reserves, size, life history stage, and resource gathering strategy affect its speed and completeness of recovery (Box E3). Animals that are in good health and have abundant energy reserves before an effect takes place will likely recover more quickly.

Animals that recover quickly and completely are unlikely to suffer reductions in their health or reproductive success, or experience changes in habitat utilization (Box F2). No population-level effects would be expected if individual animals do not suffer reductions in their lifetime reproductive success or change their habitat utilization (Box G2). Animals that do not recover quickly and fully could suffer reductions in their health and lifetime reproductive success; they could be permanently displaced or change how they use the environment; or they could die (Box F1). These long-term consequences to the individual can lead to consequences for the population (Box G1); although, population dynamics and abundance play a role in determining how many individuals would need to suffer long-term consequences before there was an effect on the population.

Long-term consequences to individuals can translate into consequences for populations dependent upon population abundance, structure, growth rate, and carry capacity. Carrying capacity describes the theoretical maximum number of animals of a particular species that the environment can support. When a population nears its carrying capacity, its growth is naturally limited by available resources and predator pressure. If one, or a few animals, in a population are removed or gather fewer resources, then other animals in the population can take advantage of the freed resources and potentially increase their health and lifetime reproductive success. Abundant populations that are near their carrying capacity (theoretical maximum abundance) that suffer consequences on a few individuals may not be affected overall. Populations that exist well below their carrying capacity may suffer greater consequences from any lasting consequences to even a few individuals. Population-level consequences can include a change in the population dynamics, a decrease in the growth rate, or a change in geographic distribution.

## REFERENCES

- Baird, R. W. (2013). False Killer Whales Around Kauai and Niihau. *Offshore Neighbors, Hawaii Fishing News*, pp. 24–25.
- Baird, R. W., D. L. Webster, S. Watwood, R. Morrissey, B. K. Rone, S. D. Mahaffy, A. M. Gorgone, D. B. Anderson, and D. J. Moretti. (2016). *Odontocete Studies on the Pacific Missile Range Facility in February 2015: Satellite-Tagging, Photo-Identification, and Passive Acoustic Monitoring. Final Report*. Olympia, WA: HDR Environmental Inc.
- Barlow, J. (2016). *Cetacean Abundance in the California Current Estimated from Ship-based Line-transect Surveys in 1991–2014*. (NOAA Administrative Report NMFS-SWFSC-LJ-1601). La Jolla, CA: Southwest Fisheries Science Center.
- Becker, E. A., K. A. Forney, P. C. Fiedler, J. Barlow, S. J. Chivers, C. A. Edwards, A. M. Moore, and J. V. Redfern. (2016). Moving Towards Dynamic Ocean Management: How Well Do Modeled Ocean Products Predict Species Distributions? *Remote Sensing*, 8(2), 149.
- Berlett, B. S., and E. R. Stadtman. (1997). Protein oxidation in aging, disease, and oxidative stress. *The Journal of Biological Chemistry*, 272(33), 20313–20316.
- Bousman, W. G., and R. M. Kufeld. (2005). *UH-60A Airloads Catalog*. Moffett Field, CA: National Aeronautics and Space Administration.
- Bradford, A. L., K. A. Forney, E. M. Oleson, and J. Barlow. (2017). Abundance estimates of cetaceans from a line-transect survey within the U.S. Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin*, 115(2), 129–142.
- Campbell, G. S., L. Thomas, K. Whitaker, A. B. Douglas, J. Calambokidis, and J. A. Hildebrand. (2015). Inter-annual and seasonal trends in cetacean distribution, density and abundance off southern California. *Deep Sea Research Part II: Topical Studies in Oceanography*, 112, 143–157.
- Carretta, J. V., E. Oleson, D. W. Weller, A. R. Lang, K. A. Forney, J. Baker, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, J. Moore, D. Lynch, L. Carswell, and R. L. Brownell. (2015). *U.S. Pacific Marine Mammal Stock Assessments: 2014* (NOAA Technical Memorandum NMFS-SWFSC-549). La Jolla, CA: Southwest Fisheries Science Center.
- Corning Incorporated. (2005). *Corning SMF-28e Optical Fiber Product Information*. Corning, NY: Corning Incorporated.
- Courbis, S., and G. Timmel. (2008). Effects of vessels and swimmers on behavior of Hawaiian spinner dolphins (*Stenella longirostris*) in Kealake'akua, Honaunau, and Kauhako bays, Hawai'i. *Marine Mammal Science*, 25(2), 430–440.
- Crum, L., and Y. Mao. (1996). Acoustically enhanced bubble growth at low frequencies and its implications for human diver and marine mammal safety. *The Journal of the Acoustical Society of America*, 99(5), 2898–2907.
- Crum, L., M. Bailey, J. Guan, P. Hilmo, S. Kargl, and T. Matula. (2005). Monitoring bubble growth in supersaturated blood and tissue *ex vivo* and the relevance to marine mammal bioeffects. *Acoustics Research Letters Online*, 6(3), 214–220.
- Eller, A. I., and R. C. Cavanagh. (2000). *Subsonic Aircraft Noise at and Beneath the Ocean Surface: Estimation of Risk for Effects on Marine Mammals*. McLean, VA: United States Air Force Research Laboratory.

- Environmental Sciences Group. (2005). *Canadian Forces Maritime Experimental and Test Ranges: Environmental Assessment Update 2005*. Kingston, Canada: Environmental Sciences Group, Royal Military College.
- Fahlman, A., P. L. Tyack, P. J. O. Miller, and P. H. Kvadsheim. (2014). How man-made interference might cause gas bubble emboli in deep diving whales. *Frontiers in Physiology*, 5(13), 1–6.
- Forney, K. A., E. A. Becker, D. G. Foley, J. Barlow, and E. M. Oleson. (2015). Habitat-based models of cetacean density and distribution in the central North Pacific. *Endangered Species Research*, 27, 1–20.
- Frisk, G. V. (2012). Noiseconomics: The relationship between ambient noise levels in the sea and global economic trends. *Scientific Reports*, 2(437), 1–4.
- HDR. (2012). *Summary Report: Compilation of Visual Survey Effort and Sightings for Marine Species Monitoring in the Hawaii Range Complex, 2005–2012*. Pearl Harbor, HI: U.S. Pacific Fleet.
- Henderson, D., E. C. Bielefeld, K. C. Harris, and B. H. Hu. (2006). The role of oxidative stress in noise-induced hearing loss. *Ear & Hearing*, 27, 1–19.
- Hennessy, M. B., J. P. Heybach, J. Vernikos, and S. Levine. (1979). Plasma corticosterone concentrations sensitively reflect levels of stimulus intensity in the rat. *Physiology and Behavior*, 22, 821–825.
- Houser, D. S., L. A. Dankiewicz-Talmadge, T. K. Stockard, and P. J. Ponganis. (2009). Investigation of the potential for vascular bubble formation in a repetitively diving dolphin. *The Journal of Experimental Biology*, 213, 52–62.
- Jefferson, T. A., M. A. Smultea, and C. E. Bacon. (2014). Southern California Bight marine mammal density and abundance from aerial survey, 2008–2013. *Journal of Marine Animals and Their Ecology*, 7(2), 14–30.
- Kujawa, S. G., and M. C. Liberman. (2009). Adding insult to injury: Cochlear nerve degeneration after "temporary" noise-induced hearing loss. *The Journal of Neuroscience*, 29(45), 14077–14085.
- Mintz, J. D., and C. L. Parker. (2006). *Vessel Traffic and Speed Around the U.S. Coasts and Around Hawaii*. Alexandria, VA: Center for Naval Analyses.
- Mintz, J. D., and R. J. Filadelfo. (2011). *Exposure of Marine Mammals to Broadband Radiated Noise (Specific Authority N0001-4-05-D-0500)*. Washington, DC: Center for Naval Analyses.
- Mintz, J. D. (2012). *Vessel Traffic in the Hawaii-Southern California and Atlantic Fleet Testing and Training Study Areas*. Alexandria, VA: Center for Naval Analyses.
- Mintz, J. D. (2016). *Characterization of Vessel Traffic in the Vicinities of HRC, SOCAL, and the Navy Operating Areas off the U.S. East Coast*. Alexandria, VA: Center for Naval Analyses.
- Mulsow, J., C. E. Schlundt, L. Brandt, and J. J. Finneran. (2015). Equal latency contours for bottlenose dolphins (*Tursiops truncatus*) and California sea lions (*Zalophus californianus*). *The Journal of the Acoustical Society of America*, 138(5), 2678.
- National Marine Fisheries Service. (2016). *Technical Guidance for Assessing the Effects of Anthropogenic Sound on Marine Mammal Hearing Underwater Acoustic Thresholds for Onset of Permanent and Temporary Threshold Shifts*. (NOAA Technical Memorandum NMFS-OPR-55). Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.

- Popper, A. N., A. D. Hawkins, R. R. Fay, D. A. Mann, S. M. Bartol, T. J. Carlson, S. Coombs, W. T. Ellison, R. L. Gentry, M. B. Halvorsen, S. Løkkeborg, P. H. Rogers, B. L. Southall, D. G. Zeddies, and W. N. Tavolga. (2014). *ASA S3/SC1.4 TR-2014 Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI*. New York, NY and London, United Kingdom: Acoustical Society of America Press and Springer Briefs in Oceanography.
- Purdy, R. (2016). Kauai's first documented green sea turtle nest of 2016 hatches: Sixty-three hatchlings emerge from Pacific Missile Range Facility. *Currents*(Fall), 24–25.
- Raytheon Company. (2015). *Airborne Mine Neutralization System (AMNS): Alternative Optical Fiber Engineering Study Final Report*. Portsmouth, RI: Integrated Defense Systems.
- Reeder, D. M., and K. M. Kramer. (2005). Stress in free-ranging mammals: Integrating physiology, ecology, and natural history. *Journal of Mammalogy*, 86(2), 225–235.
- Sies, H. (1997). Physiological Society Symposium: Impaired endothelial and smooth muscle cell function in oxidative stress-oxidative stress: Oxidants and antioxidants. *Experimental Physiology*, 82, 291–295.
- Slabbekoorn, H., and E. A. Ripmeester. (2007). Birdsong and anthropogenic noise: Implications and applications for conservation. *Molecular Ecology*, 17(1), 72–83.
- Sohn, R. A., F. Vernon, J. A. Hildebrand, and S. C. Webb. (2000). Field measurements of sonic boom penetration into the ocean. *The Journal of the Acoustical Society of America*, 107(6), 3073–3083.
- Sparrow, V. W. (2002). Review and status of sonic boom penetration into the ocean. *The Journal of the Acoustical Society of America*, 111(1), 537–543.
- St. Aubin, D., and L. A. Dierauf. (2001). Stress and Marine Mammals. In L. A. Dierauf & F. M. D. Gulland (Eds.), *Marine Mammal Medicine* (2nd ed., pp. 253–269). Boca Raton, FL: CRC Press.
- Touyz, R. M. (2004). Reactive oxygen species, vascular oxidative stress, and redox signaling in hypertension: What is the clinical significance? *Hypertension*, 44, 248–252.
- U.S. Department of the Air Force. (2000). *Supersonic Aircraft Noise At and Beneath the Ocean Surface: Estimation of Risk for Effects on Marine Mammals* (AFRL-HE-WP-TR-2000-0167). McLean, VA: United States Air Force Research Laboratory.
- U.S. Department of the Air Force. (2016). *United States Air Force F-35A Operational Beddown–Pacific Final Environmental Impact Statement*. Eielson Air Force Base, AK: United States Air Force.
- U.S. Department of the Army. (1999). *Finding of No Significant Impact for the Life Cycle Environmental Assessment for the HELLFIRE Modular Missile System*. Washington, DC: U.S. Department of Defense.
- U.S. Department of the Navy. (1975). *Explosion Effects and Properties Part I – Explosion Effects in Air*. Silver Spring, MD: White Oak Laboratory, Naval Surface Weapons Center.
- U.S. Department of the Navy. (1981). *Gun Blast Far Field Peak Overpressure Contours*. (NSWC TR 79-442). Silver Spring, MD: Naval Surface Weapons Center.
- U.S. Department of the Navy. (2001). *Sonic Boom Parametric Study*. Naval Air Station Patuxent River, MD: Applied Ordnance Technology, Inc. and Operational Environmental Planning Office
- U.S. Department of the Navy, and Department of Defense. (2007). *Finding of No Significant Harm and Final Environmental Assessment of the Naval Air Routine Training Exercises in East and Gulf*

- Coast Operation Areas and Seaward*. Norfolk, VA: Atlantic Division, Naval Facilities Engineering Command.
- U.S. Department of the Navy. (2010). *Laser System Usage in the Marine Environment: Applications and Environmental Considerations*. San Diego, CA: Space and Naval Warfare Systems Command Center Pacific.
- U.S. Department of the Navy. (2012a). *Biological Assessment for the Expeditionary Electronic Attack Squadron Realignment and Transition at Naval Air Station Whidbey Island, Oak Harbor, Washington*. Washington, DC: U.S. Department of the Navy.
- U.S. Department of the Navy. (2012b). *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis*. San Diego, CA: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2013). *Petition for Regulations Pursuant to Section 101(a)(5) of the Marine Mammal Protection Act Covering Taking of Marine Mammals Incidental to Target and Missile Launch Activities for the Period 2014–2019 at San Nicolas Island, California (50 CFR Part 216, Subpart I)*. Point Mugu, CA: Office of Protected Resources, National Marine Fisheries Service, National Oceanographic and Atmospheric Administration.
- U.S. Department of the Navy. (2014). *U.S. Navy Testing of Hypervelocity Projectiles and an Electromagnetic Railgun*. Wallops Island, VA: U.S. Department of the Navy.
- U.S. Department of the Navy. (2015a). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2015b). *Record of Decision for the Mariana Islands Training and Testing Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS)*. Washington, DC: U.S. Department of Defense.
- U.S. Department of the Navy. (2017a). *NATOPS General Flight and Operating Instructions Manual; OPNAV Instruction M-3710.7*. Washington, DC: Department of the Navy, Office of the Chief of Naval Operations.
- U.S. Department of the Navy. (2017b). *U.S. Navy Marine Species Density Database Phase III for the Hawaii-Southern California Training and Testing Study Area* (Naval Facilities Engineering Command Pacific Technical Report). Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- U.S. Department of the Navy. (2017c). *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)*. San Diego, CA: Space and Naval Warfare System Command, Pacific.
- U.S. Department of the Navy. (2018a). *U.S. Navy Marine Species Density Database Phase III for the Mariana Islands Training and Testing Study Area* (Naval Facilities Engineering Command Pacific Technical Report). Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- U.S. Department of the Navy. (2018b). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (Technical Report prepared by NUWC Division Newport, Space and Naval Warfare Systems Center Pacific, G2 Software Systems, and the National Marine Mammal Foundation). Newport, RI: Naval Undersea Warfare Center.
- U.S. Naval Research Advisory Committee. (2009). *Report on Jet Engine Noise Reduction*. Patuxent River, MD: Department of Defense.

Yagla, J., and R. Stiegler. (2003). *Gun blast noise transmission across the air-sea interface*. Paper presented at the 5th European Conference on Noise Control. Naples, Italy.

---

---

## 3.1 Sediments and Water Quality



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|         |   |        |
|---------|---|--------|
| 3.1     | Sediments and Water Quality.....          | 3.1-1  |
| 3.1.1   | Affected Environment.....                 | 3.1-1  |
| 3.1.1.1 | Existing Conditions.....                  | 3.1-1  |
| 3.1.2   | Environmental Consequences .....          | 3.1-5  |
| 3.1.2.1 | Explosives and Explosives Byproducts..... | 3.1-6  |
| 3.1.2.2 | Metals .....                              | 3.1-7  |
| 3.1.2.3 | Chemicals Other Than Explosives .....     | 3.1-8  |
| 3.1.2.4 | Other Materials.....                      | 3.1-11 |
| 3.1.3   | Public Scoping Comments .....             | 3.1-12 |

**List of Figures**

There are no figures in this section.

**List of Tables**

|              |  |       |
|--------------|--|-------|
| Table 3.1-1: | Rates of Failure and Low-Order Detonations ..... | 3.1-3 |
|--------------|--|-------|

This page intentionally left blank.

### **3.1 Sediments and Water Quality**

#### **3.1.1 Affected Environment**

The purpose of this section is to supplement the analysis of impacts on sediments and water quality as presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon de Medinilla (FDM). Information presented in the 2015 MITT Final EIS/OEIS that remains valid is noted as such and referenced in the appropriate sections. Any new or updated information describing the affected environment and analysis of impacts on sediments and water quality associated with the Proposed Action is provided in this section. Comments received from the public during scoping related to sediments and water quality are addressed in Section 3.1.3 (Public Scoping Comments) (Denton et al., 1997; U.S. Department of the Navy, 2015).

##### **3.1.1.1 Existing Conditions**

Ocean water resources, climate, and the existing water quality in the MITT Study Area were discussed in the 2015 MITT Final EIS/OEIS. The 2015 MITT Final EIS/OEIS analyzed potential impacts on sediment quality in nearshore and deep water sediments, as well as water quality on the surface and within the water column, and determined that potential impacts on sediment and water quality would not be significant. As stated in the 2015 MITT Final EIS/OEIS, some studies suggest that deep water is, in general, of higher quality than surface waters. Additionally, water quality in marine environments is determined by complex interactions between physical, chemical, and biological processes (U.S. Department of the Navy, 2015).

There is no new information since the publication of the 2015 MITT Final EIS/OEIS for marine sediments that would alter the analysis of potential impacts on water and sediment quality. New information, however, has been released that would improve the understanding of existing conditions for water quality, mainly with regards to marine debris and climate change. In addition, published results that inform the evaluation for water quality impacts in the immediate vicinity of FDM are now available and summarized below. The new information, however, does not indicate an appreciable change to the existing environmental conditions as described in the 2015 MITT Final EIS/OEIS.

##### **3.1.1.1.1 Marine Debris and Water Quality**

Richardson et al. (2016) describe the results of seine net (vertical nets that are held in place with weights and buoys) surveys in open ocean waters of the western and central Pacific Ocean within exclusive economic zones (EEZs) of 25 Pacific countries and territories, as well as in international waters. A majority of the reported purse seine (a seine net that fully encompasses an area of fish) pollution incidents are related to plastics waste. Other common pollution incidents are related to oil spillages and to abandoned, lost, or dumped fishing gear. Data analysis highlighted the need for increased monitoring, reporting, enforcement of pollution violations for all types of fishing vessels operating in the Pacific region, a regional outreach and compliance assistance program on marine pollution prevention, and improvements in Pacific port waste reception facilities. Most of the pollution incidents associated with marine debris occurred in Papua New Guinea's EEZ (approximately 45 percent), while less than 1 percent of the debris accumulations collected on the surface by Richardson et al. (2016) were within the portion of the United States (U.S.) EEZ surrounding Guam, the Commonwealth of the Northern Mariana Islands (CNMI), and other U.S. Pacific islands.

### 3.1.1.1.2 Climate Change and Water Quality

New information on the potential for climate change to impact water quality was obtained for the western Pacific region. The 2015 MITT Final EIS/OEIS identified decreasing ocean pH (i.e., increasing acidity), increasing water temperatures, and increasing storm activity as aspects of climate change that potentially impact water quality.

Rainfall and tropical cyclones are significant aspects of the climate on islands within the Study Area; however, potential impacts on rainfall and tropical cyclone patterns are difficult to estimate (Keener et al., 2015). One study for Guam predicts fewer, but more intense, storms, that would likely follow new storm tracks, and a moderate increase in daily and annual average rainfall (U.S. Marine Corps, 2014). On Saipan, an assessment of vulnerability to climate change assumed a future small increase in average rainfall, an increase in extreme rainfall, as well as more extreme wet and dry seasons. Although difficult to predict, changes in rainfall and storm intensity are generally anticipated to be harmful to ecosystems that provide ecological services beneficial to water quality within the Study Area.

Keener et al. (2015) documented a coral bleaching event off of Guam in 2013 through 2014. That event, combined with the strong associations between sea surface temperature increases and coral bleaching events throughout the Pacific (Griesser & Spillman, 2016), suggests that it is highly likely sea surface temperature increases in the Mariana Islands are at least partially to blame for coral bleaching events. Coral cover on Guam is generally similar to other southern Mariana Islands, but lower than the northern islands (Raymundo et al., 2016). Because coral distribution and coral cover on reefs is naturally patchy and heterogeneous, a single island-wide number is not a particularly useful summary of the coral community. Long-term monitoring surveys conducted by the National Oceanic and Atmospheric Administration's Coral Reef Ecosystem Division Pacific Assessment and Monitoring Program found approximately 10–15 percent coral cover overall, but the recent multi-year coral bleaching events have had dramatic, if patchy, consequences for the reef communities on Guam. For example, Raymundo et al. (2017) estimated a 53 percent decline in staghorn *Acropora* spp. on Guam. Of the 21 sites in the study, 6 are on Joint Region Marianas-administered submerged lands, including 4 in Apra Harbor. The estimated mean mortality of staghorn *Acropora* spp. was 80 percent at Big Blue Shoals, 80 percent at Western Shoals, 30 percent at Dogleg, and 90 percent at Gab (Raymundo et al., 2016). In the past several years, corals in Guam have been bleaching regularly each summer and recovery has been limited, leading to significant levels of coral mortality (Harvey, 2016; Raymundo et al., 2017).

Even though the new studies show variability in coral cover at FDM and decreases in some coral species off Guam, this information does not appreciably change the analysis presented in the 2015 MITT Final EIS/OEIS because the species composition on the reefs has not changed.

Changes in pH outside the normal range can make it difficult for marine organisms that make hard structures through calcification (e.g., shells or skeletons) to maintain their structures. Many of those creatures are at the base of the marine food chain, such as phytoplankton, so changes may cascade through the ecosystem. Rising water temperatures can be detrimental to coastal ecosystems and, by extension, coastal water quality because these ecosystems provide ecological services (e.g., sediment trapping, nutrient cycling).

### 3.1.1.1.3 Farallon de Medinilla

Range condition assessments are conducted at all operational ranges within the Mariana Islands Range Complex in accordance with Department of Defense (DoD) Instruction 4715.14, Operational Range Assessments, and the Chief of Naval Operations Range Sustainability Environmental Program Analysis

Policy. The Navy is committed to surveying the FDM coral reef environment every five years, as well as performing the routine clearance of unexploded ordnance and other range debris from the FDM impact areas. The coral reef surveys provide an indication if the waters surrounding FDM (designated Class A) are degrading in quality, as evidenced by coral health. Routine clearance of unexploded ordnance from the FDM impact areas removes potential sources of munition constituents, helping to protect CNMI's water quality. The Navy engaged with the National Marine Fisheries Service in coral consultations under the Endangered Species Act (ESA) and through the Essential Fish Habitat Assessment, relevant to all species of corals and essential fish habitats that are present in the Study Area. These consultations and regulatory conclusions were summarized in the 2015 MITT Final EIS/OEIS.

No detailed data was available for munitions expenditure during the last three decades of the 20th century on FDM, but early environmental planning documents in 1974 and 1999 provide some insight to the source loading. Delivered munitions that resulted in either a low-order detonation or a “dud” are the predominant energetic munition constituent source material on FDM. Munition constituents commonly associated with munitions such as high melting explosive (HMX) (also referred to as octogen), royal demolition explosive (RDX) (also referred to as cyclonite), dinitrotoluene (DNT), and heavy metals are likely present in small dispersed residual quantities associated with high-order detonations and in localized higher concentrations associated with duds or low-order detonations. Areas with high explosive concentrations are often found around “carcasses” of munitions that were only partly detonated. Heavily cratered areas on military ranges often have below detection or low high-explosive concentrations, suggesting that high-order detonations leave only trace amounts of explosive residues (U.S. Army Corps of Engineers, 2007).

The frequency of low-order detonations or dud rates of munitions fired into the impact zones at FDM is not known; however (MacDonald & Mendez, 2005) provided failure rates and low-order detonation rates for various munitions types, shown in Table 3.1-1.

**Table 3.1-1: Rates of Failure and Low-Order Detonations**

| Ordnance                  | Failure Rate (Percent) | Low-Order Detonation Rate (Percent) |
|---------------------------|------------------------|-------------------------------------|
| Guns/artillery            | 4.68                   | 0.16                                |
| Hand grenades             | 1.78                   | –                                   |
| Explosive ordnance        | 3.37                   | 0.09                                |
| Rockets                   | 3.84                   | –                                   |
| Submunitions <sup>1</sup> | 8.23                   | –                                   |

<sup>1</sup>Submunitions are munitions contained within and distributed by another device such as a rocket.

Sources: MacDonald and Mendez (2005); U.S. Army Corps of Engineers (2007)

Since the publication of the 2015 MITT Final EIS/OEIS, the draft analysis of multi-year dive surveys conducted in nearshore waters of FDM between 1997 and 2012 has been published (Smith & Marx, 2016). During these dive surveys, Smith and Marx (2016) provide qualitative observations of water quality and sediment quality surrounding the live-fire range. A summary of the observations is included below, and a more detailed description of the surveys and observations may be found in Section 3.1.3.1.5.3 (Farallon de Medinilla Specific Impacts) of the 2015 MITT Final EIS/OEIS. (Smith & Marx, 2016).

- **Natural causes of erosion.** Based on these direct observations of damage off the coast of FDM, the majority of disturbances to the seafloor sediments, substrates, and mass wasting of FDM can be attributed to typhoons and storm surges. Further, damage attributed to military training activities recovered within two to three years at the same rate of damage associated with natural phenomenon (Smith & Marx, 2016). As discussed in Section 3.10 (Terrestrial Species and Habitats), prior to the mid-1990s, the ordnance drops on FDM were not confined to designated impact zones, and there were no ordnance constraints in terms of net explosive weight. The vegetation loss on the island and subsequent erosion has likely decreased under current training constraints for FDM relative to the intensive range use over the decades prior to the mid-1990s.
- **Assessment of long-term impacts: military impacts.** Based on the dive surveys, there is no evidence that long-term adverse impacts on the nearshore environment have taken place as a result of military training activities. These findings are based on the number of detectable impacts (e.g., from visual observation during dive surveys), the size of those impacts, and the apparent recovery time (e.g., how long an ordnance fragment or physical damage is no longer visually apparent). Impacts on the physical environment clearly attributable to military training activities were noted in 2007, 2008, 2010, and 2012 (Smith & Marx, 2016). Indirect impacts, such as ordnance skipping or eroding off of FDM and rock and ordnance fragments blasted off of the island, were detected in every survey year. Dive surveys completed in 2005 noted that disturbed sites in 2004 showed no color differences with surrounding undamaged areas, and revealed new, small (less than 3 centimeters), scattered colonies of coral and crustose coralline algae. By 2006 and observed again through 2012, no visual evidence of abnormalities, or of damaged or diseased coral, could be detected (Smith et al., 2013). Further, no new submerged cliff blocks were observed between 2005 and 2012. Small-to-medium-sized fresh rock fragments (generally less than 1 foot [30 centimeters]) have been observed yearly, and are attributed to detonation impacts. In 2007, the first clear indication of a detonation of a bomb on the seafloor was observed. The impact area was measured to be approximately 100 square feet (9 square meters). During the subsequent survey in 2008, the impact area supported new growth of stony corals and crustose algae; by 2009, no trace of the disturbance could be detected by the surveyors (Smith & Marx, 2009). The vast majority of unexploded ordnance observed in the water lacked fins and tail assemblies, which indicates that the ordnance either skipped or ricocheted off of the island, or eroded or washed off of FDM at a later date (Smith & Marx, 2016).
- **Indicators of diminished water quality.** The dive surveys have looked for indicators of diminished water quality in waters surrounding FDM. For instance, high densities of macrobioeroders (e.g., boring sponges), bleaching of corals, surface lesions, or dead patches on stony corals or stony coral mucus production have been associated with sedimentation, pollutants, or other stressors that diminish water quality. Although a moderate bleaching event was noted in 2007, and a barnacle infestation was noted in 2012 (Smith et al., 2013), the bleaching event was regional and extended from southern Japan through the Mariana Islands and south through waters surrounding Palau, which suggests that it was not due to training events at FDM. In addition, subsequent surveys observed soft and fire corals had recovered completely, and 75 percent of the stony corals had recovered by 2008 (Smith & Marx, 2009, 2016). The dive surveys were not conducted in more recent years with bleaching events; however, the health of the marine ecosystem surrounding FDM was comparable to similar

habitats within the Mariana Archipelago, demonstrating that training activities occurring at FDM do not have an appreciable impact on the water quality (Smith & Marx, 2016).

### 3.1.2 Environmental Consequences

Section 3.1 (Sediments and Water Quality) of the 2015 MITT Final EIS/OEIS analyzed potential impacts of training and testing activities resulting from the following stressors: (1) explosives (in-air explosives and in-water explosives) and explosives byproducts, (2) metals, (3) chemicals other than explosives, and (4) a miscellaneous category of other materials. The 2015 MITT Final EIS/OEIS assessed the likelihood for these stressors to result in the following potential impacts on sediments and water quality:

- The potential release of materials into the water that subsequently disperse, react with seawater, or dissolve over time
- The potential for depositing materials on the ocean bottom and any subsequent interactions with sediments or the accumulation of such materials over time
- The potential for depositing materials on the ocean bottom and any subsequent interaction with the water column
- The potential for depositing materials on the ocean bottom and any subsequent disturbance of those sediments or their resuspension in the water column

This section evaluates how, and to what degree, potential impacts on sediments and water quality from stressors described in Section 3.0 (Introduction) may have changed since the analysis presented in the 2015 MITT Final EIS/OEIS was completed. Tables 2.5-1 and 2.5-2 in Chapter 2 (Description of Proposed Action and Alternatives) list the proposed training and testing activities, the number of times each event would be conducted annually, and the locations within the Study Area where the activity would typically occur under each alternative. The tables also present the same information for activities described in the 2015 MITT Final EIS/OEIS so that the proposed levels of training and testing under this supplement can be easily compared.

The Navy conducted a review of federal and state regulations and standards relevant to sediments and water quality, as well as a review of new literature pertaining to sediments and water quality that could inform the analysis presented in the 2015 MITT Final EIS/OEIS. Although additional information was found and described in Section 3.1.1 (Affected Environment), the new information does not indicate an appreciable change to the existing environmental conditions as described in the 2015 MITT Final EIS/OEIS. Thus, the analysis in the 2015 MITT Final EIS/OEIS remains valid. The analysis presented in this section also considers standard operating procedures, which are discussed in Section 2.3.3 (Standard Operating Procedures) of this Draft Supplemental EIS (SEIS)/OEIS, and mitigation measures that are described in Chapter 5 (Mitigation). These measures are not specifically designed to offset potential impacts on water resources; however, implementation of some of these measures designed for other resource areas discussed in this SEIS/OEIS would avoid or reduce potential impacts on sediments and water quality. For example, Table 5.4-1 (Seafloor Resource Mitigation Areas) lists several protective measures that avoid or reduce disturbance to corals and benthic habitats, as well as targeting and ordnance restrictions that would reduce runoff into FDM's nearshore habitats.

The most relevant new information used in this section is published by the Hawaii Undersea Military Munitions Assessment (HUMMA), a program administered by the DoD and the University of Hawaii at Manoa (Briggs et al., 2016; Edwards et al., 2016; Kelley et al., 2016; Koide et al., 2016; Silva & Chock, 2016; Tomlinson & De Carlo, 2016). The investigations completed as part of the program provide

quantitative information on the fate and transport of sea-disposed conventional munitions at a munitions dump site south of Oahu, including (1) the spatial extent and distribution of munitions; (2) the integrity of munitions casings; (3) whether munitions constituents could be detected in sediment, seawater, or animals near munitions; (4) whether constituent levels at munitions sites differed significantly from levels at reference control sites; (5) whether statistically significant differences in ecological population metrics could be detected between the two types of sites; and (6) whether munitions constituents or their derivatives potentially pose an unacceptable risk to human health.

### **3.1.2.1 Explosives and Explosives Byproducts**

Sources of explosives and explosives byproducts include the various munitions used during training and testing activities. Potential impacts of explosives and explosive byproducts were analyzed in Section 3.1.3.1 (Explosives and Explosives Byproducts) in the 2015 MITT Final EIS/OEIS, and that analysis remains valid.

Over 98 percent of residual explosive materials would result from ordnance failures (i.e., the munition fails to detonate and explosives remain in the casing). Ordnance failure rates for various munition types are shown in Table 3.1-4 in Section 3.1.3.1.3 (Ordnance Failure and Low-Order Detonations) of the 2015 MITT Final EIS/OEIS. The percentages for ordnance failure range from just below two percent to just over eight percent. As part of the HUMMA program, Briggs et al. (2016) sampled for explosive materials in sediments and marine invertebrates and fish, showing no detections of explosive residue chemical markers in the biological samples. In 2009, no explosive residues were located within sediments; however, in 2012, 2 of the 121 samples showed low concentrations (0.09 and 0.12 milligrams per kilogram) of an explosive residue compound, 4-nitrotoluene. These samples were collected within 50 centimeters of a munitions casing, with no detections further away from the casing (Briggs et al., 2016). Within the Study Area, ocean currents would quickly disperse leached explosive materials in the water column, and residual explosive materials would not result in water toxicity.

#### **3.1.2.1.1 Impacts from Explosives and Explosives Byproduct Stressors Under Alternative 1**

Under Alternative 1, the number of explosive munitions used during at-sea training and testing activities would decrease, compared to the number analyzed in the 2015 MITT Final EIS/OEIS (see Table 3.0-16); however, there would be an increase in the number of activities on FDM that use explosive ordnance (Table 3.0-19). The Navy conducted an analysis as part of this SEIS/OEIS to quantify the amount of ordnance used on FDM, in terms of net explosive weight, that would change compared to what was analyzed in the 2015 MITT Final EIS/OEIS. This analysis shows that the proposed increases in ordnance use on FDM would be less than 1 percent compared to levels analyzed previously.

This small increase on FDM under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS because (1) most of the explosives would be consumed during detonation; (2) the frequency of low-order detonations would be low, and therefore the frequency of releases of explosives would be low; and (3) the constituents of explosives would be subject to physical, chemical, and biological processes that would render the materials harmless or otherwise disperse them to undetectable levels. Neither state nor federal standards or guidelines would be violated. The impacts of unconsumed explosives on water and sediment quality would be long term, local, and negative. Chemical, physical, or biological changes in sediment or water quality would likely be measurable, but neither state nor federal standards or guidelines would be violated.

### **3.1.2.1.2 Impacts from Explosives and Explosives Byproduct Stressors Under Alternative 2**

As with Alternative 1, activities proposed under Alternative 2 would decrease the number of explosive munitions used during at-sea training and testing activities, compared to the number analyzed in the 2015 MITT Final EIS/OEIS (see Table 3.0-16) and increase the number on FDM. At-sea ordnance use under Alternative 2 would be greater than Alternative 1; however, the amount of ordnance use on FDM would be the same under Alternative 2 as with Alternative 1 (Table 3.0-19). Increases for at-sea activities under Alternative 2 as compared to Alternative 1 would have no appreciable change on the impact conclusions for explosives and explosives byproducts stressors as was presented in the 2015 MITT Final EIS/OEIS. Therefore, under Alternative 2, impacts on sediments and water quality from the use of explosives and generating explosives byproducts would be negligible.

### **3.1.2.1.3 Impacts from Explosives and Explosives Byproduct Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosives and explosives byproduct stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing at-sea training and testing activities would result in fewer explosives and explosive byproducts introduced into the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing at-sea training and testing activities under the No Action Alternative would lessen the potential for impacts on sediments and water quality resulting from explosives and explosive byproducts.

### **3.1.2.2 Metals**

Sources of metals introduced into the marine environment as part of training and testing activities include munitions and expended materials containing metals (i.e., lead, brass, manganese, copper, nickel, tungsten, chromium, molybdenum, vanadium, boron, selenium, columbium, or titanium). Since the publication of the 2015 MITT Final EIS/OEIS, the Navy has conducted a review of new literature pertaining to the potential impacts of metals on sediments and water quality. Although additional information was found, as described in the following paragraph, the new information does not indicate an appreciable change to the existing environmental conditions as described in the 2015 MITT Final EIS/OEIS.

As described in Section 3.1.3.1 (Explosives and Explosives Byproducts) of the 2015 MITT Final EIS/OEIS, sediment samples collected from World War II-era munitions disposal sites and heavily used Navy ranges show that metals are not impacting sediment quality despite longtime use and high concentrations of military munitions composed primarily of metal components (Briggs et al., 2016; Kelley et al., 2016; Koide et al., 2016; Smith & Marx, 2016; U.S. Department of the Navy, 2013). The concentration of munitions and other expended materials containing metals in any one location in the Study Area would be a small fraction of that from a munitions disposal site, a target island used for 45 years, or a water range in a river used for almost 100 years. Chemical, physical, or biological changes to sediments or water quality in the Study Area would not be detectable and would be similar to nearby areas without munitions or other expended materials containing metals. This conclusion is based on the following: (1) most of the metals are benign, and those of potential concern make up a small percentage of expended munitions and other metal objects; (2) metals released through corrosion would be diluted by currents or bound up and sequestered in adjacent sediments; (3) elevated concentrations of metals

in sediments would be limited to the immediate area around the expended material; and (4) the areas over which munitions and other metal components would be distributed are large.

#### **3.1.2.2.1 Impacts from Metal Stressors Under Alternative 1**

Under Alternative 1, the number of sources of metals that would be expended during training and testing would increase as compared to the 2015 MITT Final EIS/OEIS (see Table 3.0-14 through Table 3.0-17 and Table 3.0-19). Although the overall amount of metals introduced to the Study Area would increase, the analysis is not dependent on the amount of metals. Instead, the 2015 MITT Final EIS/OEIS analyzed whether or not the metals deposited from training and testing activities would impact sediments and water quality.

Since the publication of the 2015 MITT Final EIS/OEIS, the Navy has conducted a review of existing federal and local regulations and standards relevant to sediments and water quality, as well as a review of new literature pertaining to sediments and water quality. There is no new information that changes the basis of the conclusions presented for the potential impacts of metals on sediments and water quality. Therefore, the increases shown in Tables 2.5-1 and 2.5-2 for training and testing activities proposed under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

#### **3.1.2.2.2 Impacts from Metal Stressors Under Alternative 2**

Under Alternative 2, the number of sources of metals being expended would increase as compared to the 2015 MITT Final EIS/OEIS and Alternative 1 (see Tables 3.0-14 through Table 3.0-17 and Table 3.0-19). These increases would have no appreciable change on the impact conclusions for metals as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, under Alternative 2, impacts on sediments and water quality from activities that expend metals would be negligible.

#### **3.1.2.2.3 Impacts from Metal Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Metal stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing training and testing activities would result in fewer metals introduced into the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on sediments and water quality resulting from metals released during training and testing activities.

#### **3.1.2.3 Chemicals Other Than Explosives**

Chemicals other than explosives are associated with the following military expended materials: (1) solid-fuel propellants in missiles and rockets; (2) Otto Fuel II torpedo propellant and combustion byproducts; (3) polychlorinated biphenyls in target vessels used during sinking exercises; (4) other chemicals associated with explosive munitions; and (5) chemicals that simulate chemical warfare agents, referred to as "simulants."

Simulants: Simulants were not analyzed in the 2015 MITT Final EIS/OEIS. The Department of Defense uses compounds, referred to as simulants, as substitutes for chemical and biological warfare agents to test equipment intended to detect their presence. Simulants must have one or more characteristics of a real chemical or biological agent—size, density, or aerosol behavior—to effectively mimic the agent. Simulants must also pose a minimal risk to human health and the environment to be used safely in outdoor tests.

Simulants are selected using the following criteria: (1) safety to humans and the environment, and (2) the ability to trigger a response by sensors used in the detection equipment. Simulants would be relatively benign (e.g., low toxicity or effects potential) from a human health, safety, and environmental perspective. Exposure levels during testing activities would be well below concentrations associated with any adverse human health or environmental effects. The degradation products of simulants used during testing would also be harmless. Given these characteristics of simulants used during testing activities, it is reasonable to conclude that simulants would have no impact on sediments and water quality in the Study Area. Simulants are not analyzed further in this section.

#### **3.1.2.3.1 Impacts from Chemical Stressors Other than Explosives Under Alternative 1**

Under Alternative 1, the number of sources of chemicals other than explosives would increase as compared to the 2015 MITT Final EIS/OEIS (see Table 3.0-14 through Table 3.0-17 and Table 3.0-19).

The fate and transport of solid fuel propellants are described in Section 3.1.3.3.2 (Missile and Rocket Propellant – Solid Fuel) of the 2015 MITT Final EIS/OEIS. The analysis in the 2015 MITT Final EIS/OEIS concluded that, based on the small amount of residual propellant that would remain from training and testing activities using missiles or rockets, perchlorates would not occur in concentrations that would impact sediments and water quality in the Study Area. The changes in the number of missiles and rockets shown in Tables 2.5-1 and 2.5-2 for activities proposed under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

The fate and transport of Otto Fuel II torpedo propellant and combustion byproducts are described in Section 3.1.3.3.3 (Torpedo Propellant – Otto Fuel II and Combustion Byproducts) of the 2015 MITT Final EIS/OEIS. Otto Fuel II and its combustion byproducts would be released into the water column only in small amounts during combustion. Furthermore, all non-explosive torpedoes are typically recovered for reuse following training and testing activities, which removes any unconsumed fuel from the environment immediately after completion of the activity. Combustion byproducts of Otto Fuel II would be released into the water column, where they would dissolve, dissociate, or be dispersed and diluted. One combustion byproduct, hydrogen cyanide, does not normally occur in seawater; however, it is soluble in seawater and would be diluted to less than 1 micrograms per liter (1.0 part per billion)—below EPA-recommended concentrations (U.S. Environmental Protection Agency, 2010)—at a distance of approximately 18 feet from the center of the torpedo's path when first discharged. Additional dilution would occur thereafter, with the rate of dilution depending, in part, upon circulation in the water column in the vicinity of the discharge. The changes in the number of torpedoes shown in Tables 2.5-1 and 2.5-2 for activities proposed under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

The fate and transport of Polychlorinated Biphenyls (PCBs) are described in Section 3.1.3.3.4 (Polychlorinated Biphenyls in Target Vessels) of the 2015 MITT Final EIS/OEIS. Sinking exercises would decrease under Alternative 1 in this SEIS/OEIS and are therefore not analyzed further. Public comments, however, were received that concerned the potential resuspension of PCBs in the water column after

activities that use underwater explosives in Outer Apra Harbor. Figure 2.1-5 in Chapter 2 (Description of the Proposed Activities and Alternatives) shows the location of the Outer Apra Harbor Underwater Detonation (UNDET) site. The Navy's literature review found PCB measurements obtained by a University of Guam study in 1997 for PCB contamination within Apra Harbor (Denton et al., 1997). The location of the UNDET site in Outer Apra Harbor corresponds to a sediment sampling site that was considered by Denton et al. (1997) as within the "light" contamination range (1–10 nanograms/gram dry weight). PCB profiles, determined in sediments from Hotel Wharf and the Commercial Port area, closely resembled those of Aroclor 1254, a commercial PCB mixture that was once widely used as a dielectric fluid in electrical transformers (Denton et al., 1997). Another set of samples were collected in 2014 within Outer Apra Harbor. As part of this sampling regime, preliminary remediation goals were established for different types of PCBs. The location within Outer Apra Harbor that is used for underwater explosions did not exceed these preliminary remediation goal thresholds for PCBs (U.S. Department of the Navy, 2017). Because the same location is used for UNDET sites, the Navy avoids resuspension of PCBs from undisturbed benthic habitats where PCBs may have migrated. There is no information in the University of Guam study that changes the basis of the above findings. Therefore, based on the findings above, the changes in the numbers of UNDETs used within Outer Apra Harbor as shown in Tables 2.5-1 and 2.5-2 would have not appreciably changed the impacts that chemicals other than explosives would have on sediments and water quality.

The fate and transport of other chemicals associated with explosive munitions are described in Section 3.1.3.3.5 (Other Chemicals Associated with Ordnance) of the 2015 MITT Final EIS/OEIS. Residual chemical constituents associated with explosive munitions can remain in the environment after low-order (i.e., incomplete) detonations and in unconsumed explosives. These constituents, listed in Table 3.1-10 of the 2015 MITT Final EIS/OEIS, are in addition to the explosives contained in the munition. Lead azide, titanium compounds, perchlorates, barium chromate, and fulminate of mercury are not naturally constituents of seawater. Another residual constituent, lead oxide, is a rare, naturally occurring mineral (Agency for Toxic Substances and Disease Registry, 2007). As noted in Section 3.1.2.1 (Explosives and Explosives Byproducts), fewer explosive munitions would be used during training activities under Alternative 1 compared to the number of explosives proposed in the 2015 MITT Final EIS/OEIS. Some testing activities would use more explosive munitions, while others would use fewer. Based on the detailed analysis in Section 3.1.3.1 (Explosives and Explosion Byproducts) in the 2015 MITT Final EIS/OEIS and the summary of recent studies in Section 3.1.2.1 (Explosives and Explosives Byproducts) in this SEIS/OEIS, concentrations of chemical constituents associated with explosive munitions is expected to be localized to areas adjacent to the munition and similar to concentrations from nearby sites. The changes in the number of explosions shown in Tables 2.5-1 and 2.5-2 for activities proposed under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

#### **3.1.2.3.2 Impacts from Chemical Stressors Other than Explosives Under Alternative 2**

Under Alternative 2, the number of sources used that would generate chemicals other than explosives would increase as compared to Alternative 1 (see 3.0-14 through Table 3.0-17 and Table 3.0-19). As discussed in Alternative 1, increases as associated with Alternative 2 would have no appreciable change on the impact conclusions for chemicals other than explosives as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, under Alternative 2, impacts on sediments and water quality from activities that expend chemicals other than explosives would be negligible.

### **3.1.2.3.3 Impacts from Chemical Stressors Other than Explosives Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Chemical stressors other than explosives as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer releases of chemical stressors other than explosives into the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on sediments and water quality resulting from chemical stressors other than explosives.

### **3.1.2.4 Other Materials**

Other materials include marine markers and flares, chaff, towed and stationary targets, and miscellaneous components of other devices. These materials and components are made mainly of nonreactive or slowly reactive materials (e.g., glass, carbon fibers, and plastics), or they break down or decompose into benign byproducts (e.g., rubber, steel, iron, and concrete). Most of these objects would settle to the sea floor where they would (1) be exposed to seawater, (2) become lodged in or covered by seafloor sediments, (3) become encrusted (e.g., by rust) through oxidation, (4) dissolve slowly, or (5) be covered by marine organisms such as coral. Plastics may float or descend to the bottom, depending upon their buoyancy.

The various types of expended materials that would be used during training and testing activities are described in detail in Section 3.1.3.4 (Other Materials) in the 2015 MITT Final EIS/OEIS. The section describes the constituent components of marine markers, flares, and chaff as well as other items, and the fate and transport of those constituents in the marine environment. Pyrotechnic materials in marine markers and flares are largely consumed during use, and byproducts are released into the air. Chemical constituents of marine markers and flares are listed in Table 3.1-11 and the constituents of chaff are listed in Table 3.1-12 of the 2015 MITT Final EIS/OEIS.

#### **3.1.2.4.1 Impacts from Other Materials Under Alternative 1**

Under Alternative 1, the number of proposed training and testing activities that would introduce other materials, such as marine markers and flares, chaff, towed and stationary targets, and miscellaneous components would increase over levels analyzed previously in the 2015 MITT Final EIS/OEIS (see Table 3.0-17, Tables 3.0-20 through 3.0-24). Increases in training and testing activities under Alternative 1 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

#### **3.1.2.4.2 Impacts from Other Materials Under Alternative 2**

Under Alternative 2, the number of proposed training and testing activities that would introduce other materials, such as marine markers and flares, chaff, towed and stationary targets, and miscellaneous components would increase over levels analyzed previously in the 2015 MITT Final EIS/OEIS (see Table 3.0-17, Tables 3.0-20 through 3.0-24). There would also be increases under Alternative 2 in the number of training and testing activities that would likely introduce other materials into the environment, as compared to Alternative 1. As with Alternative 1, increases in training and testing activities proposed under Alternative 2 would have no appreciable change on the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

#### 3.1.2.4.3 Impacts from Other Materials Under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Other materials as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer releases of other materials within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on sediments and water quality resulting from plastics, marine markers, flares, and chaff released during training and testing activities.

#### 3.1.3 Public Scoping Comments

The public raised a number of issues during the scoping period in regard to sediments and water quality. The issues are summarized in the list below.

- **Public scoping comments concerning FDM.** Some commenters noted a lack of studies documenting the amount of ordnance debris and unexploded ordnance in waters surrounding FDM, while other comments requested that the Navy analyze potential loss of land mass associated with military training activities on FDM. The Navy has included a detailed summary of recent published studies that describe multi-year dive studies conducted by Smith and Marx (2016). The results of these surveys are included in Section 3.1.1.1.3 (Farallon de Medinilla) of this SEIS/OEIS. Throughout all dive surveys, the coral fauna at FDM were observed to be healthy and robust. The nearshore physical environment and basic habitat types at FDM have remained unchanged over the 13 years of survey activity. These conclusions are based on (1) a limited amount of physical damage, (2) very low levels of partial mortality and disease (less than 1 percent of all species observed), (3) absence of excessive mucus production, (4) good coral recruitment, (5) complete recovery by 2012 of the 2007 bleaching event, and (6) a limited number of macrobioeroders and an absence of invasive crown of thorns starfish (*Acanthaster planci*). These factors suggest that sedimentation that may result from military use of FDM is not sufficient as to adversely impact water quality, a conclusion substantiated by repeated dive surveys discussed above (Smith & Marx, 2016).
- **Public scoping comments regarding the potential loss of landmass through erosion of FDM from military use.** Some commenters have expressed concerns regarding erosion of FDM, and the potential loss of landmass. The U.S. military has used FDM as a bombing range since at least 1971. FDM's vegetation appears to have undergone significant changes since the island was leased by the DoD and the subsequent use for military training. The 2015 MITT Final EIS/OEIS compared historic aerial photographs to recent aerial imagery, which shows that the island has lost substantial forests over the decades, with the northern portion of the island with the most intact forest structure remaining on FDM (see Section 3.10, Terrestrial Species and Habitats, of the 2015 MITT Final EIS/OEIS). It is likely that the loss of vegetation over the past decades has accelerated erosion of soils and limestone weathering on the island. The current training activities that use ordnance are constrained in terms of ordnance type and target location (e.g., designated impact zones). These restrictions were put in place as part of past Section 7 ESA consultations with the U.S. Fish and Wildlife Service and would continue under the SEIS. While

these measures were specifically designed to protect ESA-listed species and habitats on FDM, the restrictions would likely reduce the rates of erosion experienced in previous decades on the island. In addition, since the 2015 MITT Final EIS/OEIS, the Navy has relocated vertical cliff targets (established on the western side of the island) to interior locations within impact zones. The target relocations were done to minimize impacts on seabird rookeries along the western side of the island. Smith and Marx (2016) also provided anecdotal observations of coral reefs surrounding the island over the course of multi-year dive surveys. These observations suggest healthy reef environments surrounding the island, without signs of sedimentation that would result from erosion of soils from the impact areas. In summary, the intensive bombing regimes of FDM in past decades likely resulted in the loss of forested areas on the island; such reductions in forests likely resulted in erosion of the upper plateau of the island. Current restrictions, however, confine the bombing activities to discrete impact zones located in the interior of the island, with additional restrictions on the types of ordnance allowed for use on the island, thereby reducing the potential for erosion and loss of land mass of FDM.

- **Public scoping comments concerning resuspension of PCBs in Outer Apra Harbor.** Some commenters were concerned about resuspension of PCBs in the water column resulting from underwater explosions within Outer Apra Harbor. Section 3.1.2.3 (Chemicals Other Than Explosives) of this SEIS/OEIS includes additional information on the potential for resuspension of PCBs in the water column, which includes sediment data collected from a site in close proximity to the Outer Apra Underwater Detonation Site. The potential for resuspension of PCBs in the water column is reduced because (1) the sediment samples collected by Denton et al. (1997) showed that this location is within the “light” concentration range (1–10 nanograms/gram dry weight), (2) additional sediment sampling from 2014 shows that the area where underwater detonations would occur contains sediments that do not exceed remediation goals for different types of PCBs, and (3) the Navy uses the same seafloor location for underwater explosions. Therefore, the Navy does not conduct this training activity in other areas of Apra Harbor identified as “moderate” or “high” concentrations. In addition, no new undisturbed benthic locations that are contaminated by PCBs would be used for underwater explosions.
- **Public scoping comments concerning general impacts on water quality in offshore marine environments.** Some commenters were concerned about the fate and transport of metal fragments as they are deposited in open ocean training locations. Section 3.1.3.2 (Metals) in the 2015 MITT Final EIS/OEIS describes the potential impacts of metals introduced into marine environments from training locations. Although Guam does not maintain screening standards for metals in sediments or water, the U.S. Environmental Protection Agency maintains “threshold” values for metals in marine environments (see Table 3.1-8 of the 2015 MITT Final EIS/OEIS). In 2014, the CNMI Bureau of Environmental and Coastal Quality established water quality standards, designating the coastal waters surrounding FDM as “Class A” waters, which are maintained for recreational and aesthetic use, with some allowable uses as long as it is compatible with the protection and propagation of fish, shellfish, and wildlife (Commonwealth of the Northern Mariana Islands Bureau of Coastal and Environmental Quality, 2014). Based on the multi-year dive surveys discussed above, there are no indications of adverse impacts on fish, shellfish, or wildlife within the coastal waters surrounding FDM, with the dive surveys showing healthy ecosystem functions and wildlife abundance within these waters. While no quantitative sampling for metals in training areas have been completed, there are a number of studies

conducted in marine training and testing locations that have attempted to measure metal content where military activities occur. In one study, the water was sampled for lead, manganese, nickel, vanadium, and zinc at a shallow bombing range in Pamlico Sound (state waters of North Carolina) immediately following a training event with non-explosive practice bombs. All water quality parameters tested, except nickel, were within the state limits. The nickel concentration was significantly higher than the state criterion, although the concentration did not differ significantly from the control site located outside the bombing range. The results suggest that bombing activities were not responsible for the elevated nickel concentrations (U.S. Department of the Navy, 2010). A recent study conducted by the U.S. Marine Corps sampled sediments and water quality for 26 different constituents related to munitions at several U.S. Marine Corps water-based training ranges. Metals included lead and magnesium. These areas were also used for bombing practice. No munitions constituents were detected above screening values used at the U.S. Marine Corps water ranges (U.S. Department of the Navy, 2010). A study by Pait et al. (2010) of previous Navy training areas at Vieques, Puerto Rico, found generally low concentrations of metals in marine sediments. Areas in which live ammunition and loaded weapons were used (“live-fire areas”) were also included in the analysis. Additional studies are summarized in the 2015 MITT Final EIS/OEIS. In no instance did metals exceed federal or state thresholds. It is unlikely that metals in sediments or the water column from military training activities would exceed federal thresholds in the Study Area, a conclusion that is consistent with other range locations and qualitative observations of ecosystem health surrounding FDM, as observed by Smith and Marx (2016).

## REFERENCES

- Agency for Toxic Substances and Disease Registry. (2007). *Toxicological Profile for Lead*. Atlanta, GA: U.S. Department of Health and Human Services.
- Briggs, C., S. M. Shjegstad, J. A. K. Silva, and M. H. Edwards. (2016). Distribution of chemical warfare agent, energetics, and metals in sediments at a deep-water discarded military munitions site. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 63–69.
- Commonwealth of the Northern Mariana Islands Bureau of Coastal and Environmental Quality. (2014). *Commonwealth of the Northern Mariana Islands Water Quality Standards*. Retrieved from <https://www.epa.gov/sites/production/files/2014-12/documents/northern-mariana-wqs.pdf>.
- Denton, G. R., H. R. Wood, L. P. Concepcion, H. G. Siegrist, V. S. Eflin, D. K. Narcis, and G. T. Pangelinan. (1997). *Analysis of In-place Contaminants in Marine Sediments from Four Harbor Locations on Guam*. Mangilao, Guam: Water and Environmental Research Institute of the Western Pacific, University of Guam.
- Edwards, M. H., S. M. Shjegstad, R. Wilkens, J. C. King, G. Carton, D. Bala, B. Bingham, M. C. Bissonnette, C. Briggs, N. S. Bruso, R. Camilli, M. Cremer, R. B. Davis, E. H. DeCarlo, C. DuVal, D. J. Fornari, I. Kaneakua-Pia, C. D. Kelley, S. Koide, C. L. Mah, T. Kerby, G. J. Kurras, M. R. Rognstad, L. Sheild, J. Silva, B. Wellington, and M. V. Woerkom. (2016). The Hawaii undersea military munitions assessment. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 4–13.
- Griesser, A., and C. Spillman. (2016). Assessing the skill and value of seasonal thermal stress forecasts for coral bleaching risk in the western Pacific. *Journal of Applied Meteorology and Climatology*, 55(7), 1565–1578.
- Harvey, C. (2016, 21 June 2018). 'I cried... right into my mask': Scientists say Guam's reefs have bleached four years straight. *The Washington Post*.
- Keener, V. W., S. B. Gingerich, and M. L. Finucane. (2015). *Climate Change in the Commonwealth of the Northern Mariana Islands*. Retrieved from [ftp://ftp.soest.hawaii.edu/coastal/USAPI\\_ClimateChangeBooklets/CNMI%20Climate%202016.pdf](ftp://ftp.soest.hawaii.edu/coastal/USAPI_ClimateChangeBooklets/CNMI%20Climate%202016.pdf).
- Kelley, C., G. Carton, M. Tomlinson, and A. Gleason. (2016). Analysis of towed camera images to determine the effects of disposed mustard-filled bombs on the deep water benthic community off south Oahu. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 34–42.
- Koide, S., J. A. K. Silva, V. Dupra, and M. Edwards. (2016). Bioaccumulation of chemical warfare agents, energetic materials, and metals in deep-sea shrimp from discarded military munitions sites off Pearl Harbor. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 53–62.
- MacDonald, J., and C. Mendez. (2005). *Unexploded Ordnance Cleanup Costs: Implications of Alternative Protocols* (MG-244-RC). Santa Monica, CA, Arlington, VA, and Pittsburgh, PA: RAND Corporation.
- Pait, A. S., A. L. Mason, D. R. Whitall, J. D. Christensen, and S. I. Hartwell. (2010). Chapter 5: Assessment of Chemical Contaminants in Sediments and Corals in Vieques. In L. J. Bauer & M. S. Kendall (Eds.), *An Ecological Characterization of the Marine Resources of Vieques, Puerto Rico* (pp. 101–150). Silver Spring, MD: National Oceanic and Atmospheric Administration.
- Raymundo, L. J., M. C. D. Malay, and A. N. Williams. (2016). *Final Report: Research on Stony Coral Health and Community Structure*. Mangilao, Guam: University of Guam Marine Laboratory.

- Raymundo, L. J., D. Burdick, V. A. Lapacek, R. J. Miller, and V. Brown. (2017). Anomalous temperatures and extreme tides: Guam staghorn *Acropora* succumb to a double threat. *Marine Ecology Progress Series*, 564, 47–55.
- Richardson, K., D. Haynes, A. Talouli, and M. Donoghue. (2016). Marine pollution originating from purse seine and longline fishing vessel operations in the Western and Central Pacific Ocean, 2003–2015. *Ambio*, 46(2), 190–200.
- Silva, J. A. K., and T. Chock. (2016). Munitions integrity and corrosion features observed during the HUMMA deep-sea munitions disposal site investigation. *Deep-Sea Research I*, 14–24.
- Smith, S. H., and D. E. Marx, Jr. (2009). *Assessment of Near Shore Marine Resources at Farallon De Medinilla: 2006, 2007 and 2008*. Captiol Hill, Commonwealth of the Northern Mariana Islands: Pacific Division, Naval Facilities Engineering Command.
- Smith, S. H., J. Marx, D. E., and L. H. Shannon. (2013). *Calendar Year 2012 Assessment of Near Shore Marine Resources at Farallon de Medinilla, Commonwealth of the Northern Mariana Islands*. Port Hueneme, CA: U.S. Department of the Navy.
- Smith, S. H., and D. E. Marx, Jr. (2016). De-facto marine protection from a Navy bombing range: Farallon de Medinilla, Mariana Archipelago, 1997 to 2012. *Marine Pollution Bulletin*, 102(1), 187–198.
- Tomlinson, M. S., and E. H. De Carlo. (2016). Occurrence and possible sources of arsenic in seafloor sediments surrounding sea-disposed munitions and chemical agents near Oahu, Hawaii. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 70–84.
- U.S. Army Corps of Engineers. (2007). *Explosives residues resulting from the detonation of common military munitions: 2002–2006*.
- U.S. Department of the Navy. (2010). *Water Range Assessment for the VACAPES Range Complex*. Norfolk, VA: Parsons.
- U.S. Department of the Navy. (2013). *Military Expended Material Shrimp Fisheries Study in the U.S. South Atlantic and Eastern Gulf of Mexico*. Newport, RI: Naval Undersea Warfare Center Division.
- U.S. Department of the Navy. (2015). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2017). *Preliminary Apra Harbor Sediment Data Package*. Washington, DC: U.S. Department of the Navy.
- U.S. Environmental Protection Agency. (2010). *Water Quality Criteria: Suspended and Bedded Sediments*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Marine Corps. (2014). *Environmental Assessment and Finding of No Significant Impact: Reconfiguration and Construction of Small Arms Ranges at the Ulupau Range Training Facility, Marine Corps Base Hawaii Kaneohe Bay*. Fort Collins, CO: Center for Environmental Management of Military Lands.

---

---

## 3.2 Air Quality



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|         |                                  |       |
|---------|----------------------------------|-------|
| 3.2     | Air Quality .....                | 3.2-1 |
| 3.2.1   | Affected Environment.....        | 3.2-1 |
| 3.2.1.1 | Climate of the Study Area.....   | 3.2-1 |
| 3.2.1.2 | Regional Emissions .....         | 3.2-2 |
| 3.2.1.3 | Existing Air Quality.....        | 3.2-2 |
| 3.2.2   | Environmental Consequences ..... | 3.2-2 |
| 3.2.2.1 | Criteria Pollutants .....        | 3.2-2 |
| 3.2.2.2 | Hazardous Pollutants.....        | 3.2-4 |
| 3.2.2.3 | Greenhouse Gases .....           | 3.2-5 |
| 3.2.3   | Public Scoping Comments .....    | 3.2-5 |

**List of Figures**

There are no figures in this section.

**List of Tables**

|  |       |
|--|-------|
| Table 3.2-1: Baseline Pollutant Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast from Aircraft, Vessels, and Ordnance (tpy) .....      | 3.2-3 |
| Table 3.2-2: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast Under Alternative 1 from Aircraft, Vessels, and Ordnance (tpy) .. | 3.2-4 |
| Table 3.2-3: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast Under Alternative 2 from Aircraft, Vessels, and Ordnance (tpy) .. | 3.2-4 |
| Table 3.2-4: Annual Greenhouse Gas Emissions Under All Three Alternatives .....  | 3.2-5 |

This page intentionally left blank.

## 3.2 Air Quality

The purpose of this section is to supplement the analysis of impacts on Air Quality presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon de Medinilla (FDM). Information presented in the 2015 Final MITT EIS/OEIS that remains valid is noted as such and referenced in the appropriate sections. Any new or updated information describing the affected environment and analysis of impacts on Air Quality associated with the Proposed Action is provided in this section. Comments received from the public during scoping related to Air Quality are addressed in Section 3.2.3 (Public Scoping Comments).

This analysis is complicated because the emitted emissions are scattered widely over multiple regions, covering thousands of square miles of ocean from the sea surface and up to over a half mile in altitude. Therefore, the analysis makes use of screening thresholds. Screening thresholds are compared against the change in annual emissions that an action would have from the baseline. Changes in emissions below a screening threshold in a region or air basin are presumed to have no significant impact on the human environment. Changes in emissions above the thresholds may be significant, such that they require a harder look to make such a determination. For purposes of analysis, screening thresholds used in this analysis are derived from legal standards that are either legislated or contained in regulations promulgated by expert agencies with the input of the public and scientific community. These are conservative as they have been developed mainly for stationary sources or individual projects. They also represent potential cumulative impact(s) in that the thresholds are smaller in areas with degraded air in order to accommodate degradation from past and present actions as well as future progress toward attainment.

The 2015 MITT Final EIS/OEIS used the General Conformity Rule's *de minimis* thresholds as a screening factor because certain training and testing activities occurred within the nonattainment areas of Piti and Tanguisson. However, this Supplemental EIS (SEIS)/OEIS does not address any activities that occur within nonattainment areas. Therefore, the General Conformity Rule is not applicable to the Proposed Action, and thereby the *de minimis* thresholds that were used as screening criterion in the 2015 MITT Final EIS/OEIS are no longer pertinent. This SEIS/OEIS will instead utilize the thresholds established by the Prevention of Significant Deterioration (PSD) program as screening factors. PSD thresholds typically apply to the construction or modification of major stationary sources within nonattainment areas and are in place to ensure that a project would not cause a violation of the National Ambient Air Quality Standards (NAAQS) within an attainment area. Although the Proposed Action does not consist of any stationary sources, the PSD thresholds are more applicable than *de minimis* thresholds due to the attainment status of the region. In addition to criteria pollutants, greenhouse gas emissions were analyzed in this SEIS/OEIS by illustrating their cumulative contribution to climate change.

### 3.2.1 Affected Environment

#### 3.2.1.1 Climate of the Study Area

Climate in the MITT Study Area was discussed in detail in the 2015 MITT Final EIS/OEIS. The climate within the Study Area has not changed since the publication of the 2015 MITT Final EIS/OEIS (Climatemps.com, 2017).

### 3.2.1.2 Regional Emissions

Regional emissions have changed since the publication of the 2015 MITT Final EIS/OEIS. Guam and Saipan still contain the majority of the stationary sources of air pollutants within the Study Area. The largest point source emitters for air pollutants were the power-generating facilities at Piti and Tanguisson. However, the power-generating facility at Tanguisson has been retired since then and an explosion and fire at the power-generating facility in Piti has left two turbines inoperable. This has reduced the amount of pollutants being released into the atmosphere from manmade sources. In addition to anthropogenic sources, volcanic activity within the Study Area naturally contributes to sulfur dioxide concentrations in the region.

### 3.2.1.3 Existing Air Quality

As noted in the 2015 MITT Final EIS/OEIS, Guam and the Commonwealth of the Northern Mariana Islands, including Farallon de Medinilla, meet all national and local ambient air quality standards except for the areas of Piti and Tanguisson, which are in nonattainment of the 1971 sulfur dioxide primary NAAQS (U.S. Environmental Protection Agency, 2017). These nonattainment areas extend in a circle with a radius of 2.2 miles from the power-generating facilities. However, the retirement of the Tanguisson facility and reduction in functionality of the Piti facility have decreased pollutant emissions and could potentially affect the attainment status for these areas. In general, the islands are considered to have excellent ambient air quality due to geographic isolation and favorable climate. Consistent winds and rain help to remove and carry away pollutants from the islands.

## 3.2.2 Environmental Consequences

The 2015 MITT Final EIS/OEIS analyzed potential impacts on air quality resulting from proposed training and testing activities. This supplemental analysis will update and consider changes to air quality resulting from proposed changes to training and testing activities conducted at sea and on FDM (see Tables 2.5-1 and 2.5-2). This SEIS/OEIS only addresses training and testing activities that would occur at-sea or on Farallon de Medinilla, therefore there is no discussion of any activities that occur within the nonattainment areas surrounding the Piti and Tanguisson facilities. Any activities associated with the 2015 MITT Final EIS/OEIS that currently occur within these nonattainment areas would not be affected by the Proposed Action. Therefore, all activities associated with this SEIS/OEIS are considered to occur in attainment areas that meet the NAAQS for all criteria pollutants. For these reasons and the reasons stated above, the PSD thresholds of 250 tons per year for criteria pollutants are used as a screening factor to determine the extent and likelihood of impacts that could arise from implementation of the Proposed Action.

### 3.2.2.1 Criteria Pollutants

Estimated emissions are compared against baseline emissions (Table 3.2-1) to determine whether PSD thresholds are exceeded for criteria pollutants. While most of the emissions would be off shore, only emissions that would be released below 3,000 feet elevation and within 3 nautical miles (NM) of the coastline are analyzed for their impacts on ambient air quality. Pollutants that would be emitted more than 3 NM offshore would be intermittent and distributed across a very large area of ocean, and would not be concentrated in any one area. Therefore, pollutants emitted beyond 3 NM are expected to disperse sufficiently enough to not be detectable, meaning that these emissions would have very little effect on the ambient air quality over the ocean and almost no impact on the ambient air quality over Guam or the Commonwealth of the Northern Mariana Islands. Emissions calculations for the Baseline, Alternative 1, and Alternative 2 can be seen in Appendix D (Air Quality Emissions Calculations). Baseline

emissions are derived from those presented in Alternative 1 of the 2015 MITT Final EIS/OEIS (Table 3.2-7). However, this SEIS/OEIS only addresses at-sea activities and activities occurring at FDM, whereas the 2015 MITT Final EIS/OEIS addressed those activities in addition to land-based activities. Therefore, only those activities that are covered under this SEIS/OEIS are considered in the baseline emissions.

**Table 3.2-1: Baseline Pollutant Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast from Aircraft, Vessels, and Ordnance (tpy)**

| Criteria Pollutant                | NO <sub>x</sub> | CO  | VOC | SO <sub>x</sub> | PM <sub>10</sub> | PM <sub>2.5</sub> |
|-----------------------------------|-----------------|-----|-----|-----------------|------------------|-------------------|
| Aircraft                          | 117             | 170 | 24  | 8               | 49               | 44                |
| Vessels                           | 365             | 41  | 97  | 288             | 54               | 48                |
| Ordnance                          | 0               | 0   | 0   | 0               | 0                | 0                 |
| Total of At-Sea and FDM Emissions | 482             | 211 | 121 | 296             | 103              | 92                |

Notes: NO<sub>x</sub> = nitrogen oxide, CO = carbon monoxide, VOC = volatile organic compounds, SO<sub>x</sub> = sulfur oxide, PM<sub>10</sub> = particulate matter less than 10 microns, PM<sub>2.5</sub> = particulate matter less than 2.5 microns, FDM = Farallon de Medinilla; tpy = tons per year. Baseline emissions are derived from those presented in Alternative 1 of the 2015 MITT Final EIS/OEIS.

#### 3.2.2.1.1 Alternative 1

Under Alternative 1, estimated pollutant emissions from aircraft, vessels, and ordnance would increase, as shown in Table 3.2-2. Criteria pollutants emitted in the Study Area within territorial waters could be transported ashore but would not affect the attainment status of the relevant air quality control regions. The amounts of air pollutants emitted in the Study Area and subsequently transported ashore would be minor because (1) the pollutants are emitted over large areas (i.e., the Study Area is an area source) and mostly beyond 3 NM, (2) the distances the air pollutants would be transported are often large, and (3) the pollutants would be substantially dispersed during transport. The criteria pollutants emitted over non-territorial waters within the Study Area would be dispersed over vast areas of open ocean and thus would not cause significant harm to environmental resources in those areas.

The changes in emissions produced under Alternative 1 above baseline conditions are compared against the PSD threshold of 250 tons per year to determine whether implementation of the Proposed Action would result in significant impacts. As shown in Table 3.2-2, Alternative 1 would not cause pollutant emissions to exceed the PSD threshold of 250 tons per year for any of the criteria pollutants, meaning this alternative would not have a significant impact on ambient air quality.

**Table 3.2-2: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast Under Alternative 1 from Aircraft, Vessels, and Ordnance (tpy)**

| Criteria Pollutant          | NO <sub>x</sub> | CO        | VOC       | SO <sub>x</sub> | PM <sub>10</sub> | PM <sub>2.5</sub> |
|-----------------------------|-----------------|-----------|-----------|-----------------|------------------|-------------------|
| Alternative 1 Emissions     | 547             | 272       | 169       | 313             | 123              | 111               |
| Baseline Emissions          | 470             | 194       | 119       | 295             | 99               | 88                |
| <b>Changes in Emissions</b> | <b>77</b>       | <b>78</b> | <b>50</b> | <b>18</b>       | <b>24</b>        | <b>23</b>         |
| PSD Thresholds              | 250             | 250       | 250       | 250             | 250              | 250               |

Notes: NO<sub>x</sub> = nitrogen oxide, CO = carbon monoxide, VOC = volatile organic compounds, SO<sub>x</sub> = sulfur oxide, PM<sub>10</sub> = particulate matter less than 10 microns, PM<sub>2.5</sub> = particulate matter less than 2.5 microns, PSD = prevention of significant deterioration, tpy = tons per year

### 3.2.2.1.2 Alternative 2

Under Alternative 2, there would be a slight increase in annual air emissions from Alternative 1 (Table 3.2-3). However, the change in emissions would not exceed the PSD thresholds and would not affect the attainment status of the relevant air quality control regions.

**Table 3.2-3: Annual Emissions for At-Sea and FDM Training and Testing Activities that Occur Within 3 Nautical Miles of the Coast Under Alternative 2 from Aircraft, Vessels, and Ordnance (tpy)**

| Criteria Pollutant      | NO <sub>x</sub> | CO        | VOC       | SO <sub>x</sub> | PM <sub>10</sub> | PM <sub>2.5</sub> |
|-------------------------|-----------------|-----------|-----------|-----------------|------------------|-------------------|
| Alternative 2 Emissions | 548             | 273       | 169       | 313             | 123              | 111               |
| Baseline Emissions      | 470             | 194       | 119       | 295             | 99               | 88                |
| <b>Difference</b>       | <b>78</b>       | <b>79</b> | <b>50</b> | <b>18</b>       | <b>24</b>        | <b>23</b>         |
| PSD Thresholds          | 250             | 250       | 250       | 250             | 250              | 250               |

Notes: NO<sub>x</sub> = nitrogen oxide, CO = carbon monoxide, VOC = volatile organic compounds, SO<sub>x</sub> = sulfur oxide, PM<sub>10</sub> = particulate matter less than 10 microns, PM<sub>2.5</sub> = particulate matter less than 2.5 microns, PSD = Prevention of Significant Deterioration, tpy = tons per year

### 3.2.2.1.3 No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Therefore, implementation of the No Action Alternative would mean that the emissions shown in Table 3.2-1 would no longer be produced, resulting in improved air quality.

### 3.2.2.2 Hazardous Pollutants

These emissions are typically one or more orders of magnitude smaller than concurrent emissions of criteria air pollutants, and only become a concern when large amounts of fuel, explosives, or other materials are consumed during a single activity or in one location. Hazardous air pollutants are analyzed qualitatively in relation to the prevalence of the sources emitting these pollutants during training and testing activities. Mobile sources operating as a result of the Proposed Action would be functioning intermittently over a large area and would produce negligible ambient hazardous air pollutants in a

localized area not located near any publicly accessible areas. For these reasons, hazardous air pollutants are not further evaluated in the analysis.

**3.2.2.3 Greenhouse Gases**

Table 3.2-4 summarizes the greenhouse gas emissions that would be generated under baseline conditions, Alternative 1, and Alternative 2. Greenhouse gas emissions would increase from the baseline by approximately 20 percent under Alternative 1 and Alternative 2. Since greenhouse gases are relevant in a global scope, they are analyzed based on the extent to which they would contribute to climate change. Implementation of Alternative 2 would generate approximately 0.0133 percent of the United States annual greenhouse gas emissions, which is less than a 0.0025 percent increase from baseline contributions. This minor increase is not expected to significantly affect the global climate.

**Table 3.2-4: Annual Greenhouse Gas Emissions Under All Three Alternatives**

|                      | Annual Greenhouse Gas Emissions (metric tons per year) |                  |                 |                   |
|----------------------|--|------------------|-----------------|-------------------|
|                      | CO <sub>2</sub>  | N <sub>2</sub> O | CH <sub>4</sub> | CO <sub>2</sub> e |
| Baseline Emissions   | 704,268  | 23               | 20              | 711,764           |
| Alternative 1        | 854,586  | 28               | 24              | 863,682           |
| Alternative 2        | 855,700  | 28               | 24              | 864,807           |
| Nationwide Emissions |  |                  |                 | 6,511,000,000     |

Notes: CO<sub>2</sub> = carbon dioxide, N<sub>2</sub>O = nitrous oxide, CH<sub>4</sub> = methane, CO<sub>2</sub>e = carbon dioxide equivalent

**3.2.3 Public Scoping Comments**

The public did not raise any issues during the scoping period in regard to air quality.

## **REFERENCES**

Climateemps.com. (2017). *Guam Climate & Temperature*. Retrieved from <http://www.mariana-islands.climateemps.com/>.

U.S. Environmental Protection Agency. (2017). *Nonattainment Areas for Criteria Pollutants (Green Book)*. Retrieved from <https://www.epa.gov/green-book>.

---

---

## 3.3 Marine Habitats



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

3.3 Marine Habitats ..... 3.3-1  
3.3.1 Affected Environment..... 3.3-1  
3.3.1.1 Existing Conditions..... 3.3-1  
3.3.2 Environmental Consequences ..... 3.3-3  
3.3.2.1 Explosive Stressors..... 3.3-3  
3.3.2.2 Physical Disturbance and Strike Stressors ..... 3.3-4  
3.3.3 Public Scoping Comments ..... 3.3-6

**List of Figures**

Figure 3.3-1: Nearshore Marine Habitats around Saipan ..... 3.3-2

**List of Tables**

There are no tables in this section.

This page intentionally left blank.

### **3.3 Marine Habitats**

#### **3.3.1 Affected Environment**

The purpose of this section is to supplement the analysis of impacts on Marine Habitats presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon De Medinilla. New information made available since the publication of the 2015 MITT Final EIS/OEIS is included below to better understand potential stressors and impacts on the nonliving (abiotic) Marine Habitats resulting from training and testing activities. Information presented in the 2015 MITT Final EIS/OEIS that remains valid is noted as such and referenced in the appropriate sections (U.S. Department of the Navy, 2015). Any new or updated information describing the affected environment and analysis of impacts on Marine Habitats associated with the Proposed Action is provided in this section. Other necessary habitats for living resources, including those that form biotic habitats such as aquatic plant beds and coral reefs, are discussed in other sections (e.g., Section 3.7, Marine Vegetation, and Section 3.8, Marine Invertebrates). Comments received from the public during scoping related to Marine Habitats are addressed in Section 3.3.3 (Public Scoping Comments).

##### **3.3.1.1 Existing Conditions**

The information and analysis on marine habitat types (i.e., soft shores, rocky shores, vegetated shores, aquatic beds, soft bottoms, hard bottoms, and artificial structures) presented in the 2015 MITT Final EIS/OEIS has not substantially changed and remains valid. In 2017, Kendall et al. (2017) mapped the benthic habitat of Saipan Lagoon. This new data was taken into consideration during the development of this Supplemental EIS (SEIS)/OEIS and is shown in Figure 3.3-1. After reviewing this data, the information and analysis on marine habitat types (i.e., soft shores, rocky shores, vegetated shores, aquatic beds, soft bottoms, hard bottoms, and artificial structures) presented in the 2015 MITT Final EIS/OEIS has not substantially changed and remains valid. The majority of the MITT Study Area lies within open-ocean areas. Relatively little of the Study Area includes intertidal and shallow subtidal areas in U.S. territory waters, where numerous habitats are exclusively present (e.g., salt/brackish marsh, mangrove, coral reefs, and seagrass beds). Intertidal abiotic habitats (e.g., beaches, tidal deltas, mudflats, rocky shores) are addressed only where intersections with military training and testing activities are reasonably likely to occur. Impacts on the water column are analyzed in Section 3.1 (Sediments and Water Quality). In addition, since the publication of the 2015 MITT Final EIS/OEIS, no critical habitat has been designated that needs to be considered here. Essential Fish Habitat (EFH) is discussed in Section 6.1.3, Magnuson-Stevens Fishery Conservation and Management Act, and is not discussed further in this section.



### 3.3.2 Environmental Consequences

The 2015 MITT Final EIS/OEIS considered training and testing activities that currently occur in the Study Area and considered potential stressors related to marine habitats. The stressors applicable to marine habitats in the Study Area for this SEIS/OEIS are the same stressors considered in the 2015 MITT Final EIS/OEIS.

- Explosive (in-air explosions and in-water explosions)
- Physical disturbance and strike (vessels, in-water devices, military expended materials, and seafloor devices)

This section evaluates how and to what degree potential impacts on marine habitats from stressors described in Section 3.0.1 (Overall Approach to Analysis) may have changed since the analysis presented in the 2015 MITT Final EIS/OEIS was completed. Proposed training and testing activities, the number of times each event would be conducted annually, and the locations within the Study Area where the activity would typically occur under each alternative are presented in Tables 2.5-1 and 2.5-2 in Chapter 2 (Description of Proposed Action and Alternatives). Information for training and testing activities proposed in the 2015 MITT Final EIS/OEIS are also included for comparison purposes.

The analysis presented in this section also considers standard operating procedures, which are discussed in Section 2.3.3 (Standard Operating Procedures) of this Draft SEIS/OEIS, and mitigation measures that are described in Chapter 5 (Mitigation). The Navy would implement these measures to avoid or reduce potential impacts on marine habitats from stressors associated with the proposed training and testing activities. Marine habitats in the remainder of this section will be referred to as marine substrates to reflect the subset of marine habitats being evaluated, similar to the 2015 MITT Final EIS/OEIS analysis.

#### 3.3.2.1 Explosive Stressors

As stated in the 2015 MITT Final EIS/OEIS, underwater detonations that occur on or near the bottom are the only explosive stressors that would impact marine substrates. All other explosive stressors (e.g., gunnery exercises, missile exercises, and air-to-surface rockets) used during training and testing activities occur on the water surface or in the water column and would not impact marine substrates. Underwater detonations that occur on or near the bottom are primarily used during various mine warfare training activities. The impacts of in-water explosions vary with the bottom substrate type. As stated in the 2015 MITT Final EIS/OEIS, mine warfare training and testing activities utilizing bottom placed detonations would only occur in the existing mine warfare underwater detonation areas at Piti, Agat, and Outer Apra Harbor. Cobble, rocky reef, and other hard bottom habitat may be scattered throughout the area, but those areas would be avoided during training and testing to the maximum extent practicable.

##### 3.3.2.1.1 Impacts from Explosive Stressors Under Alternative 1

Under Alternative 1, underwater detonations associated with training activities would increase for underwater demolition qualification/certification (Table 2.5-1). However, these activities would continue to occur in the same areas at the Agat Bay site, Piti, and Outer Apra Harbor sites, and would have no appreciable change in the impact analysis or conclusions for explosive stressors as presented in the 2015 MITT Final EIS/OEIS. There would be no increases in underwater detonations associated with testing activities. Therefore, the analysis in the 2015 MITT Final EIS/OEIS remains valid.

Mitigation measures will help the Navy avoid or reduce impacts on seafloor resources (including shallow-water coral reefs, live hard bottom, artificial reefs, and shipwrecks) from explosives during applicable activities, as described in Section 5.4.1 (Mitigation Areas for Seafloor Resources).

#### **3.3.2.1.2 Impacts from Explosive Stressors Under Alternative 2**

Under Alternative 2, the number of explosives used during training and testing events are proposed to increase compared to Alternative 1 described in this SEIS/OEIS (Table 3.0-7). Under Alternative 2, proposed increases would have no appreciable change on the impact conclusions for explosive stressors as summarized above under Alternative 1 because these activities would continue to occur in the same designated areas. Mitigation measures will help the Navy avoid or reduce impacts on seafloor resources (including shallow-water coral reefs, live hard bottom, artificial reefs, and submerged cultural resources) from explosives during applicable activities, as described in Section 5.4.1 (Mitigation Areas for Seafloor Resources).

#### **3.3.2.1.3 Impacts from Explosive Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosive stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer explosive stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for explosive impacts on marine habitat, but would not measurably improve the overall distribution or abundance of marine habitat.

#### **3.3.2.2 Physical Disturbance and Strike Stressors**

Bottom substrates could be disturbed by vessels (i.e., during amphibious landings and only in amphibious landing areas) and in-water devices, military expended materials, seafloor devices used for training and testing activities, and from walking, standing, or swimming in the nearshore waters during activities such as raids and assaults. Disturbance from vessel strikes would be avoided or reduced to the maximum extent practicable by taking precautions to avoid or reduce impacts on bottom substrates such as only performing amphibious landings in amphibious landing areas, at high tide, and where there are clear routes. Raids and assaults are planned to occur in areas that are primarily soft-bottom sandy habitat. Due to the nature of high-energy surf and shifting sands in these areas, ocean approaches would not be expected to affect marine habitats.

Seafloor devices would be located in areas that would be primarily soft-bottom and previously disturbed habitat to the greatest extent practical. The impact of seafloor devices on marine habitats is unlikely because these activities would occur over soft-bottom sediment, the items used in nearshore areas have a small footprint, and the items are retrieved. These potential impacts to bottom substrates would be minimal in size and temporary (recovery in days to weeks) to short term (recovery in weeks up to three years) in duration. Once on the seafloor, military expended material would be buried by sediment, corroded from exposure to the marine environment, or colonized by benthic organisms. As stated in the 2015 MITT Final EIS/OEIS, impacts of physical disturbance or strike resulting from training and testing activities on biogenic soft bottom (e.g., seagrasses, macroalgae, etc.) and hard bottom (e.g., corals,

sponges, tunicates, oysters, mussels, macroalgae, etc.) substrates are discussed in Sections 3.7 (Marine Vegetation) and 3.8 (Marine Invertebrates), respectively.

#### **3.3.2.2.1 Impacts from Physical Disturbance and Strike Stressors Under Alternative 1**

Under Alternative 1, there would be a slight increase in the use of towed in-water devices (Table 3.0-13). The increase in the number of in-water devices is unlikely to change the impact conclusion presented in the 2015 MITT Final EIS/OEIS. As stated in the 2015 MITT Final EIS/OEIS, the impact of vessels and in-water devices on marine habitats would remain inconsequential because (1) vessel and in-water activities that could come into contact with marine substrates would be located in previously disturbed areas (i.e., nearshore shallow waters), (2) military expended materials could be colonized by benthic organisms, and (3) seafloor devices would be used in predominantly soft bottom previously disturbed areas and therefore would not be expected to affect marine substrates.

Amphibious assault and raid activities that could involve walking, standing, and swimming from nearshore waters to shore would not increase under Alternative 1 on Tinian or Guam within the Mariana Islands Range Complex. These activities would cause minor and temporary increases in suspended sediments in soft bottom habitats, similar to impacts that occur on beaches that are open to the public (i.e., where people walk around and swim). Hard bottom substrates would be impacted by walking and standing in cobble-laid or reef areas. Contact with hard bottom substrate in nearshore waters, such as coral reefs, would be avoided or reduced to the greatest extent possible.

Under Alternative 1, the number of military expended materials used for training and testing events that have the potential to impact marine habitats would generally increase (see Tables 3.0-14 through 3.0-17). Military expended materials are very small relative to marine habitat and would not change the quality or type of habitat present. Therefore, these increases are not expected to pose a risk to marine habitats.

Under Alternative 1, the number of seafloor devices used in shallow-water habitats during training and testing events would decrease slightly from the number presented in the 2015 MITT Final EIS/OEIS (Table 3.0-18). Seafloor devices would pose a negligible risk to marine habitat for the same reason described above for military expended materials.

Any impacts on marine habitats incurred by vessel movements and in-water devices or military expended materials to soft bottom substrates would be minimal and temporary. Physical disturbance and strike of live hard bottom substrates would be permanent but minimal, and would be avoided or reduced through implementation of standard operating procedures as described in Section 2.3.3 (Standard Operating Procedures) and mitigation measures as described in Section 5.4.1 (Mitigation Areas for Seafloor Resources).

In addition, potential impacts on bottom substrates would be localized and temporary (recovery in days to weeks) to short-term (recovery in weeks up to three years) in duration. Artificial structures should not be adversely affected by the use of seafloor devices.

Mitigation measures will help the Navy avoid or reduce impacts on shallow-water coral reefs, live hard bottom, artificial reefs, and submerged cultural resources from precision anchoring and military expended materials during applicable activities, as described in Section 5.4.1 (Mitigation Areas for Seafloor Resources).

### 3.3.2.2.2 Impacts from Physical Disturbance and Strike Stressors Under Alternative 2

Under Alternative 2, the combined number of proposed training and testing events involving vessels and in-water devices (Table 3.0-12 and Table 3.0-13) are proposed to increase slightly from those presented in the 2015 MITT Final EIS/OEIS. Military expended materials (Table 3.0-14, Table 3.0-15, and Table 3.0-16) combined are proposed to increase, and seafloor devices (Table 3.0-18) are proposed to decrease slightly from the number in the 2015 MITT Final EIS/OEIS. Proposed increases in some physical disturbance and strike stressors such as military expended materials could increase the impact risk on marine habitats but does not appreciably change the analysis or impact conclusions presented in the 2015 MITT Final EIS/OEIS.

### 3.3.2.2.3 Impacts from Physical Disturbance and Strike Stressors Under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Physical disturbance and strike stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer physical disturbance and strike stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for physical disturbance and strike impacts on marine habitat, but would not measurably improve the overall distribution or abundance of marine habitat.

### 3.3.3 Public Scoping Comments

The public raised a number of issues during the scoping period in regard to marine habitats. The issues are summarized in the list below.

- **Impact of unexploded ordnance on the ocean floor** – Unexploded ordnance is not part of the military expended materials proposed to be expended under the Proposed Action.
- **Destruction of habitat** – Proposed training and testing activities are not expected to cause the destruction of marine habitat in the Study Area. Any impacts on marine habitats incurred by vessel movements and in-water devices or military expended materials to soft- and hard bottom substrates would be minimal. Explosive impacts on hard bottom habitat are determined to be permanent but minimal throughout the Study Area because they only occur in three existing mine warfare in-water detonation locations, which are made up mostly of soft bottom substrate (Piti, Agat, and Outer Apra Harbors). Furthermore, the implementation of mitigation measures helps to avoid or reduce impacts on live hard bottom, as defined in Chapter 5 (Mitigation). Impacts on the soft bottom substrate are determined to be short term and minimal due to the mobile nature of soft bottom substrates (i.e., sandy bottoms can be stirred up and settle relatively quickly when compared to impacts on hard bottom substrates).
- **Recommend consideration of temporal mitigation and habitat avoidance mitigation** – Temporal mitigation and habitat avoidance mitigation were considered, and mitigation areas are discussed in Section 5.4 (Mitigation Areas to be Implemented) of Chapter 5 (Mitigation) as well as Appendix I (Geographic Mitigation Assessment).

- **Impacts on EFH from training activities (deposition and resuspension of sediments, erosion and sedimentation, and impacts from unexploded ordnance)** – Because training activities would have adverse effects on EFH, the Navy is undergoing supplemental consultation with NMFS addressing activities that have changed (i.e., increased) as a result of the Proposed Action, see Section 6.1.3, Magnuson-Stevens Fishery Conservation and Management Act, for further discussion. To avoid or reduce adverse impacts on hard bottom habitat, the Navy created mitigation measures to protect the resource. As shown in Table 5.4-1 of Chapter 5 (Mitigation), shallow-water coral reefs, live hard bottom, artificial reefs, and submerged cultural resources are areas of focus for protection from explosives and physical disturbance and strike stressors. Mitigation area requirements to reduce impacts on live hard bottom substrate are listed in Table 5.4-1.

## **REFERENCES**

- Kendall, M. S., B. Costa, S. McKagan, L. Johnston, and D. Okano. (2017). *Benthic habitat maps of Saipan Lagoon* (NOAA Technical Memorandum NOS-NCCOS-229). Silver Spring, MD: National Oceanic and Atmospheric Administration.
- U.S. Department of the Navy. (2015). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.

---

---

## 3.4 Marine Mammals



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|          |  |        |
|----------|--|--------|
| 3.4      | Marine Mammals.....  | 3.4-1  |
| 3.4.1    | Affected Environment.....  | 3.4-1  |
| 3.4.1.1  | General Background .....   | 3.4-1  |
| 3.4.1.2  | Species Unlikely to Be Present in the Study Area.....              | 3.4-3  |
| 3.4.1.3  | Group Size.....  | 3.4-3  |
| 3.4.1.4  | Habitat Use .....  | 3.4-4  |
| 3.4.1.5  | Dive Behavior.....   | 3.4-4  |
| 3.4.1.6  | Hearing and Vocalization .....                                     | 3.4-4  |
| 3.4.1.7  | General Threats .....  | 3.4-8  |
| 3.4.1.8  | Blue Whale ( <i>Balaenoptera musculus</i> ).....                   | 3.4-15 |
| 3.4.1.9  | Bryde’s Whale ( <i>Balaenoptera edeni</i> ).....                   | 3.4-16 |
| 3.4.1.10 | Fin Whale ( <i>Balaenoptera physalus</i> ).....                    | 3.4-18 |
| 3.4.1.11 | Humpback Whale ( <i>Megaptera novaeangliae</i> ) .....             | 3.4-18 |
| 3.4.1.12 | Minke Whale ( <i>Balaenoptera acutorostrata</i> ) .....            | 3.4-22 |
| 3.4.1.13 | Omura’s Whale ( <i>Balaenoptera omurai</i> ) .....                 | 3.4-23 |
| 3.4.1.14 | Sei Whale ( <i>Balaenoptera borealis</i> ) .....                   | 3.4-23 |
| 3.4.1.15 | Blainville’s Beaked Whale ( <i>Mesoplodon densirostris</i> ).....  | 3.4-24 |
| 3.4.1.16 | Common Bottlenose Dolphin ( <i>Tursiops truncatus</i> ) .....      | 3.4-25 |
| 3.4.1.17 | Cuvier’s Beaked Whale ( <i>Ziphius cavirostris</i> ).....          | 3.4-27 |
| 3.4.1.18 | Dwarf Sperm Whale ( <i>Kogia sima</i> ).....                       | 3.4-28 |
| 3.4.1.19 | False Killer Whale ( <i>Pseudorca crassidens</i> ) .....           | 3.4-29 |
| 3.4.1.20 | Fraser’s Dolphin ( <i>Lagenodelphis hosei</i> ) .....              | 3.4-30 |
| 3.4.1.21 | Ginkgo-Toothed Beaked Whale ( <i>Mesoplodon ginkgodens</i> ) ..... | 3.4-31 |
| 3.4.1.22 | Killer Whale ( <i>Orcinus orca</i> ) .....                         | 3.4-32 |
| 3.4.1.23 | Longman’s Beaked Whale ( <i>Indopacetus pacificus</i> ) .....      | 3.4-34 |
| 3.4.1.24 | Melon-Headed Whale ( <i>Peponocephala electra</i> ) .....          | 3.4-35 |

|          |   |         |
|----------|---|---------|
| 3.4.1.25 | Pantropical Spotted Dolphin ( <i>Stenella attenuata</i> ) .....                   | 3.4-36  |
| 3.4.1.26 | Pygmy Killer Whale ( <i>Feresa attenuata</i> ) .....                              | 3.4-37  |
| 3.4.1.27 | Pygmy Sperm Whale ( <i>Kogia breviceps</i> ) .....                                | 3.4-38  |
| 3.4.1.28 | Risso’s Dolphin ( <i>Grampus griseus</i> ) .....                                  | 3.4-39  |
| 3.4.1.29 | Rough-Toothed Dolphin ( <i>Steno bredanensis</i> ).....                           | 3.4-40  |
| 3.4.1.30 | Short-Finned Pilot Whale ( <i>Globicephala macrorhynchus</i> ) .....              | 3.4-41  |
| 3.4.1.31 | Sperm Whale ( <i>Physeter macrocephalus</i> ).....                                | 3.4-42  |
| 3.4.1.32 | Spinner Dolphin ( <i>Stenella longirostris</i> ).....                             | 3.4-44  |
| 3.4.1.33 | Striped Dolphin ( <i>Stenella coeruleoalba</i> ).....                             | 3.4-45  |
| 3.4.2    | Environmental Consequences .....  | 3.4-46  |
| 3.4.2.1  | Acoustic Stressors.....   | 3.4-50  |
| 3.4.2.2  | Explosive Stressors.....  | 3.4-175 |
| 3.4.2.3  | Energy Stressors .....  | 3.4-245 |
| 3.4.2.4  | Physical Disturbance and Strike Stressors .....                                   | 3.4-247 |
| 3.4.2.5  | Entanglement Stressors.....   | 3.4-249 |
| 3.4.2.6  | Ingestion Stressors.....  | 3.4-250 |
| 3.4.2.7  | Secondary Stressors.....  | 3.4-252 |
| 3.4.3    | Summary of Potential Impacts on Marine Mammals.....                               | 3.4-253 |
| 3.4.3.1  | Combined Impacts of All Stressors Under Alternative 1 .....                       | 3.4-255 |
| 3.4.3.2  | Combined Impacts of All Stressors Under Alternative 2 .....                       | 3.4-255 |
| 3.4.3.3  | Combined Impacts of All Stressors Under the No Action Alternative<br>.....        | 3.4-255 |
| 3.4.3.4  | Summary of Monitoring and Observations During Navy Activities Since<br>2015 ..... | 3.4-255 |
| 3.4.4    | Marine Mammal Protection Act Determinations.....                                  | 3.4-258 |
| 3.4.5    | Endangered Species Act Determinations .....                                       | 3.4-259 |
| 3.4.6    | Public Scoping Comments .....   | 3.4-259 |

### List of Figures

|               |  |        |
|---------------|--|--------|
| Figure 3.4-1: | Composite Audiograms for Hearing Groups Likely Found in the Study Area ..... | 3.4-7  |
| Figure 3.4-2: | Two Hypothetical Threshold Shifts.....                                       | 3.4-55 |
| Figure 3.4-3: | Odontocete Critical Ratios .....   | 3.4-62 |
| Figure 3.4-4: | Critical Ratios for Different Noise Types.....                               | 3.4-63 |
| Figure 3.4-5: | Navy Auditory Weighting Functions for all Species Groups .....               | 3.4-93 |

Figure 3.4-6: TTS and PTS Exposure Functions for Sonar and Other Transducers.....3.4-94

Figure 3.4-7: Behavioral Response Function for Odontocetes .....3.4-96

Figure 3.4-8: Behavioral Response Function for Mysticetes .....3.4-96

Figure 3.4-9: Behavioral Response Function for Beaked Whales .....3.4-97

Figure 3.4-10: Relative Likelihood of a Response Being Significant Based on the Duration and Severity of Behavioral Reactions.....3.4-98

Figure 3.4-11: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-115

Figure 3.4-12: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-116

Figure 3.4-13: Bryde’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-117

Figure 3.4-14: Bryde’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-118

Figure 3.4-15: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-119

Figure 3.4-16: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-120

Figure 3.4-17: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-121

Figure 3.4-18: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-122

Figure 3.4-19: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-123

Figure 3.4-20: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-124

Figure 3.4-21: Omura’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-125

Figure 3.4-22: Omura’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-126

Figure 3.4-23: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-127

Figure 3.4-24: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-129

Figure 3.4-25: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-134

Figure 3.4-26: Cuvier’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-135

---

|   |         |
|---|---------|
| Figure 3.4-27: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 ..... | 3.4-136 |
| Figure 3.4-28: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....      | 3.4-137 |
| Figure 3.4-29: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....   | 3.4-138 |
| Figure 3.4-30: Cuvier’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....       | 3.4-139 |
| Figure 3.4-31: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 ..... | 3.4-140 |
| Figure 3.4-32: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....      | 3.4-141 |
| Figure 3.4-33: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....          | 3.4-142 |
| Figure 3.4-34: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....          | 3.4-143 |
| Figure 3.4-35: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....           | 3.4-144 |
| Figure 3.4-36: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....           | 3.4-145 |
| Figure 3.4-37: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....           | 3.4-146 |
| Figure 3.4-38: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....           | 3.4-147 |
| Figure 3.4-39: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....          | 3.4-148 |
| Figure 3.4-40: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....          | 3.4-149 |
| Figure 3.4-41: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....            | 3.4-150 |
| Figure 3.4-42: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....            | 3.4-151 |
| Figure 3.4-43: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....                | 3.4-153 |
| Figure 3.4-44: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....                | 3.4-154 |
| Figure 3.4-45: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....          | 3.4-155 |
| Figure 3.4-46: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....          | 3.4-156 |

Figure 3.4-47: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-157

Figure 3.4-48: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-158

Figure 3.4-49: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-159

Figure 3.4-50: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-160

Figure 3.4-51: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-161

Figure 3.4-52: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-162

Figure 3.4-53: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-163

Figure 3.4-54: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-164

Figure 3.4-55: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-165

Figure 3.4-56: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-166

Figure 3.4-57: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-167

Figure 3.4-58: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-168

Figure 3.4-59: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-169

Figure 3.4-60: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-170

Figure 3.4-61: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1 .....3.4-171

Figure 3.4-62: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2 .....3.4-172

Figure 3.4-63: Navy Phase III Weighting Functions for All Species Groups .....3.4-188

Figure 3.4-64: Navy Phase III Behavioral, TTS and PTS Exposure Functions for Explosives.....3.4-189

Figure 3.4-65: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-205

Figure 3.4-66: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-206

Figure 3.4-67: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-208

Figure 3.4-68: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-209

Figure 3.4-69: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-210

Figure 3.4-70: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-211

Figure 3.4-71: Omura’s Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-212

Figure 3.4-72: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-214

Figure 3.4-73: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-215

Figure 3.4-74: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-218

Figure 3.4-75: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-219

Figure 3.4-76: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-220

Figure 3.4-77: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-221

Figure 3.4-78: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-223

Figure 3.4-79: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-224

Figure 3.4-80: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-225

Figure 3.4-81: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-226

Figure 3.4-82: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-228

Figure 3.4-83: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-229

Figure 3.4-84: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-231

Figure 3.4-85: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-232

Figure 3.4-86: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-233

Figure 3.4-87: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-234

Figure 3.4-88: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-236

Figure 3.4-89: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-237

Figure 3.4-90: Short-Finned Pilot Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-239

Figure 3.4-91: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-241

Figure 3.4-92: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-242

Figure 3.4-93: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1 .....3.4-243

Figure 3.4-94: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2 .....3.4-244

### List of Tables

Table 3.4-1: Marine Mammal Occurrence within the Study Area .....3.4-2

Table 3.4-2: Species within Marine Mammal Hearing Groups Likely Found in the Study Area .....3.4-6

Table 3.4-3: Cutoff Distances for Moderate Source Level, Single Platform Training and Testing Events and for All Other Events with Multiple Platforms or Sonar with Source Levels at or Exceeding 215 dB re 1  $\mu$ Pa at 1 m .....3.4-97

Table 3.4-4: Range to Permanent Threshold Shift for Five Representative Sonar Systems .....3.4-101

Table 3.4-5: Ranges to Temporary Threshold Shift for Sonar Bin HF4 over a Representative Range of Environments within the Study Area .....3.4-102

Table 3.4-6: Ranges to Temporary Threshold Shift for Sonar Bin LF4 over a Representative Range of Environments within the Study Area .....3.4-102

Table 3.4-7: Ranges to Temporary Threshold Shift for Sonar Bin MF1 over a Representative Range of Environments within the Study Area .....3.4-103

Table 3.4-8: Ranges to Temporary Threshold Shift for Sonar Bin MF4 over a Representative Range of Environments within the Study Area .....3.4-103

Table 3.4-9: Ranges to Temporary Threshold Shift for Sonar Bin MF5 over a Representative Range of Environments within the Study Area .....3.4-104

Table 3.4-10: Ranges to a Potentially Significant Behavioral Response for Sonar Bin HF4 over a Representative Range of Environments within the Study Area.....3.4-104

Table 3.4-11: Ranges to a Potentially Significant Behavioral Response for Sonar Bin LF4 over a Representative Range of Environments within the Study Area.....3.4-105

---

|  |         |
|--|---------|
| Table 3.4-12: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF1 over a Representative Range of Environments within the Study Area..... | 3.4-106 |
| Table 3.4-13: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF4 over a Representative Range of Environments within the Study Area..... | 3.4-107 |
| Table 3.4-14: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF5 over a Representative Range of Environments within the Study Area..... | 3.4-108 |
| Table 3.4-15: Criteria to Quantitatively Assess Non-Auditory Injury Due to Underwater Explosions   | 3.4-187 |
| Table 3.4-16: Navy Phase III Weighted Sound Exposure Thresholds for Underwater Explosive Sounds .....  | 3.4-190 |
| Table 3.4-17: Ranges to Non-Auditory Injury (in meters) for All Marine Mammal Hearing Groups..   | 3.4-192 |
| Table 3.4-18: Ranges to Mortality (in meters) for All Marine Mammal Hearing Groups as a Function of Animal Mass .....  | 3.4-193 |
| Table 3.4-19: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for High-Frequency Cetaceans .....                                 | 3.4-194 |
| Table 3.4-20: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for High-Frequency Cetaceans .....   | 3.4-195 |
| Table 3.4-21: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Low-Frequency Cetaceans .....                                  | 3.4-197 |
| Table 3.4-22: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Low-Frequency Cetaceans .....  | 3.4-198 |
| Table 3.4-23: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Mid-Frequency Cetaceans .....                                  | 3.4-200 |
| Table 3.4-24: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Mid-Frequency Cetaceans .....  | 3.4-201 |

### 3.4 Marine Mammals

#### 3.4.1 Affected Environment

The purpose of this section is to supplement the analysis of impacts on marine mammals presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon de Medinilla (FDM). Information presented in the 2015 MITT Final EIS/OEIS that remains valid is noted as such and referenced in the appropriate sections. Any new or updated information describing the affected environment and analysis of impacts on marine mammals associated with the Proposed Action is provided in this section. Comments received from the public during scoping related to marine mammals are addressed in Section 3.4.6 (Public Scoping Comments).

The complete analysis and summary of potential impacts of the Proposed Action on marine mammals are found in Section 3.4.2 (Environmental Consequences) and Section 3.4.3 (Summary of Potential Impacts on Marine Mammals). For additional information, also see the 2015 MITT Final EIS/OEIS, Section 3.4 (Marine Mammals) (U.S. Department of the Navy, 2015a).

##### 3.4.1.1 General Background

Marine mammals are a diverse group of approximately 130 species. Most live predominantly in the marine habitat, although some species, such as seals, spend time in terrestrial habitats, and other species such as manatees and certain dolphins spend time in freshwater habitats (Rice, 1998; U.S. Department of the Navy, 2007). The exact number of formally recognized marine mammal species changes periodically with new scientific understanding or findings (Rice, 1998). For a list of current species classifications, see the formal list of *Marine Mammal Species and Subspecies* maintained online by the Society for Marine Mammalogy. In this document, the Navy follows the naming conventions presented by National Marine Fisheries Service (NMFS) in the applicable annual Stock Assessment Reports (SAR) for the Pacific and Alaska<sup>1</sup> regions covering the marine mammals present in the MITT Study Area (Carretta et al., 2017d; Carretta et al., 2018a; Muto et al., 2017b).

All marine mammals in the United States are protected under the Marine Mammal Protection Act (MMPA), and some species receive additional protection under the Endangered Species Act (ESA). The MMPA defines a marine mammal “stock” as “a group of marine mammals of the same species or smaller taxon in a common spatial arrangement that interbreed when mature” (16 United States Code [U.S.C.] section 1362; for further details, see Oleson et al. (2013). As provided by NMFS guidance, “for purposes of management under the MMPA a stock is recognized as being a management unit that identifies a demographically independent biological population.” (Carretta et al., 2017c; National Marine Fisheries Service, 2016b). However, in practice, recognized management stocks may fall short of this ideal for various reasons, including a lack of information, and, in some cases, may even include multiple distinct population segments in a management unit, such as with the Western North Pacific humpback whale stock (Bettridge et al., 2015).

---

<sup>1</sup> Some stocks in the Pacific and the Mariana Islands, such as the Northeast Pacific stocks of sperm whales and fin whales, and the Western North Pacific Stock of humpback whales, which include individuals that may spend the summer season foraging in Alaska waters, are covered in the Alaska Stock Assessment Report.

The ESA provides for listing species, subspecies, or distinct population segments of species, all of which are referred to as “species” under the ESA. The *Interagency Policy Regarding the Recognition of Distinct Vertebrate Population Segments Under the ESA* defines a distinct population segment as, “any subspecies of fish or wildlife or plants, and any distinct population segment of any species of vertebrate fish or wildlife which interbreeds when mature” (61 Federal Register [FR] 4722, February 7, 1996). If a population meets the criteria to be identified as a distinct population segment, it is eligible for listing under the ESA as a separate species (National Marine Fisheries Service, 2016b).

Twenty-six cetacean marine mammal species are known to exist in the Study Area, including 7 mysticetes (baleen whales) and 19 odontocetes (dolphins and toothed whales) (U.S. Department of the Navy, 2005, 2018b). The species expected to be present in the Study Area are provided in Table 3.4-1 and listed alphabetically within the two suborder groupings. The information presented in this Supplemental EIS (SEIS)/OEIS incorporates data from the U.S. Pacific and the Alaska Marine Mammal Stock Assessments (Carretta et al., 2017d; Muto et al., 2017b), which cover some of those species present in the Study Area and incorporate the best available science, including monitoring data from Navy marine mammal research efforts. For those few species for which stock information exists in the region, relevant data are included in the species-specific Status and Management summaries provided subsequently in this section.

**Table 3.4-1: Marine Mammal Occurrence within the Study Area**

| Common Name <sup>1</sup>  | Scientific Name                   | ESA Status | Occurrence in Study Area |                  |             |
|---------------------------|-----------------------------------|------------|--------------------------|------------------|-------------|
|                           |                                   |            | Mariana Islands          | Transit Corridor | Apra Harbor |
| <b>Mysticetes</b>         |                                   |            |                          |                  |             |
| Blue whale                | <i>Balaenoptera musculus</i>      | Endangered | Seasonal                 | Seasonal         | -           |
| Bryde’s whale             | <i>Balaenoptera edeni</i>         | n/a        | Regular                  | Regular          | -           |
| Fin whale                 | <i>Balaenoptera physalus</i>      | Endangered | Rare                     | Rare             | -           |
| Humpback whale            | <i>Megaptera novaeangliae</i>     | Endangered | Seasonal                 | Seasonal         | -           |
| Minke whale               | <i>Balaenoptera acutorostrata</i> | n/a        | Seasonal                 | Seasonal         | -           |
| Omura’s whale             | <i>Balaenoptera omurai</i>        | n/a        | Rare                     | Rare             | -           |
| Sei whale                 | <i>Balaenoptera borealis</i>      | Endangered | Seasonal                 | Seasonal         | -           |
| <b>Odontocetes</b>        |                                   |            |                          |                  |             |
| Blainville’s beaked whale | <i>Mesoplodon densirostris</i>    | n/a        | Regular                  | Regular          | -           |
| Common bottlenose dolphin | <i>Tursiops truncatus</i>         | n/a        | Regular                  | Regular          | -           |
| Cuvier’s beaked whale     | <i>Ziphius cavirostris</i>        | n/a        | Regular                  | Regular          | -           |
| Dwarf sperm whale         | <i>Kogia sima</i>                 | n/a        | Regular                  | Regular          | -           |

**Table 3.4-1: Marine Mammal Occurrence within the Study Area (continued)**

| Common Name <sup>1</sup>    | Scientific Name                   | ESA Status | Occurrence in Study Area |                  |             |
|-----------------------------|-----------------------------------|------------|--------------------------|------------------|-------------|
|                             |                                   |            | Mariana Islands          | Transit Corridor | Apra Harbor |
| <b>Odontocetes</b>          |                                   |            |                          |                  |             |
| False killer whale          | <i>Pseudorca crassidens</i>       | n/a        | Regular                  | Regular          | -           |
| Fraser's dolphin            | <i>Lagenodelphis hosei</i>        | n/a        | Regular                  | Regular          | -           |
| Ginkgo-toothed beaked whale | <i>Mesoplodon ginkgodens</i>      | n/a        | Regular                  | Regular          | -           |
| Killer whale                | <i>Orcinus orca</i>               | n/a        | Regular                  | Regular          | -           |
| Longman's beaked whale      | <i>Indopacetus pacificus</i>      | n/a        | Regular                  | Regular          | -           |
| Melon-headed whale          | <i>Peponocephala electra</i>      | n/a        | Regular                  | Regular          | -           |
| Pantropical spotted dolphin | <i>Stenella attenuata</i>         | n/a        | Regular                  | Regular          | -           |
| Pygmy killer whale          | <i>Feresa attenuata</i>           | n/a        | Regular                  | Regular          | -           |
| Pygmy sperm whale           | <i>Kogia breviceps</i>            | n/a        | Regular                  | Regular          | -           |
| Risso's dolphin             | <i>Grampus griseus</i>            | n/a        | Regular                  | Regular          | -           |
| Rough-toothed dolphin       | <i>Steno bredanensis</i>          | n/a        | Regular                  | Regular          | -           |
| Short-finned pilot whale    | <i>Globicephala macrorhynchus</i> | n/a        | Regular                  | Regular          | -           |
| Sperm whale                 | <i>Physeter macrocephalus</i>     | Endangered | Regular                  | Regular          | -           |
| Spinner dolphin             | <i>Stenella longirostris</i>      | n/a        | Regular                  | Regular          | -           |
| Striped dolphin             | <i>Stenella coeruleoalba</i>      | n/a        | Regular                  | Regular          | -           |

<sup>1</sup> If available for the species, information on stocks is included in the species-specific Status and Management summaries.

Notes: n/a = status is not applicable for those species that are not listed under ESA; Regular = a species that occurs as a regular or usual part of the fauna of the area, regardless of how abundant or common it is; Rare = a species that occurs in the area only sporadically; Seasonal = species is only seasonally present in the Study Area. Additional details regarding presence in the Study Area are provided in the species-specific subsections.

#### 3.4.1.2 Species Unlikely to Be Present in the Study Area

Consistent with the analysis provided in the 2015 MITT Final EIS/OEIS, the species carried forward for analysis in this SEIS/OEIS are those likely to be found in the Study Area based on the most recent sighting, survey, and habitat modeling data available. The analysis does not include species that may have once inhabited or transited the area, but have not been sighted in recent years (e.g., species which no longer occur in an area due to factors such as 19th-century commercial exploitation). These species include the North Pacific right whale (*Eubalaena japonica*), the western subpopulation of gray whale (*Eschrichtius robustus*), short-beaked common dolphin (*Delphinus delphis*), Indo-Pacific bottlenose dolphin (*Tursiops aduncus*), northern elephant seal (*Mirounga angustirostris*), and dugong (*Dugong dugon*). Details regarding the reasons for these exclusions are explained in detail in the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015a).

#### 3.4.1.3 Group Size

Many species of marine mammals, particularly odontocetes, are highly social animals that spend much of their lives living in groups called "pods." The size and structures of these groups are dynamic and depending on the species, can range from several to several thousand individuals. Similarly,

aggregations of mysticete whales may form during particular breeding or foraging seasons, although they do not persist through time as a social unit. Marine mammals that live or travel in groups are more likely to be detected by observers, and group size characteristics are incorporated into the many density and abundance calculations. Group size characteristics are also incorporated into acoustic effects modeling to represent a more realistic patchy distribution for a given density. The behavior of aggregating into groups is also important for the purposes of mitigation and monitoring since animals that occur in larger groups have an increased probability of being detected. A comprehensive and systematic review of relevant literature and data was conducted for available published and unpublished literature, including journals, books, technical reports, cruise reports, and raw data from cruises, theses, and dissertations. The results of this review were compiled into a Technical Report, which includes tables of group size information by species along with relevant citations.

#### **3.4.1.4 Habitat Use**

Many factors influence the distribution of marine mammals in the Study Area, primarily patterns of major ocean currents, bottom relief, and water temperature, which, in turn, affect prey distribution and productivity. The continuous movement of water from the ocean bottom to the surface creates a nutrient-rich, highly productive environment for marine mammal prey in upwelling zones (Jefferson et al., 2015); the equatorial upwelling in the western Pacific is one such area (Di Lorenzo et al., 2010; Helber & Weisberg, 2001). While most baleen whales (such as humpback whales) are migratory, some species such as Bryde's whales and Omura's whales are thought to be present within the Study Area year round. Many of the toothed whales do not migrate in the strictest sense, but some do undergo seasonal shifts in distribution both within and outside of the Study Area.

#### **3.4.1.5 Dive Behavior**

All marine mammals, with the exception of polar bears, spend part of their lives underwater while traveling or feeding. Some species of marine mammals have developed specialized adaptations to allow them to make deep dives lasting over an hour, primarily for the purpose of foraging on deep-water prey such as squid. Other species spend the majority of their lives close to the surface, and make relatively shallow dives. The diving behavior of a particular species or individual has implications for the ability to visually detect them for mitigation and monitoring. In addition, their relative distribution through the water column based on diving behavior is an important consideration when conducting acoustic effects modeling. Information and data on diving behavior for each species of marine mammal were compiled and summarized in a technical report (U.S. Department of the Navy, 2018a) that provides the detailed summary of time at depth.

#### **3.4.1.6 Hearing and Vocalization**

The typical terrestrial mammalian ear (which is ancestral to that of marine mammals) consists of an outer ear that collects and transfers sound to the tympanic membrane and then to the middle ear (Fay & Popper, 1994; Rosowski, 1994). The middle ear contains ossicles that amplify and transfer acoustic energy to the sensory cells (called hair cells) in the cochlea, which transforms acoustic energy into electrical neural impulses that are transferred by the auditory nerve to high levels in the brain (Møller, 2013). All marine mammals display some degree of modification to the terrestrial ear; however, there are differences in the hearing mechanisms of marine mammals with an amphibious ear versus those with a fully aquatic ear (Wartzok & Ketten, 1999). Marine mammals with an amphibious ear include the marine carnivores: pinnipeds, sea otters, and polar bears (Ghoul & Reichmuth, 2014; Owen & Bowles, 2011; Reichmuth et al., 2013). Outer ear adaptations in this group include external pinnae (ears) that are

reduced or absent, and in the pinnipeds, cavernous tissue, muscle, and cartilaginous valves seal off water from entering the auditory canal when submerged (Wartzok & Ketten, 1999). Marine mammals with the fully aquatic ear (cetaceans and sirenians) use bone and fat channels in the head to conduct sound to the ear; while the auditory canal still exists it is narrow and sealed with wax and debris (Ketten, 1998).


The most accurate means of determining the hearing capabilities of marine mammal species are direct measures that assess the sensitivity of the auditory system (Nachtigall et al., 2000; Supin et al., 2001). Studies using these methods produce audiograms — plots describing hearing threshold (the quietest sound a listener can hear) as a function of frequency. Marine mammal audiograms, like those of terrestrial mammals, typically have a “U-shape,” with a frequency region of best hearing sensitivity and a progressive decrease in sensitivity outside of the range of best hearing (Fay, 1988; Mooney et al., 2012; Nedwell et al., 2004; Reichmuth et al., 2013). The “gold standard” for producing audiograms is the use of behavioral (psychophysical) methods, where marine mammals are trained to respond to acoustic stimuli (Nachtigall et al., 2000). For species that are untrained for behavioral psychophysical procedures, those that are difficult to house under human care, or in stranding rehabilitation and temporary capture contexts, auditory evoked potential methods are increasingly used to measure hearing sensitivity (e.g., Castellote et al., 2014; Finneran et al., 2009; Montie et al., 2011; Mulsow et al., 2011; Nachtigall et al., 2007; Nachtigall et al., 2008; Supin et al., 2001).

These auditory evoked potential methods, which measure electrical potentials generated by the auditory system in response to sound and do not require the extensive training of psychophysical methods, can provide an efficient estimate of behaviorally measured sensitivity (Finneran & Houser, 2006; Schlundt et al., 2007; Yuen et al., 2005). The thresholds provided by auditory evoked potential methods are, however, typically elevated above behaviorally measured thresholds, and auditory evoked potential methods are not appropriate for estimating hearing sensitivity at frequencies much lower than the region of best hearing sensitivity (Finneran, 2015; Finneran et al., 2016). For marine mammal species for which access is limited and therefore psychophysical or auditory evoked potential testing is impractical (e.g., mysticete whales and rare species), some aspects of hearing can be estimated from anatomical structures, frequency content of vocalizations, and extrapolations from related species.

Direct measurements of hearing sensitivity exist for approximately 25 of the nearly 130 species of marine mammals. Table 3.4-2 summarizes hearing capabilities for marine mammal species in the Study Area. For this analysis, marine mammals are arranged into the following functional hearing groups based on their generalized hearing sensitivities: high-frequency (HF) cetaceans (HF group: porpoises, Kogia whales), mid-frequency (MF) cetaceans (MF group: delphinids, beaked whales, sperm whales), and low-frequency (LF) cetaceans (LF group: mysticetes). Note that the designations of high-, mid-, and low-frequency cetaceans are based on relative differences of sensitivity between groups, as opposed to conventions used to describe active sonar systems. For Phase III analyses, a single representative composite audiogram (Figure 3.4-1) was created for each functional hearing group using audiograms from published literature. For discussion of all marine mammal functional hearing groups and their derivation see technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a). The mid-frequency cetacean composite audiogram is consistent with recently published behavioral audiograms of killer whales (Branstetter et al., 2017a).

**Table 3.4-2: Species within Marine Mammal Hearing Groups Likely Found in the Study Area**

| <i>Hearing Group</i>     | <i>Species within the Study Area</i> |
|--------------------------|--------------------------------------|
| High-frequency cetaceans | Dwarf sperm whale                    |
|                          | Pygmy sperm whale                    |
| Mid-frequency cetaceans  | Blainville's beaked whale            |
|                          | Common bottlenose dolphin            |
|                          | Cuvier's beaked whale                |
|                          | False killer whale                   |
|                          | Fraser's dolphin                     |
|                          | Ginkgo-toothed beaked whale          |
|                          | Killer whale                         |
|                          | Longman's beaked whale               |
|                          | Melon-headed whale                   |
|                          | Northern right whale dolphin         |
|                          | Pantropical spotted dolphin          |
|                          | Pygmy killer whale                   |
|                          | Risso's dolphin                      |
|                          | Rough-toothed dolphin                |
|                          | Short-finned pilot whale             |
|                          | Sperm whale                          |
|                          | Spinner dolphin                      |
| Striped dolphin          |                                      |
| Low-frequency cetaceans  | Blue whale                           |
|                          | Bryde's whale                        |
|                          | Fin whale                            |
|                          | Humpback whale                       |
|                          | Minke whale                          |
|                          | Omura's whale                        |
|                          | Sei whale                            |



Source: (U.S. Department of the Navy, 2017a)

Notes: For hearing in the water; LF = low-frequency, MF = mid-frequency, HF = high-frequency

**Figure 3.4-1: Composite Audiograms for Hearing Groups Likely Found in the Study Area**

Similar to the diversity of hearing capabilities among species, the wide variety of acoustic signals used in marine mammal communication (including biosonar or echolocation) is reflective of the diverse ecological characteristics of cetacean species (see Avens, 2003; Richardson et al., 1995b). This makes a succinct summary difficult (see Richardson et al., 1995b; Wartzok & Ketten, 1999 for thorough reviews); however, a division can be drawn between lower frequency communication signals that are used by marine mammals in general, and the specific, high-frequency biosonar signals that are used by odontocetes to sense their environment.

Non-biosonar communication signals span a wide frequency range, primarily having energy up into the tens of kilohertz (kHz) range. Of particular note are the very low-frequency calls of mysticete whales that range from tens of hertz (Hz) to several kilohertz, and have source levels of 150–200 decibels referenced to 1 micropascal (dB re 1  $\mu$ Pa) (Cummings & Thompson, 1971; Edds-Walton, 1997; Širović et al., 2007; Stimpert et al., 2007; Wartzok & Ketten, 1999). These calls most likely serve social functions such as mate attraction, but may serve an orientation function as well (Green, 1994; Green et al., 1994; Richardson et al., 1995b). Humpback whales are a notable exception within the mysticetes, with some calls exceeding 10 kHz (Zoidis et al., 2008).

Odontocete cetaceans use underwater communicative signals that, while not as low in frequency as those of many mysticetes, likely serve similar functions. The acoustic characteristics of these signals are quite diverse among species, but can be generally classified as having dominant energy at frequencies below 20 kHz (Richardson et al., 1995b; Wartzok & Ketten, 1999).

Odontocete cetaceans generate short-duration (500–200  $\mu$ s), specialized clicks used in biosonar with peak frequencies between 10 and 200 kHz to detect, localize, and characterize underwater objects such as prey (Au, 1993; Wartzok & Ketten, 1999). These clicks are often more intense than other communicative signals, with reported source levels as high as 229 dB re 1  $\mu$ Pa peak-to-peak (Au et al., 1974). The echolocation clicks of high-frequency cetaceans are narrower in bandwidth (i.e., the difference between the upper and lower frequencies in a sound) and higher in frequency than those of mid-frequency cetaceans (Madsen et al., 2005; Villadsgaard et al., 2007).

In general, frequency ranges of vocalization lie within the audible frequency range for an animal (i.e., animals vocalize within their audible frequency range); however, auditory frequency range and vocalization frequencies do not perfectly align. The frequency range of vocalization in a species can therefore be used to infer some characteristics of their auditory system; however, caution must be taken when considering vocalization frequencies alone in predicting the hearing capabilities of species for which no data exist (i.e., mysticetes). It is important to note that aspects of vocalization and hearing sensitivity are subject to evolutionary pressures that are not solely related to detecting communication signals. For example, hearing plays an important role in detecting threats (e.g., Deecke et al., 2002), and high-frequency hearing is advantageous to animals with small heads in that it facilitates sound localization based on differences in sound levels at each ear (Heffner & Heffner, 1982). This may be partially responsible for the difference in best hearing thresholds and dominant vocalization frequencies in some species of marine mammals (e.g., Steller sea lions) (Mulsow & Reichmuth, 2010).

#### **3.4.1.7 General Threats**

Marine mammal populations can be influenced by various natural factors as well as human activities. There can be direct effects from disease, hunting, and whale watching, or indirect effects such as through reduced prey availability or lowered reproductive success of individuals. Research presented in Twiss and Reeves (1999) and National Marine Fisheries Service (2011a, 2011b, 2011c, 2011d) provides a general discussion of marine mammal conservation and the threats they face. As detailed in National Marine Fisheries Service (2011e), investigations of stranded marine mammals are undertaken to monitor threats to marine mammals (Simeone et al., 2015). Investigations into the cause of death for stranded animals can also provide indications of the general threats to marine mammals in a given location (Bradford & Forney, 2017; Carretta et al., 2017b; Helker et al., 2017). The causes for strandings include infectious disease, parasite infestation, climate change reducing prey availability and leading to starvation, pollution exposure, trauma (e.g., injuries from ship strikes or fishery entanglements), sound (human-generated or natural), harmful algal blooms and associated biotoxins, tectonic events such as underwater earthquakes, and ingestion of or interaction with marine debris (for more information see NMFS Marine Mammal Stranding Response Fact Sheet (National Marine Fisheries Service, 2016d). Since 1963, Guam Department of Agriculture Division of Aquatic and Wildlife Resources has conducted aerial surveys twice every month (weather permitting) of the coastal margin around Guam at a distance of approximately 200–300 meters (m) offshore of the outer reef margin (Martin et al., 2016). Therefore, the Navy assumes any animals stranded on Guam are likely to have been identified; see also Mobley (2007). For a general discussion of strandings and their causes as well as strandings in association with U.S. Navy activity, see the technical report titled *Strandings Associated with U.S. Navy Activity* (U.S. Department of the Navy, 2017c).

##### **3.4.1.7.1 Water Quality**

Chemical pollution and impacts on ocean water quality are of great concern, although the effects on marine mammals are just starting to be understood (Bachman et al., 2014; Bachman et al., 2015;

Desforges et al., 2016; Foltz et al., 2014; Godard-Codding et al., 2011; Hansen et al., 2015; Jepson & Law, 2016; Law, 2014; Peterson et al., 2014; Peterson et al., 2015; Ylitalo et al., 2005; Ylitalo et al., 2009). Oil and other chemical spills are a specific type of ocean contamination that can have damaging effects on some marine mammal species directly through exposure to oil or chemicals and indirectly due to pollutants' impacts on prey and habitat quality (Engelhardt, 1983; Marine Mammal Commission, 2010; Matkin et al., 2008).

On a broader scale ocean contamination resulting from chemical pollutants inadvertently introduced into the environment by industrial, urban, and agricultural use is also a concern for marine mammal conservation and has been the subject of numerous studies (Desforges et al., 2016; Fair et al., 2010; Krahn et al., 2007; Krahn et al., 2009; Moon et al., 2010; Ocean Alliance, 2010). For example, the chemical components of pesticides used on land flow as runoff into the marine environment, which can accumulate in the bodies of marine mammals, and be transferred to nursing young through the mother's milk (Fair et al., 2010). The presence of these chemicals in marine mammals has been assumed to put those animals at greater risk for adverse health effects and potential impact on their reproductive success given toxicology studies and results from laboratory animals (Fair et al., 2010; Godard-Codding et al., 2011; Krahn et al., 2007; Krahn et al., 2009; Peterson et al., 2014; Peterson et al., 2015). Desforges et al. (2016) have suggested that exposure to chemical pollutants may act in an additive or synergistic manner with other stressors resulting in significant population level consequences. Although the general trend has been a decrease in chemical pollutants in the environment following their regulation, chemical pollutants remain important given their potential to impact marine mammals and marine life in general (Bonito et al., 2016; Jepson & Law, 2016; Law, 2014).

#### **3.4.1.7.2 Commercial Industries**

Human impacts on marine mammals through fisheries interactions have received much attention in recent decades, and include bycatch (accidental or incidental catch), gear entanglement, and indirect effects from takes of prey species; noise pollution; marine debris (ingestion and entanglement); hunting (both commercial and native practices); vessel strikes; entrainment into power plant water intakes; increased ocean acidification; and general habitat deterioration or destruction.

#### **3.4.1.7.3 Bycatch**

Fishery bycatch is likely the most impactful threat to marine mammal individuals and populations and may account for the deaths of more marine mammals than any other cause (Bradford & Forney, 2017; Carretta et al., 2016b; Carretta et al., 2017b; Geijer & Read, 2013; Hamer et al., 2010; Helker et al., 2017; Lent & Squires, 2017; National Marine Fisheries Service, 2016c; Northridge, 2009; Read, 2008; Song, 2017). In 1994, the MMPA was amended to formally address bycatch in the United States. The amendment required the development of a take reduction plan when bycatch exceeds a level considered sustainable and lead to marine mammal population decline. In addition, NMFS develops and implements take reduction plans that help recover and prevent the depletion of strategic stocks of marine mammals that interact with certain fisheries.

At least in part as a result of the amendment, estimates of bycatch in the Pacific by U.S. fisheries declined by a total of 96 percent from 1994 to 2006 (Geijer & Read, 2013). Information on bycatch associated with non-U.S. fishery activities is generally not available in the Study Area. It has been argued that the bycatch of marine mammals by Japan and South Korea is more like an unregulated commercial hunt than an incidental or illegal fishery given the products from bycatch whales can be sold openly on commercial markets in both countries (Baker et al., 2006b; Lukoschek et al., 2009). For example, in 2008

the reported bycatch in Japan and South Korea totaled 214 minke whales (Lukoschek et al., 2009) and in nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that 230 products contained minke whale meat in addition to that of other species (Baker et al., 2006b). A total of 717 cetacean bycatches involving South Korean fish vessels in the East Sea that involved 10 species were reported in a two-year period between 2013 and 2014 (Song, 2017).

#### **3.4.1.7.4 Other Fishery Interactions**

Fishery interactions other than bycatch include entanglement in abandoned or partial nets, fishing line, hooks, and the ropes and lines connected to fishing gear (Bradford & Lyman, 2015; Bradford & Forney, 2016, 2017; Carretta, 2013; Carretta et al., 2014; Carretta et al., 2016b; Carretta et al., 2017b; Helker et al., 2015; Helker et al., 2017; National Oceanic and Atmospheric Administration, 2017; Richardson et al., 2016; Saez et al., 2013). The National Oceanic and Atmospheric Administration Marine Debris Program (2014b) reports that abandoned, lost, or otherwise discarded fishing gear constitutes the vast majority of mysticete entanglements. For the identified sources of entanglement, none included Navy-expended materials. Identified species entangled in the Pacific in 2015 and 2016 included humpback, gray, blue, fin and killer whales with a total of 133 entanglements in the two-year period (National Oceanic and Atmospheric Administration, 2017).

In waters off Alaska where humpback whales from the Study Area may forage in the summer season, passive acoustic monitoring efforts since 2009 have documented the routine use of non-military explosives at sea (Baumann-Pickering et al., 2013; Rice et al., 2015). Based on the acoustic spectral properties of the recorded sounds and their correspondence with known fishing seasons or activity, the source of these explosions has been linked to the use of explosive marine mammal deterrents, which as a group are commonly known as “seal bombs” (Baumann-Pickering et al., 2013). Seal bombs are intended to be used by commercial fishers to deter marine mammals, particularly pinnipeds, from preying upon their catch and to prevent marine mammals from interacting and potentially becoming entangled with fishing gear (National Marine Fisheries Service, 2015c). The prevalent and continued use of seal bombs in Alaska seems to indicate that, while a potential threat, their use has had no significant effect on populations of marine mammals given that it is likely at least some individuals, if not larger groups of marine mammals, have been repeatedly exposed to this explosive stressor.

#### **3.4.1.7.5 Noise**

In some locations, especially where urban or industrial activities or commercial shipping is intense, anthropogenic noise can be a potential habitat-level stressor (Dunlop, 2016; Dyndo et al., 2015; Erbe et al., 2014; Frisk, 2012; Gedamke et al., 2016; Hermannsen et al., 2014; Li et al., 2015; McKenna et al., 2012; Melcón et al., 2012; Miksis-Olds & Nichols, 2015; Nowacek et al., 2015; Pine et al., 2016; Williams et al., 2014b). Noise is of particular concern to marine mammals because many species use sound as a primary sense for navigating, finding prey, avoiding predators, and communicating with other individuals. Noise may cause marine mammals to leave a habitat, impair their ability to communicate, or cause physiological stress (Courbis & Timmel, 2008; Erbe, 2002; Erbe et al., 2016; Hildebrand, 2009; Rolland et al., 2012; Tyack et al., 2011; Tyne et al., 2017; Williams et al., 2013). Noise can cause behavioral disturbances, mask other sounds including their own vocalizations, may result in injury, and in some cases may result in behaviors that ultimately lead to death (Erbe et al., 2014; Erbe et al., 2016; National Research Council, 2003, 2005; Nowacek et al., 2007a; Southall et al., 2009; Tyack, 2009; Würsig & Richardson, 2009). Anthropogenic noise is generated from a variety of sources including commercial shipping, oil and gas exploration and production activities, commercial and recreational fishing (including fish-finding sonar, fathometers, and acoustic deterrent and harassment devices), foreign

navies, recreational boating and whale watching activities, offshore power generation, and research (including sound from air guns, sonar, and telemetry).

Vessel noise in particular is a major contributor to noise in the ocean (Southall et al., 2018). Commercial shipping's contribution to ambient noise in the ocean increased by as much as 12 decibels (dB) between approximately the 1960s and 2005 (Hildebrand, 2009; McDonald et al., 2008). Frisk (2012) confirmed the trend and reported that between 1950 and 2007 ocean noise in the 25–50 Hz frequency range has increased 3.3 dB per decade, resulting in a cumulative increase of approximately 19 dB over a baseline of 52 dB. The increase in noise is associated with an increase in commercial shipping, which correlates with global economic growth (Frisk, 2012). Subsequently, Miksis-Olds and Nichols (2015) have demonstrated that the trends for low-frequency ocean sound levels no longer show a uniform increase across the globe.

Although Guam and the Mariana Islands lack a major port, many thousands of trans-Pacific shipments to and from Asia occur as part of the global shipping transportation network and pass in proximity to the Mariana Islands and throughout the Study Area (Kaluza et al., 2010). Redfern et al. (2017) found that shipping channels leading to and from the ports of Los Angeles and Long Beach may have degraded the habitat for blue, fin, and humpback whales due to the loss of communication space where important habitat for these species overlaps with elevated noise from commercial vessel traffic, and similar impacts are also likely in the Study Area on great circle routes and other shortest point-to-point shipping routes.

In many areas of the world, including the Study Area, oil and gas seismic exploration in the ocean is undertaken using a group of air guns towed behind large research vessels. The airguns convert high pressure air into very strong shock wave impulses that are designed to return information off the various buried layers of sediment under the seafloor. Seismic exploration surveys last many days and cover vast overlapping swaths of the ocean area being explored. Most of the impulse energy produced by these airguns is heard as low-frequency sound, which can travel long distances and has the potential to impact marine mammals. NMFS routinely issues permits for the taking of marine mammals associated with these commercial activities.

#### **3.4.1.7.6 Hunting**

Commercial hunting, as in whaling and sealing operations, provided the original impetus for marine mammal management efforts and has driven much of the early research on cetaceans and pinnipeds (Twiss & Reeves, 1999). With the enactment of the MMPA and the 1946 International Convention for the Regulation of Whaling, hunting-related mortality has decreased over the last 40 years. Unregulated harvests are still considered as direct threats. However, since passage of the MMPA, there have been relatively few serious calls for culls of marine mammals in the United States compared to other countries, including Canada (Roman et al., 2013). Review of uncovered Union of Soviet Socialist Republics catch records in the North Pacific Ocean indicate extensive illegal whaling activity between 1948 and 1979, with a harvest totaling 195,783 whales. Of these, only 169,638 were reported by the Union of Soviet Socialist Republics to the International Whaling Commission (Ilyashenko et al., 2013; Ilyashenko et al., 2014; Ilyashenko & Chapham, 2014; Ilyashenko et al., 2015).

#### **3.4.1.7.7 Vessel Strike**

Ship strikes are also a growing issue for most marine mammals, especially for large whales as populations recover from widespread commercial whaling. Some of the largest ports in the world are located to the west of the Study Area, and a substantial portion of the world's commercial vessel traffic

from Asia and Japan transits through the Study Area heading south to ports of call along the coast of eastern Australia and to New Zealand, in addition to goods shipped into the Mariana Islands.

Since 1995, the U.S. Navy and U.S. Coast Guard have reported all known or suspected vessel collisions with whales to NMFS. There are no known collisions between Navy vessels and whales in the MITT Study Area associated with any of the proposed training or testing activities. The assumed under-reporting of whale collisions by vessels other than U.S. Navy or U.S. Coast Guard makes any comparison of data involving vessel strikes between Navy vessels and other vessels heavily biased. This under-reporting is recognized by NMFS; for example, in the Technical Memorandum providing the analysis of the impacts from vessel collisions with whales in Hawaii (Bradford & Lyman, 2015), NMFS takes into account unreported vessel strikes by civilian vessels.

#### **3.4.1.7.8 Disease and Parasites**

As with humans, marine mammals, especially the young, old, and weak, are susceptible to disease. For example, the first case of morbillivirus (a virus related to measles in humans) in the central Pacific was documented for a stranded juvenile male Longman's beaked whale discovered in 2010 in Hawaii (West et al., 2012; West et al., 2015) and subsequently in 2011 *brucella* (a bacterial pathogen) and *morbillivirus* were discovered in a sperm whale that stranded on Oahu (West et al., 2015); both these species are present in the Study Area. Occasionally disease epidemics can also injure or kill a large percentage of a marine mammal population (Keck et al., 2010; Paniz-Mondolfi & Sander-Hoffmann, 2009; Simeone et al., 2015). Recent review of odontocetes stranded along the California coast from 2000 to 2015 found evidence for morbilliviral infection in 9 of the 212 animals examined, therefore indicating this disease may be a contributor to mortality in cetaceans stranding along the California coast (Serrano et al., 2017).

Mass die-offs of some marine mammal species have been linked to toxic algal blooms, which occur as larger organisms consume multiple prey containing those toxins and thereby accumulating fatal doses. A comprehensive study that sampled over 900 marine mammals across 13 species, including several mysticetes, odontocetes, pinnipeds, and mustelids in Alaska, found detectable concentrations of domoic acid in all 13 species and saxitoxin, a toxin absorbed from ingesting dinoflagellates, in 10 of the 13 species (Lefebvre et al., 2016). Algal toxins may have contributed to the stranding and mortality of 30 whales found around the islands in the western Gulf of Alaska and the southern shoreline of the Alaska Peninsula starting in May 2015 (National Oceanic and Atmospheric Administration, 2016; Rosen, 2015; Savage et al., 2017; Summers, 2017). These findings from studies in Alaska are relevant to the Study Area given that the fin and humpback whales from the Mariana Islands migrate to Alaska waters in the summer to feed.

Additionally, all marine mammals have parasites that, under normal circumstances, probably do little overall harm, but under certain conditions, can cause serious health problems or even death (Bull et al., 2006; Fauquier et al., 2009; Jepson et al., 2005). Parasitic toxoplasmosis from feral cats (introduced into the ocean from urban runoff) has been found in two stranded spinner dolphins and eight monk seals in Hawaii (Rogers, 2016; West, 2018). In 2011, a Cuvier's beaked whale stranded at Saipan, when necropsied, it was found to have abnormalities and a level of parasitism that resulted in "the worst example of kidneys" the stranding coordinator said he had ever seen (Saipan Tribune, 2011).

#### **3.4.1.7.9 Climate Change**

The global climate is warming and is having impacts on some populations of marine mammals (National Marine Fisheries Service, 2015e; National Oceanic and Atmospheric Administration, 2018a; Salvadeo et al., 2010; Shirasago-Germán et al., 2015; Simmonds & Elliott, 2009). Climate change can affect marine

mammal species directly by causing them to shift their distribution to match physiological tolerance under changing environmental conditions (Doney et al., 2012; Silber et al., 2017), which may or may not result in net habitat loss (some can experience habitat gains). Climate change can also affect marine mammals indirectly via impacts on prey, changes in prey distributions and locations, and changes in water temperature (Giorli & Au, 2017). Changes in prey can impact marine mammal foraging success, which in turn affects reproductive success and survival. Researchers in July 2016 shifted the location of blue, fin, and humpback whale satellite tagging efforts from Southern California to Central California, following sightings of thin and apparently unwell whales. The whales' conditions were thought to be the result of a change in the distribution of their prey away from traditional foraging areas. (Oregon State University, 2017) In Central California waters, the researchers identified good numbers of blue, fin, and humpback whales in better condition and indicative of a good feeding area that was likely to be sustained that season (Oregon State University, 2017).

Harmful algal blooms may become more prevalent in warmer ocean waters with increased salinity levels such that blooms will begin earlier, last longer, and cover a larger geographical range (Edwards, 2013; Moore, 2008). Warming ocean waters have been linked to the spread of harmful algal blooms into the North Pacific where waters had previously been too cold for most of these algae to thrive. Most of the mysticetes found in the Study Area spend part of the year in the North Pacific. The spread of the algae and associated blooms has led to disease in marine mammals in locations where algae-caused diseases had not been previously known (Lefebvre et al., 2016).

Climate change may indirectly influence marine mammals through changes in human behavior, such as increased shipping and oil and gas extraction, which benefit from sea ice loss (Alter et al., 2010). Ultimately impacts from global climate change may result in an intensification of current and on-going threats to marine mammals (Edwards, 2013).

Marine mammals are influenced by climate-related phenomena, such as typhoons and shifts in extreme weather patterns such as the 2015–2016 El Niño in the ocean off the U.S. West Coast. Generally, not much is known about how large storms and other weather patterns affect marine mammals, other than that mass strandings (when two or more marine mammals become beached or stuck in shallow water) sometimes coincide with hurricanes, typhoons, and other tropical storms (Bradshaw et al., 2006; Marsh, 1989; Rosel & Watts, 2008; Zellar et al., 2017), or other oceanographic conditions. Indirect impacts of climate change may include altered water chemistry in estuaries (low dissolved oxygen or increased nutrient loading) causing massive fish kills (Burkholder et al., 2004), which changes prey distribution and availability for cetaceans (Stevens et al., 2006). Human responses to extreme weather events may indirectly affect behavior and reproductive rates of marine mammals. For example, Miller et al. (2010) reported an increase in reproductive rates in bottlenose dolphins after Hurricane Katrina in the Mississippi Sound, presumably resulting from an increase in fish abundance due to a reduction in fisheries landings, a decrease in recreational and commercial boat activities (National Marine Fisheries Service, 2007b), and an increase in the number of reproductively active females available during the breeding seasons following the storm. Smith et al. (2013a) supplemented the findings from this study and documented a marked increase in foraging activity in newly identified foraging areas that were observed during the two-year study period after the storm.

Habitat deterioration and loss is a major factor for almost all coastal and inshore species of marine mammals and may include such factors as depleting a habitat's prey base and the complete loss of habitat (Ayres et al., 2012; Kemp, 1996; Pine et al., 2016; Rolland et al., 2012; Smith et al., 2009; Veirs et al., 2015; Williams et al., 2014a). Many researchers predict that if oceanic temperatures continue to rise

with an associated effect on marine habitat and prey availability, then either changes in foraging or life history strategies, including poleward shifts in many marine mammal species distributions, should be anticipated (Alter et al., 2010; Fleming et al., 2016; Ramp et al., 2015; Salvadeo et al., 2015; Sydeman & Allen, 1999). Poloczanska et al. (2016) analyzed climate change impact data that integrates multiple climate-influenced changes in ocean conditions (e.g., temperature, acidification, dissolved oxygen, and rainfall) to assess anticipated changes to a number of key ocean fauna across representative areas. Their results predict a northward expansion in the distribution of zooplankton, fish, and squid, all of which are prey for many marine mammal species.

Marine mammals as a whole are subject to the various influences and factors delineated in this section. If specific threats to individual species in the Study Area are known, those threats are described below in individual species accounts.

#### **3.4.1.7.10 Marine Debris**

The majority of marine debris in the ocean comes from land-based sources (Jambeck et al., 2015; Thiel et al., 2018). Without improved waste management and infrastructure in undeveloped coastal countries worldwide, the cumulative quantity of plastic waste available to enter the ocean from land is predicted to increase by an order of magnitude by 2025 (Jambeck et al., 2015). Marine debris is a global threat to marine mammals (National Oceanic and Atmospheric Administration Marine Debris Program, 2014a; Puig-Lozano et al., 2018). A literature review by Baulch and Perry (2014) found that 56 percent of cetacean species are documented as having ingested marine debris and that it can be a significant source of injury and mortality. A Cuvier's beaked whale that stranded at Saipan in 2011 was found to have an approximate 1-inch-diameter circular piece of plastic in its stomach (Saipan Tribune, 2011). Attributing cause of death to marine debris ingestion is difficult (Laist, 1997), but ingestion of plastic bags and Styrofoam has been identified as the cause of injury or death of minke whales (De Pierrepont et al., 2005) and deep-diving odontocetes, including beaked whales (Baulch & Perry, 2014), pygmy sperm whales (Sadove & Morreale, 1989; Stamper et al., 2006; Tarpley & Marwitz, 1993), and sperm whales (Jacobsen et al., 2010; Sadove & Morreale, 1989; Unger et al., 2016).

Without improved waste management and infrastructure in undeveloped coastal countries worldwide, the cumulative quantity of plastic waste available to enter the ocean from land is predicted to increase by an order of magnitude by 2025 (Jambeck et al., 2015). There have been not marine debris surveys in the Mariana Islands, so information from the Hawaiian Islands may be relevant. Since 1996, NOAA has removed 848 metric tons of derelict fishing nets and debris from the Northwest Hawaiian Islands and has estimated that an additional 52 tons of debris collects on the shallow coral reefs and shores there every year (National Oceanic and Atmospheric Administration Fisheries, 2018). From April 2013 to April 2016 in the waters around Lanai and channels between Lanai, Maui, and Kahoolawe, surveys were conducted to quantify the presence of marine mammals and floating marine debris (Currie et al., 2017). The surveys encountered, collected, and categorized 1,027 pieces of marine debris. Items categorized as "plastic" were the predominant type of debris encountered, accounting for 86 percent of total debris and consisting mainly of plastic bottles, tubs, baskets, foamed polystyrene disposable plates, cups, fragments, plastic bags, and other soft plastic films. A smaller portion of the plastic debris (13 percent; 11 percent of the total debris) was fishing-related and included items such as buoys, netting, rope, and fishing lines; milled lumber and rubber accounted for 10 percent of debris, with the remaining 4 percent attributed to metal, glass, and clothing/fabric. Similar findings have been documented for other locations in the Pacific region (Choy & Drazen, 2013; Horton et al., 2017; Smith, 2012).

Although there are no similar records for the Study Area, for some of the same cetacean species in Hawaii between 2007 and 2012, there were 48 humpback whales, a sperm whale, a bottlenose dolphin, 3 spinner dolphins, and a pantropical spotted dolphin found entangled in marine debris (Bradford & Lyman, 2015). One humpback whale was known to be injured, and it is believed that interaction with debris led to the mortality of a second humpback whale and a spinner dolphin (Bradford & Lyman, 2015). Marine mammals migrating from the Mariana Islands to Alaska also encounter threats outside the Study Area. In Alaska from 2011 through 2015, records of approximately 3,700 human-marine mammal interactions were reviewed by NMFS and determined to have resulted in 440 entanglement/entrapment-related marine mammal serious injury or mortality to various species (Helker et al., 2017).

An estimated 75 percent or more of marine debris consists of plastic (Derraik, 2002; Hardesty & Wilcox, 2017). High concentrations of floating plastic have been reported in the central areas of the Pacific Ocean (Cozar et al., 2014). Plastic pollution found in the oceans is primarily dominated by particles smaller than 1 centimeter, commonly referred to as microplastics (Hidalgo-Ruz et al., 2012). Other researchers have defined microplastics as particles with a diameter ranging from a few micrometers up to 5 millimeters (mm) that are not readily visible to the naked eye (Andrady, 2015). Microplastic fragments and fibers found throughout the oceans result from the breakdown of larger items, such as clothing, packaging, and rope and have accumulated in the pelagic zone and sedimentary habitats (Thompson et al., 2004). Results from the investigation by Browne et al. (2011) have also suggested that microplastic fibers are discharged in sewage effluent resulting from the washing of synthetic fiber clothes. DeForges et al. (2014) sampled the Northeast Pacific Ocean in areas in and near the coastal waters of British Columbia, Canada, and found microplastics (those 62–5,000 micrometers in size) were abundant in all samples with elevated concentrations near urban centers. This finding should be applicable to all urban centers in the Pacific, including those in the Study Area. Besseling et al. (2015) documented the first occurrence of microplastics in the intestines of a humpback whale; while the primary cause of the stranding was not determined, the researchers found multiple types of microplastics ranging in sizes from 1 mm to 17 centimeters. There is still a large knowledge gap about possible negative effects of microplastics, but it remains a concern (Besseling et al., 2015). Specifically, the propensity of plastics to absorb and concentrate dissolved pollutant chemicals, such as persistent organic pollutants, is a concern because microfauna may be able to digest plastic nanoparticles, facilitating the delivery of dissolved pollutant chemicals across trophic levels and making them bioavailable to larger marine organisms, such as marine mammals (Andrady, 2015).

Marine mammals as a whole are subject to the various influences and factors delineated in this section above. If specific threats to individual species in the Study Area are known, those threats are described below in individual species accounts.

## ***Mysticetes***

### **3.4.1.8 Blue Whale (*Balaenoptera musculus*)**

#### **3.4.1.8.1 Status and Management**

The blue whale is listed as endangered under the ESA and as depleted under the MMPA throughout its range, but there is no designated critical habitat for this species. NMFS has determined that more research is still needed to rigorously and specifically define the features that make habitat important to blue whales (National Marine Fisheries Service, 2018b). Although the designated Central North Pacific Stock of blue whales are present in winter in “lower latitudes in the western and central Pacific,

including Hawaii,” blue whales in the Study Area have not been assigned to a stock in either the Alaska or Pacific SARs (Carretta et al., 2017d; Muto et al., 2017b).

#### **3.4.1.8.2 Geographic Range and Distribution**

Blue whales inhabit all oceans and typically occur in both nearshore and deep oceanic waters. Blue whales belonging to the Central Pacific Stock feed in summer in the Pacific south of the Aleutian Islands and in the Gulf of Alaska, and then migrate to lower latitudes in the winter. There are no recent sighting records for blue whales in the Study Area (Fulling et al., 2011; Hill et al., 2017a; Uyeyama, 2014). Although rare, acoustic detections from passive monitoring devices deployed at Saipan and Tinian have recorded the presence of blue whales over short periods of time (a few days) (Oleson et al., 2015). However, since blue whale calls can travel up to 621 miles (mi.) (1,000 kilometers [km]), it is unknown whether the animals were actually within the Study Area. Blue whales would be most likely to occur in the Study Area during the winter and are expected to be few in number.

#### **3.4.1.8.3 Population and Abundance**

Widespread whaling over the last century was believed to have decreased the global blue whale population to approximately 1 percent of its pre-whaling population size (Branch, 2007; Monnahan, 2013; Monnahan et al., 2014; Rocha et al., 2014; Širović et al., 2004). The most current information suggests that following the cessation of commercial whaling in 1971, the population in the North Pacific may have recovered and since the 1990s has been at a stable level despite the impacts of ship strikes, interactions with fishing gear, and increased levels of ambient sound in the Pacific Ocean (Campbell et al., 2015; Carretta et al., 2017c; Carretta et al., 2018b; International Whaling Commission, 2016; Monnahan, 2013; Monnahan et al., 2014; Monnahan et al., 2015; Rockwood et al., 2017; Širović et al., 2015). For the portion of the population present in the eastern Pacific, findings have suggested that the population is now near the environment’s carrying capacity and that the rate of change of the population size has declined as a result (Carretta et al., 2018b; International Whaling Commission, 2016; Monnahan et al., 2014; Monnahan et al., 2015).

#### **3.4.1.8.4 Predator-Prey Interactions**

Blue whales feed almost exclusively on various types of zooplankton, especially krill (Jefferson et al., 2015). Blue whales with data recording tags have been recorded feeding from the surface to depths approaching 300 m (Goldbogen et al., 2013a; Goldbogen et al., 2013b).

Blue whales have been documented to be preyed on by killer whales, and 25 percent of photo-identified whales in the Gulf of California carry rake scars from killer whale attacks (Jefferson et al., 2015; Pitman et al., 2007; Sears & Perrin, 2009).

#### **3.4.1.8.5 Species-Specific Threats**

Blue whales are susceptible to entanglement in fishing gear and ship strikes (Berman-Kowalewski et al., 2010; Calambokidis et al., 2009a; Calambokidis, 2012; Carretta et al., 2013; Carretta et al., 2016b; Laggner, 2009; Monnahan et al., 2015; National Marine Fisheries Service, 2011e; Rockwood et al., 2017).

### **3.4.1.9 Bryde’s Whale (*Balaenoptera edeni*)**

#### **3.4.1.9.1 Status and Management**

Bryde’s whales are not listed as endangered under the ESA. There is currently no biological basis for defining separate stocks of Bryde’s whales in the western or central North Pacific. (Carretta et al., 2017c; Carretta et al., 2017d; Constantine et al., 2018). NMFS recognizes two stocks of Bryde’s whales in the

Pacific with one for Hawaiian waters and the other for the Gulf of California and waters off California (Carretta et al., 2017c; Carretta et al., 2017d); none of the ranges described for these stocks include the Study Area.

#### **3.4.1.9.2 Geographic Range and Distribution**

Data suggest that winter and summer grounds partially overlap in the central North Pacific (Murase et al., 2015). Bryde's whales are distributed in the central North Pacific in summer; the southernmost summer distribution of Bryde's whales inhabiting the central North Pacific is about 20° N (Kishiro, 1996). Some whales remain in higher latitudes (around 25° N) in both winter and summer, but are not likely to move poleward of 40° N (Jefferson et al., 2015; Kishiro, 1996). Bryde's whales are expected to be present in the Study Area based on sighting records (Fulling et al., 2011; Hill et al., 2017a; Mobley, 2007; Oleson & Hill, 2010a; Uyeyama, 2014). Bryde's whales were detected in the Transit Corridor between the Study Area and Hawaii during a NMFS survey in January 2010 (Oleson & Hill, 2010a) and regularly encountered during the 2007 survey of the MITT Study Area<sup>2</sup> (Fulling et al., 2011). Bryde's whales were encountered during a NMFS cetacean survey in May–June 2015 off Pagan, Alamagan, and south of Guam (Hill et al., 2018b; Oleson, 2017), and later that same year off Rota during a small boat survey in August–September 2015 (Hill et al., 2017a). In May 2017 a single individual was encountered in deep water off the west side of Saipan (Hill et al., 2018a).

#### **3.4.1.9.3 Population and Abundance**

Based on the best available science, there are an estimated 233 (Coefficient of Variation [CV] = 0.45) Bryde's whales present in the Study Area (Fulling et al., 2011).

#### **3.4.1.9.4 Predator-Prey Interactions**

Bryde's whales primarily feed on schooling fish and are lunge feeders. Prey includes anchovy, sardine, mackerel, herring, krill, and other invertebrates, such as pelagic red crab (Baker & Madon, 2007; Nemoto & Kawamura, 1977). Bryde's whales have been observed using "bubble nets" to herd prey (Kato & Perrin, 2009). Bubble nets are used in a feeding strategy where the whales dive and release bubbles of air that float up in a column and trap prey inside where whales lunge through the column to feed.

Bryde's whale is known to be prey for killer whales, as evidenced by an aerial observation of 15 killer whales attacking a Bryde's whale in the Gulf of California (Weller, 2009).

#### **3.4.1.9.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Bettridge et al., 2015; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that four products contained

---

<sup>2</sup> The Mariana Islands Sea Turtle and Cetacean Survey (MISTCS) undertaken in 2007 covered an area of approximately 301,300 square kilometers within the larger MITT Study Area, which encompasses approximately 1,300,000 square kilometers; see Chapter 2 (Description of Proposed Action and Alternatives) for more details with regard to the extent of the MITT Study Area. The MISTCS abundance estimates from the 2007 survey as reported in Fulling et al. (2011) and cited throughout this section, thus represent the number of marine mammals estimated to be present in the approximately 24 percent of the MITT Study Area covered by the MISTCS survey.

Bryde's whale meat (Baker et al., 2006a). There has been one recorded stranding of a Bryde's whale (at Tinian in 2005) within the Study Area (Trianni & Tenorio, 2012).

#### **3.4.1.10 Fin Whale (*Balaenoptera physalus*)**

##### **3.4.1.10.1 Status and Management**

Fin whales are listed as endangered under the ESA, but there is no designated critical habitat for this species. The stock structure of fin whales remains uncertain (Mizroch et al., 2009), and fin whales in the Study Area have not been assigned to a stock in the current SARs (Carretta et al., 2017d; Muto et al., 2017b). NMFS recognizes three stocks of fin whales in the North Pacific (Carretta et al., 2017d; Muto et al., 2017b), and none of the ranges described for these stocks include the Study Area.

##### **3.4.1.10.2 Geographic Range and Distribution**

Fin whales prefer temperate and polar waters; they are rarely seen in warm, tropical waters and are not expected south of 20°N latitude (Miyashita et al., 1996; Reeves et al., 2002). There are no sighting records for fin whales in the Study Area (Fulling et al., 2011; Hill et al., 2017a; Oleson et al., 2015; Uyeyama, 2014). Based on acoustic detections, fin whales are expected to be seasonally present in the Study Area although few in number. Acoustic detections from passive monitoring devices deployed at Saipan and Tinian have recorded the presence of fin whales over short (a few days) periods of time (Oleson et al., 2015), and fin whale vocalizations were detected in January 2010 in the Transit Corridor between Hawaii and Guam (Oleson & Hill, 2010a). Fin whales were not, however, detected in the Transit Corridor using the same equipment and methods in May of that year (Oleson & Hill, 2010a).

##### **3.4.1.10.3 Population and Abundance**

There is no current abundance estimate available for fin whales in the Study Area (Carretta et al., 2017d; Muto et al., 2017b). There were approximately 50,000 reported fin whales killed during commercial whaling in the North Pacific from 1911 to 1985 (C. Allison, pers. comm. as provided in Mizroch et al. (2009), and it is assumed the population is still recovering.

##### **3.4.1.10.4 Predator-Prey Interactions**

Fin whale prey vary by region and may include krill, small invertebrates such as copepods, squid, and schooling fish such as capelin, herring, pollock, and mackerel (Goldbogen et al., 2006; Jefferson et al., 2015; Mizroch et al., 2009).

The fin whale is not known to have a significant number of predators. However, in regions where killer whales are abundant, some fin whales exhibit attack scars on their flippers, flukes, and flanks, suggesting possible predation by killer whales (Aguilar de Soto et al., 2008).

##### **3.4.1.10.5 Species-Specific Threats**

Fin whales are susceptible to entanglement in fishing gear (Carretta et al., 2013; Carretta et al., 2016b; Carretta et al., 2017a; Helker et al., 2015). Given this and as discussed in Section 3.4.1.7.3 (Bycatch) for other species, entanglement risk includes fishing activities out of Japan and South Korea.

#### **3.4.1.11 Humpback Whale (*Megaptera novaeangliae*)**

##### **3.4.1.11.1 Status and Management**

Humpback whales in the Study Area are indirectly addressed in the Alaska SAR given that the historic range of humpbacks in the "Asia wintering area" includes the Mariana Islands. The detected presence of humpbacks in the Mariana Islands (Fulling et al., 2011; Hill et al., 2016a; Hill et al., 2017a; Hill et al.,

2018a; Klinck et al., 2016a; Munger et al., 2014; Oleson et al., 2015; Uyeyama, 2014) are consistent with the Study Area as a plausible migratory destination for humpback whales from Alaska (Muto et al., 2017a; Muto et al., 2017b).

Effective as of October 11, 2016, NMFS changed the status of all humpback whales from an endangered species to a specific status for each of 14 newly identified distinct population segments (DPS) (81 FR 62259). For the Study Area, the Navy believes it is likely that humpback whales in the Mariana Islands are part of the endangered Western North Pacific DPS based on the available science (Bettridge et al., 2015; Calambokidis et al., 2008; Calambokidis et al., 2010; Carretta et al., 2017c; Carretta et al., 2017d; Hill et al., 2017b; Muto et al., 2017a; Muto et al., 2017b; National Marine Fisheries Service, 2016e; National Oceanic and Atmospheric Administration, 2015a, 2018c; Wade et al., 2016). Humpback whales from the winter range of the Western North Pacific DPS (including the Study Area) that feed in the summer off Russia and Alaska have been designated by NMFS as the Western North Pacific Stock (Muto et al., 2017a; Muto et al., 2017b). As part of the Western North Pacific Stock, the population is considered depleted under the MMPA (Muto et al., 2017a; Muto et al., 2017b).

There has been no critical habitat designated for the Western North Pacific Distinct Population Segment.

#### **3.4.1.11.2 Geographic Range and Distribution**

Between 1948 and 1979, Soviet Union commercial whaling alone took 7,344 humpback whales from the North Pacific (Ilyashenko & Chapham, 2014). It is therefore likely that humpback whales in the western North Pacific are still recovering and will remain rare in parts of their former range. Researchers have reported that it is not clear whether humpback whales use the Mariana Islands as a winter breeding and calving area or as a corridor from one or more wintering areas when moving to summertime feeding area locations, which are also unknown (Hill et al., 2016a).

For purposes of the analysis presented in this SEIS/OEIS, the Navy assumes humpback whales in the Study Area are part of the endangered Western North Pacific DPS.<sup>3</sup> This population segment is based on a known breeding group of individuals found off Okinawa and Ogasawara Islands (approximately 1,230 nautical miles [NM] north of Guam) in Japan waters and in Philippine waters (approximately 1,350 NM west of Guam), as identified by photographic identification of individuals (Calambokidis et al., 2008; Calambokidis et al., 2010), in addition to an “unknown breeding group” from a location in the western North Pacific that remained unidentified until recently (National Oceanic and Atmospheric Administration, 2018c). Humpback whales found off Okinawa, Ogasawara, the Philippines, and the unknown area were combined to form the Western North Pacific population (Bettridge et al., 2015). This “unknown area” corresponds to the historical range for the western North Pacific that included waters extending from the South China Sea east through the Philippines, the Ryukyu Islands, Mariana

---

<sup>3</sup> There is reference to a “Second West Pacific DPS” in the latest NMFS humpback whale status report (Bettridge et al., 2015), although that terminology did not carry over into the rule-making establishing the 14 distinct population segments. As a result, although the humpback whales in the Study Area may exactly fit the parameters of the intended “Second West Pacific DPS,” in this SEIS/OEIS the Navy has assumed that humpback whales in the Study Area are part of the Western North Pacific Distinct Population Segment consistent with the determinations presented in 81 FR 62259, the range for the Western North Pacific Stock as presented in the Alaska Stock Assessment Report (Muto et al., 2017a; Muto et al., 2017b; Muto et al., 2018), and the findings based on a series of small boat surveys in the Mariana Islands (National Marine Fisheries Service, 2018c).

Islands, and Marshall Islands and from there, north to the Arctic (Muto et al., 2017a; Muto et al., 2017b; Rice, 1998). Genetic and photographic data collected during Navy-funded small boat surveys has provided matches to individuals identified many years previously off the Ogasawara Islands and the Western North Pacific DPS (Hill et al., 2017a; Hill et al., 2017b; Hill et al., 2018a).

Navy aerial monitoring surveys occurring at Farallon de Medinilla conducted monthly from 1997 to 2009 and irregularly thereafter documented the occasional presence of humpback whales, including mother-calf pairs and other adult individuals (Uyeyama, 2014). Shipboard survey in the MITT Study Area in February 2007 acoustically detected and subsequently sighted an estimated group of eight humpbacks at Marpi Reef north of Saipan (Fulling et al., 2011; Norris et al., 2012a, 2014). Small boat surveys in 2010 and 2014 off Guam, Saipan, Tinian, Aguijan, and Rota did not encounter humpback whales (Hill et al., 2014). The next documented observations of humpback whales in the Mariana Islands occurred from February 26 to March 8, 2015, when four mother/calf pairs and four other individual humpback whales were observed at Chalan Kanoa Reef off Saipan (Hill et al., 2015a; Hill et al., 2016b). During the subsequent NMFS Mariana Archipelago Cetacean Survey (two months later; May 8 to June 6, 2015), survey transects sampling all the Mariana Islands out to 50 NM from shore detected no humpback whales visually or acoustically in the Mariana Islands (Hill et al., 2018b; Oleson, 2017). Humpback whales were seen again off Saipan during Navy-funded surveys in 2016, 2017, and 2018 (Hill et al., 2017a; Hill et al., 2017b; Hill et al., 2018a; National Oceanic and Atmospheric Administration, 2018c); see Appendix I (Geographic Mitigation Assessment) for additional details regarding all humpback whale sightings in the MITT Study Area. These Navy-funded small boat survey investigations have included photo-identification and genetic sampling and have resulted in the documentation of mother-calf pairs, competitive groups, and 35 additional photo-identified non-calf whales (Fulling et al., 2011; Hill et al., 2017a; Hill et al., 2017b; Hill et al., 2018a; National Oceanic and Atmospheric Administration, 2018c; Norris et al., 2012a, 2014). The presence of newborn calves and competitive groups documented during the aforementioned small boat surveys confirm the Mariana Islands are serving as a breeding location for Western North Pacific DPS humpback whales (National Oceanic and Atmospheric Administration, 2018c). Investigations in Hawaii over consecutive breeding seasons between 1997 through 2008 in the Maui Basin found a preference by individual mother-calf pairs for both water depth and sea-bed terrain type, with the pair moving into deeper water and rougher terrain as a calf matured (Pack et al., 2017); this habitat preference by mother-calf pairs may also be present in the Mariana Islands.

Based on a compendium of all detections, humpback whales have been sighted in the Study Area in the months of January through March (U.S. Department of the Navy, 2005; Uyeyama, 2014), male humpback songs have been recorded from December through April, and humpback whale sounds were infrequently detected at Tinian during June to October (Hill et al., 2017a; Klinck et al., 2016a; Munger et al., 2014; Norris et al., 2014; Oleson et al., 2015). Humpback whales were not observed or acoustically detected in the Transit Corridor during a May 2010 survey (Oleson & Hill, 2010a), which is consistent with the presumption that, except when migrating to or from summer feeding areas, humpback whales will most likely be present in relative shallow water locations in the vicinity of the Mariana Islands.

Humpback whales from the Western North Pacific, Hawaii, and Mexico DPSs overlap to some extent on feeding grounds off Alaska (Bettridge et al., 2015; Muto et al., 2017a; Muto et al., 2017b; National Marine Fisheries Service, 2016a; Titova et al., 2017; Wade et al., 2016). Photographic identification data have also documented the presence of at least one whale seen multiple years off Ogasawara (Japan) later seen feeding off British Columbia (Darling et al., 1996), indicating there may be greater overlap of DPSs in the summer feeding areas than has been characterized in the SARs for Alaska and the Pacific

(Carretta et al., 2017c; Carretta et al., 2017d; Carretta et al., 2018a; Muto et al., 2017a; Muto et al., 2017b). Comparison of photographic identification data from Russian waters (where the Western North Pacific DPS humpback whales may also feed) has found 35 individual whales that were also documented in Hawaii and 11 that were from the Mexican breeding grounds (Titova et al., 2017).

#### **3.4.1.11.3 Population and Abundance**

Based on photographic identifications off Okinawa and Ogasawara gathered previously and conclusions reached in 2008 (Calambokidis et al., 2008), the abundance of humpback whales in the Western North Pacific population was estimated to be approximately 1,000 individuals (Bettridge et al., 2015; Muto et al., 2017a). From that same data set, the growth rate of the Western North Pacific Distinct Population Segment was estimated to be 6.9 percent (Bettridge et al., 2015; Calambokidis et al., 2008). This can be viewed in context of the North Pacific population, which has been increasing at a rate of between 5.5 percent and 6.0 percent per year, approximately doubling every 10 years (Bettridge et al., 2015; Muto et al., 2017a; Wade et al., 2016). The inclusion of more recent data from photographic identifications off Okinawa has documented the presence of at least 1,402 unique individuals in the Western North Pacific Distinct Population Segment (Kobayashi et al., 2016). Additional information from Navy-funded surveys and passive acoustic hydrophone recordings in the Mariana Islands has confirmed the presence of mother-calf pairs, non-calf whales, and singing males in the Study Area (Fulling et al., 2011; Hill et al., 2016a; Munger et al., 2014; Munger et al., 2015; National Oceanic and Atmospheric Administration, 2018c; Norris et al., 2014; Oleson & Hill, 2010a; Oleson et al., 2015; Uyeyama et al., 2012). The NMFS Alaska SAR provides a population estimate for humpbacks in Ogasawara Islands, Okinawa, and the Philippines of 1,107 animals, with a minimum population of 865, noting that these are likely to be an underestimate of the Western North Pacific Stock's true abundance (Muto et al., 2017a; Muto et al., 2017b). Although not specific to the Study Area, the overall abundance of humpback whales in the North Pacific was recently estimated at 21,808 individuals (Carretta et al., 2017c; Carretta et al., 2017d), confirming that this population of humpback whales has continued to increase and is now greater than some pre-whaling abundance estimates (Barlow et al., 2011; Bettridge et al., 2015; Muto et al., 2017a; Muto et al., 2017b; Wade et al., 2016).

#### **3.4.1.11.4 Predator-Prey Interactions**

When on the summer feeding grounds in Alaska, humpback whales from the Study Area feed on a wide variety of invertebrates and small schooling fishes. The most common invertebrate prey are krill (tiny crustaceans); the most common fish prey are herring, mackerel, sand lance, sardines, anchovies, and capelin (Clapham & Mead, 1999). Feeding occurs both at the surface and in deeper waters, wherever prey is abundant. Humpback whales are the only species of baleen whale that show strong evidence of cooperation when they feed in large groups (D'Vincent et al., 1985).

This species is known to be attacked by both killer whales and false killer whales as evidenced by tooth rake scars on their bodies and fins (Jefferson et al., 2015; Steiger et al., 2008).

#### **3.4.1.11.5 Species-Specific Threats**

Based on data from Alaska (including the Western North Pacific Stock), Hawaii, and the U.S. Pacific coast (Bradford & Lyman, 2015; Carretta et al., 2016c; Carretta et al., 2017b; Helker et al., 2017), humpback whales are subject to risk from entanglement in marine debris and active fishing gear; most often recorded is pot/trap fishery gear. As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under

Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that three products contained humpback whale meat (Baker et al., 2006a).

The mean vessel collision mortality and serious injury rate in Alaska is 4.3 humpback whales annually (Muto et al., 2017a), but that rate reflects Southeast Alaska waters where the presence of the Western North Pacific Stock is less likely.

#### **3.4.1.12 Minke Whale (*Balaenoptera acutorostrata*)**

##### **3.4.1.12.1 Status and Management**

Minke whales are not listed as endangered under the ESA. The stock structure for minke whales remains uncertain in the Pacific, and minke whales in the Study Area have not been assigned to a stock in the current SARs (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). NMFS recognizes three stocks of minke whales in the North Pacific: (1) the Hawaii Stock, (2) the California/Oregon/Washington Stock, and (3) the Alaska Stock (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b).

##### **3.4.1.12.2 Geographic Range and Distribution**

Surveys employing towed hydrophone arrays and sonobuoys, and long-term monitoring efforts using fixed passive acoustic recording devices, have routinely detected the presence of minke whales in the Study Area (Klinck et al., 2016b; Norris et al., 2017; Oleson & Hill, 2010a; Oleson et al., 2015). Minke whales have not been visually detected in the Study Area during any survey efforts within approximately the last decade although they are the most common acoustically detected mysticete in the area (Fulling et al., 2011; Hill et al., 2011; Hill et al., 2013a; Hill et al., 2014; Hill et al., 2015b; Hill et al., 2017a; Mobley, 2007; Norris et al., 2014; Oleson & Hill, 2010a; Tetra Tech Inc., 2014; Uyeyama, 2014).

##### **3.4.1.12.3 Population and Abundance**

No estimates have been made for the number of minke whales in the North Pacific (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). Acoustic data collected during a Navy-funded 2007 line-transect survey employing a towed hydrophone array in the Mariana Islands were used to estimate a minimum abundance of calling minke whales (Norris et al., 2017). Abundance was estimated using two different methodologies, resulting in minimum estimates of 80 or 91 animals in the surveyed area (a density of 0.13 and 0.15 animals per 1,000 km<sup>2</sup>, respectively; CV = 0.34) (Norris et al., 2017). This study provided the first abundance and density estimates for calling minke whales and the first minimum estimates by which the number of minke whales in the Mariana Islands region could be derived.

##### **3.4.1.12.4 Predator-Prey Interactions**

##### **3.4.1.12.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk for the population of minke whales includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Dalebout et al., 2002a; Lukoschek et al., 2009). For example in 2008, the reported bycatch in Japan and South Korea totaled 214 minke whales (Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that 230 products contained minke whale meat (Baker et al.,

2006a). In the two-year period between 2013 and 2014, the total bycatch by South Korean fisheries in the East Sea totaled a reported 48 minke whales (Song, 2017).

#### **3.4.1.13 Omura's Whale (*Balaenoptera omurai*)**

##### **3.4.1.13.1 Status and Management**

Omura's whale is not listed under the ESA and is not mentioned in the Pacific or Alaska SARs (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). There is no managed stock or population within U.S. waters pursuant to the MMPA, but the species is protected under that statute nonetheless, as are all marine mammals.

##### **3.4.1.13.2 Geographic Range and Distribution**

The species was first described in 2003 based on eight specimens taken by Japanese research whaling vessels in the Sea of Japan, the Solomon Sea, and the eastern Indian Ocean (Wada et al., 2003). Records of the species from Philippines shore-based whaling provide additional indication of a broad distribution that includes the western Pacific (Cerchio et al., 2015). Given the documented occurrence of the species, it is assumed the species may be present in the Study Area. Recent well-documented sightings have occurred in nearshore waters off Madagascar and off Sri Lanka, indicating in those cases a preference for relatively shallow water less than approximately 200 m in depth (Cerchio et al., 2015; de Vos, 2017).

##### **3.4.1.13.3 Population and Abundance**

There are no data available to estimate abundance for Omura's whale in the Study Area.

##### **3.4.1.13.4 Predator-Prey Interactions**

Observations of feeding Omura's whales in waters off Madagascar suggested the animals were skim feeding on zooplankton given there was an absence of fish or other observable prey (Cerchio et al., 2015).

##### **3.4.1.13.5 Species-Specific Threats**

An individual Omura's whale observed in Sri Lanka waters showed evidence of an entanglement scar on the left side of its upper jaw, indicating that entanglement is a potential threat for this species (Cerchio et al., 2015; de Vos, 2017). One Omura's whale was reported struck by a fishing boat in the Philippines (Obusan et al., 2016).

#### **3.4.1.14 Sei Whale (*Balaenoptera borealis*)**

##### **3.4.1.14.1 Status and Management**

Sei whales are listed as endangered under the ESA, but there is no designated critical habitat for this species. The stock structure for sei whales is uncertain in the Pacific (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes three stocks of sei whales in the North Pacific: (1) the Hawaii Stock, (2) the California/Oregon/Washington Stock, and (3) the Alaska Stock (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). The western Pacific and waters within the Study Area have not been addressed by NMFS, and sei whales in the Study Area have not been assigned to a stock (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b).

##### **3.4.1.14.2 Geographic Range and Distribution**

In a January-February survey in 1972, a single group of approximately 13 sei whales were sighted during a survey of the Mariana Islands and Ogasawara (Masaki, 1972). In the 2007 survey of the Mariana Islands (Fulling et al., 2011), a total of 16 sei whales were sighted. Sei whale calls documented during the

2007 survey indicated a greater variability in the vocal repertoire of sei whales than documented elsewhere (Norris et al., 2014), which may have contributed to the lack of acoustic detections in the three-year record from 2010 to 2013 (Oleson et al., 2015). Sei whales were also visually detected in the Transit Corridor between the Study Area and Hawaii during a NMFS survey in January 2010 (Oleson & Hill, 2010a).

#### **3.4.1.14.3 Population and Abundance**

During a 2007 systematic survey of the Study Area, sei whales were sighted on 16 occasions with a resulting abundance estimate of 166 individuals (CV = 0.49) (Fulling et al., 2011).

#### **3.4.1.14.4 Predator-Prey Interactions**

In the North Pacific, sei whales feed on a diversity of prey, including copepods, krill, fish [specifically sardines and anchovies], and cephalopods [squids, cuttlefish, octopuses] (Horwood, 1987; Horwood, 2009; Nemoto & Kawamura, 1977). Feeding occurs primarily around dawn, which appears to be correlated with vertical migrations of prey species (Horwood, 2009). As with other mysticetes, sei whales have been observed lunging and gulping dense concentrations of prey, but more often tend to obtain prey by skimming (Horwood, 2009). Sei whales, like other large baleen whales, are likely subject to occasional attacks by killer whales.

#### **3.4.1.14.5 Species-Specific Threats**

Based on the discovery of a sei whale entangled in rope and fishing gear in Hawaii that presumably came from Alaska (Bradford & Lyman, 2015), sei whales may be subject to entanglement from fishery activity taking place in the western Pacific, including the Study Area. Based on the statistics of other large whales along the U.S. Pacific coast and Alaska (Carretta et al., 2016b; Carretta et al., 2017b; Helker et al., 2017), it is likely that ship strikes also pose a threat to sei whales in the Study Area from commercial vessels transiting that area.

### ***Odontocetes***

#### **3.4.1.15 Blainville's Beaked Whale (*Mesoplodon densirostris*)**

##### **3.4.1.15.1 Status and Management**

Blainville's beaked whale is not listed under the ESA. The stock structure for Blainville's beaked whales remains uncertain in the western Pacific, and Blainville's beaked whales in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes a single stock of Blainville's beaked whales in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.15.2 Geographic Range and Distribution**

Blainville's beaked whales are one of the most widely distributed of the toothed whales within the *Mesoplodon* genus, occurring in temperate and tropical deep waters areas in all oceans (Jefferson et al., 2015; MacLeod, 2000; MacLeod & Mitchell, 2006). In Hawaii, some populations have been documented to be long-term residents to particular areas (Baird et al., 2009b; Baird, 2011; Baird et al., 2015; McSweeney et al., 2007). There were two *Mesoplodon* whale sightings during the 2007 survey of the Study Area, over the West Mariana Ridge, but they were not identified to the species level (Fulling et al., 2011). During the 2015 NMFS survey of the Mariana Islands, two groups of Blainville's beaked whales were identified and photographed in addition to seven other beaked whale sightings identified only as *Mesoplodon* beaked whales (Hill et al., 2018b; Oleson, 2017). During Navy-funded 2010–2018 small boat

surveys in the Mariana Islands, five *Mesoplodon* beaked whales were encountered on two occasions in a median depth of approximately 1,140 m and median approximate distance from shore of 15 km (Hill et al., 2013a; Hill et al., 2017a; Hill et al., 2018b). It could not be determined if these were Blainville's beaked whales or ginkgo-toothed beaked whales, both of which belong to the genus *Mesoplodon* and are believed to be present in the Study Area. Acoustic monitoring has indicated that Blainville's beaked whales occur regularly and year-round in the Study Area (Klinck et al., 2016b; Oleson et al., 2015; Tetra Tech Inc., 2014). Although there is no record of similar occurrences in the Mariana Islands, it has been suggested that the Philippines are a stranding "hot spot" for Blainville's beaked whales in Asia (Bachara & Blatchley, 2018).

#### **3.4.1.15.3 Population and Abundance**

There are no abundance estimates for Blainville's beaked whales in the Study Area.

#### **3.4.1.15.4 Predator-Prey Interactions**

All beaked whales probably feed at or close to the bottom in deep oceanic waters, taking suitable prey opportunistically or as locally abundant, typically by suction feeding (Heyning & Mead, 1996; Jefferson et al., 2015; Werth, 2006a, 2006b). Feeding may also occur at mid-water as shown by tagging data from Blainville's beaked whales (Baird et al., 2005; Baird et al., 2006c). Blainville's beaked whales are known to echolocate in groups when they are on foraging dives, which makes them more easily detectable by passive acoustic means (Moretti & Baird, 2015). *Mesoplodon* beaked whales have been observed being actively preyed upon by killer whales (Wellard et al., 2016).

#### **3.4.1.15.5 Species-Specific Threats**

There were five observed interactions between an unidentified beaked whale, unidentified Mesoplodont beaked whale, or Blainville's beaked whale and longline fishing activities in Hawaiian waters between 2010 and 2014 (Bradford & Forney, 2016, 2017). As similar information for U.S. fishing vessels or foreign fishing vessels in the Study Area is unavailable, this data from Hawaii provides information regarding the species interactions with fishing activities in general. As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated at least one product contained Blainville's beaked whale meat (Baker et al., 2006a).

#### **3.4.1.16 Common Bottlenose Dolphin (*Tursiops truncatus*)**

##### **3.4.1.16.1 Status and Management**

Bottlenose dolphin is not listed under the ESA. The stock structure for bottlenose dolphin remains uncertain in the western Pacific and the Mariana Islands (Martien et al., 2014b), and bottlenose dolphins in the Study Area have not been assigned to a stock in the current Pacific SAR (Carretta et al., 2017c; Carretta et al., 2017d). Other than small and resident Main Hawaiian island-associated populations of bottlenose dolphins, NMFS recognizes a single pelagic stock of bottlenose dolphin in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.16.2 Geographic Range and Distribution**

Multiple fishery interactions with bottlenose dolphins in the western North Pacific (Miyashita, 1993b) indicated their presence beginning approximately 400 NM north of the Study Area. It is possible that

bottlenose dolphins do not occur in great numbers in the Mariana Island chain, but they have been frequently sighted, although in small numbers. In the main Hawaiian Islands, data suggest that bottlenose dolphins exhibit site fidelity (Baird et al., 2009a; Baird et al., 2013c; Martien et al., 2012). Gannier (2002) noted that large densities of bottlenose dolphins do not occur at the Marquesas Islands and attributed this to the area's lack of a significant shelf component, which would be similar to the MITT Study Area.

Common bottlenose dolphins are generally found in coastal and continental shelf waters of tropical and temperate regions of the world and are known to occur in small enclosed bays or harbors (Martien et al., 2012; Rossman et al., 2015; Wells & Scott, 2009), but they have not been detected in any such enclosed water in the Study Area (such as Apra Harbor). During the 2007 survey of the Mariana Islands, there were three sightings of bottlenose dolphins to the east of Saipan in deep waters near the Mariana Trench (Fulling et al., 2011). Bottlenose dolphins were not detected during the 2010 survey of the Mariana Islands and the Transit Corridor (Oleson & Hill, 2010a), but were detected on three occasions during the 2015 NMFS cetacean survey of the Mariana Islands (Hill et al., 2018b; Oleson, 2017). In total during Navy-funded 2010–2018 small boat surveys in the Mariana Islands, bottlenose dolphins were encountered on 36 occasions in a median depth of approximately 700 m and median approximate distance from shore of 7 km (Hill et al., 2017a; Hill et al., 2018a; Hill et al., 2018b). On multiple occasions, encounters with bottlenose dolphins in the Mariana Islands have involved a mixed-species aggregation with one other species that have included short-finned pilot whales, pantropical spotted dolphins, false killer whales, spinner dolphins, or rough-toothed dolphins (Hill et al., 2011; Hill et al., 2018b). In the 2017 follow-on small boat survey, three bottlenose dolphin groups consisting of two to eight individuals were encountered off Saipan in waters with a median depth of 306 m and a median distance from shore of 5.4 km (Hill et al., 2018a).

#### **3.4.1.16.3 Population and Abundance**

In some regions of the Pacific, “inshore” and “offshore” or pelagic species differ genetically and morphologically (Baird et al., 2009a; Baird et al., 2013c; Tezanos-Pinto et al., 2009), but this has not been demonstrated for the Mariana Islands (Martien et al., 2014b). A total of 4,610 photos taken during small boat surveys between 2011 and 2014 were analyzed to identify individual bottlenose dolphins. A total of 47 individuals were identified with 30 individuals (64 percent) re-encountered and the remaining 17 of those individuals (36 percent) encountered three or more times (Hill et al., 2017a). These re-encounters occurred between all islands and may be similar to the site fidelity present for some of the island-associated populations present in the Hawaiian Islands (Baird et al., 2009a). Genetic samples from 21 bottlenose dolphins encountered off Guam and Saipan in 2007 suggest a history of hybridization with Fraser's dolphin (Martien et al., 2014b). The Mariana Islands samples shared DNA haplotypes with individuals from the Philippines, South Korea, Taiwan, and the main Hawaiian Islands but precluded determination of any small populations associated with specific locations in the Mariana Islands similar to what has been found in Hawaii (Martien et al., 2014b).

A bottlenose dolphin abundance estimate of 31,700 animals was made for the area approximately 400 NM north of the Mariana Islands (Miyashita, 1993b). There were three sightings of bottlenose dolphin during a 2007 systematic survey of the Study Area, resulting in an abundance estimate of 122 animals (CV = 0.992) (Fulling et al., 2011).

#### 3.4.1.16.4 Predator-Prey Interactions

Bottlenose dolphins are opportunistic feeders, taking a wide variety of fishes, cephalopods, and shrimps (Wells & Scott, 1999, 2009), and using a variety of feeding strategies (Shane, 1990). In addition to using echolocation, bottlenose dolphins detect and orient fish prey by listening for the sounds their prey produce (i.e., passive listening) (Gannon et al., 2005). Nearshore bottlenose dolphins prey predominantly on coastal fish and cephalopods, while offshore individuals prey on open ocean cephalopods and a large variety of near-surface and mid-water fish species (Mead & Potter, 1995). Throughout their range bottlenose dolphins are known to be preyed on by killer whales and sharks (Ferguson et al., 2012; Heithaus, 2001a; Heithaus, 2001b; Sprogis et al., 2018; Wells & Scott, 1999).

#### 3.4.1.16.5 Species-Specific Threats

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least two products contained bottlenose dolphin meat, reflecting an estimated 23 bottlenose dolphins (Baker et al., 2006a). The stranding of a single bottlenose dolphin in 2013 near Tumon, Guam, is the only known stranding for this species in the Mariana Islands area (Uyeyama, 2014).

#### 3.4.1.17 Cuvier's Beaked Whale (*Ziphius cavirostris*)

##### 3.4.1.17.1 Status and Management

Cuvier's beaked whale is not listed under the ESA. The stock structure for Cuvier's beaked whales remains uncertain in the western Pacific, and Cuvier's beaked whales in the Study Area have not been assigned to a stock in the current SARs (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). With the exception of the U.S. West Coast, NMFS only recognizes a stock of Cuvier's beaked whale in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d) and in the "eastern North Pacific" and Alaskan waters (Muto et al., 2017a; Muto et al., 2017b), whose distribution does not extend to the Study Area.

##### 3.4.1.17.2 Geographic Range and Distribution

Cuvier's beaked whales have an extensive range that includes all oceans, from the tropics to the polar waters of both hemispheres (Ferguson et al., 2006a; Ferguson et al., 2006b; Jefferson et al., 2008; Pitman et al., 1988). Worldwide, beaked whales normally inhabit continental slope and deep oceanic waters. They are commonly sighted around seamounts, escarpments, and canyons (MacLeod et al., 2004). Cuvier's beaked whales are generally sighted in waters with a bottom depth greater than 655 feet (ft.) (200 m) and are frequently recorded in waters with bottom depths greater than 3,280 ft. (1,000 m) (Falcone et al., 2009; Jefferson et al., 2008). While there are indications of potential seasonal redistribution of Cuvier's beaked whales and documented satellite tag movements in Southern California waters (Falcone & Schorr, 2014; Moretti, 2017; Schorr et al., 2014; Schorr et al., 2018), no such research findings are available from the Mariana Islands. A study spanning 21 years off the west coast of the Island of Hawaii suggests that this species may show long-term site fidelity in certain areas (McSweeney et al., 2007).

During aerial surveys conducted in August 2007 covering 2,352 km of linear effort, a single Cuvier's beaked whale was observed about 65 NM south of Guam at the edge of the Mariana Trench (Mobley,

2007). One ziphiid whale (the taxon that Cuvier's beaked whales belong to) was observed in deep water during the 2007 shipboard survey of the Study Area but was not identified to the species level (Fulling et al., 2011). A single Cuvier's beaked whale was sighted and others acoustically detected during an August 2013 survey at Pagan Island (Tetra Tech Inc., 2014). A year's duration of acoustic monitoring at Saipan and at Tinian recorded vocalizing Cuvier's beaked whales (Oleson et al., 2015). These vocalizations were detected in all months having sufficient samples to detect their presence in the Study Area, suggesting there is no seasonal aspect to the Cuvier's beaked whale's distribution.

#### **3.4.1.17.3 Population and Abundance**

There are no abundance estimates for Cuvier's beaked whales in the Study Area.

#### **3.4.1.17.4 Predator-Prey Interactions**

Cuvier's beaked whales, similar to other beaked whale species, are deepwater feeders. Stomach content analyses show that they feed mostly on deep-sea squid, fish, and crustaceans (Hickmott, 2005; Santos et al., 2007). They apparently use suction to swallow prey (Jefferson et al., 2008; Werth, 2006a).

Cuvier's beaked whales may be preyed upon by killer whales (Heyning & Mead, 2009; Jefferson et al., 2008).

#### **3.4.1.17.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least one product contained Cuvier's beaked whale meat (Baker et al., 2006a).

Cuvier's beaked whales were found stranded on Saipan in 2007, on Rizal Beach, Guam, in 2008 (*Ziphius* sp.), and two were found (one alive and one dead) on Micro Beach, Saipan, in 2011 (Uyeyama, 2014). A necropsy conducted on one of the 2011 stranded animals revealed abnormalities in the animal's kidneys and intestines (Hawaii Pacific University, 2012; Saipan Tribune, 2011). Two strandings of Cuvier's beaked whales occurred in 2015; one in March and one in July. The individual that stranded in March 2015 had a severe *Crassicauda* (nematode) infestation in its kidneys (West, 2018). The individual stranded in July 2015 was not accessible for transport until days after the stranding and, because of the resulting decomposition, necropsy and histopathology was not conducted for this whale (West, 2018).

#### **3.4.1.18 Dwarf Sperm Whale (*Kogia sima*)**

##### **3.4.1.18.1 Status and Management**

Dwarf sperm whale is not listed under the ESA. The stock structure for dwarf sperm whales remains uncertain in the western Pacific, and dwarf sperm whales in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). Other than for waters along the U.S. West Coast, NMFS recognizes a single stock of dwarf sperm whale in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.18.2 Geographic Range and Distribution**

Records of this species have been documented from the western Pacific (Taiwan and Japan) (Sylvestre, 1988; Wang et al., 2001; Wang & Yang, 2006), and there have been four known dwarf sperm whale strandings in the Mariana Islands (Trianni & Tenorio, 2012; Uyeyama, 2014).

There were no species of *Kogia* sighted during the 2007 shipboard survey of the Study Area, although this cryptic species is difficult to detect, particularly in the high sea states that are normally present in the Mariana Islands (Fulling et al., 2011). Aerial surveys in August 2007 covering 2,352 km of linear effort encountered three dwarf sperm whales (Mobley, 2007). In total during Navy-funded 2010–2016 small boat surveys in the Mariana Islands, five dwarf sperm whales have been encountered on four occasions in a median depth of approximately 750 m and at a median distance of approximately 3 km from shore (Hill et al., 2017a).

#### **3.4.1.18.3 Population and Abundance**

There are no abundance estimates for dwarf sperm whales in the Study Area.

#### **3.4.1.18.4 Predator-Prey Interactions**

Dwarf sperm whales feed on cephalopods and, less often, on deep sea fishes and shrimps (Caldwell & Caldwell, 1989; Sekiguchi et al., 1992). Dwarf sperm whales are believed to generally forage near the seafloor (McAlpine, 2009).

Killer whales are predators of dwarf sperm whales (Dunphy-Daly et al., 2008).

#### **3.4.1.18.5 Species-Specific Threats**

Based on data collected in Hawaiian waters, dwarf sperm whales are susceptible to injury or mortality from fisheries interactions (Bradford & Forney, 2014, 2017). It is assumed that fishery activities in the Study Area pose a similar threat to the species.

#### **3.4.1.19 False Killer Whale (*Pseudorca crassidens*)**

##### **3.4.1.19.1 Status and Management**

The population of false killer whales in the Mariana Islands is not listed under the ESA. The stock structure for false killer whales remains uncertain in the western Pacific (Chivers et al., 2007; Martien et al., 2014a), and false killer whales in the Study Area have not been assigned to a stock in the current SAR for the Pacific (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes multiple stocks of false killer whale in the Pacific within the U.S. Exclusive Economic Zone in Hawaiian waters, at Palmyra Atoll, and waters around American Samoa (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.19.2 Geographic Range and Distribution**

The false killer whale is an oceanic species, occurring in deep waters of the North Pacific (Miyashita et al., 1996; Wang et al., 2001) but also known to occur close to shore near oceanic islands (Baird, 2012). In Hawaii, false killer whales have been seen in groups of up to 100 over a wide range of depths and distance from shore (Baird et al., 2003; Baird et al., 2013a; Bradford et al., 2014; Bradford et al., 2015; Oleson et al., 2013). False killer whales are not considered a migratory species, although seasonal shifts in density likely occur. Seasonal movements in the western North Pacific may be related to prey distribution (Odell & McClune, 1999). Satellite-tracked individuals around the Hawaiian islands indicate that false killer whales can move extensively among different islands and also sometimes move from an island coast to as far as 60 mi. (96.6 km) offshore (Baird, 2009b).

During the 2007 survey of the Study Area, there were 10 false killer whale sightings in deep water offshore locations with group sizes ranging from 2 to 26 individuals (Fulling et al., 2011). During the 2010 NMFS survey, one sighting of a pod containing five false killer whales was made approximately midway between Guam and Hawaii in the Transit Corridor (Oleson & Hill, 2010a). During the NMFS 2015 survey of the Mariana Islands, false killer whales were encountered on only two occasions, once off Asuncion

Island and once off Alamagan Island, with estimated group sizes of 6–31 individuals (Hill et al., 2018b; Oleson, 2017). In small boat surveys in the Study Area conducted between 2010 and 2018, false killer whales were encountered only on six occasions (Hill et al., 2014; Hill et al., 2017a; Hill et al., 2018b). Three reported false killer whale strandings have been reported between 1963 and 2013; these occurred in 2000, 2003, and 2007 (Trianni & Tenorio, 2012; Uyeyama, 2014).

#### **3.4.1.19.3 Population and Abundance**

There are estimated to be about 6,000 false killer whales in the North Pacific (starting approximately 50 NM off the Study Area from 25° N to 39° N latitude) based on fishery interaction data (Miyashita, 1993b). Based on sighting data from the 2007 survey, there were an estimated 637 (CV = 0.74) false killer whales in the Study Area (Fulling et al., 2011).

#### **3.4.1.19.4 Predator-Prey Interactions**

False killer whales feed primarily on deep-sea cephalopods and fish (Baird et al., 2008; Koen-Alonso et al., 1999; Odell & McClune, 1999). Four false killer whales found stranded in Hawaii from 2010 through 2016 had stomach contents that included prey items from various squid, yellowfin tuna, mahi mahi, jack, marlin, and bonefish (West, 2016).

False killer whales have been observed to attack other cetaceans, including dolphins and large whales, such as humpback and sperm whales (Baird, 2009a, 2009b). They are known to behave aggressively toward small cetaceans in tuna purse seine nets. Unlike other whales or dolphins, false killer whales frequently pass prey back and forth among individuals before they start to eat the fish, in what appears to be a way of affirming social bonds (Baird et al., 2010b). This species is believed to be preyed on by large sharks and killer whales (Baird, 2009a).

#### **3.4.1.19.5 Species-Specific Threats**

Based on a historic decline in the number of false killer whales in Hawaii, which is believed to have been the result of various factors that include incidental take by commercial fisheries (Bradford et al., 2014; Bradford & Forney, 2017; Oleson et al., 2010; Reeves et al., 2009), it should be assumed that foreign and the limited domestic commercial longline fishing in the Study Area may also pose a similar threat to false killer whales in the Mariana Islands. (Allen & Amesbury, 2012; National Marine Fisheries Service, 2015d, 2018a; National Oceanic and Atmospheric Administration, 2018b). Necropsy results from four stranded false killer whales in Hawaii documented stomach contents that included fishing gear (hooks, leaders, and line) in two of the four animals (West, 2016). As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least 19 products contained false killer whale meat (Baker et al., 2006a). In the two-year period between 2013 and 2014, the total bycatch by South Korean fisheries in the East Sea totaled one false killer whale (Song, 2017).

#### **3.4.1.20 Fraser's Dolphin (*Lagenodelphis hosei*)**

##### **3.4.1.20.1 Status and Management**

Fraser's dolphin is not listed under the ESA. The stock structure for Fraser's dolphin remains uncertain in the western Pacific, and Fraser's dolphin in the Study Area have not been assigned to a stock in the

current SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes a single stock of Fraser's dolphin in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

#### **3.4.1.20.2 Geographic Range and Distribution**

Fraser's dolphin is a tropical oceanic species, except where deep water approaches the coast (Dolar, 2009). This species has been found off the Pacific coast of Japan (Amano et al., 1996). Fraser's dolphin does not appear to be a migratory species (Jefferson & Leatherwood, 1994). In Hawaiian waters, Fraser's dolphin was one of the most abundant species offshore, having large pod group sizes with an observed mean of 283 animals (Bradford et al., 2017).

#### **3.4.1.20.3 Population and Abundance**

There are no abundance estimates for Fraser's dolphin in the Study Area. Genetic samples from 21 bottlenose dolphins encountered off Guam and Saipan in 2007 suggests a history of hybridization with Fraser's dolphin (Martien et al., 2014b).

#### **3.4.1.20.4 Predator-Prey Interactions**

Fraser's dolphin feeds on mid-water fish, squid, and shrimp (Jefferson & Leatherwood, 1994; Mignucci-Giannoni et al., 1999; Perrin et al., 1994a; Watkins et al., 1994).

#### **3.4.1.20.5 Species-Specific Threats**

There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in other locations. There is a report of a Fraser's dolphin being taken as a result of a fishery interaction in the Philippines (Obusan et al., 2016). Fraser's dolphin has been subjected to predation by killer whales in the Bahamas (Dunn et al., 2007).

### **3.4.1.21 Ginkgo-Toothed Beaked Whale (*Mesoplodon ginkgodens*)**

#### **3.4.1.21.1 Status and Management**

Ginkgo-toothed beaked whale is not listed under the ESA. Due to the difficulty in distinguishing the different *Mesoplodon* beaked whale species during visual surveys, ginkgo-toothed beaked whales are combined with all other *Mesoplodon* species that occur off the U.S. West Coast and are managed by NMFS as a species guild (Carretta et al., 2017c; Carretta et al., 2017d). The stock structure for ginkgo-toothed beaked whale remains uncertain in the western Pacific, and ginkgo-toothed beaked whales present in the Study Area or the remainder of the Pacific have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d).

#### **3.4.1.21.2 Geographic Range and Distribution**

Worldwide, beaked whales normally inhabit continental slope and deep ocean waters (greater than 200 m) and are only occasionally reported in waters over the continental shelf (Cañadas et al., 2002; Ferguson et al., 2006a; MacLeod & D'Amico, 2006; Pitman, 2009). Acoustic monitoring at sites around the North Pacific have encountered the "BWC type" beaked whale vocalizations, which are assumed to be produced by ginkgo-toothed beaked whales (Baumann-Pickering et al., 2012; Oleson et al., 2015). Strandings of ginkgo-toothed beaked whales are not common anywhere, but the largest number of records are from Japan (Baumann-Pickering et al., 2012); there have been no known strandings of the species in the Mariana Islands.

In total during Navy-funded 2010–2018 small boat surveys in the Mariana Islands, five *Mesoplodon* beaked whales have been encountered on two occasions in a median depth of approximately 1,140 m and median approximate distance from shore of 15 km (Hill et al., 2017a; Hill et al., 2018b); it could not be determined if these were ginkgo-toothed beaked whales, which are believed to be present in the Study Area; or Blainville's beaked whales, which have been observed elsewhere in the Mariana Islands (Hill et al., 2018b; Oleson, 2017).

A year of acoustic monitoring at Saipan and at Tinian recorded the BWC type beaked whale vocalizations assumed to be produced by ginkgo-toothed beaked whales (Oleson et al., 2015). These vocalizations were detected in all months having sufficient samples to detect their presence in the Study Area, suggesting there is no seasonal aspect to their distribution. This correlates with the findings reported from a previous acoustic monitoring site off Saipan where this same signal type was encountered during 24 percent of days sampled (Baumann-Pickering et al., 2012).

#### **3.4.1.21.3 Population and Abundance**

There are no abundance estimates for ginkgo-toothed beaked whale in the Study Area.

#### **3.4.1.21.4 Predator-Prey Interactions**

All beaked whales probably feed at or close to the bottom in deep oceanic waters, taking suitable prey opportunistically or as locally abundant, typically by suction feeding (Heyning & Mead, 1996; Jefferson et al., 2015; Werth, 2006a, 2006b). Feeding may also occur at mid-water as shown from tagging data from Blainville's beaked whale habits documented in Hawaii (Baird et al., 2005; Baird et al., 2006c). *Mesoplodon* beaked whales have been observed being actively preyed upon by killer whales (Wellard et al., 2016).

#### **3.4.1.21.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of the Philippines, Japan, and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009; Obusan et al., 2016). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated at least one product contained Blainville's beaked whale meat (Baker et al., 2006a), suggesting the same risk may be present for ginkgo-toothed beaked whales. There were five observed interactions between an unidentified beaked whale, unidentified Mesoplodont beaked whale, or Blainville's beaked whale and longline fishing activities in Hawaiian waters between 2010 and 2014 (Bradford & Forney, 2016, 2017). There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in Hawaii.

#### **3.4.1.22 Killer Whale (*Orcinus orca*)**

##### **3.4.1.22.1 Status and Management**

The stock structure for killer whales remains uncertain in the western Pacific, and killer whales present in the Study Area have not been assigned to a stock in the current SARs (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b). NMFS recognizes eight stocks of killer whales for the Pacific, but none of the identified ranges are within the Study Area (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b).

Under the ESA, the Southern Resident Distinct Population Segment of killer whales is the only species listed as endangered, but those animals do not venture beyond the North American nearshore waters. Killer whales in the Study Area are not listed pursuant to the ESA.

#### **3.4.1.22.2 Geographic Range and Distribution**

Killer whales are found in all marine habitats from inland and nearshore coastal areas, to the deep mid-ocean, and from equatorial regions to the polar pack ice zones of both hemispheres. Forney and Wade (2006) found that killer whale densities increased by one to two orders of magnitude from the tropics to the poles.

There are accounts of killer whales off the coast of Japan (Kasuya, 1971). Japanese whaling and whaling sighting vessels indicate that concentrations of killer whales occurred north of the Northern Mariana Islands (Miyashita et al., 1995), and the species has been reported in the tropical waters around Guam, Yap, and Palau (Rock, 1993). Between 1987 and 2017 in the Mariana Islands, killer whales in pods of three to five individuals were observed on only six occasions (Eldredge, 1991; Uyeyama, 2014). There was also a badly decomposed killer whale found stranded on Guam in August 1981 (Kami, 1982). There were no sightings of the species during a 2007 systematic line-transect survey (Fulling et al., 2011) or a 2010 survey of the area (Oleson & Hill, 2010a). In May 2010, a group of approximately five killer whales, including one calf, were observed about 20 NM south of Farallon de Medinilla (Uyeyama, 2014; Wenninger, 2010). The Navy-funded small boat surveys between 2010 and 2016 in the Mariana Islands did not encounter any killer whales (Hill et al., 2014; Hill et al., 2017a). Vocalizations from killer whales were detected on three occasions south of Guam by passive acoustic recorders aboard an underwater glider survey in 2014 (Klinck et al., 2016b).

#### **3.4.1.22.3 Population and Abundance**

There are no abundance estimates for killer whales in the Study Area.

#### **3.4.1.22.4 Predator-Prey Interactions**

Killer whales feed on a variety of prey, including bony fishes, elasmobranchs (a class of fish composed of sharks, skates, and rays), cephalopods, seabirds, sea turtles, and other marine mammals (Fertl et al., 1996; Ford et al., 2013; Ford et al., 2014; Jefferson et al., 2015). In May 2010 during the routine Navy aerial survey of Farallon de Medinilla about 20 mi. (32 km) south of the island, a group of approximately five killer whales, including one calf, were observed feeding on a large whale carcass (Uyeyama, 2014; Wenninger, 2010). The killer whale has no known natural predators; it is considered to be the top predator of the oceans (Ford, 2008).

#### **3.4.1.22.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least two products contained killer whale meat (Baker et al., 2006a).

### 3.4.1.23 Longman's Beaked Whale (*Indopacetus pacificus*)

#### 3.4.1.23.1 Status and Management

Longman's beaked whale is not listed under the ESA. Only one stock has been identified for the Pacific for the population present in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017b). The stock structure for Longman's beaked whale remains uncertain in the western Pacific, and the species in the Study Area has not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d).

#### 3.4.1.23.2 Geographic Range and Distribution

Longman's beaked whales are found in warm tropical waters, and most sightings occur in waters with sea surface temperatures warmer than 78°F (26°C) (Anderson et al., 2006; MacLeod et al., 2006; MacLeod & D'Amico, 2006). Based on systematic survey data collected from 1986 to 2005 in the eastern Pacific, all Longman's beaked whale sightings were south of 25° N (Hamilton et al., 2009). Sighting records of this species in the Indian Ocean showed that Longman's beaked whales are typically found in waters over deep bathymetric slopes reaching 200–2,000 m or greater (Anderson et al., 2006).

Although the full extent of this species' distribution is not fully understood, there have been many recorded sightings at various locations in tropical waters of the Pacific and Indian Oceans (Afsal et al., 2009; Dalebout et al., 2002b; Dalebout et al., 2003; Moore, 1972). In the Pacific, records of this species indicate presence in the eastern, central, and western Pacific, including waters off the coast of Mexico and Hawaii. Longman's beaked whales have not been observed or detected acoustically in the Study Area, although it is assumed they are present in the area. In Hawaii, there was a single sighting of approximately 18 Longman's beaked whales during a NMFS 2002 survey (Barlow, 2006). During the follow-on 2010 survey, there were three sightings of Longman's beaked whales, with group sizes ranging from approximately 32 to 99 individuals (Bradford et al., 2017). It is assumed that Longman's beaked whales would have similar grouping behavior in the Study Area.

#### 3.4.1.23.3 Population and Abundance

There are no abundance estimates for Longman's beaked whales in the Study Area.

#### 3.4.1.23.4 Predator-Prey Interactions

All beaked whales probably feed at or close to the bottom in deep oceanic waters, taking suitable prey opportunistically or as locally abundant, typically by suction feeding (Heyning & Mead, 1996; Jefferson et al., 2015; Werth, 2006a, 2006b). Feeding may also occur at mid-water as shown by tagging data from Cuvier's and from Blainville's beaked whales in Hawaii (Baird et al., 2005; Baird et al., 2006c). *Mesoplodon* beaked whales have been observed being actively preyed upon by killer whales (Wellard et al., 2016); it is assumed this may also be the case with Longman's beaked whales.

#### 3.4.1.23.5 Species-Specific Threats

Disease may be relatively common in Longman's beaked whales since morbillivirus was documented in a juvenile male Longman's beaked whale that stranded in Hawaii in 2010 (West et al., 2012) and in five individuals stranded in New Caledonia (Garrigue et al., 2016), and also given the small sample sizes involved in those findings.

There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in Hawaii. There were two observed interactions between unidentified beaked whales and longline fishing activities in

Hawaiian waters between 2009 and 2013 (Bradford & Forney, 2016), so it is assumed that interactions with fishing activities in the Mariana Islands may also occur. As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence from three different species of beaked whale were identified (Baker et al., 2006a).

#### **3.4.1.24 Melon-Headed Whale (*Peponocephala electra*)**

##### **3.4.1.24.1 Status and Management**

Melon-headed whale is not listed under the ESA. The stock structure for melon-headed whales remains uncertain in the western Pacific, and melon-headed whales in the Study Area have not been assigned to a stock in the current Pacific SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes two stocks of melon-headed whales in the Pacific associated with Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.24.2 Geographic Range and Distribution**

Melon-headed whales are found worldwide in tropical and subtropical waters, but movement patterns for this species are poorly understood. It has been suggested that melon-headed whales near oceanic islands rest near shore during the day and feed in deeper waters at night (Brownell et al., 2009a; Gannier, 2002; Woodworth et al., 2012). In surveys around the main Hawaiian Islands, melon-headed whales showed no clear pattern in depth use (Baird, 2013). Melon-headed whales are also known to enter shallow water areas on occasion, although these are generally characterized as animals being “out of habitat” or “mass strandings.” Such out-of-habitat events, each involving a few hundred melon-headed whales, have occurred at Sasanhaya Bay, Rota (Jefferson et al., 2006); and in Hawaii (Fromm et al., 2006; Mobley et al., 2007; Southall et al., 2006) on the same day in 2004. Similar numbers did so twice in the Philippines entering Manila Bay in February 2009 and the bay at Odiongan, Romblon in March of 2009 (Aragones et al., 2010; Obusan et al., 2016).

There were two sightings of melon-headed whales during the 2007 survey of the Study Area, with group sizes of 80–109 individuals (Fulling et al., 2011). There was one sighting of approximately 53 individuals southeast of Guam and two mid-ocean sightings (pods sizes of 43 and 72) in the Transit Corridor portion of the Study Area during the large vessel Pacific Islands Fisheries Science Center survey (Oleson & Hill, 2010a). During small boat surveys in 2012 and 2014, melon-headed whales in large pods numbering between 85 and 325 individuals were sighted off Guam and Tinian/Saipan (HDR, 2012; Hill et al., 2014). The NMFS 2015 month-long survey of the Mariana Islands encountered melon-headed whales on four occasions, in offshore waters and in large pods estimated to number between 90 and 268 individuals (Hill et al., 2018b; Oleson, 2017).

There was a live stranding of a melon-headed whale on the beach at Inarajan Bay, Guam in April 1980 (Donaldson, 1983; Kami, 1982), and four individuals at Orote in 2009 (Uyeyama, 2014).

##### **3.4.1.24.3 Population and Abundance**

Based on sighting data from a systematic survey in 2007, there were an estimated 2,455 (CV = 0.70) melon-headed whales in the Study Area (Fulling et al., 2011).

#### **3.4.1.24.4 Predator-Prey Interactions**

Melon-headed whales prey on squid, pelagic fishes, and occasionally crustaceans. Most of the fish and squid families eaten by this species consist of mid-water forms found in waters up to 1,500 m deep, suggesting that feeding takes place deep in the water column (Baird et al., 2010a; Jefferson & Barros, 1997).

#### **3.4.1.24.5 Species-Specific Threats**

Melon-headed whales are believed to be preyed on by killer whales and have been observed fleeing from killer whales in Hawaiian waters (Baird et al., 2006a). The 2016 Pacific SAR (Carretta et al., 2017c; Carretta et al., 2017d) suggests that melon-headed whales may be particularly sensitive to impacts from anthropogenic sounds; see the U.S. Navy's Technical Report (U.S. Department of the Navy, 2017c) for a general discussion of strandings potentially related to the use of sonar and other anthropogenic sound.

#### **3.4.1.25 Pantropical Spotted Dolphin (*Stenella attenuata*)**

##### **3.4.1.25.1 Status and Management**

The pantropical spotted dolphin is not listed under the ESA. The stock structure for pantropical spotted dolphin remains uncertain in the western Pacific, and pantropical spotted dolphins in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes a single pelagic stock and three Hawaiian Island-associated stocks of pantropical spotted dolphin in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d). Results from genetic analyses of pantropical spotted dolphin populations, including the Indo-Pacific and eastern tropical Pacific Ocean (including eight samples from Guam and the Northern Mariana Islands), support the current taxonomy and indicate very close genetic relationships among the Indo-Pacific populations (Leslie & Morin, 2018).

##### **3.4.1.25.2 Geographic Range and Distribution**

A survey of the Mariana Islands in 2007 encountered 17 groups of pantropical spotted dolphins, ranging in size from 1 to 115 individuals (Fulling et al., 2011). Aerial surveys in August 2007 covering 2,352 km of linear effort encountered a single pod of 30 pantropical spotted dolphins (Mobley, 2007). In total during the Navy-funded 2010 to 2016 small boat surveys in the Mariana Islands, pantropical spotted dolphins were encountered on 30 occasions in group sizes of 4–70 individuals at a median approximate distance from shore of 6 km (Hill et al., 2014; Hill et al., 2017a). Approximate satellite tag locations from a pantropical spotted dolphin in 2016 demonstrated wide-ranging use of the waters at a median of 6.1 km offshore of Guam (Hill et al., 2017a).

##### **3.4.1.25.3 Population and Abundance**

Based on sighting data from the 2007 systematic survey of the Mariana Islands, the estimated abundance for pantropical spotted dolphins in the Study Area is 12,981 (CV = 0.704) (Fulling et al., 2011).

##### **3.4.1.25.4 Predator-Prey Interactions**

Pantropical spotted dolphins prey on near-surface fish, squid, and crustaceans and on some mid-water species (Perrin & Hohn, 1994). Results from various tracking and feeding studies suggest that pantropical spotted dolphins in the eastern tropical Pacific and off Hawaii feed primarily at night on surface and mid-water species that rise with the deep scattering layer toward the water's surface after dark (Baird et al., 2001; Silva et al., 2016).

#### 3.4.1.25.5 Species-Specific Threats

There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in other locations. Pantropical spotted dolphins in Hawaii and Samoa have been observed interacting with the longline fishery, resulting in injury (Bradford & Forney, 2014), and there was one case of serious injury to a spotted dolphin observed entangled in fishing line (Bradford & Lyman, 2015). Given the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

Pantropical spotted dolphins may be preyed on by killer whales and sharks and have been observed fleeing killer whales in Hawaiian waters (Baird et al., 2006b). Other predators may include the pygmy killer whale, false killer whale, and occasionally the short-finned pilot whale (Perrin, 2009b).

#### 3.4.1.26 Pygmy Killer Whale (*Feresa attenuata*)

##### 3.4.1.26.1 Status and Management

The pygmy killer whale is not listed under the ESA. The stock structure for pygmy killer whale remains uncertain in the western Pacific, and pygmy killer whales in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d).

##### 3.4.1.26.2 Geographic Range and Distribution

This species has been documented in the western Pacific (Taiwan and Japan) (Sylvestre, 1988; Wang et al., 2001; Wang & Yang, 2006). There was only one pygmy killer whale sighting of a group of six animals during the 2007 systematic survey of the Study Area (Fulling et al., 2011). The sighting was made near the Mariana Trench, south of Guam, where the bottom depth was 14,564 ft. (4,413 m). This is consistent with the known habitat preference of this species for deep, oceanic waters. During small boat surveys between 2010 and 2016, there was a single pygmy killer whale sighting northeast of Saipan in 2011 and then single sightings in 2013 and 2014 off Guam; group sizes were from six to nine individuals (Hill et al., 2014; Hill et al., 2017a).

##### 3.4.1.26.3 Population and Abundance

Based on a single sighting during the 2007 survey of the Study Area, pygmy killer whale abundance was estimated at 78 individuals (CV = 0.881) (Fulling et al., 2011).

##### 3.4.1.26.4 Predator-Prey Interactions

Pygmy killer whales feed predominantly on fish and squid. They have been known to attack other dolphin species, apparently as prey, although this is not common (Jefferson et al., 2015; Perryman & Foster, 1980; Ross & Leatherwood, 1994).

##### 3.4.1.26.5 Species-Specific Threats

Fisheries interactions in the Study Area are likely given documented evidence from fishery activities in Hawaii (Bradford & Forney, 2017; Carretta et al., 2017c). Based on the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

The 2016 Pacific SAR (Carretta et al., 2017c; Carretta et al., 2017d) suggests that two mass strandings of pygmy killer whales (that occurred in 2004 and 2005) on Taiwan were, "...possibly associated with offshore naval training exercises" based on the citation to Wang and Yang (2006). Wang and Yang (2006) only speculatively suggested that, "...naval sonar and live ammunition exercises are two of many plausible causes that need to be investigated" given there was a lack of necessary information (such as if sonar was even in use) regarding relatively contemporaneous and distant events involving the U.S. Navy, People's Republic of China Navy, Taiwan's Republic of China Navy, Japanese Navy, and oil and gas seismic exploration occurring in the eastern Pacific. Further, between 1995 and 2005 there were a total of six pygmy killer whale Mass Stranding Events and three milling events involving the same species in Taiwan (Brownell et al., 2009b; Yang et al., 2008), confounding the identification of a specific cause for these particular stranding events. The suggestion that sonar, underwater detonations, or seismic oil and gas exploration may have caused the 2004 and 2005 strandings has remained speculative with researchers pointing to the need for further investigation (Brownell et al., 2009b; Wang & Yang, 2006; Yang et al., 2008). The technical report from the U.S. Department of the Navy (U.S. Department of the Navy, 2017c) provides a general discussion of strandings potentially related to the use of sonar and other anthropogenic sound.

The pygmy killer whale has no documented predators (Weller, 2009), although it may be subject to predation by killer whales.

#### **3.4.1.27 Pygmy Sperm Whale (*Kogia breviceps*)**

##### **3.4.1.27.1 Status and Management**

Pygmy sperm whale is not listed under the ESA. The stock structure for pygmy sperm whales remains uncertain in the western Pacific, and pygmy sperm whales in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). Other than for waters along the U.S. West Coast, NMFS recognizes a single stock of pygmy sperm whale in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.27.2 Geographic Range and Distribution**

During marine mammal monitoring for Valiant Shield 07, a group of three *Kogia* (dwarf or pygmy sperm whales) was observed about 8 NM east of Guam (Mobley, 2007). The stranding of a pygmy sperm whale in 1997 (Trianni & Tenorio, 2012), is the only other confirmed occurrence of this species in the Study Area.

##### **3.4.1.27.3 Population and Abundance**

There are no abundance estimates for pygmy sperm whale in the Study Area.

##### **3.4.1.27.4 Predator-Prey Interactions**

Pygmy sperm whales feed on cephalopods and, less often, on deep sea fishes and shrimps (Caldwell & Caldwell, 1989; Sekiguchi et al., 1992; West et al., 2009). Pygmy sperm whales are believed to generally forage near the seafloor (McAlpine, 2009).

Killer whales and white sharks are documented predators of pygmy sperm whales (Dunphy-Daly et al., 2008; Long, 1991; Tirard et al., 2010).

##### **3.4.1.27.5 Species-Specific Threats**

Based on data collected in Hawaiian waters, pygmy sperm whales are susceptible to injury or mortality from fisheries interactions (Bradford & Forney, 2014, 2017). It is assumed the fishery activities in the

Study Area pose a similar threat. Given the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

#### **3.4.1.28 Risso's Dolphin (*Grampus griseus*)**

##### **3.4.1.28.1 Status and Management**

Risso's dolphin is not listed under the ESA. The stock structure for Risso's dolphin remains uncertain in the western Pacific, and Risso's dolphins in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). Other than for waters along the U.S. West Coast, NMFS recognizes a single stock of Risso's dolphins in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### **3.4.1.28.2 Geographic Range and Distribution**

Occurrence of this species is deep open ocean waters off Hawaii and in other locations in the Pacific (Au & Perryman, 1985; Bradford et al., 2017; Leatherwood et al., 1980; Miyashita et al., 1996; Wang et al., 2001). Fishery interaction data determined the species occurrence west of the International Date Line extended as far north as 40° N, but the southern extent of the range could not be determined (Miyashita, 1993a). Aerial surveys in August 2007 covering 2,352 km of linear effort encountered a single pod of eight Risso's dolphins (Mobley, 2007). During the NMFS survey of 2010, there was a single Risso's dolphin sighting of three individuals approximately 60 NM north of FDM (Oleson & Hill, 2010a). The 2015 NMFS month-long survey of the Mariana Islands encountered Risso's dolphins only twice and in small pods with a median group size of three (Hill et al., 2018b; Oleson, 2017). The species has not been detected in any other surveys efforts in the Study Area (Fulling et al., 2011; Hill et al., 2014; Hill et al., 2017a; Hill et al., 2018a; Hill et al., 2018b). Vocalizations from Risso's dolphins were also detected south of Guam by passive acoustic recorders aboard an underwater glider survey in 2014 (Klinck et al., 2016b).

##### **3.4.1.28.3 Population and Abundance**

There are no abundance estimates for Risso's dolphin in the Study Area.

##### **3.4.1.28.4 Predator-Prey Interactions**

Cephalopods and crustaceans are the primary prey for Risso's dolphins (Clarke, 1996), which feed mainly at night (Fernandez et al., 2017; Jefferson et al., 2015; Perrin et al., 2009). This dolphin may be preyed on by both killer whales and sharks, although there are no documented reports of predation by either species (Weller, 2009).

##### **3.4.1.28.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least one product contained Risso's dolphin meat (Baker et al., 2006a).

### **3.4.1.29 Rough-Toothed Dolphin (*Steno bredanensis*)**

#### **3.4.1.29.1 Status and Management**

The rough-toothed dolphin is not listed under the ESA. The stock structure for rough-toothed dolphins remains uncertain in the western Pacific, and rough-toothed dolphins in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes a single stock of rough-toothed dolphins in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

#### **3.4.1.29.2 Geographic Range and Distribution**

Rough-toothed dolphins were sighted twice during a 2007 survey; once as nine individuals in a mixed group of short-finned pilot whales and bottlenose dolphins, and once in a pod of nine individuals with calves present (Fulling et al., 2011). A pod of eight rough-toothed dolphins was also sighted approximately 175 km south of Guam during a 2007 aerial survey (Mobley, 2007). There were no rough-toothed dolphins identified in the broad offshore survey in 2010 (Oleson & Hill, 2010a). The species was encountered only three times during the month-long 2015 NMFS survey of the islands, twice in a group with another cetacean species (Oleson, 2017). Annual small boat surveys conducted from 2010 to 2018 (Hill et al., 2011; Hill et al., 2013a) have encountered rough-toothed dolphins on seven occasions, and again all but one of those encounters were in a group with other cetaceans (Hill et al., 2014; Hill et al., 2018b). Four of the same photo-identified rough-toothed dolphins encountered in 2013 have been seen multiple times since in the same general location to the west of Saipan off CK Reef (Hill et al., 2014; Hill et al., 2017a). One group of rough-toothed dolphins was sighted in 2014, but none were encountered in 2015, 2016, or 2017 (Hill et al., 2017a; Hill et al., 2018a).

#### **3.4.1.29.3 Population and Abundance**

During the 2007 systematic line-transect survey of the Study Area, there was only one on-effort sighting of rough-toothed dolphin that was used to derive an abundance estimate of 166 animals (CV = 0.892) (Fulling et al., 2011). Given the very limited sample size (a single sighting), this estimate is considered highly uncertain. In July 2004, there was a sighting of an undetermined smaller number of rough-toothed dolphins mixed in with a school of an estimated 500–700 melon-headed whales off Rota in Sasanhayan Bay (Jefferson et al., 2006).

#### **3.4.1.29.4 Predator-Prey Interactions**

Rough-toothed dolphin prey includes fish and cephalopods. They are known to feed on large fish species, such as mahi mahi (Miyazaki & Perrin, 1994; Pitman & Stinchcomb, 2002), and have been observed feeding during the day on near-surface fishes, including flying fishes (Gannier & West, 2005). They may also prey on reef fish, as Perkins and Miller (1983) noted that parts of reef fish had been found in the stomachs of stranded rough-toothed dolphins in Hawaii, although the stomach contents of a stranded animal may not be representative of the species.

#### **3.4.1.29.5 Species-Specific Threats**

There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in Hawaii. In Hawaii from 2010 to 2014, two rough-toothed dolphins were observed injured during deep-set and shallow-set fisheries in the Exclusive Economic Zone (Bradford & Forney, 2017). Given the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive

created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

Although this species has not been documented as prey by other species, it may be subject to predation from killer whales.

### **3.4.1.30 Short-Finned Pilot Whale (*Globicephala macrorhynchus*)**

#### **3.4.1.30.1 Status and Management**

Short-finned pilot whale is not listed under the ESA. The stock structure for short-finned pilot whales remains uncertain in the western Pacific, and short-finned pilot whales in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). With the exception of the U.S. West Coast, NMFS recognizes a single stock of short-finned pilot whales in the Pacific in Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

#### **3.4.1.30.2 Geographic Range and Distribution**

In the 2007 survey of the Mariana Islands, short-finned pilot whales were encountered five times in groups ranging in size from 5 to 43 animals (Fulling et al., 2011). During the 2010 NMFS survey there was a single sighting of 23 short-finned pilot whales in the northern portion of the Study Area (Oleson & Hill, 2010a). Closer to the islands, there have been numerous incidental sightings of short-finned pilot whales occurring between 1977 and 2013 (Uyeyama, 2014). During the Navy-funded 2010–2017 small boat surveys in the Mariana Islands, short-finned pilot whale groups were encountered on 16 occasions in a median depth of approximately 700 m and median approximate distance from shore of 5 km, including one pod of 35 individuals off Marpi Reef north of Saipan (Hill et al., 2014; Hill et al., 2017a; Hill et al., 2018a). Satellite tag locations from one short-finned pilot whale in 2016 appeared to indicate a position inside the mouth of Apra Harbor (there were no prior or subsequent positions on that day) (Hill et al., 2017a). However, it should be considered uncertain if the animal was in Apra Harbor, due to the limited precision (error range) of even high-quality Argos satellite fixes, and in particular with regard to reduced longitudinal precision, given the Argos satellites are in polar orbits (Boyd & Brightsmith, 2013; Vincent et al., 2002). Based on the locations from the 2013 to 2016 satellite tagged individuals in May–August timeframe, the combined data has suggested that the northwest side of Guam is a frequently used area for pilot whales during that time of the year (Hill et al., 2017a).

#### **3.4.1.30.3 Population and Abundance**

The estimated abundance for short-finned pilot whales in the Study Area is 909 (CV = 0.677), based on sighting data from the 2007 systematic survey of the Mariana Islands (Fulling et al., 2011). Genetic samples taken during small boat surveys between 2010 and 2014 found evidence of genetic differentiation for short-finned pilot whales between the Mariana Islands, although they possess haplotypes also common in the South Pacific, North Atlantic, Indian Ocean, and off of southern Japan (Martien et al., 2014b).

#### **3.4.1.30.4 Predator-Prey Interactions**

Pilot whales feed primarily on squid but also take fish (Bernard & Reilly, 1999). They are generally well adapted to feeding on squid (Jefferson et al., 2015; Werth, 2006a, 2006b). Analysis of satellite tagging data from pilot whales in Hawaii correlated with certain environmental parameters, suggesting that the deep mesopelagic boundary community serves as prey for these whales (Abecassis et al., 2015).

Pilot whales are not generally known to prey on other marine mammals, but records from the eastern tropical Pacific suggest that the short-finned pilot whale does occasionally chase and attack, and may eat, dolphins during fishery operations (Olson, 2009; Perryman & Foster, 1980).

#### **3.4.1.30.5 Species-Specific Threats**

As discussed in Section 3.4.1.7.3 (Bycatch), entanglement risk includes fishing activities out of Japan and South Korea, with the threat of mortality from any such interaction being high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009). In nine market samples from South Korea between 2003 and 2005, molecular (DNA) evidence indicated that at least two products contained short-finned pilot whale meat (Baker et al., 2006a).

This species is not known to have any predators (Weller, 2009), although it may be subject to predation by killer whales.

#### **3.4.1.31 Sperm Whale (*Physeter macrocephalus*)**

##### **3.4.1.31.1 Status and Management**

The sperm whale is listed as endangered under the ESA, but there is no designated critical habitat for this species. The stock structure for sperm whales remains uncertain in the Pacific (Mesnick et al., 2011; Mizroch & Rice, 2013; National Marine Fisheries Service, 2015a), and sperm whales in the Study Area have not been assigned to a stock in the current Pacific SAR (Carretta et al., 2017c; Carretta et al., 2017d). Except for waters off the U.S. West Coast, NMFS recognizes two stocks of sperm whales, one in the central Pacific (in Hawaiian waters) and one in the North Pacific (in Alaskan waters) (Carretta et al., 2017c; Carretta et al., 2017d; Muto et al., 2017a; Muto et al., 2017b).

##### **3.4.1.31.2 Geographic Range and Distribution**

Based on whaling data and discovery tag movement data for the North Pacific, it has been argued that the distribution of sperm whales encompasses the entire Pacific Ocean basin, with concentrations in the arctic and subtropical areas (Ilyashenko et al., 2014; Mizroch & Rice, 2013). The Study Area is south of the locations where the majority of sperm whales were encountered during whaling (Mizroch & Rice, 2013; Townsend, 1935), although during a 1972 survey of the Ogasawara and Mariana Island regions two large groups totaling 90 sperm whales were reported (Masaki, 1972). Sperm whales have been routinely sighted in the Study Area and detected in acoustic monitoring records. Acoustic recordings in August 2013 at Pagan Island indicated the presence of sperm whales within 20 NM of the island (Tetra Tech Inc., 2014). Although it has been reported that sperm whales are generally found far offshore in deep water (Mizroch & Rice, 2013), sightings in the Study Area have included animals close to shore in relatively shallow water as well as in areas near steep bathymetric relief (Fulling et al., 2011; Hill et al., 2017a; Uyeyama, 2014). A total of 23 sperm whale sightings and 93 acoustic encounters were made during the 2007 survey in water depths between approximately 400 and 1,000 m depth (Fulling et al., 2011; Yack et al., 2016). There were three encounters with sperm whales during the NMFS 2015 cetacean survey of the Mariana Islands (Hill et al., 2018b; Oleson, 2017). During the Navy-funded 2010–2018 small boat surveys in the Mariana Islands, six sperm whales were encountered on three occasions in a median depth of approximately 1,200 m and median approximate distance from shore of 12 km (Hill et al., 2017a; Hill et al., 2018b). Vocalizations from sperm whales were also detected on 20 occasions to the east and south of Guam by passive acoustic recorders during an underwater glider survey in 2014 (Klinck et al., 2016b).

### 3.4.1.31.3 Population and Abundance

It is assumed the Pacific population is still recovering, given whaling by the Soviet Union from 1948 to 1979 in the North Pacific took 157,680 sperm whales (Ilyashenko et al., 2014). NMFS has reported that for the Pacific Ocean,<sup>4</sup> the population is estimated between 26,300 and 32,100 for the North Pacific and between 14,800 and 34,600 for the eastern tropical Pacific, while the population of the Hawaii Stock is estimated between 2,539 and 3,354 (National Marine Fisheries Service, 2015a). NMFS has not explicitly stated if the western North Pacific and the Mariana Islands are included in the range for the population of sperm whales considered the North Pacific Stock (Muto et al., 2017a; Muto et al., 2017b; National Marine Fisheries Service, 2015a), although that may be the most logical assignment for those animals in the Study Area. The most recent Alaska SAR provides that there is no current abundance data available for sperm whale of the North Pacific Stock (Muto et al., 2017a; Muto et al., 2017b).

During the 2007 systematic line-transect survey of the Mariana Islands, 11 on-effort sperm whale sightings were used to derive an abundance estimate of 705 animals (CV = 0.604) for the Study Area (Fulling et al., 2011). Passive acoustic monitoring was also conducted during the 2007 survey, and 93 acoustic encounters from vocalizing sperm whales were used to develop a habitat-based density model for this species (Yack et al., 2016). The model provided spatially explicit density estimates for the Study Area, and daily model predictions indicated that sperm whale abundance varied temporally over the period of the 2007 survey (January 15 to April 10). Average Study Area abundance derived from the habitat model was similar to the line-transect estimate based on visual sightings; 700 animals (CV = 0.436) based on a model using sounds typically produced by mature males, females, and juveniles (i.e., “regular clicks”), and 637 animals (CV = 0.447) based on a model using both the regular clicks and “slow clicks” that are only produced by mature males (Yack et al., 2016).

### 3.4.1.31.4 Predator-Prey Interactions

Sperm whales are known to occur in groups for both predator defense and foraging purposes. Sperm whales feed on squid, other cephalopods, and bottom-dwelling fish and invertebrates (Davis et al., 2007; Marcoux et al., 2007; Rice, 1989). False killer whales, pilot whales, and killer whales have been documented harassing and on occasion attacking sperm whales (Arnbom et al., 1987; Baird, 2009b; Palacios & Mate, 1996; Pitman et al., 2001).

### 3.4.1.31.5 Species-Specific Threats

Sperm whales are susceptible to injury or mortality from vessel strike (Bradford & Lyman, 2015; Carretta et al., 2016b; Carretta et al., 2017c; Fulling et al., 2017). Sperm whales in the Pacific have been documented as susceptible to entanglement and other interactions with fishing gear (Bradford & Lyman, 2015; Carretta et al., 2016b; Carretta et al., 2017c; Helker et al., 2017). Sperm whales have also been documented as having ingested marine debris, resulting in mortality (Garibaldi & Podesta, 2014; Jacobsen et al., 2010), and as with all most marine mammals, are susceptible to disease (West et al., 2015).

---

<sup>4</sup>The “Pacific Ocean” estimates provided did not address or otherwise specifically include the western Pacific Ocean that would include the Study Area.

### 3.4.1.32 Spinner Dolphin (*Stenella longirostris*)

#### 3.4.1.32.1 Status and Management

The spinner dolphin is not listed under the ESA. The stock structure for spinner dolphins remains uncertain in the western Pacific, and spinner dolphins in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). NMFS recognizes seven stocks of island- or atoll-associated spinner dolphin populations in the Pacific in Hawaii and American Samoa waters (Carretta et al., 2017c; Carretta et al., 2017d), which are all at locations well to the east of the Study Area.

#### 3.4.1.32.2 Geographic Range and Distribution

Spinner dolphins traveling among the Mariana Islands chain are expected to occur throughout the Mariana Islands, having been observed from Uracas in the north to Guam in the south (Fulling et al., 2011; Hill et al., 2017a; Jefferson et al., 2006; Oleson, 2017; Oleson & Hill, 2010b; Tetra Tech Inc., 2014; Trianni & Kessler, 2002; Uyeyama, 2014; Vogt, 2008). Spinner dolphins have been the most frequently encountered species during small boat reconnaissance surveys conducted in the nearshore waters of the Mariana Islands since 2010 but were uncommon offshore (Fulling et al., 2011; HDR, 2011a; HDR EOC, 2012; Hill et al., 2013a; Hill et al., 2014; Hill et al., 2015b; Hill et al., 2016b; Hill et al., 2017a; Hill et al., 2018a; Ligon et al., 2011; Oleson, 2017; Oleson & Hill, 2010b). Previously reported spinner dolphin high-use areas nearshore at Guam include Bile Bay, Tumon Bay, Double Reef, north Agat Bay, and off Merizo (Cocos Lagoon area), where these animals congregate during the day to rest (Amesbury et al., 2001; Eldredge, 1991). More recently, high-use areas have included Agat Bay; the Merizo channel, tucked into the several small remote bays between Merizo and Facpi Point; Piti Bay; Hagatna; Tumon Bay; and Pugua Point (Ligon et al., 2011). There have been no documented sightings within Apra Harbor. The locations where spinner dolphins have been documented resting in Agat Bay have been considered for geographic mitigation, as detailed in Appendix I (Geographic Mitigation Assessment).

During the Navy-funded 2010–2017 small boat surveys in the Mariana Islands, 129 spinner dolphins have been encountered on 15 occasions in a median depth of approximately 20 m and median approximate distance from shore of 1 km (Hill et al., 2017a; Hill et al., 2018a). During a survey in August 2013 at Pagan Island, spinner dolphins calves and juveniles were encountered; although sighting rates were low relative to other island areas, re-sightings of four individual spinner dolphins on subsequent days were suggested to be consistent with residency patterns seen elsewhere (Tetra Tech Inc., 2014), which would be similar to behaviors seen in Hawaii (Heenehan et al., 2017b; Lammers, 2004; Marten & Psarakos, 1999; Norris et al., 1994; Tyne et al., 2015; Tyne et al., 2017).

#### 3.4.1.32.3 Population and Abundance

Spinner dolphins were sighted only once during the 2007 broad area line-transect survey of the Mariana Islands (Fulling et al., 2011). As noted previously, spinner dolphins have been the most commonly encountered species in nearshore waters less than 1 km from shore and have been encountered in group sizes of up to 124 individuals in a pod (HDR, 2011a; HDR EOC, 2012; Hill et al., 2011; Hill et al., 2013a; Hill et al., 2013b; Hill et al., 2014; Hill et al., 2015b; Hill et al., 2016b; Hill et al., 2017a; Hill et al., 2018a; Ligon et al., 2011; Oleson & Hill, 2010b). Genetic samples (n = 93) from spinner dolphins encountered during small boat surveys off Guam and Saipan between 2010 and 2014 suggest the population has high haplotypic diversity similar to that observed in the Society Islands of French Polynesia and that spinner dolphins around the Mariana Islands are much less isolated than those around the Hawaiian Islands (Martien et al., 2014b).

#### 3.4.1.32.4 Predator-Prey Interactions

Spinner dolphins feed primarily on small mid-water fishes, squids, and shrimp, and dive to at least 200–300 m (Benoit-Bird & Au, 2003; Perrin & Gilpatrick, 1994). They forage primarily at night, when the mid-water community migrates toward the surface and the shore (Benoit-Bird et al., 2001; Benoit-Bird, 2004; Benoit-Bird & Au, 2009; Tyne et al., 2017). Spinner dolphins track the horizontal and vertical migrations of their prey (Benoit-Bird & Au, 2003), allowing for foraging efficiencies (Benoit-Bird et al., 2001; Benoit-Bird & Au, 2003, 2004; Benoit-Bird & Au, 2009). Foraging behavior has also been linked to lunar phases in scattering layers off of Hawaii (Benoit-Bird & Au, 2004).

#### 3.4.1.32.5 Species-Specific Threats

There is no information available regarding marine mammal interactions with fishing activities in the Study Area, but the threat is presumed to be similar to what has been documented in Hawaii. In Hawaiian waters from 2008 to 2012 there were three observed serious injuries (leading to death) to spinner dolphins (Bradford & Lyman, 2015). Two of these injuries were fishing related, and one involved marine debris preventing the individual's mouth from opening. Given the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

Spinner dolphins are also at risk if ecotourism and whale-watching activities result in chronic disturbance in their resting habitats (Courbis & Timmel, 2008; Heenehan et al., 2016; Heenehan et al., 2017a; Tyne et al., 2014; Tyne, 2015; Tyne et al., 2015; Tyne et al., 2017; Tyne et al., 2018). Courbis (2008) found changes in spinner dolphin aerial behaviors and suggested it was likely that vessel and swimmer activity was at least synergistically involved in causing these changes, but whether the behavioral changes affected the survival and fitness of spinner dolphins remains unknown.

Spinner dolphins have stranded at Saipan (Trianni & Kessler, 2002). Spinner dolphins may be preyed on by sharks, killer whales, pygmy killer whales, and short-finned pilot whales (Perrin, 2009a).

#### 3.4.1.33 Striped Dolphin (*Stenella coeruleoalba*)

##### 3.4.1.33.1 Status and Management

The striped dolphin is not listed under the ESA. The stock structure for striped dolphins remains uncertain in the western Pacific, and striped dolphins in the Study Area have not been assigned to a stock in the current SAR (Carretta et al., 2017c; Carretta et al., 2017d). Other than along the U.S. West Coast, NMFS recognizes only a single stock of striped dolphins that is present within the 200-mile Exclusive Economic Zone defining Hawaiian waters (Carretta et al., 2017c; Carretta et al., 2017d).

##### 3.4.1.33.2 Geographic Range and Distribution

Striped dolphins are generally restricted to oceanic regions and are seen close to shore only where deep water approaches the coast. In the eastern tropical Pacific, striped dolphins inhabit areas with large seasonal changes in surface temperature and thermocline depth, as well as seasonal upwelling (Au & Perryman, 1985; Reilly, 1990). The observed northern limits for the species are the Sea of Japan off Hokkaido, off Washington State in the eastern Pacific, or roughly along 40° N latitude across the western and central Pacific (Reeves et al., 2002).

Prior to the 2007 survey of the Study Area (Fulling et al., 2011), striped dolphins were only known to occur in the area from two strandings, one recorded in July 1985 (Eldredge, 1991, 2003) and a second in 1993 off Saipan (Trianni & Tenorio, 2012). However, striped dolphins were sighted throughout the Study Area during the 2007 survey (Fulling et al., 2011). There was at least one sighting over the Mariana Trench, southeast of Saipan. Group sizes ranged from 7 to 44 individuals, and several sightings included calves. In early April 2010, during an oceanographic survey of waters in Micronesia and the Commonwealth of the Northern Mariana Islands, there were two striped dolphin sightings (pod sizes of 6 and 12) in waters to the south of Guam (Oleson & Hill, 2010a). Striped dolphins have not been reported during more recent non-systematic surveys in the Study Area involving small boats operating close to shore (Hill et al., 2011; Hill et al., 2013a; Hill et al., 2014; Hill et al., 2015b; Hill et al., 2017a; Hill et al., 2018a).

#### **3.4.1.33.3 Population and Abundance**

The population of striped dolphins south of 30° N in the western Pacific (which would include the Study Area) was estimated to be around 52,600 dolphins (Miyashita, 1993b). Based on the 2007 survey data from the Mariana Islands sightings, there were an estimated 3,531 (CV = 0.54) striped dolphins in the survey area (Fulling et al., 2011).

#### **3.4.1.33.4 Predator-Prey Interactions**

Striped dolphins often feed in open sea or sea bottom zones along the continental slope or just beyond it in oceanic waters. Most of their prey possess light-emitting organs (lanternfishes), suggesting that striped dolphins may be feeding at great depths, possibly diving to 200–700 m, and may feed at night in order to take advantage of the deep scattering layer's diurnal vertical movements, including small mid-water fishes and squids (Archer & Perrin, 1999; Perrin et al., 1994b). This species has been documented to be preyed upon by sharks (Ross & Bass, 1971). It may also be subject to predation by killer whales.

#### **3.4.1.33.5 Species-Specific Threats**

Striped dolphins have been taken as bycatch by the tuna purse seine fishery in the eastern tropical Pacific and are susceptible to entanglement in fishing gear in other areas (Carretta et al., 2017c; Carretta et al., 2017d). There are no specific fisheries interactions or other information on species-specific threats available for this species in the Study Area. Given the information provided in Section 3.4.1.7.3 (Bycatch), entanglement risk may include fishing activities out of Japan and South Korea (Miyashita, 1993b). The threat of mortality from any such interaction is high given the incentive created by the commercial sale of whale meat/products allowed under Japanese and South Korean law (Baker et al., 2006a; Lukoschek et al., 2009).

### **3.4.2 Environmental Consequences**

Under the Proposed Action for this SEIS/OEIS, there have been some modifications to the quantity and type of acoustic stressors under the two action alternatives. Additionally, there is one new sub-stressor (high-energy lasers) being analyzed because of the potential to affect marine species, as detailed in Section 3.0.4.3.2.2 (High-Energy Lasers).

In the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015a), the Navy considered all potential stressors associated with ongoing military readiness in the Mariana Islands and then analyzed their potential impacts on marine mammals in the Study Area. In addition, NMFS also reviewed the Navy's analysis and detailed their findings with regard to requirements under the MMPA (80 FR 46112) and

pursuant to the ESA for the Navy's Proposed Action in the Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015b).

In general, there have been no substantial changes to the activities analyzed as the Proposed Action the 2015 MITT Final EIS/OEIS that would change the conclusions reached regarding populations of marine mammals in the Study Area. Use of acoustic stressors (sonar and other transducers) and use of explosives have occurred in the Mariana Islands for decades and were last authorized by the 2015 completion of the MITT Record of Decision, MMPA Authorization, and ESA Biological Opinion. There have been no known impacts on populations of marine mammals or adverse effects to ESA-listed marine mammal species that were not otherwise previously analyzed or accounted for in the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015b), the NMFS MMPA Authorization (80 FR 46112), or the NMFS Biological Opinion pursuant to ESA (National Oceanic and Atmospheric Administration, 2015a) with regard to acoustic or explosive stressors.

In this SEIS/OEIS, the Navy has reviewed the analysis of impacts from these ongoing activities and additionally analyzed the new or changing training and testing activities as projected into the reasonably foreseeable future. The projected future actions are based on evolving operational requirements, including those associated with any anticipated new platforms or systems not previously analyzed. The Navy has compiled, thoroughly reviewed, and incorporated the best available emergent marine mammal science since 2015 that is relevant to the analysis of environmental impacts from the proposed activities as presented in the 2015 MITT Final EIS/OEIS. Where there has been no substantive or otherwise meaningful change in the action, science, or regulations, the Navy will rely on the 2015 MITT Final EIS/OEIS analysis. Where there has been substantive change in the action, science, or regulations, the information and analysis provided in this SEIS/OEIS will supplement the 2015 MITT Final EIS/OEIS to support environmental compliance with applicable environmental statutes for marine mammals (the MMPA and ESA).

The 2015 MITT Final EIS/OEIS considered all training and testing activities proposed to occur in the Study Area that may have the potential to result in the MMPA-defined take of marine mammals or to affect ESA-listed marine mammal species. The stressors applicable to marine mammals in the Study Area for this SEIS/OEIS include a new stressor (high-energy laser) and the same stressors considered in the 2015 MITT Final EIS/OEIS:

- **Acoustic** (sonar and other transducers, vessel noise, aircraft noise, weapon noise)
- **Explosives** (in-air explosions and in-water explosions)
- **Energy** (in-water electromagnetic devices, high-energy lasers)
- **Physical disturbance and strike** (vessels and in-water devices, military expended materials, seafloor devices)
- **Entanglement** (wires and cables, decelerators/parachutes)
- **Ingestion** (military expended materials – munitions, military expended materials other than munitions)
- **Secondary** (impacts on habitat, impacts on prey availability)

This section of this SEIS/OEIS evaluates how and to what degree potential impacts on marine mammals from stressors described in Section 3.0 (Introduction) may have changed since the analysis presented in the 2015 MITT Final EIS/OEIS was completed. Table 2.5-1 and Table 2.5-2 in Chapter 2 (Description of Proposed Action and Alternatives) list the proposed training and testing activities and include the number of times each activity would be conducted annually and the locations within the Study Area

where the activity would typically occur under each alternative. The tables also present the same information for activities described in the 2015 MITT Final EIS/OEIS so that the proposed levels of training and testing under this SEIS/OEIS can be easily compared. The analysis in this SEIS/OEIS includes consideration of the Navy's standard operating procedures and mitigation that the Navy will implement to avoid or reduce potential impacts on marine mammals from acoustic, explosive, and physical disturbance and strike stressors. Mitigation for marine mammals will be coordinated with NMFS through the MMPA and ESA consultation processes, and is detailed in Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment) of this SEIS/OEIS.

In 2015, the Navy and NMFS determined that within the Study Area only acoustic stressors and explosive stressors could potentially result in harassment and/or the incidental taking of marine mammals from Navy training and testing activities (80 FR 46112) and that none of the other stressors would result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b).

There has been no emergent science that would necessitate changes to conclusions reached by Navy or NMFS (as a cooperating agency) in the 2015 MITT Final EIS/OEIS regarding those other dismissed stressors as having a negligible or discountable impact on marine mammal populations or species. As detailed in Chapter 2 (Description of Proposed Action and Alternatives) of this SEIS/OEIS, there are no changes to proposed training and testing activities that would necessitate re-analysis of any of the activities associated with those stressors for which NMFS has previously determined did not rise to the level of a take under MMPA. The analysis presented in this section of the SEIS/OEIS also considers standard operating procedures, which are discussed in Section 2.3.3 (Standard Operating Procedures) of this Draft SEIS/OEIS, and mitigation measures that are described in Chapter 5 (Mitigation). The Navy would implement these measures to avoid or reduce potential impacts on marine mammals from stressors associated with the proposed training and testing activities. Mitigation for marine mammals will be coordinated with NMFS through the ESA consultation process. In addition, in the Draft SEIS/OEIS the Navy has developed Appendix I (Geographic Mitigation Assessment), which details consideration of specific mitigation areas identified by the public during the scoping process. In short, Appendix I contains the background information for each area being considered and lays out the methodology used by the Navy in its scientific and operational analysis for assessing and developing proposed geographic mitigation areas within the MITT Study Area to further avoid or reduce potential impacts on marine mammals in areas that may be of particular biological importance.

As presented in Section 3.0 (Introduction), since completion of the 2015 MITT Final EIS/OEIS there have been refinements made in the modeling of potential impacts from sonar and other transducers and in-water explosives. These changes have been incorporated into the re-analysis of acoustic and explosive stressors presented in this SEIS/OEIS. In addition to the new effects criteria, weighting functions, and thresholds for multiple species, new information for marine mammals includes the integration of new marine mammal density data based on new survey data (Bradford et al., 2017) and the integration of data from acoustic monitoring (Norris et al., 2017; Yack et al., 2016).

There have been no changes to the MITT Study Area, existing conditions, species life histories, or any new information available since 2015 that would otherwise substantively change the conclusions<sup>5</sup> presented in the 2015 MITT Final EIS/OEIS. What is new since 2015 are refinements to the Navy Acoustic Effects Model. This SEIS/OEIS, therefore, focuses on a re-analysis of potential impacts on marine mammals from acoustic stressors involving use of sonar and other transducers, and the use of in-water explosives. The following paragraphs provide details on refinements to the Navy's acoustic modeling since 2015. Most important is the information found in Section 3.4.3.4 (Summary of Monitoring and Observations During Navy Activities Since 2015) regarding scientific data gathered on marine mammals in locations where the Navy has been training and testing, which serves as an empirical basis for the marine mammal impact assessment presented in this SEIS/OEIS.

### **New effects criteria, weighting functions, and thresholds**

A detailed description of the Phase III acoustic and explosive criteria and threshold development regarding marine mammals is included in the supporting technical report *Criteria and Thresholds for U.S Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a). In summary, the update to the acoustic impact criteria has largely been predicated on a series of behavioral studies (often sponsored by the U.S. Navy), which have led to a new understanding of how some marine mammals react to sonar and other sound sources (Baird et al., 2017; Crowell et al., 2016; Curé et al., 2016; Dunlop et al., 2016; Friedlaender et al., 2016; Graham et al., 2017; Harris et al., 2018; Henderson et al., 2015a; Kvasdheim et al., 2017; Pirotta et al., 2016; Sabet et al., 2016; Sivle et al., 2015; Southall et al., 2016; Visser et al., 2016). As a result of that new understanding, the previous behavioral response functions for estimating alterations in behavior have been refined to accurately reflect studies undertaken both in the ocean and in well controlled studies done in laboratory settings. Additional studies have also provided information allowing for the refinement of the previous auditory weighting functions (Finneran et al., 2015; Houser et al., 2016; Houser et al., 2017; Kastelein et al., 2015b; Kastelein et al., 2015c; Kastelein et al., 2015d; Kastelein et al., 2016; Kastelein et al., 2017; Mulsow et al., 2015), and has led to a new methodology to predict these functions for each hearing group along with the accompanying hearing loss thresholds. These criteria for estimating hearing loss in marine mammals was largely adopted by NMFS for species within their purview (81 FR 51693) and is used in the analysis for impacts on marine mammals presented in this SEIS/OEIS.

The majority of the changes in the results of the impact analyses presented in this SEIS/OEIS pursuant to requirements of the MMPA arise from changes in the model input; specifically, more accurate marine mammal density data, revised acoustic impact criteria, and more comprehensive computer modeling of predicted effects on marine mammals. Assessment of likely long-term consequences to populations of

---

<sup>5</sup> Conclusions in this regard refer to the findings reached by Navy and NMFS on the two previous sets of analyses for the continuation of training and testing in Study Area. NMFS has recently re-considered analysis of Navy training and testing for many of the same for actions elsewhere (83 FR 10954 and 83 FR 29872) and for a third time reaffirmed their earlier conclusions regarding Navy military readiness activities. The Navy and NMFS have found that there would not be significant impacts on populations of marine mammals resulting from the continuation of training and testing. Under ESA, the Proposed Action may affect certain ESA-listed marine mammal species, but were not likely to jeopardize the continued existence of the continued existence of those species.

marine mammals are provided by empirical data gathered from areas where the Navy routinely trains and tests. Substantial Navy-funded marine mammal survey data, monitoring data, and scientific research have been completed since 2007. These empirical data are beginning to provide insight on the qualitative analysis of the actual (as opposed to model predicted numerical) impact on marine mammals resulting from Navy training and testing activities based on observations of marine mammals generally in and around Navy Range Complexes.

The following subsections of this SEIS/OEIS present the potential environmental consequences based on an updated modeling methodology and the scientific observations and investigations made over 12 years of monitoring training and testing activities in the Pacific and elsewhere that are representative of the type of activities proposed in this SEIS/OEIS.

#### **3.4.2.1 Acoustic Stressors**

Assessing whether a sound may disturb or injure a marine mammal involves understanding the characteristics of the acoustic sources, the marine mammals that may be present in the vicinity of the sources, and the effects that sound may have on the physiology and behavior of those marine mammals. Although it is known that sound is important for marine mammal communication, navigation, and foraging (National Research Council, 2003, 2005), there are many unknowns in assessing impacts, such as the potential interaction of different effects and the significance of responses by marine mammals to sound exposures (Nowacek et al., 2007b; Southall et al., 2007). Many other factors besides just the received level of sound may affect an animal's reaction, such as the duration of the sound-producing activity, the animal's physical condition, prior experience with the sound, activity at the time of exposure (e.g., feeding, traveling, resting), the context of the exposure (e.g., in a semi-enclosed bay versus open ocean), and proximity to the source of the sound.

The ways in which an acoustic exposure could result in immediate effects or long-term consequences for an animal are explained in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities). The following background section discusses what is currently known about acoustic effects to marine mammals. These effects could hypothetically extend from physical injury or trauma to a behavioral or stress response that may or may not be detectable. Injury (physical trauma) can occur to organs or tissues of an animal (Section 3.4.2.1.1.1, Injury). Hearing loss (Section 3.4.2.1.1.2, Hearing Loss) is a noise-induced decrease in hearing sensitivity, which can be either temporary or permanent. Masking (Section 3.4.2.1.1.4, Masking) can occur when the perception of a biologically important sound (i.e., signal) is interfered with by a second sound (i.e., noise). Physiological stress (Section 3.4.2.1.1.3, Physiological Stress) is an adaptive process that helps an animal cope with changing conditions; however, too much stress can potentially result in additional physiological effects. Behavioral response (Section 3.4.2.1.1.5, Behavioral Reactions) ranges from brief distractions to avoidance of a sound source to prolonged flight. Extreme behavioral or physiological responses can lead to stranding (Section 3.4.2.1.1.6, Stranding). Long-term consequences (Section 3.4.2.1.1.7, Long-Term Consequences) are those impacts, or accumulation of impacts, that can result in decreases in individual fitness or population changes. To avoid or reduce potential impacts to the maximum extent practicable, the Navy will implement marine mammal mitigation measures during applicable training and testing activities that generate acoustic stressors (see Chapter 5, Mitigation; and Appendix I, Geographic Mitigation Assessment).

The Navy will rely on the previous 2015 MITT Final EIS/OEIS for the analysis of vessel noise, aircraft noise, and weapon noise; new applicable and emergent science in regard to these sub-stressors is presented in the sections that follow. Due to new acoustic impact criteria, marine mammal densities,

and revisions to the acoustic effects model, the analysis provided in Section 3.4.2.1.2 (Impacts from Sonar and Other Transducer Stressors) of this SEIS/OEIS supplants the 2015 MITT Final EIS/OEIS for marine mammals and changes estimated impacts for some species since the 2015 MITT Final EIS/OEIS.

#### **3.4.2.1.1 Background**

##### **3.4.2.1.1.1 Injury**

Injury (i.e., physical trauma) refers to the effects on the tissues or organs of an animal due to exposure to pressure waves. Injury due to exposure to non-explosive acoustic stressors such as sonar is discussed below. Moderate- to low-level sound sources, including vessel and aircraft noise, would not cause any injury. Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) provides additional information on injury (i.e., physical trauma) and the framework used to analyze this potential impact.

Several mechanisms of acoustically induced tissue damage (non-auditory) have been proposed and are discussed below.

#### **Injury Due to Sonar-Induced Acoustic Resonance**

An object exposed to its resonant frequency will tend to amplify its vibration at that frequency, a phenomenon called acoustic resonance. Acoustic resonance has been proposed as a mechanism by which a sonar or sources with similar operating characteristics could damage tissues of marine mammals. In 2002, NMFS convened a panel of government and private scientists to investigate the potential for acoustic resonance to occur in marine mammals (National Oceanic and Atmospheric Administration, 2002). They modeled and evaluated the likelihood that Navy mid-frequency sonar caused resonance effects in beaked whales that eventually led to their stranding. The conclusions of the group were that resonance in air-filled structures was not likely to have caused the Bahamas stranding in 2000. The frequency at which resonance was predicted to occur in the animals' lungs was 50 Hz, well below the frequencies used by the mid-frequency sonar systems associated with the Bahamas event. Furthermore, air cavity vibrations, even at resonant frequencies, were not considered to be of sufficient amplitude to cause tissue damage, even under the unrealistic scenario in which air volumes would be undamped (unrestrained) by surrounding tissues and the amplitude of the resonant response would be greatest. These same conclusions would apply to other training and testing activities involving acoustic sources. Therefore, the Navy concludes that acoustic resonance would not occur under realistic conditions during training and testing activities, and this type of impact is not considered further in this analysis.

#### **Nitrogen Decompression**

Marine mammals are thought to deal with nitrogen loads in their blood and other tissues, caused by gas exchange from the lungs under conditions of high ambient pressure during diving, through anatomical, behavioral, and physiological adaptations (Hooker et al., 2012).

Although not a direct injury, variations in marine mammal diving behavior or avoidance responses have been hypothesized to result in nitrogen off-gassing in super-saturated tissues, possibly to the point of deleterious vascular and tissue bubble formation (Hooker et al., 2012; Jepson et al., 2003; Saunders et al., 2008) with resulting symptoms similar to decompression sickness (also known as "the bends"). The process has been under debate in the scientific community (Hooker et al., 2012; Saunders et al., 2008), although analyses of bycaught and drowned animals have demonstrated that nitrogen bubble formation can occur in animals that no longer exchange gas with the lungs (drowned) and which are brought to the

surface where tissues become supersaturated with nitrogen due to the reduction in hydrostatic pressure (Bernaldo de Quiros et al., 2013b; Moore et al., 2009). Deep-diving whales, such as beaked whales, have been predicted to have higher nitrogen loads in body tissues for certain modeled changes in dive behavior, which might make them more susceptible to decompression (Fahlman et al., 2014b; Fernandez et al., 2005; Hooker et al., 2012; Jepson et al., 2003).

Researchers have examined how dive behavior affects tissue supersaturation conditions that could put an animal at risk of gas bubble embolism. An early hypothesis was that if exposure to a startling sound elicits a rapid ascent to the surface, tissue gas saturation sufficient for the evolution of nitrogen bubbles might result (Fernandez et al., 2005; Jepson et al., 2003). However, modeling suggested that even unrealistically rapid rates of ascent from normal dive behaviors are unlikely to result in supersaturation to the extent that bubble formation would be expected in beaked whales (Zimmer & Tyack, 2007). Instead, emboli observed in animals exposed to mid-frequency active sonar (Fernandez et al., 2005; Jepson et al., 2003) could stem from a behavioral response that involves repeated dives, shallower than the depth of lung collapse (Aguilar de Soto et al., 2006; Hooker et al., 2012; Tyack et al., 2006; Zimmer & Tyack, 2007). Longer times spent diving at mid-depths above lung collapse would allow gas exchange from the lungs to continue under high hydrostatic pressure conditions, increasing potential for supersaturation; below the depth of lung collapse, gas exchange from the lungs to the blood would likely not occur (Fahlman et al., 2014b). To examine the potential for gas bubble formation, a bottlenose dolphin was trained to dive repetitively to depths shallower than lung collapse to elevate nitrogen saturation to the point that asymptomatic nitrogen bubble formation was predicted to occur. However, inspection of the vascular system of the dolphin via ultrasound did not demonstrate the formation of any nitrogen gas bubbles (Houser et al., 2009). To estimate risk of decompression sickness, Kvadsheim et al. (2012) modeled gas exchange in the tissues of sperm, pilot, killer, and beaked whales based on actual dive behavior during exposure to sonar in the wild. Results indicated that venous supersaturation was within the normal range for these species, which have naturally high levels of nitrogen loading.

Still, little is known about respiratory physiology of deep-diving breath-hold animals. Costidis and Rommel (2016) suggest that gas exchange may continue to occur across the tissues of air-filled sinuses in deep-diving odontocetes below the depth of lung collapse, if hydrostatic pressures are high enough to drive gas exchange across into non-capillary veins, contributing to tissue gas loads. Researchers have also considered the role of carbon dioxide accumulation produced during periods of high activity by an animal, theorizing that accumulating carbon dioxide, which cannot be removed by gas exchange below the depth of lung collapse, may facilitate the formation of bubbles in nitrogen-saturated tissues (Bernaldo de Quiros et al., 2012; Fahlman et al., 2014b). Garcia Parraga et al. (2018) suggest that diving marine mammals have physiological and anatomical adaptations to control gas uptake above the depth of lung collapse, favoring oxygen uptake while minimizing nitrogen uptake. Under the hypothesis of Garcia Parraga et al. (2018), elevated activity due to a strong evasive response could lead to increased uptake of nitrogen, resulting in an increased risk of nitrogen decompression.

Modeling has suggested that the long, deep dives performed regularly by beaked whales over a lifetime could result in the saturation of long-halftime tissues (i.e., tissues that take longer to give off nitrogen, such as fat and bone lipid) to the point that they are supersaturated when the animals are at the surface (Fahlman et al., 2014b; Hooker et al., 2009; Saunders et al., 2008). The presence of osteonecrosis (bone death due to reduced blood flow) in deep-diving sperm whales has been offered as evidence of chronic supersaturation (Moore & Early, 2004). Proposed adaptations for prevention of bubble formation under conditions of persistent tissue saturation have been suggested (Fahlman et al., 2006; Hooker et al.,

2009), while the condition of supersaturation required for bubble formation in these tissues has been demonstrated in marine mammals drowned at depth as fisheries bycatch and brought to the surface (Moore et al., 2009). For beaked whale strandings associated with sonar use, one theory is that observed bubble formation might be caused by long periods of compromised blood flow caused by the stranding itself (which reduces ability to remove nitrogen from tissues) following rapid ascent dive behavior that does not allow for typical management of nitrogen in supersaturated, long-half-time tissues (Houser et al., 2009).

A fat embolic syndrome (out-of-place fat particles, typically in the bloodstream) was identified by Fernández et al. (2005) coincident with the identification of bubble emboli in stranded beaked whales. The fat embolic syndrome was the first pathology of this type identified in marine mammals and was thought to possibly arise from the formation of bubbles in fat bodies, which subsequently resulted in the release of fat emboli into the blood stream. Although rare, similar findings have been found in the Risso's dolphin, another deep-diving species, but with presumably non-anthropogenic causes (Fernandez et al., 2017).

Dennison et al. (2012) reported on investigations of dolphins stranded in 2009–2010 and, using ultrasound, identified gas bubbles in kidneys of 21 of the 22 live-stranded dolphins and in the livers of two of the 22. The authors postulated that stranded animals were unable to recompress by diving, and thus retained bubbles that would have otherwise re-absorbed in animals that continued to dive. The researchers concluded that the minor bubble formation observed could be tolerated since the majority of stranded dolphins released did not re-strand.

The appearance of extensive bubble and fat emboli in beaked whales was unique to a small number of strandings associated with certain high-intensity sonar events; the phenomenon has not been observed to the same degree in other stranded marine mammals, including other beaked whale strandings not associated with sonar use. It is uncertain as to whether there is some more easily triggered mechanism for this phenomenon specific to beaked whales or whether the phenomenon occurs only following rapidly occurring stranding events (i.e., when whales are not capable of sufficiently decompressing). Nevertheless, based on the rarity of observations of bubble pathology, the potential for nitrogen decompression sickness, or “the bends,” is considered discountable.

#### **Acoustically Induced Bubble Formation Due to Sonars**

A suggested cause of injury to marine mammals is rectified diffusion (Crum & Mao, 1996), the process of increasing the size of a microscopic gas bubble by exposing it to a sound field. The process is dependent upon a number of factors, including the sound pressure level (SPL) and duration. Under this hypothesis, microscopic bubbles assumed to exist in the tissues of marine mammals may experience one of three things: (1) bubbles grow to the extent they become emboli or cause localized tissue trauma, (2) bubbles develop to the extent that a complement immune response is triggered or the nervous tissue is subjected to enough localized pressure that pain or dysfunction occurs (a stress response without injury), or (3) the bubbles are cleared by the lung without negative consequence to the animal.

Rectified diffusion is facilitated if the environment in which the ensonified bubbles exist is supersaturated with gas. As discussed above, repetitive diving by marine mammals can cause the blood and some tissues to become supersaturated (Ridgway & Howard, 1979). The dive patterns of some marine mammals (e.g., beaked whales) are predicted to induce greater supersaturation (Houser et al., 2001). If rectified diffusion were possible in marine mammals exposed to high-level sound, conditions of tissue supersaturation could theoretically speed the rate and increase the size of bubble growth.

Subsequent effects due to tissue trauma and emboli would presumably mirror those observed in humans suffering from decompression sickness.

It is unlikely that the short duration of sonar pulses would be long enough to drive bubble growth to any substantial size, if such a phenomenon occurs. However, an alternative but related hypothesis has also been suggested: stable microbubbles could be destabilized by high-level sound exposures such that bubble growth then occurs through static diffusion of gas out of supersaturated tissues. In such a scenario, the marine mammal would need to be in a gas-supersaturated state for a long enough time for bubbles to become a problematic size. The phenomena of bubble growth due to a destabilizing exposure was shown by Crum et al. (2005) by exposing highly supersaturated ex vivo bovine tissues to a 37 kHz source at 214 dB re 1  $\mu$ Pa. Although bubble growth occurred under the extreme conditions created for the study, these conditions would not exist in the wild because the levels of tissue supersaturation in the study (as high as 400–700 percent) are substantially higher than model predictions for marine mammals (Fahlman et al., 2009; Fahlman et al., 2014b; Houser et al., 2001; Saunders et al., 2008), and such high exposure level would only occur in very close proximity to the most powerful sonars. It is improbable that this mechanism is responsible for stranding events or traumas associated with beaked whale strandings.

There has been considerable disagreement among scientists as to the likelihood of this phenomenon (Evans & Miller, 2003; Piantadosi & Thalmann, 2004). Although it has been argued that traumas from beaked whale strandings are consistent with gas emboli and bubble-induced tissue separations (Fernandez et al., 2005; Jepson et al., 2003), nitrogen bubble formation as the cause of the traumas has not been verified. The presence of bubbles postmortem, particularly after decompression, is not necessarily indicative of bubble pathology (Bernaldo de Quiros et al., 2012; Bernaldo de Quiros et al., 2013a; Bernaldo de Quiros et al., 2013b; Dennison et al., 2012; Moore et al., 2009).

#### **3.4.2.1.1.2 Hearing Loss**


Exposure to intense sound may result in noise-induced hearing loss that persists after cessation of the noise exposure. Hearing loss may be temporary or permanent, depending on factors such as the exposure frequency, received sound pressure level, temporal pattern, and duration. The frequencies affected by hearing loss will vary depending on the frequency of the fatiguing noise, with frequencies at and above the noise frequency most strongly affected. The amount of hearing loss may range from slight to profound, depending on the ability of the individual to hear at the affected frequencies.

Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) provides additional information on hearing loss and the framework used to analyze this potential impact. Hearing loss has only been studied in a few species of marine mammals, although hearing studies with terrestrial mammals are also informative.

Hearing loss is typically quantified in terms of threshold shift (TS)—the amount (in dB) that hearing thresholds at one or more specified frequencies are elevated, compared to their pre-exposure values, at some specific time after the noise exposure. The amount of TS measured usually decreases with increasing recovery time—the amount of time that has elapsed since a noise exposure. If the TS eventually returns to zero (i.e., the hearing threshold returns to the pre-exposure value), the threshold shift is called a temporary threshold shift (TTS). If the TS does not completely recover (the threshold remains elevated compared to the pre-exposure value), the remaining TS is called a permanent threshold shift (PTS). Figure 3.4-2 shows two hypothetical TSs: one that completely recovers, a TTS, and one that does not completely recover, leaving some PTS. By definition, TTS is a function of the recovery

time, therefore comparing the severity of noise exposures based on the amount of induced TTS can only be done if the recovery times are also taken into account. For example, a 20 dB TTS measured 24 hours post-exposure indicates a more hazardous exposure than one producing 20 dB of TTS measured only two minutes after exposure; if the TTS is 20 dB after 24 hours, the TTS measured after two minutes would have likely been much higher. Conversely, if 20 dB of TTS was measured after two minutes, the TTS measured after 24 hours would likely have been much smaller.

Studies have revealed that intense noise exposures may also cause auditory system injury that does not result in PTS (i.e., hearing thresholds return to normal after the exposure, but there is injury nonetheless). Kujawa and Liberman (2009) found that noise exposures sufficient to produce a TTS of 40 dB, measured 24 hours post-exposure using electro-physiological methods, resulted in acute loss of nerve terminals and delayed degeneration of the cochlear nerve in mice. Lin et al. (2011) found a similar result in guinea pigs, that a TTS in auditory evoked potential of up to approximately 50 dB, measured 24 hours post-exposure, resulted in neural degeneration. These studies demonstrate that PTS should not be used as the sole indicator of auditory injury, since exposures producing high levels of TTS (40 to 50 dB measured 24 hours after exposure)—but no PTS—may result in auditory injury.



Notes: TTS = Temporary Threshold Shift, TS = Threshold Shift, PTS = Permanent Threshold Shift

**Figure 3.4-2: Two Hypothetical Threshold Shifts**

There are no simple functional relationships between TTS and the occurrence of PTS or other auditory injury (e.g., neural degeneration). However, TTS and PTS are, by definition, mutually exclusive. An exposure that produces TTS cannot also produce PTS in the same individual; conversely, if an initial threshold shift only partially recovers, resulting in some amount of PTS, the difference between the initial TS and the PTS is not called TTS. As TTS increases, the likelihood that additional exposure SPL or duration will result in PTS or other injury also increases. Exposure thresholds for the occurrence of PTS or other auditory injury can therefore be defined based on a specific amount of TTS; that is, although an exposure has been shown to produce only TTS, we assume that any additional exposure may result in some PTS or other injury. The specific upper limit of TTS is based on experimental data showing amounts of TTS that have not resulted in PTS or injury. In other words, we do not need to know the exact functional relationship between TTS and PTS or other injury, we only need to know the upper limit for TTS before some PTS or injury is possible.

A variety of human and terrestrial mammal data indicate that threshold shifts up to 40 to 50 dB may be induced without PTS, and that 40 dB is a reasonable upper limit for allowable threshold shift to prevent PTS (e.g., Kryter et al., 1965; Miller et al., 1963; Ward et al., 1958; Ward et al., 1959; Ward, 1960). It is

reasonable to assume the same relationship would hold for marine mammals, since there are many similarities between the inner ears of marine and terrestrial mammals, and experiments with marine mammals have revealed similarities to terrestrial mammals for features such as TTS, age-related hearing loss, drug-induced hearing loss, masking, and frequency selectivity (Finneran et al., 2005a; Finneran, 2015; Ketten, 2000). Therefore, we assume that sound exposures sufficient to produce 40 dB of TTS measured approximately four minutes after exposure represent the limit of a non-injurious exposure (i.e., higher-level exposures have the potential to cause auditory injury). Exposures sufficient to produce a TTS of 40 dB, measured approximately four minutes after exposure, therefore represent the threshold for auditory injury. The predicted injury could consist of either hair cell damage/loss resulting in PTS or other auditory injury, such as the delayed neural degeneration identified by Kujawa and Liberman (2009) and Lin et al. (2011) that may not result in PTS.

Numerous studies have directly examined noise-induced hearing loss in marine mammals (see Finneran, 2015). In these studies, hearing thresholds were measured in marine mammals before and after exposure to intense sounds. The difference between the pre-exposure and post-exposure thresholds was then used to determine the amount of TTS at various post-exposure times. The major findings from these studies include the following:

- The method used to test hearing may affect the resulting amount of measured TTS, with neurophysiological measures producing larger amounts of TTS compared to psychophysical measures (Finneran et al., 2007; Finneran, 2015).
- The amount of TTS varies with the hearing test frequency. As the exposure SPL increases, the frequency at which the maximum TTS occurs also increases (Kastelein et al., 2014b). For high-level exposures, the maximum TTS typically occurs one-half to one octave above the exposure frequency (Finneran et al., 2007; Mooney et al., 2009a; Nachtigall et al., 2004; Popov et al., 2011; Popov et al., 2013; Schlundt et al., 2000). The overall spread of TTS from tonal exposures can therefore extend over a large frequency range (i.e., narrowband exposures can produce broadband [greater than one octave] TTS).
- The amount of TTS increases with exposure SPL and duration and is correlated with sound exposure level (SEL), especially if the range of exposure durations is relatively small (Kastak et al., 2007; Kastelein et al., 2014b; Popov et al., 2014). As the exposure duration increases, however, the relationship between TTS and SEL begins to break down. Specifically, duration has a more significant effect on TTS than would be predicted on the basis of SEL alone (Finneran et al., 2010a; Kastak et al., 2005; Mooney et al., 2009a). This means if two exposures have the same SEL but different durations, the exposure with the longer duration (thus lower SPL) will tend to produce more TTS than the exposure with the higher SPL and shorter duration. In most acoustic impact assessments, the scenarios of interest involve shorter duration exposures than the marine mammal experimental data from which impact thresholds are derived; therefore, use of SEL tends to over-estimate the amount of TTS. Despite this, SEL continues to be used in many situations because it is relatively simple, more accurate than SPL alone, and lends itself easily to scenarios involving multiple exposures with different SPL.
- The amount of TTS depends on the exposure frequency. Sounds at low frequencies, well below the region of best sensitivity, are less hazardous than those at higher frequencies, near the region of best sensitivity (Finneran & Schlundt, 2013). The onset of TTS—defined as the exposure level necessary to produce 6 dB of TTS (i.e., clearly above the typical variation in threshold measurements)—also varies with exposure frequency. At low frequencies onset-TTS exposure levels are higher compared to those in the region of best sensitivity.

- TTS can accumulate across multiple exposures, but the resulting TTS will be less than the TTS from a single, continuous exposure with the same SEL (Finneran et al., 2010a; Kastelein et al., 2014b; Kastelein et al., 2015b; Mooney et al., 2009b). This means that TTS predictions based on the total, cumulative SEL will overestimate the amount of TTS from intermittent exposures such as sonars and impulsive sources.
- The amount of observed TTS tends to decrease with increasing time following the exposure; however, the relationship is not monotonic (i.e., increasing exposure does not always increase TTS). The time required for complete recovery of hearing depends on the magnitude of the initial shift; for relatively small shifts recovery may be complete in a few minutes, while large shifts (e.g., approximately 40 dB) may require several days for recovery. Under many circumstances TTS recovers linearly with the logarithm of time (Finneran et al., 2010a, 2010b; Finneran & Schlundt, 2013; Kastelein et al., 2012a; Kastelein et al., 2012b; Kastelein et al., 2013a; Kastelein et al., 2014b; Kastelein et al., 2014c; Popov et al., 2011; Popov et al., 2013; Popov et al., 2014). This means that for each doubling of recovery time, the amount of TTS will decrease by the same amount (e.g., 6 dB recovery per doubling of time).

Nachtigall et al. (2018) and Finneran (2018) describe the measurements of hearing sensitivity of multiple odontocete species (bottlenose dolphin, harbor porpoise, beluga, and false killer whale) when a relatively loud sound was preceded by a warning sound. These captive animals were shown to reduce hearing sensitivity when warned of an impending intense sound. Based on these experimental observations of captive animals, the authors suggest that wild animals may dampen their hearing during prolonged exposures or if conditioned to anticipate intense sounds. Finneran recommends further investigation of the mechanisms of hearing sensitivity reduction in order to understand the implications for interpretation of some existing temporary threshold shift data obtained from captive animals, notably for considering TTS due to short duration, unpredictable exposures. No modification of analysis of auditory impacts is currently suggested, as the Phase III auditory impact thresholds are based on best available data for both impulsive and non-impulsive exposures to marine mammals.

Due to the higher exposure levels or longer exposure durations required to induce hearing loss, only a few types of human-made sound sources have the potential to cause a threshold shift to a marine mammal in the wild. These include some sonars and other transducers and impulsive sound sources such as air guns and impact pile driving, neither of which will be used in the Study Area.

#### **Threshold Shift due to Sonar and Other Transducers**

TTS in mid-frequency cetaceans exposed to non-impulsive sound has been investigated in multiple studies (Finneran et al., 2005b; Finneran et al., 2010b; Finneran & Schlundt, 2013; Mooney et al., 2009a; Mooney et al., 2009b; Nachtigall et al., 2003; Nachtigall et al., 2004; Popov et al., 2013; Popov et al., 2014; Schlundt et al., 2000) from two species, bottlenose dolphins and beluga whales. Two high-frequency cetacean species have been studied for TTS due to non-impulsive sources: the harbor porpoise (Kastelein et al., 2012b) and the finless porpoise (*Neophocaena phocaenoides*) (Popov et al., 2011). TTS from non-impulsive sounds has also been investigated in three pinniped species: harbor seal (*Phoca vitulina*), California sea lion (*Zalophus californianus*), and Northern elephant seal (*Mirounga angustirostris*) (e.g., Kastak et al., 2005; Kastelein et al., 2012a). These data are reviewed in detail in Finneran (2015) as well as the *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* technical report (U.S. Department of the Navy, 2017a), and the major findings are summarized above.

### 3.4.2.1.1.3 Physiological Stress

The growing field of conservation physiology relies in part on the ability to monitor stress hormones in populations of animals, particularly those that are threatened or endangered. The ability to make predictions from stress hormones about impacts on individuals and populations exposed to various forms of stressors, natural and human-caused, relies on understanding the linkages between changes in stress hormones and resulting physiological impacts. At this time, the sound characteristics that correlate with specific stress responses in marine mammals are poorly understood, as are the ultimate consequences due to these changes. Navy-funded efforts are underway to try to improve the understanding of and the ability to predict how stressors ultimately affect marine mammal populations (e.g., King et al., 2015; New et al., 2013a; New et al., 2013b; Pirotta et al., 2015a). With respect to acoustically induced stress, this includes not only determining how and to what degree various types of anthropogenic sound cause stress in marine mammals, but what factors can mitigate those responses. Factors potentially affecting an animal's response to a stressor include the mammal's life history stage, sex, age, reproductive status, overall physiological and behavioral plasticity, and whether they are naïve or experienced with the sound [e.g., prior experience with a stressor may result in a reduced response due to habituation (Finneran & Branstetter, 2013; St. Aubin & Dierauf, 2001a)]. Because there are many unknowns regarding the occurrence of acoustically induced stress responses in marine mammals, the Navy assumes in its effects analysis that any physiological response (e.g., hearing loss or injury) or significant behavioral response is also associated with a stress response.

Marine mammals naturally experience stressors within their environment and as part of their life histories. Changing weather and ocean conditions, exposure to disease and naturally occurring toxins, lack of prey availability, and interactions with predators all contribute to the stress a marine mammal experiences (Atkinson et al., 2015). Breeding cycles, periods of fasting, and social interactions with members of the same species are also stressors, although they are natural components of an animal's life history. Anthropogenic activities have the potential to provide additional stressors beyond those that occur naturally (Fair et al., 2014; Meissner et al., 2015; Rolland et al., 2012). Anthropogenic stressors potentially include such things as fishery interactions, pollution, tourism, and ocean noise.

The stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor (Moberg & Mench, 2000). However, if the magnitude and duration of the stress response is too great or too long, then it can have negative consequences to the organism (e.g., decreased immune function, decreased reproduction). The generalized stress response is classically characterized by the release of cortisol, a hormone that has many functions including elevation of blood sugar, suppression of the immune system, and alteration of the biochemical pathways that affect fat, protein, and carbohydrate metabolism. However, it is now known that the endocrine response (glandular secretions of hormones into the blood) to a stressor can extend to other hormones. For instance, thyroid hormones can also vary under the influence of certain stressors, particularly food deprivation. These types of responses typically occur on the order of minutes to days. The "fight or flight" response, an acute stress response, is characterized by the very rapid release of hormones that stimulate glucose release, increase heart rate, and increase oxygen consumption.

What is known about the function of the various stress hormones is based largely upon observations of the stress response in terrestrial mammals. The endocrine response of marine mammals to stress may not be the same as that of terrestrial mammals because of the selective pressures marine mammals faced during their evolution in an ocean environment (Atkinson et al., 2015). For example, due to the necessity of breath-holding while diving and foraging at depth, the physiological role of epinephrine and

norepinephrine (the catecholamines) in marine mammals might be different than in other mammals. Catecholamines increase during breath-hold diving in seals, co-occurring with a reduction in heart rate, peripheral vasoconstriction (constriction of blood vessels), and an increased reliance on anaerobic metabolism during extended dives (Hance et al., 1982; Hochachka et al., 1995; Hurford et al., 1996); the catecholamine increase is not associated with an increased heart rate, glycemic release, and increased oxygen consumption typical of terrestrial mammals. Other hormone functions might also be different, such as aldosterone, which has been speculated to not only contribute to electrolyte balance, but possibly also the maintenance of blood pressure during periods of vasoconstriction (Houser et al., 2011). In marine mammals, aldosterone is thought to play a particular role in stress mediation because of its pronounced increase in response to handling stress (St. Aubin & Dierauf, 2001a; St. Aubin & Geraci, 1989).

Relatively little information exists on the linkage between anthropogenic sound exposure and stress in marine mammals, and even less information exists on the ultimate consequences of sound-induced stress responses (either acute or chronic). Most studies to date have focused on acute responses to sound either by measuring catecholamines or by measuring heart rate as an assumed proxy for an acute stress response. Belugas demonstrated no catecholamine response to the playback of oil drilling sounds (Thomas et al., 1990b) but showed a small but statistically significant increase in catecholamines following exposure to impulsive sounds produced from a seismic water gun (Romano et al., 2004). A bottlenose dolphin exposed to the same seismic water gun signals did not demonstrate a catecholamine response, but did demonstrate a statistically significant elevation in aldosterone (Romano et al., 2004), albeit the increase was within the normal daily variation observed in this species (St. Aubin et al., 1996). Increases in heart rate were observed in bottlenose dolphins to which known calls of other dolphins were played, although no increase in heart rate was observed when background tank noise was played back (Miksis et al., 2001). Unfortunately, in this study, it cannot be determined whether the increase in heart rate was due to stress or an anticipation of being reunited with the dolphin to which the vocalization belonged. Similarly, a young beluga's heart rate was observed to increase during exposure to noise, with increases dependent upon the frequency band of noise and duration of exposure, and with a sharp decrease to normal or below normal levels upon cessation of the exposure (Lyamin et al., 2011). Spectral analysis of heart rate variability corroborated direct measures of heart rate (Bakhchina et al., 2017). This response might have been in part due to the conditions during testing, the young age of the animal, and the novelty of the exposure; a year later the exposure was repeated at a slightly higher received level and there was no heart rate response, indicating the beluga whale had potentially acclimated to the noise exposure. Kvalsheim et al. (2010) measured the heart rate of captive hooded seals during exposure to sonar signals and found an increase in the heart rate of the seals during exposure periods versus control periods when the animals were at the surface. When the animals dove, the normal dive-related bradycardia (decrease in heart rate) was not impacted by the sonar exposure. Similarly, Thompson et al. (1998) observed a rapid but short-lived decrease in heart rates in harbor and grey seals exposed to seismic air guns (cited in Gordon et al., 2003). Williams et al. (2017) recently monitored the heart rates of narwhals released from capture and found that a profound dive bradycardia persisted, even though exercise effort increased dramatically as part of their escape response following release. Thus, although some limited evidence suggests that tachycardia might occur as part of the acute stress response of animals that are at the surface, the dive bradycardia persists during diving and might be enhanced in response to an acute stressor.

Whereas a limited amount of work has addressed the potential for acute sound exposures to produce a stress response, almost nothing is known about how chronic exposure to acoustic stressors affects

stress hormones in marine mammals, particularly as it relates to survival or reproduction. In what is probably the only study of chronic noise exposure in marine mammals associating changes in a stress hormone with changes in anthropogenic noise, Rolland et al. (2012) compared the levels of cortisol metabolites in North Atlantic right whale feces collected before and after September 11, 2001. Following the events of September 11, shipping was significantly reduced in the region where fecal collections were made, and regional ocean background noise declined. Fecal cortisol metabolites significantly decreased during the period of reduced ship traffic and ocean noise (Rolland et al., 2012). Considerably more work has been conducted in an attempt to determine the potential effect of boating on smaller cetaceans, particularly killer whales (Bain, 2002; Erbe, 2002; Lusseau, 2006; Noren et al., 2009; Pirotta et al., 2015b; Rolland et al., 2012; Skarke et al., 2014; Williams et al., 2006; Williams et al., 2009; Williams et al., 2013; Williams et al., 2014b). Most of these efforts focused primarily on estimates of metabolic costs associated with altered behavior or inferred consequences of boat presence and noise, but did not directly measure stress hormones. However, Ayres et al. (2012) investigated southern resident killer whale fecal thyroid hormone and cortisol metabolites to assess two potential threats to the species recovery: lack of prey (salmon) and impacts from exposure to the physical presence of vessel traffic (but without measuring vessel traffic noise). Ayres et al. (2012) concluded from these stress hormone measures that the lack of prey overshadowed any population-level physiological impacts on southern resident killer whales due to vessel traffic. Collectively, these studies indicate the difficulty in teasing out factors that are dominant in exerting influence on the secretion of stress hormones, including the separate and additive effects of vessel presence and vessel noise. Nevertheless, although the reduced presence of the ships themselves cannot be ruled out as potentially contributing to the reduction in fecal cortisol metabolites in North Atlantic right whales, the work of Rolland et al. (2012) represents the most provocative link between ocean noise and cortisol in cetaceans to date.

Navy-funded efforts are underway to try and improve our understanding and ability to predict how stressors ultimately affect marine mammal populations (e.g., King et al., 2015; New et al., 2013a; New et al., 2013c; Pirotta et al., 2015a), and to determine whether a marine mammal being naïve or experienced with the sound (e.g., prior experience with a stressor) may result in a reduced response due to habituation (St. Aubin & Dierauf, 2001b).


#### **3.4.2.1.1.4 Masking**

Masking occurs when one sound, distinguished as the “noise,” interferes with the detection, discrimination, or recognition of another sound. The quantitative definition of masking is the amount in decibels an auditory detection, discrimination, or recognition threshold is raised in the presence of a masker (Erbe et al., 2016). As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), masking can effectively limit the distance over which a marine mammal can communicate, detect biologically relevant sounds, and echolocate (odontocetes). Masking only occurs in the presence of the masking noise and does not persist after the cessation of the noise. Masking can lead to vocal changes (e.g., Lombard effect, increasing amplitude, or changing frequency) and behavior changes (e.g., cessation of foraging, leaving an area) to both signalers and receivers, in an attempt to compensate for noise levels (Erbe et al., 2016).

Critical ratios are the lowest signal-to-noise ratio in which detection occurs (Finneran & Branstetter, 2013; Johnson et al., 1989; Southall et al., 2000). When expressed in dB, critical ratios can easily be calculated by subtracting the noise level (in dB re 1  $\mu\text{Pa}^2/\text{Hz}$ ) from the signal level (in dB re 1  $\mu\text{Pa}$ ) at threshold. Critical ratios have been measured for pinnipeds (Southall et al., 2000, 2003; Thomas et al., 1990a), odontocetes (see Figure 3.4-3) (Au & Moore, 1990; Branstetter et al., 2017b; Johnson et al.,

1989; Kastelein & Wensveen, 2008; Lemonds et al., 2011), manatees (Gaspard et al., 2012), and sea otters (Ghoul & Reichmuth, 2014). Critical ratios are directly related to the bandwidth of auditory filters; as a result, critical ratios increase as a function of signal frequency (Au & Moore, 1990; Lemonds et al., 2011). Higher frequency noise is more effective at masking higher frequency signals. Composite critical ratio functions have been estimated for odontocetes, which allows predictions of masking if the spectral density of noise is known (Branstetter et al., 2017b). Although critical ratios are typically estimated in controlled laboratory conditions using Gaussian (white) noise, critical ratios can vary considerably (see Figure 3.4-4) depending on the noise type (Branstetter et al., 2013; Branstetter et al., 2017b; Trickey et al., 2010)


Clark et al. (2009) developed a model for estimating masking effects on communication signals for low-frequency cetaceans, including calculating the cumulative impact of multiple noise sources. For example, the model estimates that a right whale's optimal communication space (around 20 km) is decreased by 84 percent when two commercial ships pass through it. Similarly, Aguilar de Soto et al. (2006) found that a 15 dB increase in background noise due to vessels led to a communication range of only 18 percent of its normal value for foraging beaked whales. This method relies on empirical data on source levels of calls (which is unknown for many species) and requires many assumptions such as pre-industrial ambient noise conditions and simplifications of animal hearing and behavior, but it is an important step in determining the impact of anthropogenic noise on animal communication. Erbe (2016) developed a model with a noise source-centered view of masking to examine how a call may be masked from a receiver by a noise as a function of caller, receiver, and noise-source location; distance relative to each other; and received level of the call.



Source: Branstetter et al. (2017b)

Notes: (A) Odontocete critical ratios and composite model:  $CR = a[\log_{10}(f)]^b + c$ , where  $a$ ,  $b$ , and  $c$  are model coefficients and  $f$  is the signal frequency in Hz. Equation 1 was fit to aggregate data for all odontocetes. (B) *T. truncatus* critical ratios and composite model. (C) *P. phocoena* critical ratios and composite model. Parameter values for composite models are displayed in the lower right of each panel

**Figure 3.4-3: Odontocete Critical Ratios**



Source: Branstetter et al. (2013)

Notes: CM = comodulated, SS = snapping shrimp, RN = rain noise, G = Gaussian, PS = pile saw, BT = boat engine noise, and IS = ice squeaks

**Figure 3.4-4: Critical Ratios for Different Noise Types**

Vocal changes in response to anthropogenic noise can occur across the repertoire of sound production modes used by marine mammals, such as whistling, echolocation click production, calling, and singing. Vocalization changes include increasing the source level, modifying the frequency, increasing the call repetition rate of vocalizations, or ceasing to vocalize in the presence of increased noise (Hotchkiss & Parks, 2013). In cetaceans, vocalization changes were reported from exposure to anthropogenic noise sources such as sonar, vessel noise, and seismic surveying (Gordon et al., 2003; Holt et al., 2008; Holt et al., 2011; Lesage et al., 1999; McDonald et al., 2009; Rolland et al., 2012) as well as changes in the natural acoustic environment (Dunlop et al., 2014). Vocal changes can be temporary, or can be persistent, as seen in the increase in starting frequency for the North Atlantic right whale upcall over the last 50 years (Tennessen & Parks, 2016). Model simulation suggests that the frequency shift resulted in increased detection ranges between right whales; the frequency shift, coupled with an increase in call intensity by 20 dB, led to a call detectability range of less than 3 km to over 9 km (Tennessen & Parks, 2016). In some cases, these vocal changes may have fitness consequences, such as an increase in metabolic rates and oxygen consumption, as was found for bottlenose dolphins when increasing their call amplitude (Holt et al., 2015). A switch from vocal communication to physical, surface-generated sounds such as pectoral fin slapping or breaching was observed for humpback whales in the presence of increasing natural background noise levels, indicating that adaptations to masking may also move beyond vocal modifications (Dunlop et al., 2010). These changes all represent possible tactics by the sound-producing animal to reduce the impact of masking. The receiving animal can also reduce masking by using active listening strategies such as orienting to the sound source, moving to a quieter location, or reducing self-noise from hydrodynamic flow by remaining still. The temporal structure of noise (e.g., amplitude modulation) may also provide a considerable release from masking through comodulation masking release (a reduction of masking that occurs when broadband noise, with a frequency spectrum wider than an animal's auditory filter bandwidth at the frequency of interest, is amplitude modulated)

(Branstetter & Finneran, 2008; Branstetter et al., 2013). Signal type (e.g., whistles, burst-pulse, sonar clicks) and spectral characteristics (e.g., frequency modulated with harmonics) may further influence masked detection thresholds (Branstetter et al., 2016; Cunningham et al., 2014).

### **Informational Masking**

Much emphasis has been placed on signal detection in noise, and as a result, most masking studies and communication space models have focused on masked detection thresholds. However, from a fitness perspective, signal detection is almost meaningless without the ability to determine the sound source location and recognize “what” is producing the sound. Marine mammals use sound to recognize conspecifics, prey, predators, or other biologically significant sources (Branstetter et al., 2016). Masked recognition thresholds (often called informational masking) for whistle-like sounds have been measured for bottlenose dolphins (Branstetter et al., 2016), and are approximately 4 dB above detection thresholds (energetic masking) for the same signals. It should be noted that the term “threshold” typically refers to the listener’s ability to detect or recognize a signal 50 percent of the time. For example, human speech communication where only 50 percent of the words are recognized would result in poor communication (Branstetter et al., 2016). Likewise, recognition of a conspecific call or the acoustic signature of a predator at only the 50 percent level could have severe negative impacts. If “quality communication” is arbitrarily set at 90 percent recognition (which may be more appropriately related to animal fitness), the output of communication space models (which are based on 50 percent detection) would likely result in a significant decrease in communication range (Branstetter et al., 2016).

Marine mammals use sound to recognize predators (Allen et al., 2014; Cummings & Thompson, 1971; Curé et al., 2015; Fish & Vania, 1971). Auditory recognition may be reduced in the presence of a masking noise, particularly if it occurs in the same frequency band. Therefore, the occurrence of masking may prevent marine mammals from responding to the acoustic cues produced by their predators. Whether this is a possibility depends on the duration of the masking and the likelihood of encountering a predator during the time that detection and recognition of predator cues are impeded. For example, harbor seals that reside in the coastal waters off British Columbia are frequently targeted by mammal-eating killer whales. The seals acoustically discriminate between the calls of mammal-eating and fish-eating killer whales (Deecke et al., 2002), a capability that should increase survivorship while reducing the energy required to attend to all killer whale calls. Similarly, sperm whales (Curé et al., 2016; Isojunno et al., 2016), long-finned pilot whales (Visser et al., 2016), and humpback whales (Curé et al., 2015) changed their behavior in response to killer whale vocalization playbacks; these findings indicate that some recognition of predator cues could be missed if the killer whale vocalizations were masked.

### **Masking by Sonar and Other Transducers**

Masking by low-frequency or mid-frequency active sonar with relatively low-duty cycles is unlikely for most cetaceans and pinnipeds as sonar signals occur over a relatively short duration, and narrow bandwidth that does not overlap with vocalizations for most marine mammal species. While dolphin whistles and mid-frequency active sonar are similar in frequency, masking is unlikely due to the low-duty cycle of most sonars. Low-frequency active sonar could also overlap with mysticete vocalizations (e.g., minke and humpback whales). For example, in the presence of low-frequency active sonar, humpback whales were observed to increase the length of their songs (Fristrup et al., 2003; Miller et al., 2000), possibly due to the overlap in frequencies between the whale song and the low-frequency active sonar.

Newer high-duty cycle or continuous active sonars have more potential to mask vocalizations, particularly for delphinids and other mid-frequency cetaceans. These sonars transmit more frequently

(greater than 80 percent duty cycle) than traditional sonars, but at a substantially lower source level. Similarly, high-frequency acoustic sources such as pingers that operate at higher repetition rates (e.g., 2–10 kHz with harmonics up to 19 kHz, 76 to 77 pings per minute (Culik et al., 2001), also operate at lower source levels. While the lower source levels limit the range of impact compared to traditional systems, animals close to the sonar source are likely to experience masking on a much longer time scale than those exposed to traditional sonars. The frequency range at which high-duty cycle systems operate overlaps the vocalization frequency of many mid-frequency cetaceans. Continuous noise at the same frequency of communicative vocalizations may cause disruptions to communication, social interactions, and acoustically mediated cooperative behaviors such as foraging or reproductive activities. Similarly, because the systems are mid-frequency, there is the potential for the sonar signals to mask important environmental cues like predator vocalizations (e.g., killer whales), possibly affecting survivorship for targeted animals. While there are currently no available studies of the impacts of high-duty cycle sonars on marine mammals, masking due to these systems is likely analogous to masking produced by other continuous sources (e.g., vessel noise and low-frequency cetaceans), and would likely have similar short-term consequences, though longer in duration due to the duration of the masking noise. These may include changes to vocalization amplitude and frequency (Brumm & Slabbekoorn, 2005; Hotchkiss & Parks, 2013) and behavioral impacts such as avoidance of the area and interruptions to foraging or other essential behaviors (Gordon et al., 2003). Long-term consequences could include changes to vocal behavior and vocalization structure (Foote et al., 2004; Parks et al., 2007), abandonment of habitat if masking occurs frequently enough to significantly impair communication (Brumm & Slabbekoorn, 2005), a potential decrease in survivorship if predator vocalizations are masked (Brumm & Slabbekoorn, 2005), and a potential decrease in recruitment if masking interferes with reproductive activities or mother-calf communication (Gordon et al., 2003).

### **Masking by Vessel Noise**

Masking is more likely to occur in the presence of broadband, relatively continuous noise sources such as vessels. For example, right whales were observed to shift the frequency content of their calls upward while reducing the rate of calling in areas of increased anthropogenic noise (Parks et al., 2007) as well as increasing the amplitude (intensity) of their calls (Parks, 2009; Parks et al., 2011). Right whales also had their communication space reduced by up to 84 percent in the presence of vessels (Clark et al., 2009). However, Cholewiak et al. (2018) found that right whale gunshot calls had the lowest loss of communication space in Stellwagen National Sanctuary (5 percent), while fin and humpback whales lost up to 99 percent of their communication space with increased ambient noise and shipping noise combined. Although humpback whales off Australia did not change the frequency or duration of their vocalizations in the presence of ship noise, their source levels were lower than expected based on source level changes to wind noise, potentially indicating some signal masking (Dunlop, 2016).

Multiple delphinid species have also been shown to increase the minimum or maximum frequencies of their whistles in the presence of anthropogenic noise (Papale et al., 2015). More specifically, Williams et al. (2013) found that in median noise conditions in Haro Strait, killer whales lose 62 percent of their acoustic communication space due to vessel traffic noise, and in peak traffic hours lose up to 97 percent of that space. Holt et al. (2008; 2011) showed that southern resident killer whales in the waters surrounding the San Juan Islands increased their call source level as vessel noise increased. Hermanssen et al. (2014) estimated that broadband vessel noise could extend up to 160 kHz at ranges from 60 to 1,200 m, and that the higher frequency portion of that noise might mask harbor porpoise clicks. However, this may not be an issue as harbor porpoises may avoid vessels and may not be close enough

to have their clicks masked (Dyndo et al., 2015; Polacheck & Thorpe, 1990; Sairanen, 2014). Furthermore, Hermannsen et al. (2014) estimated that a 6 dB elevation in noise would decrease the hearing range of a harbor porpoise by 50 percent, and a 20 dB increase in noise would decrease the hearing range by 90 percent. Dugong vocalizations were recorded in the presence of passing boats, and although the call rate, intensity, and frequency of the calls did not change, the duration of the vocalizations was increased, as was the presence of harmonics. This may indicate more energy was being used to vocalize in order to maintain the same received level (Ando-Mizobata et al., 2014). Gervaise et al. (2012) estimated that beluga whales in the St. Lawrence Marine Park had their communication space reduced to 30 percent during average vessel traffic. During peak traffic, communication space was further reduced to 15 percent. Lesage et al. (1999) found belugas in the St. Lawrence River estuary reduced overall call rates but increased the production of certain call types when ferry and small outboard motor boats were approaching. Furthermore, these belugas increased the vocalization frequency band when vessels were in close proximity. Liu et al. (2017) found that broadband shipping noise could cause masking of humpback dolphin whistles within 1.5–3 km, and masking of echolocation clicks within 0.5–1.5 km.

#### **3.4.2.1.1.5 Behavioral Reactions**

As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), any stimulus in the environment can cause a behavioral response in marine mammals. These stimuli include noise from anthropogenic sources such as vessels, sonar, or aircraft, but could also include the physical presence of a vessel or aircraft. However, these stimuli could also influence how or if a marine mammal responds to a sound such as the presence of predators, prey, or conspecifics. Furthermore, the response of a marine mammal to an anthropogenic sound may depend on the frequency, duration, temporal pattern and amplitude of the sound as well as the animal's prior experience with the sound and their behavioral state (i.e., what the animal is doing and their energetic needs at the time of the exposure) (Ellison et al., 2011). The distance from the sound source and whether it is approaching or moving away can also affect the way an animal responds to a sound (Wartzok et al., 2003).

For marine mammals, a review of responses to anthropogenic sound was first conducted by Richardson et al. (1995b). Other reviews (Nowacek et al., 2007a; Southall et al., 2007) addressed studies conducted since 1995 and focused on observations where the received sound level of the exposed marine mammal(s) was known or could be estimated, and also examined the role of context. Southall et al. (2007) synthesized data from many past behavioral studies and observations to determine the likelihood of behavioral reactions at specific sound levels. While in general, the louder the sound source the more intense the behavioral response, it was clear that the proximity of a sound source and the animal's experience, motivation, and conditioning were also critical factors influencing the response (Southall et al., 2007; Southall et al., 2016). Ellison et al. (2011) outlined an approach to assessing the effects of sound on marine mammals that incorporates these contextual-based factors. They recommend considering not just the received level of sound, but also in what activity the animal is engaged, the nature and novelty of the sound (i.e., is this a new sound from the animal's perspective), and the distance between the sound source and the animal. They submit that this "exposure context," as described, greatly influences the type of behavioral response exhibited by the animal (see technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a)). Forney et al. (2017) also point out that an apparent lack of response (e.g., no displacement or avoidance of a sound source) may not necessarily mean there is no cost to the

individual or population, as some resources or habitats may be of such high value that animals may choose to stay, even when experiencing stress or hearing loss. Forney et al. (2017) recommend considering both the costs of remaining in an area of noise exposure such as TTS, PTS, or masking, which could lead to an increased risk of predation or other threats or a decreased capability to forage, and the costs of displacement, including potential increased risk of vessel strike or bycatch, increased risks of predation or competition for resources, or decreased habitat suitable for foraging, resting, or socializing.

Behavioral reactions could result from a variety of sound sources such as sonar and other active acoustic sources (e.g., pingers), vessel noise, and aircraft noise. There is data on the reactions of some species in different behavioral states, providing evidence on the importance of context in gauging a behavioral response. However, for most species, little or no data exist on behavioral responses to any sound source, and so all species have been grouped into broad taxonomic groups from which general response information can be inferred (see technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a)).

### **Behavioral Reactions to Sonar and Other Transducers**

Sonar and other transducers can range in frequency from less than 1 kHz (e.g., low-frequency active sonar) to over 200 kHz (e.g., fish finders), with duty cycles that range from one ping per minute to an almost continuous sound. Although very high-frequency sonars are out of the hearing range of most marine mammals, some of these sources may contain artifacts at lower frequencies that could be detected (Deng et al., 2014; Hastie et al., 2014). High-duty cycle sonar systems operate at lower source levels, but with a more continuous sound output. These sources can be stationary, or on a moving platform, and there can be more than one source present at a time. Guan et al. (2017) also found that sound levels in the mid-frequency sonar bandwidth remained elevated at least 5 dB above background levels for the first 7–15 seconds (within 2 km) after the emission of a sonar ping; depending on the length of the sonar ping and the inter-ping interval, this reverberation could increase cumulative SEL estimates during periods of active sonar. This variability in parameters associated with sonar and other transducers makes the estimation of behavioral responses to these sources difficult, with observed responses ranging from no apparent change in behavior to more severe responses that could lead to some costs to the animal. As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) and Section 3.4.2.1.1.5 (Behavioral Reactions), responses may also occur in the presence of different contextual factors regardless of received level, including the proximity and number of vessels, the behavioral state and prior experience of an individual, and even characteristics of the signal itself or the propagation of the signal through the environment.

In order to explore this complex question, behavioral response studies have been conducted through the collaboration of various research and government organizations in Bahamian, United States (off Southern California), Mediterranean, Australian, and Norwegian waters. These studies have attempted to define and measure responses of beaked whales and other cetaceans to controlled exposures of sonar and other sounds to understand better their potential impacts. While controlling for as many variables as possible (e.g., the distance and movement of the source), these studies also introduce additional variables that do not normally occur in a real Navy training or testing activity, including the tagging of whales, following the tagged animals with multiple vessels, and continually approaching the animal to create a dose escalation. In addition, distances of the sound source from the whales during behavioral response studies were always within 1–8 km. Some of these studies have suggested that ramping up a source from a lower source level would act as a mitigation measure to protect against higher order (e.g., TTS or PTS) impacts of some active sonar sources; however, this practice may only be

effective for more responsive animals, and for short durations (e.g., five minutes) of ramp-up (von Benda-Beckmann et al., 2014; von Benda-Beckmann et al., 2016). Therefore, while these studies have provided the most information to date on behavioral responses of marine mammals to sonar, there are still many contextual factors to be teased apart, and determining what might produce a significant behavioral response is not a trivial task. Additional information about active sonar ramp-up procedures, including why the Navy will not implement them as mitigation under the Proposed Action, is provided in Section 5.6.1 (Active Sonar).

Passive acoustic monitoring and visual observational behavioral response studies have also been conducted on Navy ranges, taking advantage of the existing seafloor hydrophones and real training and testing activity and associated sources to assess behavioral responses (Deakos & Richlen, 2015; Henderson et al., 2016; Manzano-Roth et al., 2016; Martin et al., 2015; McCarthy et al., 2011; Mobley & Deakos, 2015; Moretti et al., 2014; Tyack et al., 2011). In addition, extensive aerial, visual, and passive acoustic monitoring have been conducted before, during, and after training events to watch for behavioral responses during training and look for injured or stranded animals after training (Falcone et al., 2017; Farak et al., 2011; HDR, 2011b; Henderson et al., 2016; Manzano-Roth et al., 2016; Norris et al., 2012b; Smultea & Mobley, 2009; Smultea et al., 2009; Trickey et al., 2015; U.S. Department of the Navy, 2011a, 2013b, 2014a, 2015b). During all of these monitoring efforts, very few behavioral responses were observed, and no injured or dead animal was observed that was directly related to a training event (some dead animals were observed but typically before the event or appeared to have been deceased prior to the event; e.g., Smultea et al., 2011). While passive acoustic studies are limited to observations of vocally active marine mammals, and visual studies are limited to what can be observed at the surface, these study types have the benefit of occurring in the absence of some of the added contextual variables in the controlled exposure studies. Furthermore, when visual and passive acoustic data collected during a training event are combined with ship movements and sonar use, and with tagged animal data when possible, they provide a unique and realistic scenario for analysis, as in Falcone et al. (2017), Manzano-Roth et al. (2016) or Baird et al. (2017). In addition to these types of observational behavioral response studies, Harris & Thomas (2015) highlighted additional research approaches that may provide further information on behavioral responses to sonars and other transducers beyond behavior response type studies or passive acoustic monitoring, including conducting controlled exposures on captive animals with scaled (smaller sized and deployed at closer proximity) sources, on wild animals with both scaled and real but directed sources, and predator playback studies, all of which will be discussed below.

The above behavioral response studies and observations have been conducted on a number of mysticete and odontocete species, which can be extrapolated to other similar species in these taxonomic groups. No field studies of pinniped behavioral responses to sonar have been conducted; however, there are several captive studies on some pinniped and odontocete species that can provide insight into how these animals may respond in the wild. The captive studies typically represent a more controlled approach, which allow researchers to better estimate the direct impact of the received level of sound leading to behavioral responses, and to potentially link behavioral to physiological responses. However, there are still contextual factors that must be acknowledged, including previous training to complete tasks and the presence of food rewards upon completion. There are no corresponding captive studies on mysticete whales, therefore some of the responses to higher-level exposures must be extrapolated from odontocetes.

### *Mysticetes*

The responses of mysticetes to sonar and other duty-cycled tonal sounds are highly dependent upon the characteristics of the signal, the behavioral state of the animal, the particular sensitivity and previous experience of an individual, and other contextual factors including distance of the source, movement of the source, and the physical presence of vessels in addition to the sonar (Goldbogen et al., 2013b; Harris et al., 2015; Martin et al., 2015; Sivle et al., 2015). Behavioral response studies have been conducted over a variety of contextual and behavioral states, helping to identify which contextual factors may lead to a response beyond just the received level of the sound. Observed reactions during behavioral response studies have not been consistent across individuals based on received sound levels alone, and likely were the result of complex interactions between these contextual factors.

Surface feeding blue whales did not show a change in behavior in response to mid-frequency simulated and real sonar sources with received levels between 90 and 179 dB re 1  $\mu$ Pa, but deep feeding and non-feeding whales showed temporary reactions including cessation of feeding, reduced initiation of deep foraging dives, generalized avoidance responses, and changes to dive behavior (DeRuiter et al., 2017; Goldbogen et al., 2013b; Sivle et al., 2015). Similarly, while the rates of foraging lunges decreased in humpback whales due to sonar exposure, there was variability in the response across individuals, with one animal ceasing to forage completely and another animal starting to forage during the exposure (Sivle et al., 2016). In addition, lunges decreased (although not significantly) during a no-sonar control vessel approach prior to the sonar exposure, and lunges decreased less during a second sonar approach than during the initial approach, possibly indicating some response to the vessel and some habituation to the sonar and vessel after repeated approaches. In the same experiment, most of the non-foraging humpback whales did not respond to any of the approaches (Sivle et al., 2016). These humpback whales also showed variable avoidance responses, with some animals avoiding the sonar vessel during the first exposure but not the second, while others avoided the sonar during the second exposure, and only one avoided both. In addition, almost half of the animals that avoided were foraging before the exposure but the others were not; the animals that avoided while not feeding responded at a slightly lower received level and greater distance than those that were feeding (Wensveen et al., 2017). These findings indicate that the behavioral state of the animal plays a role in the type and severity of a behavioral response. In fact, when the prey field was mapped and used as a covariate in similar models looking for a response in the same blue whales, the response in deep-feeding behavior by blue whales was even more apparent, reinforcing the need for contextual variables to be included when assessing behavioral responses (Friedlaender et al., 2016). However, even when responses did occur the animals quickly returned to their previous behavior after the sound exposure ended (Goldbogen et al., 2013b; Sivle et al., 2015). In another study, humpback whales exposed to a 3 kHz pinger meant to act as a net alarm to prevent entanglement did not respond or change course, even when within 500 m (Harcourt et al., 2014). However, five out of six North Atlantic right whales exposed to an acoustic alarm interrupted their foraging dives; in this case, the alarm was comprised of a mixture of signals with frequencies from 500 to 4500 Hz, was long in duration (lasting several minutes), and was purposely designed to elicit a reaction from the animals as a prospective means to protect them from ship strikes (Nowacek et al., 2004). Although the animals' received SPL was similar in the latter two studies (133–150 dB re 1  $\mu$ Pa<sup>2</sup>s), the frequency, duration, and temporal pattern of signal presentation were different.

Humpback whales in another behavioral response experiment in Australia also responded to a 2 kHz tone stimulus by changing their course during migration to move more offshore and surfaced more frequently, but otherwise did not respond (Dunlop et al., 2013). Humpback whales in the Norwegian

behavioral response study may have habituated slightly between the first and second sonar exposure (Sivle et al., 2015), and actually responded more severely to killer whale vocalization playbacks than they did to the sonar playbacks. Several humpback whales have been observed during aerial or visual surveys during Navy training events involving sonar; no avoidance or other behavioral responses were ever noted, even when the whales were observed within 5 km of a vessel with active (or possibly active) sonar and maximum received levels were estimated to be between 135 and 161 dB re 1  $\mu$ Pa (Mobley & Millette, 2010; Mobley, 2011; Mobley & Pacini, 2012; Mobley et al., 2012; Smultea et al., 2009). In fact, one group of humpback whales approached a vessel with active sonar so closely that the sonar was shut down and the vessel slowed; the animals continued approaching and swam under the bow of the vessel (U.S. Department of the Navy, 2011b). Another group of humpback whales continued heading towards a vessel with active sonar as the vessel was moving away for almost 30 minutes, with an estimated median received level of 143 dB re 1  $\mu$ Pa. This group was observed producing surface active behaviors such as pec slaps, tail slaps, and breaches; however, these are very common behaviors in competitive pods during the breeding season and were not considered to have occurred in response to the sonar (Mobley et al., 2012).

The strongest baleen whale response in any behavioral response study was observed in a minke whale in the 3S2 study, which responded at 146 dB re 1  $\mu$ Pa by strongly avoiding the sound source (Kvadsheim et al., 2017; Sivle et al., 2015). Although the minke whale increased its swim speed, directional movement, and respiration rate, none of these were greater than rates observed in baseline behavior, and its dive behavior remained similar to baseline dives. A minke whale tagged in the Southern California behavioral response study also responded by increasing its directional movement, but maintained its speed and dive patterns, and so did not demonstrate as strong of a response (Kvadsheim et al., 2017). In addition, the 3S2 minke whale demonstrated some of the same avoidance behavior during the controlled ship approach with no sonar, indicating at least some of the response was to the vessel (Kvadsheim et al., 2017). Martin et al. (2015) found that the density of calling minke whales was reduced during periods of Navy training involving sonar relative to the periods before training, and increased again in the days after training was completed. The responses of individual whales could not be assessed, so in this case it is unknown whether the decrease in calling animals indicated that the animals left the range, or simply ceased calling. Similarly, minke whale detections made using Marine Acoustic Recording Instruments off Jacksonville, FL, were reduced or ceased altogether during periods of sonar use (Simeone et al., 2015; U.S. Department of the Navy, 2013b), especially with an increased ping rate (Charif et al., 2015). Two minke whales also stranded in shallow water after the U.S. Navy training event in the Bahamas in 2000, although these animals were successfully returned to deep water with no physical examinations; therefore, no final conclusions were drawn on whether the sonar led to their stranding (Filadelfo et al., 2009a; Filadelfo et al., 2009b; U.S. Department of Commerce & U.S. Department of the Navy, 2001).

Baleen whales have also been exposed to lower frequency sonars, with the hypothesis that these whales may react more strongly to lower frequency sounds that overlap with their vocalization range. One series of studies was undertaken in 1997–1998 pursuant to the Navy’s Low-Frequency Sound Scientific Research Program. The frequency bands of the low-frequency sonars used were between 100 and 500 Hz, with received levels between 115 and 150 dB re 1  $\mu$ Pa, and the source was always stationary. Fin and blue whales were targeted on foraging grounds, singing humpback whales were exposed on breeding grounds, and gray whales were exposed during migratory behavior. These studies found only short-term responses to low-frequency sound by some fin and humpback whales, including changes in vocal activity and avoidance of the source vessel, while other fin, humpback, and blue whales did not

respond at all. When the source was in the path of migrating gray whales they changed course up to 2 km to avoid the sound, but when the source was outside their path, little response was observed, although received levels were similar (Clark & Fristrup, 2001; Croll et al., 2001; Fristrup et al., 2003; Miller et al., 2000; Nowacek et al., 2007a). Low-frequency signals of the Acoustic Thermometry of Ocean Climate sound source were also not found to affect dive times of humpback whales in Hawaiian waters (Frankel & Clark, 2000).

Opportunistic passive acoustic based studies have also detected behavioral responses to sonar, although definitive conclusions are harder to draw. Blue whales exposed to mid-frequency sonar in the Southern California Bight were less likely to produce low-frequency calls usually associated with feeding behavior, beginning at received levels of 110–120 dB re 1  $\mu$ Pa (Melcón et al., 2012); however, without visual observations it is unknown whether there was another factor that contributed to the reduction in foraging calls, such as the presence of conspecifics. In another example, Risch et al. (2012, 2014) determined that humpback whale song produced in the Stellwagen Bank National Marine Sanctuary was reduced, and since the timing was concurrent with an Ocean Acoustic Waveguide Remote Sensing experiment occurring 200 km away, they concluded that the reduced song was a result of the Ocean Acoustic Waveguide Remote Sensing. However, Gong et al. (2014) analyzed the same data set while also looking at the presence of herring in the region, and found that the singing humpbacks were actually located on nearby Georges Bank and not on Stellwagen, and that the song rate in their data did not change in response to Ocean Acoustic Waveguide Remote Sensing, but could be explained by natural causes.

Although some strong responses have been observed in mysticetes to sonar and other transducers (e.g., the single minke whale), for the most part mysticete responses appear to be fairly moderate across all received levels. While some responses such as cessation of foraging or changes in dive behavior could carry short-term impacts, in all cases behavior returned to normal after the signal stopped. Mysticete responses also seem to be highly mediated by behavioral state, with no responses occurring in some behavioral states, and contextual factors and signal characteristics having more impact than received level alone. Many of the contextual factors resulting from the behavioral response studies (e.g., close approaches by multiple vessels or tagging) would never be introduced in real Navy training and testing scenarios. While data are lacking on behavioral responses of mysticetes to continuously active sonars, these species are known to be able to habituate to novel and continuous sounds (Nowacek et al., 2004), suggesting that they are likely to have similar responses to high-duty cycle sonars. Therefore, mysticete behavioral responses to Navy sonar would likely be a result of the animal's behavioral state and prior experience rather than external variables such as ship proximity; thus, if significant behavioral responses occur they would likely be short term. In fact, no significant behavioral responses such as panic, stranding, or other severe reactions have been observed during monitoring of actual training exercises (Smultea et al., 2009; U.S. Department of the Navy, 2011a, 2014b; Watwood et al., 2012).

#### ***Odontocetes***

Behavioral response studies have been conducted on odontocete species since 2007, with a focus on beaked whale responses to active sonar transmissions or controlled exposure playback of simulated sonar on various military ranges (Claridge et al., 2009; Defence Science and Technology Laboratory, 2007; Falcone et al., 2017; Henderson et al., 2015b; Henderson et al., 2016; Manzano-Roth et al., 2016; McCarthy et al., 2011; Moretti et al., 2009; Southall et al., 2011; Southall et al., 2012; Southall et al., 2013; Southall et al., 2014; Southall et al., 2015; Tyack et al., 2011). Through analyses of these behavioral response studies, a preliminary overarching effect of greater sensitivity to most

anthropogenic exposures was seen in beaked whales compared to the other odontocetes studied (Southall et al., 2009).

Observed reactions by Blainville's, Cuvier's, and Baird's beaked whales to mid-frequency sonar sounds have included cessation of clicking, termination of foraging dives, changes in direction to avoid the sound source, slower ascent rates to the surface, longer deep and shallow dive durations, and other unusual dive behavior (Boyd et al., 2008; Defence Science and Technology Laboratory, 2007; DeRuiter et al., 2013b; Miller et al., 2015; Southall et al., 2011; Stimpert et al., 2014; Tyack et al., 2011). A similar response was observed in a northern bottlenose whale, which conducted the longest and deepest dive on record for that species after the sonar exposure and continued swimming away from the source for over seven hours (Miller et al., 2015). Responses occurred at received levels between 95 and 150 dB re 1  $\mu$ Pa, although all of these exposures occurred within 1–8 km of the focal animal, within a few hours of tagging the animal, and with one or more boats within a few kilometers to observe responses and record acoustic data. One Cuvier's beaked whale was also incidentally exposed to real Navy sonar located over 100 km away, and the authors did not detect similar responses at comparable received levels. Received levels from the mid-frequency active sonar signals from the controlled and incidental exposures were calculated as 84–144 and 78–106 dB re 1  $\mu$ Pa, respectively, indicating that context of the exposures (e.g., source proximity, controlled source ramp-up) may have been a significant factor in the responses to the simulated sonars (DeRuiter et al., 2013b). Falcone et al. (2017) modeled deep and shallow dive durations, surface interval durations, and inter-deep dive intervals of Cuvier's beaked whales against predictor values that included helicopter-dipping, mid-power mid-frequency active sonar and hull-mounted, high-power mid-frequency active sonar along with other, non-mid-frequency active sonar predictors. They found both shallow and deep dive durations to increase as the proximity to both mid- and high-powered sources decreased, and found surface intervals and inter-deep dive intervals to also increase in the presence of both types of sonars, although surface intervals shortened during periods of no mid-frequency active sonar. The responses to the mid-power mid-frequency active sonar at closer ranges were comparable to the responses to the higher SL ship sonar, again highlighting the importance of proximity. This study also supports context as a response factor, as helicopter-dipping sonars are shorter duration and randomly located, so more difficult for beaked whales to predict or track and therefore potentially more likely to cause a response, especially when they occur at closer distances (6–25 km in this study). Watwood et al. (2017) found that helicopter-dipping events occurred more frequently but with shorter durations than periods of hull-mounted sonar, and also found that the longer the duration of a sonar event, the greater reduction in detected Cuvier's beaked whale group dives. Therefore, when looking at the number of detected group dives, there was a greater reduction during periods of hull-mounted sonar than during helicopter-dipping sonar. Long-term tagging work has demonstrated that the longer duration dives considered a behavioral response by DeRuiter et al. (2013b) fell within the normal range of dive durations found for eight tagged Cuvier's beaked whales on the Southern California Offshore Range (Schorr et al., 2014). However, the longer inter-deep dive intervals found by DeRuiter et al. (2013b), which were among the longest found by Schorr et al. (2014) and Falcone et al. (2017), could indicate a response to sonar. In addition, Williams et al. (2017) note that in normal deep dives or during fast swim speeds, beaked whales and other marine mammals use strategies to reduce their stroke rates, including leaping or wave surfing when swimming, and interspersing glides between bouts of stroking when diving. They determined that in the post-exposure dives by the tagged Cuvier's beaked whales described in DeRuiter et al. (2013b), the whales ceased gliding and swam with almost continuous strokes. This change in swim behavior was calculated to increase metabolic costs about 30.5 percent and increase the amount of energy expending on fast swim

speeds from 27 to 59 percent of their overall energy budget. This repartitioning of energy was detected in the model up to 1.7 hours after the single sonar exposure. Therefore, while the overall post-exposure dive durations were similar, the metabolic energy calculated by Williams et al. (2017) was higher.

On Navy ranges, Blainville's beaked whales located on the range appear to move off-range during sonar use and return only after the sonar transmissions have stopped, sometimes taking several days to do so (Claridge et al., 2009; Henderson et al., 2015b; Manzano-Roth et al., 2016; McCarthy et al., 2011; Moretti et al., 2009; Tyack et al., 2011). However, Blainville's beaked whales remain on the range to forage throughout the rest of the year (Henderson et al., 2016), possibly indicating that this a preferred foraging habitat regardless of the effects of the noise, or it could be that there are no long-term consequences of the sonar activity. Similarly, photo-identification studies in the Southern California Range Complex have identified approximately 100 individual Cuvier's beaked whale individuals, with 40 percent having been seen in one or more prior years, with re-sightings up to seven years apart, indicating a possibly resident population on the range (Falcone et al., 2009; Falcone & Schorr, 2014).

Beaked whales may respond similarly to shipboard echosounders, commonly used for navigation, fisheries, and scientific purposes, with frequencies ranging from 12 to 400 kHz and source levels up to 230 dB re 1  $\mu$ Pa but typically a very narrow beam (Cholewiak et al., 2017). During a scientific cetacean survey, an array of echosounders was used in a one-day-on, one-day-off paradigm. Beaked whale acoustic detections occurred predominantly (96 percent) when the echosounder was off, with only four detections occurring when it was on. Beaked whales were sighted fairly equally when the echosounder was on or off, but sightings were further from the ship when the echosounder was on (Cholewiak et al., 2017). These findings indicate that the beaked whales may be avoiding the area and may cease foraging near the echosounder.

Tyack et al. (2011) hypothesized that beaked whale responses to sonar may represent an anti-predator response. To test this idea, vocalizations of a potential predator—a killer whale—were also played back to a Blainville's beaked whale. This exposure resulted in a similar but more pronounced reaction than that elicited by sonar playback, which included longer inter-dive intervals and a sustained straight-line departure of more than 20 km from the area (Allen et al., 2014; Tyack et al., 2011). This anti-predator hypothesis was also tested by playing back killer whale vocalizations to pilot whales, sperm whales, and even other killer whales, to determine responses by both potential prey and conspecifics (Miller et al., 2011; Miller, 2012). Results varied, from no response by killer whales to an increase in group size and attraction to the source in pilot whales (Curé et al., 2012).

While there has been a focus on beaked whale responses to sonar, other species have been studied during behavioral response studies as well, including pilot whales, killer whales, and sperm whales. Responses by these species have also included horizontal avoidance, changes in behavioral state, and changes in dive behavior (Antunes et al., 2014; Miller et al., 2011; Miller, 2012; Miller et al., 2014). Additionally, separation of a killer whale calf from its group during exposure to mid-frequency sonar playback was observed (Miller et al., 2011). Received level thresholds at the onset of avoidance behavior were generally higher for pilot whales (mean 150 dB re 1  $\mu$ Pa) and sperm whales (mean 140 dB re 1  $\mu$ Pa) than killer whales (mean 129 dB re 1  $\mu$ Pa) (Antunes et al., 2014; Miller, 2012; Miller et al., 2014). A close examination of the tag data from the Norwegian groups showed that responses seemed to be behaviorally or signal frequency mediated. For example, killer whales only changed their dive behavior when doing deep dives at the onset of 1–2 kHz sonar (sweeping across frequencies), but did not change their dive behavior if they were deep diving during 6–7 kHz sonar (sweeping across frequencies). Nor did they change their dive behavior if they were conducting shallow dives at the onset of either type of

sonar. Similarly, pilot whales and sperm whales performed normal deep dives during 6–7 kHz sonar, while during 1–2 kHz sonar the pilot whales conducted fewer deep dives and the sperm whales performed shorter and shallower dives (Sivle et al., 2012). In addition, pilot whales were also more likely to respond to lower received levels when non-feeding than feeding during 6–7 kHz sonar exposures, but were more likely to respond at higher received levels when non-feeding during 1–2 kHz sonar exposures. Furthermore, pilot whales exposed to a 38 kHz downward-facing echosounder did not change their dive and foraging behavior during exposure periods, although the animals' heading variance increased and fewer deep dives were conducted (Quick et al., 2017). In contrast, killer whales were more likely to respond to either sonar type when non-feeding than when feeding (Harris et al., 2015). These results again demonstrate that the behavioral state of the animal mediates the likelihood of a behavioral response, as do the characteristics (e.g., frequency) of the sound source itself.

Other responses during behavioral response studies included the synchronization of pilot whale surfacings with sonar pulses during one exposure, possibly as a means of mitigating the sound (Wensveen et al., 2015), and mimicry of the sonar with whistles by pilot whales (Alves et al., 2014), false killer whales (DeRuiter et al., 2013b) and Risso's dolphins (Smultea et al., 2012). In contrast, in another study melon-headed whales had "minor transient silencing" (a brief, non-lasting period of silence) after each 6–7 kHz signal, and (in a different oceanographic region) pilot whales had no apparent response (DeRuiter et al., 2013a). The probability of detecting delphinid vocalizations (whistles, clicks, and buzzes) increased during periods of sonar relative to the period prior to sonar in a passive acoustic study using Marine Autonomous Recording Units in the Jacksonville Range Complex, while there was no impact of sonar to the probability of detecting sperm whale clicks (Charif et al., 2015; U.S. Department of the Navy, 2013c).

In addition, killer whale sighting data from the same region in Norway as the behavioral response study was used to compare the presence or absence of whales from other years against the period with sonar. The authors found a strong relationship between the presence of whales and the abundance of herring, and only a weak relationship between the whales and sonar activity (Kuningas et al., 2013). Baird et al. (2013b; 2014; 2017) also tagged four shallow-diving odontocete species (rough-toothed dolphins, pilot whales, bottlenose dolphins, and false killer whales) in Hawaii off the Pacific Missile Range Facility before Navy training events. None of the tagged animals demonstrated a large-scale avoidance response to the sonar as they moved on or near the range, in some cases even traveling towards areas of higher noise levels, while estimated received SPLs varied from 130 to 168 dB re 1  $\mu$ Pa and distances from sonar sources ranged between 3.2 and 94.4 km. However, one pilot whale did have reduced dive rates (from 2.6 dives per hour before to 1.6 dives per hour during) and deeper dives (from a mean of 124 m to 268 m) during a period of sonar exposure. Baird et al. (2016) also tagged four short-finned pilot whales from both the resident island-associated population and from the pelagic population. The core range for the pelagic population was over 20 times larger than for the resident population, leading Baird et al. (2016) to hypothesize that that likelihood of exposure to mid-frequency active sonar, and therefore the potential for response, would be very different between the two populations. These diverse examples demonstrate that responses can be varied, are often context- and behavior-driven, and can be species and even exposure specific.

Other opportunistic observations of behavioral responses to sonar have occurred as well, although in those cases it is difficult to attribute observed responses directly to the sonar exposure, or to know exactly what form the response took. For example, both sperm and pilot whales potentially ceased sound production during the Heard Island feasibility test, with transmissions centered at 57 Hz and up to

220 dB re 1  $\mu$ Pa (Bowles et al., 1994), although it could not be determined whether the animals ceased sound production or left the area. In May 2003, killer whales in Haro Strait, Washington, exhibited what were believed by some observers to be aberrant behaviors, during which time the USS Shoup was in the vicinity and engaged in mid-frequency active sonar operations. Sound fields modeled for the USS Shoup transmissions (Fromm, 2009; National Marine Fisheries Service, 2005; U.S. Department of the Navy, 2003) estimated a mean received SPL of approximately 169 dB re 1  $\mu$ Pa at the location of the killer whales at the closest point of approach between the animals and the vessel (estimated SPLs ranged from 150 to 180 dB re 1  $\mu$ Pa). However, attributing the observed behaviors to any one cause is problematic given there were six nearby whale watch vessels surrounding the pod, and subsequent research has demonstrated that “Southern Residents modify their behavior by increasing surface activity (breaches, tail slaps, and pectoral fin slaps) and swimming in more erratic paths when vessels are close” (National Oceanic and Atmospheric Administration Fisheries, 2014). Several odontocete species, including bottlenose dolphins, Risso’s dolphins, Pacific white-sided dolphins, and common dolphins have been observed near the Southern California Offshore Range during periods of mid-frequency active sonar; responses included changes in or cessation of vocalizations, changes in behavior, and departures from the area, and at the highest received levels animals were not present in the area at all (Henderson et al., 2014). However, these observations were conducted from a vessel off-range, and so any observed responses could not be attributed to the sonar with any certainty. Research on sperm whales in the Caribbean in 1983 coincided with the U.S. intervention in Grenada, where animals were observed scattering and leaving the area in the presence of military sonar, presumably from nearby submarines (Watkins & Schevill, 1975; Watkins et al., 1985). The authors did not report received levels from these exposures and reported similar reactions from noise generated by banging on their boat hull; therefore, it was unclear if the sperm whales were reacting to the sonar signal itself or to a potentially new unknown sound in general.

During aerial and visual monitoring of Navy training events involving sonar, rough-toothed dolphins and unidentified dolphins were observed approaching the vessel with active sonar as if to bow ride, while spotted dolphins were observed nearby but did not avoid or approach the vessel (Mobley, 2011; U.S. Department of the Navy, 2011b; Watwood et al., 2012). During small boat surveys near the Southern California Offshore Range in southern California, more dolphins were encountered in June compared to a similar survey conducted the previous November after seven days of mid-frequency sonar activity; it was not investigated if this change was due to the sonar activity or was a seasonal difference that was also observed in other years (Campbell et al., 2010). There were also fewer passive acoustic dolphin detections during and after longer sonar activities in the Mariana Islands Range Complex, with the post-activity absence lasting longer than the mean dolphin absence of two days when sonar was not present (Munger et al., 2014; Munger et al., 2015).

Acoustic harassment devices and acoustic deterrent devices, that transmit sound into the acoustic environment similar to Navy sonar, have been used to deter marine mammals from fishing gear both to prevent entanglement and to reduce depredation (taking fish). These devices have been used successfully to deter harbor porpoises and beaked whales from getting entangled in fishing nets. For example, Kyhn et al. (2015) tested two types of pingers, one with a 10 kHz tone and one with a broadband 30–160 kHz sweep. Porpoise detection rates were reduced by 65 percent for the sweep and 40 percent for the tone, and while there was some gradual habituation after the first two to four exposures, longer-term exposures (over 28 days) showed no evidence of additional habituation. Additionally, sperm whales in the Caribbean stopped vocalizing when presented with sounds from nearby acoustic pingers (Watkins & Schevill, 1975). However, acoustic harassment devices used to deter

marine mammals from depredating long lines or aquaculture enclosures have proven less successful. For example, Tixier et al. (2014) used a 6.5 kHz pinger with a source level of 195 dB re 1  $\mu$ Pa on a longline to prevent depredation by killer whales, and although two groups of killer whales fled over 700 m away during the first exposure, they began depredating again after the third and seventh exposures, indicating rapid habituation. In a review of marine mammal deterrents, Schakner & Blumstein (2013) point out that both the characteristics of deterrents and the motivation of the animal play a role in the effectiveness of acoustic harassment devices. Deterrents that are strongly aversive or simulate a predator or are otherwise predictive of a threat are more likely to be effective, unless the animal habituates to the signal or learns that there is no true threat associated with the signal. In some cases net pingers may create a “dinner bell effect,” where marine mammals have learned to associate the signal with the availability of prey (Jefferson & Curry, 1996; Schakner & Blumstein, 2013). This may be why net pingers have been more successful at reducing entanglements for harbor porpoise and beaked whales since these species are not depredating from the nets but are getting entangled when foraging in the area and are unable to detect the net (Carretta et al., 2008; Schakner & Blumstein, 2013). Similarly, a 12 kHz acoustic harassment device intended to scare seals was ineffective at deterring seals but effectively caused avoidance in harbor porpoises out to over 500 m from the source, highlighting different species- and device-specific responses (Mikkelsen et al., 2017). Additional behavioral studies have been conducted with captive harbor porpoises using acoustic alarms, such as those used on fishing nets to help deter marine mammals from becoming caught or entangled (Kastelein et al., 2006; Kastelein et al., 2001). These studies have found that high-frequency sources with varied duration, interval, and sweep characteristics can prove to be effective deterrents for harbor porpoises (Kastelein et al., 2017). Van Beest et al. (2017) modeled the long-term, population-level impacts of fisheries bycatch, pinger deterrents, and time-area closures on a population of harbor porpoises. They found that when pingers were used alone (in the absence of gillnets or time-area closures), the animals were deterred from the area often enough to cause a population level reduction of 21 percent, greater even than the modeled level of current bycatch impacts. However, when the pingers were coupled with gillnets in the model, and time-area closures were also used (allowing a net- and pinger-free area for the porpoises to move into while foraging), the population only experienced a 0.8 percent decline even with current gillnet use levels. This demonstrates that, when used correctly, pingers can successfully deter porpoises from gillnets without leading to any negative impacts.

Controlled experiments have also been conducted on captive animals to estimate received levels at which behavioral responses occur. In one study, bottlenose dolphin behavioral responses were recorded when exposed to 3 kHz sonar-like tones between 115 and 185 dB re 1  $\mu$ Pa (Houser et al., 2013), and in another study bottlenose dolphins and beluga whales were presented with one-second tones up to 203 dB re 1  $\mu$ Pa to measure TTS (Finneran et al., 2001; Finneran et al., 2003a; Finneran & Schlundt, 2004; Finneran et al., 2005b; Schlundt et al., 2000). During these studies, responses included changes in respiration rate, fluke slaps, and a refusal to participate or return to the location of the sound stimulus. This refusal included what appeared to be deliberate attempts to avoid a sound exposure or to avoid the location of the exposure site during subsequent tests (Finneran et al., 2002; Schlundt et al., 2000). In the behavioral response experiment, bottlenose dolphins demonstrated a 50 percent probability of response at 172 dB re 1  $\mu$ Pa over 10 trials, and in the TTS study bottlenose dolphins exposed to one-second intense tones exhibited short-term changes in behavior above received sound levels of 178–193 dB re 1  $\mu$ Pa, while beluga whales did so at received levels of 180–196 dB re 1  $\mu$ Pa and above. In some instances, animals exhibited aggressive behavior toward the test apparatus (Ridgway et al., 1997; Schlundt et al., 2000). While animals were commonly reinforced with food during these studies, the

controlled environment and ability to measure received levels provide insight on received levels at which animals would behaviorally respond to noise sources.

Behavioral responses to a variety of sound sources have been studied in captive harbor porpoises, including acoustic alarms (Kastelein et al., 2006; Kastelein et al., 2001); emissions for underwater data transmission (Kastelein et al., 2005); and tones, including 1–2 kHz and 6–7 kHz sweeps with and without harmonics (Kastelein et al., 2014d), 25 kHz with and without sidebands (Kastelein et al., 2015d; Kastelein et al., 2015e), and mid-frequency sonar tones at 3.5–4.1 kHz at 2.7 percent and 96 percent duty cycles (e.g., one tone per minute versus a continuous tone for almost a minute) (Kastelein et al., 2018). Responses include increased respiration rates, more jumping, or swimming further from the source, but responses were different depending on the source. For example, harbor porpoises responded to the 1–2 kHz upsweep at 123 dB re 1  $\mu$ Pa, but not to the downsweep or the 6–7 kHz tonal at the same level (Kastelein et al., 2014d). When measuring the same sweeps for a startle response, the 50 percent response threshold was 133 and 101 dB re 1  $\mu$ Pa for 1–2 kHz and 6–7 kHz sweeps, respectively, when no harmonics were present, and decreased to 90 dB re 1  $\mu$ Pa for 1–2 kHz sweeps with harmonics present (Kastelein et al., 2014d). Harbor porpoises did not respond to the low-duty cycle mid-frequency tones at any received level, but one did respond to the high-duty cycle signal with more jumping and increased respiration rates (Kastelein et al., 2018). Harbor porpoises responded to seal scarers with broadband signals up to 44 kHz with a slight respiration response at 117 dB re 1  $\mu$ Pa and an avoidance response at 139 dB re 1  $\mu$ Pa, but another scarer with a fundamental (strongest) frequency of 18 kHz did not have an avoidance response until 151 dB re 1  $\mu$ Pa (Kastelein et al., 2014a). Exposure of the same acoustic pinger to a striped dolphin under the same conditions did not elicit a response (Kastelein et al., 2006), again highlighting the importance in understanding species differences in the tolerance of underwater noise, although sample sizes in these studies was small so these could reflect individual differences as well.

Behavioral responses by odontocetes to sonar and other transducers appear to range from no response at all to responses that could potentially lead to long-term consequences for individual animals (e.g., mother-calf separation). This is likely in part due to the fact that this taxonomic group is so broad and includes some of the most sensitive species (e.g., beaked whales and harbor porpoise) as well as some of the least sensitive species (e.g., bottlenose dolphins). This is also the only group for which both field behavioral response studies and captive controlled exposure experiments have been conducted, leading to the assessment of both contextually driven responses as well as dose-based responses. This wide range in both exposure situations and individual- and species-sensitivities makes reaching general conclusions difficult. However, it does appear as though exposures in close proximity, with multiple vessels that approach the animal, lead to higher-level responses in most odontocete species regardless of received level or behavioral state. In contrast, in more “real-world” exposure situations, with distant sources moving in variable directions, behavioral responses appear to be driven by behavioral state, individual experience, or species-level sensitivities. These responses may also occur more in line with received level such that the likelihood of a response would increase with increased received levels. However, these “real-world” responses are more likely to be short-term, lasting the duration of the exposure or even shorter as the animal assesses the sound and (based on prior experience or contextual cues) determines a threat is unlikely. Therefore, while odontocete behavioral responses to Navy sonar would vary across species, populations, and individuals, they are not likely to lead to long-term consequences or population-level effects.

## Behavioral Reactions to Vessels

Sound emitted from large vessels, such as cargo ships, is the principal source of low-frequency noise in the ocean today, and marine mammals are known to react to or be affected by that noise (Hatch & Wright, 2007; Hildebrand, 2005; Richardson et al., 1995b). For example, Erbe et al. (2012) estimated the maximum annual underwater SEL from vessel traffic near Seattle was 215 dB re 1  $\mu\text{Pa}^2\text{-s}$ , and Bassett et al. (2010) measured mean SPLs at Admiralty Inlet from commercial shipping at 117 dB re 1  $\mu\text{Pa}$  with a maximum exceeding 135 dB re 1  $\mu\text{Pa}$  on some occasions. Similarly, Veirs et al. (2015) found average broadband noise levels in Haro Strait to be 110 dB re 1  $\mu\text{Pa}$  that extended up to 40 kHz, well into the hearing range of odontocetes.

Many studies of behavioral responses by marine mammals to vessels have been focused on the short- and long-term impacts of whale watching vessels. In short-term studies, researchers noted changes in resting and surface behavior states of cetaceans to whale-watching vessels (Acevedo, 1991; Aguilar de Soto et al., 2006; Arcangeli & Crosti, 2009; Au & Green, 2000; Christiansen et al., 2010; Erbe, 2002; Noren et al., 2009; Stockin et al., 2008; Williams et al., 2009). Received levels were often not reported, so it is difficult to distinguish responses to the presence of the vessel from responses to the vessel noise. Most studies examined the short-term response to vessel sound and vessel traffic (Magalhães et al., 2002; Richardson et al., 1995b; Watkins, 1981), with behavioral and vocal responses occurring when received levels were over 20 dB greater than ambient noise levels. Other research has attempted to quantify the effects of whale watching using focused experiments (Meissner et al., 2015; Pirota et al., 2015b).

The impact of vessel noise has received increased consideration, particularly as whale watching and shipping traffic has risen (McKenna et al., 2012; Pirota et al., 2015b; Veirs et al., 2015). Odontocetes and mysticetes in particular have received increased attention relative to vessel noise and vessel traffic, with pinnipeds less so. Still, not all species in all taxonomic groups have been studied, and so results do have to be extrapolated across these broad categories in order to assess potential impacts.

### *Mysticetes*

Baleen whales demonstrate a variety of responses to vessel traffic and noise, from not responding at all to both horizontal (swimming away) and vertical (increased diving) avoidance (Baker et al., 1983; Gende et al., 2011; Watkins, 1981). Other common responses include changes in vocalizations, surface time, swimming speed, swimming angle or direction, respiration rates, dive times, feeding behavior, and social interactions (Au & Green, 2000; Richter et al., 2003; Williams et al., 2002a).

The likelihood of response may be driven by the distance or speed of the vessel, the animal's behavioral state, or by the prior experience of the individual or population. For example, in one study fin and humpback whales largely ignored vessels that remained 100 m or more away (Watkins, 1981). In another study, minke whales in the Antarctic did not show any apparent response to a survey vessel moving at normal cruising speeds (about 12 knots) at a distance of 5.5 NM. However, when the vessel drifted or moved at very slow speeds (about 1 knot), many whales approached it (Leatherwood et al., 1982). Similarly, Bernasconi et al. (2012) observed the reactions of six individual baleen whales of unknown species at distances of 50–400 m from a fishing vessel conducting an acoustic survey of pelagic fisheries, with only a slight change in swim direction when the vessel began moving around the whales. Gray whales were likely to continue feeding when approached by a vessel in areas with high motorized vessel traffic, but in areas with less motorized vessel traffic they were more likely to change behaviors, either indicating habituation to vessels in high traffic area, or indicating possible startle reactions to

close-approaching non-motorized vessels (e.g., kayaks) in quieter areas (Sullivan & Torres, 2018). Changes in behavior of humpback whales when vessels came within 500 m were also dependent on behavioral state such that they would keep feeding but were more likely to start traveling if they were surface active when approached; and changes in behavior were also affected by time of day or season (Di Clemente et al., 2018). Sei whales have been observed ignoring the presence of vessels entirely and even passing close to the vessel (Reeves et al., 1998), and North Atlantic right whales tend not to respond to the sounds of oncoming vessels and continue to use habitats in high vessel traffic areas (Nowacek et al., 2004). Studies show that North Atlantic right whales demonstrate little if any reaction to sounds of vessels approaching or the presence of the vessels themselves. This lack of response may be due to habituation to the presence and associated noise of vessels in right whale habitat, or may be due to propagation effects that may attenuate vessel noise near the surface (Nowacek et al., 2004; Terhune & Verboom, 1999).

When baleen whales do respond to vessels, responses can be as minor as a change in breathing patterns (e.g., Baker et al., 1983; Jahoda et al., 2003), or can be evidenced by a decrease in overall presence, as was observed during a construction project in the United Kingdom, when fewer minke whales were observed as vessel traffic increased (Anderwald et al., 2013). Avoidance responses can be as simple as an alteration in swim patterns or direction by increasing speed and heading away from the vessel (Jahoda et al., 2003), or by increasing swim speed, changing direction to avoid, and staying submerged for longer periods of time (Au & Green, 2000). For example, in the presence of approaching vessels, blue whales perform shallower dives accompanied by more frequent surfacing but otherwise do not exhibit strong reactions (Calambokidis et al., 2009b). In another study in Hawaii, humpback whales exhibited two forms of behavioral avoidance: horizontal avoidance (changing direction or speed) when vessels were between 2,000 m and 4,000 m away, and vertical avoidance (increased dive times and change in diving pattern) when vessels were less than 2,000 m away (Baker et al., 1983). Similarly, humpback whales in Australia demonstrated variable responses to whale watching vessels, including both horizontal avoidance, approaching, and changes in dive and surface behavior (Stamation et al., 2009). Humpback whales avoided a Navy vessel by increasing their dive times and decreasing respiration rates at the surface (Smultea et al., 2009). Williamson et al. (2016) specifically looked at close approaches to humpback whales by small research boats for the purposes of tagging. They found that while dive behavior did not change for any groups, some groups did increase their speed and change their course during or right after the approach, but resumed pre-approach speed and heading shortly thereafter. Only mother-calf groups were found to increase their speed during the approach and maintain the increased speed for longer after the approach, but these groups too resumed normal swim speeds after about 40 minutes. It should be noted that there were no responses by any groups that were approached closely but with no attempts at tagging, indicating that the responses were not due to the vessel presence but to the tagging attempt. In addition, none of the observed changes in behavior were outside the normal range of swim speeds or headings for these migrating whales.

Mysticetes have been shown to both increase and decrease calling behavior in the presence of vessel noise. Based on passive acoustic recordings and in the presence of sounds from passing vessels, Melcón et al. (2012) reported that blue whales had an increased likelihood of producing certain types of calls. An increase in feeding call rates and repetition by humpback whales in Alaskan waters is associated with vessel noise (Doyle et al., 2008), while decreases in singing activity have been noted near Brazil due to boat traffic (Sousa-Lima & Clark, 2008). Frequency parameters of fin whale calls also decreased in the presence of increasing background noise due to shipping traffic (Castellote et al., 2012). Bowhead whales avoided the area around icebreaker ship noise and increased their time at the surface and

number of blows (Richardson et al., 1995a). Right whales increase the amplitude or frequency of their vocalizations or call at a lower rate in the presence of increased vessel noise (Parks et al., 2007; Parks et al., 2011), and these vocalization changes may persist over long periods if background noise levels remained elevated.

The long-term consequences of vessel noise are not well understood (see Section 3.4.2.1.1.7, Long-Term Consequences). In a short-term study, minke whales on feeding grounds in Iceland responded to increased whale-watching vessel traffic with a decrease in foraging, both during deep dives and at the surface (Christiansen et al., 2013). They also increased their avoidance of the boats while decreasing their respiration rates, likely leading to an increase in their metabolic rates. Christiansen and Lusseau (2015) and Christiansen et al. (2014) followed up this study by modeling the cumulative impacts of whale-watching boats on minke whales, but found that although the boats cause temporary feeding disruptions, there were not likely to be long-term consequences as a result. This suggests that short-term responses may not lead to long-term consequences and that over time animals may habituate to the presence of vessel traffic. However, in an area of high whale-watch activity, vessels were within 2,000 m of blue whales 70 percent of the time, with a maximum of eight vessels observed within 400 m of one whale at the same time. This study found reduced surface time, fewer breaths at the surface, and shorter dive times when vessels were within 400 m (Lesage et al., 2017). Since blue whales in this area forage 68 percent of the time, and their foraging dive depths are constrained by the location of prey patches, these reduced dive durations may indicate reduced time spent foraging by over 36 percent. In the short term this reduction may be compensated for, but prolonged exposure to vessel traffic could lead to long-term consequences. Using historical records, Watkins (1986) showed that the reactions of four species of mysticetes to vessel traffic and whale-watching activities in Cape Cod had changed over the 25-year period examined (1957–1982). Reactions of minke whales changed from initially more positive reactions, such as coming towards the boat or research equipment to investigate, to more uninterested reactions towards the end of the study. Fin whales, the most numerous species in the area, showed a trend from initially more negative reactions, such as swimming away from the boat with limited surfacing, to more uninterested reactions (ignoring) allowing boats to approach within 30 m. Right whales showed little change over the study period, with a roughly equal number of reactions judged to be negative and uninterested; no right whales were noted as having positive reactions to vessels. Humpback whales showed a trend from negative to positive reactions with vessels during the study period. The author concluded that the whales had habituated to the human activities over time (Watkins, 1986).

Overall baleen whale responses to vessel noise and traffic are varied but are generally minor, and habituation or disinterest seems to be the predominant long-term response. When baleen whales do avoid ships they do so by altering their swim and dive patterns to move away from the vessel, but no strong reactions have been observed. In fact, in many cases the whales do not appear to change their behavior at all. This may result from habituation by the whales, but may also result from reduced received levels near the surface due to propagation, or due to acoustic shadowing of the propeller cavitation noise by the ship's hull. Although a lack of response in the presence of a vessel may minimize potential disturbance from passing ships, it does increase the whales' vulnerability to vessel strike, which may be of greater concern for baleen whales than vessel noise (see Section 3.4.4.4, Physical Disturbance and Strike Stressors in the 2015 MITT Final EIS/OEIS).

### *Odontocetes*

Most odontocetes react neutrally to vessels, although both avoidance and attraction behavior have been observed (Hewitt, 1985; Würsig et al., 1998). Würsig et al. (1998) found that Kogia whales and beaked whales were the most sensitive species to vessels, and reacted by avoiding marine mammal survey vessels in 73 percent of sightings, more than any other odontocetes. Avoidance reactions include a decrease in resting behavior or change in travel direction (Bejder et al., 2006a). Incidents of attraction include common, rough-toothed, and bottlenose dolphins bow riding and jumping in the wake of a vessel (Norris & Prescott, 1961; Ritter, 2002; Shane et al., 1986; Würsig et al., 1998). A study of vessel reactions by dolphin communities in the eastern tropical Pacific found that populations that were often the target of tuna purse-seine fisheries (spotted, spinner, and common dolphins) show evasive behavior when approached; however, populations that live closer to shore (within 100 NM; coastal spotted and bottlenose dolphins) that are not set on by purse-seine fisheries tend to be attracted to vessels (Archer et al., 2010). The presence of vessels has also been shown to interrupt feeding behavior in delphinids (Meissner et al., 2015; Pirotta et al., 2015b).

Short-term displacement of dolphins due to tourist boat presence has been documented (Carrera et al., 2008), while longer-term or repetitive/chronic displacement for some dolphin groups due to chronic vessel noise has been noted (Haviland-Howell et al., 2007). Delphinid behavioral states also change in the presence of tourist boats that often approach animals, with travel increasing and foraging decreasing (Cecchetti et al., 2017; Meissner et al., 2015). Most studies of the behavioral reactions to vessel traffic of bottlenose dolphins have documented at least short-term changes in behavior, activities, or vocalization patterns when vessels are near, although the distinction between vessel noise and vessel movement has not been made clear (Acevedo, 1991; Arcangeli & Crosti, 2009; Berrow & Holmes, 1999; Gregory & Rowden, 2001; Janik & Thompson, 1996; Lusseau, 2004; Mattson et al., 2005; Scarpaci et al., 2000). Steckenreuter (2011) found bottlenose dolphin groups to feed less, become more tightly clustered, and have more directed movement when approached to 50 m than groups approached to 150 m or approached in a controlled manner. Guerra et al. (2014) demonstrated that bottlenose dolphins subjected to chronic noise from tour boats responded to boat noise by alterations in group structure and in vocal behavior but also found the dolphins' reactions varied depending on whether the observing research vessel was approaching or moving away from the animals being observed. This demonstrates that the influence of the sound exposure cannot be decoupled from the physical presence of a surface vessel, thus complicating interpretations of the relative contribution of each stimulus to the response. Indeed, the presence of surface vessels, their approach, and speed of approach, seemed to be significant factors in the response of the Indo-Pacific humpback dolphins (Ng & Leung, 2003).

The effects of tourism and whale watching have highly impacted killer whales, such as the Northern and Southern Resident populations. These animals are targeted by numerous small whale-watching vessels in the Pacific Northwest and, from 1998 to 2012 during the viewing season, have had an annual monthly average of nearly 20 vessels of various types within 0.5 mile of their location during daytime hours (Clark, 2015; Eisenhardt, 2014; Erbe et al., 2014). These vessels have source levels that ranged from 145 to 169 dB re 1  $\mu$ Pa and produce broadband noise up to 96 kHz and 116 dB re 1  $\mu$ Pa. While new regulations on the distance boats had to maintain were implemented, there did not seem to be a concurrent reduction in the received levels of vessel noise, and noise levels were found to increase with more vessels and faster-moving vessels (Holt et al., 2017). These noise levels have the potential to result in behavioral disturbance, interfere with communication, and affect the killer whales' hearing capabilities via masking (Erbe, 2002; Veirs et al., 2015). Killer whales foraged significantly less and

traveled significantly more when boats were within 100 m of the whales (Kruse, 1991; Lusseau et al., 2009; Trites & Bain, 2000; Williams et al., 2002a; Williams et al., 2002b; Williams et al., 2009). These short-term feeding activity disruptions may have important long-term population-level effects (Lusseau et al., 2009; Noren et al., 2009). As with other delphinids, the reaction of the killer whales to whale-watching vessels may be in response to the vessel pursuing them rather than to the noise of the vessel itself, or to the number of vessels in their proximity. Williams et al. (2013) modeled behavioral responses of killer whales to vessel traffic by looking at their surface behavior relative to the received level of three large classes of ships. The authors found that the severity of the response was largely dependent on seasonal data (e.g., year and month) as well as the animal's prior experience with vessels (e.g., age and sex), and the number of other vessels present, rather than the received level of the larger ships (Williams et al., 2013).

Sperm whales generally react only to vessels approaching within several hundred meters; however, some individuals may display avoidance behavior, such as quick diving (Magalhães et al., 2002; Würsig et al., 1998) or a decrease in time spent at the surface (Isojunno & Miller, 2015). One study showed that after diving, sperm whales showed a reduced timeframe before they emitted the first click than prior to a vessel interaction (Richter et al., 2006). Smaller whale-watching and research vessels generate more noise in higher frequency bands and are more likely to approach odontocetes directly, and to spend more time near an individual whale. Azzara et al. (2013) also found a reduction in sperm whale clicks while a vessel was passing, as well as up to a half hour after the vessel had passed. It is unknown whether the whales left the area, ceased to click, or surfaced during this period. However, some of the reduction in click detections may be due to masking of the clicks by the vessel noise, particularly during the closest point of approach.

Little information is available on the behavioral impacts of vessels or vessel noise on beaked whales (Cox et al., 2006), although it seems most beaked whales react negatively to vessels by quick diving and other avoidance maneuvers (Würsig et al., 1998). Limited evidence suggests that beaked whales respond to vessel noise, anthropogenic noise in general, and mid-frequency sonar at similar sound levels (Aguilar de Soto et al., 2006; Tyack et al., 2011; Tyack, 2009). An observation of vocal disruption of a foraging dive by a Cuvier's beaked whale when a large, noisy vessel passed suggests that some types of vessel traffic may disturb foraging beaked whales (Aguilar de Soto et al., 2006). Tyack et al. (2011) noted the result of a controlled exposure to pseudorandom noise suggests that beaked whales would respond to vessel noise at similar received levels to those noted previously for mid-frequency sonar. Pirotta et al. (2012) found that while the distance to a vessel did not change the duration of a foraging dive, the proximity of the vessel may have restricted the movement of the group. The maximum distance at which this change was significant was 5.2 km, with an estimated received level of 135 dB re 1  $\mu$ Pa.

Small dolphins and porpoises may also be more sensitive to vessel noise. Both finless porpoises (Li et al., 2008) and harbor porpoises (Polacheck & Thorpe, 1990) routinely avoid and swim away from large motorized vessels, and harbor porpoises may click less when near large ships (Sairanen, 2014). A resident population of harbor porpoise in Swansea Bay are regularly near vessel traffic, but only 2 percent of observed vessels had interactions with porpoises in one study (Oakley et al., 2017). Of these, 74 percent of the interactions were neutral (no response by the porpoises) while vessels were 10 m–1 km away. Of the 26 percent of interactions in which there was an avoidance response, most were observed in groups of 1–2 animals to fast-moving or steady plane-hulling motorized vessels. Larger groups reacted less often, and few responses were observed to non-motorized or stationary vessels. Another study found that when vessels were within 50 m, harbor porpoises had an 80 percent

probability of changing their swimming direction when vessels were fast moving; this dropped to 40 percent probability when vessels were beyond 400 m (Akkaya Bas et al., 2017). These porpoises also demonstrated a reduced proportion of feeding and shorter behavioral bout durations in general, if vessels were in close proximity, 62 percent of the time. Although most vessel noise is constrained to lower frequencies below 1 kHz, at close range vessel noise can extend into mid- and high-frequencies (into the tens of kHz) (Hermannsen et al., 2014; Li et al., 2015); these frequencies are what harbor porpoises are likely responding to, at M-weighted received SPLs with a mean of 123 dB re 1  $\mu$ Pa (Dyndo et al., 2015). Foraging harbor porpoises also have fewer prey capture attempts and have disrupted foraging when vessels pass closely and noise levels are higher (Wisniewska et al., 2018).

Odontocetes have been shown to make short-term changes to vocal parameters such as intensity as an immediate response to vessel noise, as well as to increase the pitch, frequency modulation, and length of whistling (May-Collado & Wartzok, 2008), with whistle frequency increasing in the presence of low-frequency noise and whistle frequency decreasing in the presence of high-frequency noise (Gospić & Picciulin, 2016). For example, bottlenose dolphins in Portuguese waters decrease their call rates and change the frequency parameters of whistles in the presence of boats (Luís et al., 2014), while dolphin groups with calves increase their whistle rates when tourist boats are within 200 m and when the boats increase their speed (Guerra et al., 2014). Likewise, modification of multiple vocalization parameters was shown in belugas residing in an area known for high levels of commercial traffic. These animals decreased their call rate, increased certain types of calls, and shifted upward in frequency content in the presence of small vessel noise (Lesage et al., 1999). Another study detected a measurable increase in the amplitude of their vocalizations when ships were present (Scheifele et al., 2005). Killer whales are also known to modify their calls during increased noise. For example, the source level of killer whale vocalizations was shown to increase with higher background noise levels associated with vessel traffic (the Lombard effect) (Holt et al., 2008). In addition, calls with a high-frequency component have higher source levels than other calls, which may be related to behavioral state, or may reflect a sustained increase in background noise levels (Holt et al., 2011). On the other hand, long-term modifications to vocalizations may be indicative of a learned response to chronic noise, or of a genetic or physiological shift in the populations. This type of change has been observed in killer whales off the northwestern coast of the United States between 1973 and 2003. This population increased the duration of primary calls once a threshold in observed vessel density (e.g., whale watching) was reached, which is suggested as being a long-term response to increased masking noise produced by the vessels (Foote et al., 2004).

The long-term and cumulative implications of ship sound on odontocetes is largely unknown (National Academies of Sciences Engineering and Medicine, 2017; National Marine Fisheries Service, 2007a), although some long-term consequences have been reported (Lusseau & Bejder, 2007). Repeated exposure to acoustic and other anthropogenic stimuli has been studied in several cases, especially as related to vessel traffic and whale watching. Common dolphins in New Zealand responded to dolphin-watching vessels by interrupting foraging and resting bouts, and took longer to resume behaviors in the presence of the vessel (Stockin et al., 2008). The authors speculated that repeated interruptions of the dolphins' foraging behaviors could lead to long-term implications for the population. Bejder et al. (2006a) studied responses of bottlenose dolphins to vessel approaches and found stronger and longer lasting reactions in populations of animals that were exposed to lower levels of vessel traffic overall. The authors indicated that lesser reactions in populations of dolphins regularly subjected to high levels of vessel traffic could be a sign of habituation, or it could be that the more sensitive animals in this population previously abandoned the area of higher human activity.

Similar to mysticetes, odontocete responses to vessel noise are varied, although many odontocete species seem to be more sensitive to vessel presence and vessel noise, and these two factors are difficult to tease apart. Some species, in particular killer whales and porpoises, may be sensitized to vessels and respond at further distances and lower received levels than other delphinids. In contrast, many odontocete species also approach vessels to bow ride, indicating either that these species are less sensitive to vessels, or that the behavioral drive to bow ride supersedes any impact of the associated noise. With these broad and disparate responses, it is difficult to assess the impacts of vessel noise on odontocetes.

### **Behavioral Reactions to Aircraft Noise**

The following paragraphs summarize what is known about the reaction of various marine mammal species to overhead flights of many types of fixed-wing aircraft and rotary-wing aircraft (i.e., helicopters), as well as unmanned aerial systems. Thorough reviews of the subject and available information is presented in Richardson et al. (1995b) and elsewhere (e.g., Efroymson et al., 2001; Holst et al., 2011; Luksenburg & Parsons, 2009; Smith et al., 2016). The most common responses of cetaceans to overflights were short surfacing durations, abrupt dives, and percussive behavior (breaching and tail slapping) (Nowacek et al., 2007a). Other behavioral responses such as flushing and fleeing the area of the source of the noise have also been observed (Holst et al., 2011; Manci et al., 1988). Richardson et al. (1995b) noted that marine mammal reactions to aircraft overflight largely consisted of opportunistic and anecdotal observations lacking clear distinction between reactions potentially caused by the noise of the aircraft and the visual cue an aircraft presents. In addition, it was suggested that variations in the responses noted were due to generally other undocumented factors associated with overflights (Richardson et al., 1995b). These factors could include aircraft type (e.g., single engine, multi engine, jet turbine), flight path (altitude, centered on the animal, off to one side, circling, level and slow), environmental factors (e.g., wind speed, sea state, cloud cover), and locations where native subsistence hunting continues and animals are more sensitive to anthropogenic impacts, including the noise from aircraft. Christiansen et al. (2016b) measured the in-air and underwater noise levels of two unmanned aerial vehicles, and found that in-air the broadband source levels were around 80 dB re 20  $\mu$ Pa, while at a meter underwater received levels were 95–100 dB re 1  $\mu$ Pa when the vehicle was only 5–10 m above the surface, and were not quantifiable above ambient noise levels when the vehicle was higher. Therefore, if an animal is near the surface and the unmanned aerial vehicle is low, it may be detected, but in most cases these vehicles are operated at much higher altitudes (e.g., over 30 m) and so are not likely to be heard.

The impact of aircraft overflights is one of the least well-known sources of potential behavioral response by any species or taxonomic group, and so many generalities must be made based on the little data available. There is some data for each taxonomic group; taken together it appears that in general, marine mammals have varying levels of sensitivity to overflights depending on the species and context.

#### ***Mysticetes***

Mysticetes either ignore or occasionally dive in response to aircraft overflights (Koski et al., 1998). Richardson (1985; 1995b) found no evidence that single or occasional aircraft flying above mysticetes causes long-term displacement of these mammals.

Bowhead whales in the Beaufort Sea exhibited a transient behavioral response to fixed-wing aircraft and vessels. Reactions were frequently observed at less than 1,000 ft. (304.8 m.) above sea level, infrequently observed at 1,500 ft. (457.2 m.), and not observed at all at 2,000 ft. (609.6 m.) (Richardson

et al., 1985). Bowhead whales reacted to helicopter overflights by diving, breaching, changing direction or behavior, and altering breathing patterns. Behavioral reactions decreased in frequency as the altitude of the helicopter increased to 150 m or higher. The bowheads exhibited fewer behavioral changes than did the odontocetes in the same area (Patenaude et al., 2002). It should be noted that bowhead whales in this study may have more acute responses to anthropogenic activity than many other marine mammals since these animals were presented with restricted egress due to limited open water between ice floes. Additionally, these animals are hunted by Alaska Natives, which could lead to animals developing additional sensitivity to human noise and presence.

A pilot study was conducted on the use of unmanned aerial systems to observe bowhead whales; flying at altitudes between 120 and 210 m above the surface, no behavioral responses were observed in any animals (Koski et al., 1998; Koski et al., 2015). Similarly, Christiansen et al. (2016a) did not observe any responses to an unmanned aerial vehicle flown 30–120 m above the water when taking photos of humpback whales to conduct photogrammetry and assess fitness. Acevedo-Whitehouse et al. (2010) successfully maneuvered a remote-controlled helicopter over large baleen whales to collect samples of their blows, with no more avoidance behavior than noted for typical photo-identification vessel approaches. These vehicles are much smaller and quieter than typical aircraft and so are less likely to cause a behavioral response, although they may fly at much lower altitudes (Smith et al., 2016).

#### ***Odontocetes***

Variable responses to aircraft have been observed in toothed whales, though overall little change in behavior has been observed during flyovers. Some toothed whales dove, slapped the water with their flukes or flippers, or swam away from the direction of the aircraft during overflights; others did not visibly react (Richardson et al., 1995b). Würsig et al. (1998) found that beaked whales were the most sensitive cetacean and reacted by avoiding marine mammal survey aircraft in 89 percent of sightings and at more than twice the rate as Kogia whales, which was the next-most reactive of the odontocetes in 39 percent of sightings; these are the same species that were sensitive to vessel traffic.

During standard marine mammal surveys at an altitude of 750 ft., some sperm whales remained on or near the surface the entire time the aircraft was in the vicinity, while others dove immediately or a few minutes after being sighted. Other authors have corroborated the variability in sperm whales' reactions to fixed-wing aircraft or helicopters (Green et al., 1992; Richter et al., 2003; Richter et al., 2006; Smultea et al., 2008; Würsig et al., 1998). In one study, sperm whales showed no reaction to a helicopter until they encountered the downdrafts from the rotors (Richardson et al., 1995b). A group of sperm whales responded to a circling aircraft (altitude of 800 to 1,100 ft.) by moving closer together and forming a defensive fan-shaped semicircle, with their heads facing outward. Several individuals in the group turned on their sides, apparently to look up toward the aircraft (Smultea et al., 2008). Whale-watching aircraft (fixed-wing airplanes and helicopters) apparently caused sperm whales to turn more sharply but did not affect blow interval, surface time, time to first click, or the frequency of aerial behavior (Richter et al., 2003).

Smaller delphinids generally react to overflights either neutrally or with a startle response (Würsig et al., 1998). The same species that show strong avoidance behavior to vessel traffic (Kogia whales and beaked whales) show similar reactions to aircraft (Würsig et al., 1998). Beluga whales reacted to helicopter overflights by diving, breaching, changing direction or behavior, and altering breathing patterns to a greater extent than mysticetes in the same area (Patenaude et al., 2002). These reactions increased in frequency as the altitude of the helicopter dropped below 150 m. A change in travel direction was noted in a group of pilot whales as the aircraft circled while conducting monitoring (State of Hawaii, 2015).

Much like mysticetes, odontocetes have demonstrated no responses to unmanned aerial systems. For example, Durban et al. (2015) conducted photogrammetry studies of killer whales using a small helicopter flown 35–40 m above the animals with no disturbance noted. However, odontocete responses may increase with reduced altitude, due either to noise or the shadows created by the vehicle (Smith et al., 2016). Bottlenose dolphins responded to a small portion of unmanned aerial vehicles by briefly orienting when the vehicle was relatively close (10–30 m high), but in most cases didn't respond at all (Ramos et al., 2018).

#### 3.4.2.1.1.6 Stranding

Marine mammals are subjected to a variety of natural and anthropogenic factors, acting alone or in combination, which may cause a marine mammal to strand (Geraci et al., 1999; Geraci & Lounsbury, 2005). When a marine mammal (alive or dead) swims or floats onto shore and becomes beached or incapable of returning to sea, the event is termed a “stranding” (Geraci et al., 1999; Geraci & Lounsbury, 2005; Perrin & Geraci, 2002). A stranding can also occur away from the shore if the animal is unable to cope in its present situation (e.g., disabled by a vessel strike, out of habitat) (Geraci & Lounsbury, 2005). Specifically, under U.S. law, a stranding is an event in the wild in which: “(A) a marine mammal is dead and is (i) on a beach or shore of the United States; or (ii) in waters under the jurisdiction of the United States (including any navigable waters); or (B) a marine mammal is alive and is (i) on a beach or shore of the United States and is unable to return to the water; (ii) on a beach or shore of the United States and, although able to return to the water, is in need of medical attention; or (iii) in the waters under the jurisdiction of the United States (including any navigable waters), but is unable to return to its natural habitat under its own power or without assistance” (16 U.S.C. section 1421h).

Natural factors related to strandings include limited food availability or following prey inshore, predation, disease, parasitism, natural toxins, echolocation disturbance, climatic influences, and aging (Bradshaw et al., 2006; Culik, 2004; Geraci et al., 1999; Geraci & Lounsbury, 2005; Huggins et al., 2015; National Research Council, 2006; Perrin & Geraci, 2002; Walker et al., 2005). Anthropogenic factors include pollution (Hall et al., 2006; Jepson et al., 2005), vessel strike (Geraci & Lounsbury, 2005; Laist et al., 2001), fisheries interactions (Read et al., 2006), entanglement (Baird & Gorgone, 2005; Saez et al., 2012; Saez et al., 2013), human activities (e.g., feeding, gunshot) (Geraci & Lounsbury, 2005; Dierauf & Gulland, 2001), and noise (Cox et al., 2006; National Research Council, 2003; Richardson et al., 1995b). For some stranding events, environmental factors (e.g., ocean temperature and wind speed and geographic conditions) can be utilized in predictive models to aid in understanding why marine mammals strand in certain areas more than others (Berini et al., 2015). In most instances, even for the more thoroughly investigated strandings involving post-stranding data collection and necropsies, the cause (or causes) for strandings remains undetermined.

Several mass strandings (strandings that involve two or more individuals of the same species, excluding a single mother-calf pair) that have occurred over the past two decades have been associated with anthropogenic activities that introduced sound into the marine environment, such as naval operations and seismic surveys. An in-depth discussion of strandings is in the Navy's technical report titled *Marine Mammal Strandings Associated with U.S. Navy Sonar Activities* (U.S. Department of the Navy, 2017c).

Sonar use during exercises involving the U.S. Navy has been identified as a contributing cause or factor in five specific mass stranding events: Greece in 1996; the Bahamas in March 2000; Madeira Island, Portugal in 2000; the Canary Islands in 2002, and Spain in 2006 (Cox et al., 2006; Fernandez, 2006; U.S. Department of the Navy, 2017c). These five mass strandings resulted in about 40 known cetacean

deaths consisting mostly of beaked whales and with close linkages to mid-frequency active sonar activity. In these circumstances, exposure to non-impulsive acoustic energy was considered a possible indirect cause of death of the marine mammals (Cox et al., 2006). Strandings of other marine mammal species have not been as closely linked to sonar exposure, but rather, have typically been attributed to natural or other anthropogenic factors. The Navy has reviewed training requirements, standard operating procedures, and potential mitigation measures, and has implemented changes to reduce the potential for acoustic related strandings to occur in the future. Discussions of procedures associated with these and other training and testing events are presented in Chapter 5 (Mitigation).

Multiple hypotheses regarding the relationship between non-impulsive sound exposure and stranding have been proposed. These range from direct impact of the sound on the physiology of the marine mammal, to behavioral reactions contributing to altered physiology (e.g., “gas and fat embolic syndrome”) (Fernandez et al., 2005; Jepson et al., 2003; Jepson et al., 2005), to behaviors directly contributing to the stranding (e.g., beaching of fleeing animals). Unfortunately, without direct observation of not only the event but also the underlying process, and given the potential for artefactual evidence (e.g., chronic condition, previous injury) to complicate conclusions from the post-mortem analyses of stranded animals (Cox et al., 2006), it has not been possible to determine with certainty the exact mechanism underlying these strandings.

Historically, stranding reporting and response efforts have been inconsistent, although they have improved considerably over the last 25 years. Although reporting forms have been standardized nationally, data collection methods, assessment methods, detail of reporting and procedures vary by region and are not yet standardized across the United States. Conditions such as weather, time, location, and decomposition state may also affect the ability to thoroughly examine a specimen (Carretta et al., 2016b; Moore et al., 2013). Because of this, the current ability to interpret long-term trends in marine mammal stranding is limited. While the investigation of stranded animals provides insight into the types of threats marine mammal populations face, investigations are only conducted on a small fraction of the total number of strandings that occur, limiting our understanding of the causes of strandings (Carretta et al., 2016a). Although many marine mammals likely strand due to natural or anthropogenic causes, the majority of reported type of occurrences in marine mammal strandings in the Pacific include fisheries interactions, entanglement, vessel strike, and predation (Carretta et al., 2017b; Helker et al., 2017).

#### **3.4.2.1.1.7 Long-Term Consequences**

Long-term consequences to a population are determined by examining changes in the population growth rate. Physical effects that could lead to a reduction in the population growth rate include mortality or injury, which could remove animals from the reproductive pool, and permanent hearing impairment or chronic masking, which could impact navigation, foraging, predator avoidance, or communication. The long-term consequences due to individual behavioral reactions and short-term or chronic instances of physiological stress are especially difficult to predict because individual experience over time can create complex contingencies, especially for long-lived animals like marine mammals. For example, a lost reproductive opportunity could be a measurable cost to the individual, or for very small populations to the population as a whole; however, short-term costs may be recouped during the life of an otherwise healthy individual. These factors are taken into consideration when assessing risk of long-term consequences. It is more likely that any long-term consequences to an individual would be a result of costs accumulated over a season, year, or life stage due to multiple behavioral or stress responses resulting from exposure to many sound-producing activities over significant periods.

Marine mammals exposed to high levels of human activities may leave the area, habituate to the activity, or tolerate the disturbance and remain in the area (Wartzok et al., 2003). Highly resident or localized populations may also stay in an area of disturbance because the cost of displacement may be higher than the cost of remaining (Forney et al., 2017). Longer-term displacement can lead to changes in abundance or distribution patterns of the species in the affected region (Bejder et al., 2006b; Blackwell et al., 2004; Teilmann et al., 2006). Gray whales in Baja California abandoned a historical breeding lagoon in the mid-1960s due to an increase in dredging and commercial shipping operations. However, whales did repopulate the lagoon after shipping activities had ceased for several years (Bryant et al., 1984). Mysticetes in the northeast tended to adjust to vessel traffic over a number of years, trending towards more neutral responses to passing vessels (Watkins, 1986), indicating that some animals may habituate or otherwise learn to cope with high levels of human activity. Bejder et al. (2006a) studied responses of bottlenose dolphins to vessel approaches and found that lesser reactions in populations of dolphins regularly subjected to high levels of vessel traffic could be a sign of habituation, or it could be that the more sensitive animals in this population previously abandoned the area of higher human activity.

Moore and Barlow (2013) noted a decline in the overall beaked whale population in a broad area of the Pacific Ocean along the U.S. West Coast. Moore and Barlow (2013) provide several hypotheses for the decline of beaked whales in those waters, one of which is anthropogenic sound including the use of sonar by the U.S. Navy; however, new data have been published that raise uncertainties over whether a decline in the beaked whale population occurred off the U.S. West Coast between 1996 and 2014 (Barlow, 2016). Moore and Barlow (2017) have since incorporated information from the entire 1991 to 2014 time series, which suggests an increasing abundance trend and a reversal of the declining trend along the U.S. West Coast that had been noted in their previous (2013) analysis.

In addition, studies on the Atlantic Undersea Test and Evaluation Center instrumented range in the Bahamas have shown that some Blainville's beaked whales may be resident during all or part of the year in the area. Individuals may move off the range for several days during and following a sonar event, but return within a few days (McCarthy et al., 2011; Tyack et al., 2011). Photo-identification studies in the Southern California Range Complex have identified approximately 100 individual Cuvier's beaked whale individuals, with 40 percent having been seen in one or more prior years and re-sightings up to seven years apart (Falcone et al., 2009; Falcone & Schorr, 2014). These results indicate long-term residency by individuals in an intensively used Navy training and testing area, which may suggest a lack of long-term consequences as a result of exposure to Navy training and testing activities, but could also be indicative of high-value resources that exceed the cost of remaining in the area. Long-term residency does not mean there has been no impact on population growth rates, and there are no data existing on the reproductive rates of populations inhabiting the Navy range area around San Clemente Island as opposed to beaked whales from other areas. In that regard however, recent results from photo-identifications are beginning to provide critically needed calving and weaning rate data for resident animals on the Navy's Southern California range. Three adult females that had been sighted with calves in previous years were again sighted in 2016, one of these was associated with her second calf, and a fourth female that was first identified in 2015 without a calf, was sighted in 2016 with a calf (Schorr et al., 2017). Resident females documented with and without calves from year to year will provide the data for this population that can be applied to future research questions.

Research involving three tagged Cuvier's beaked whales in the Southern California Range Complex reported on by Falcone and Schorr (2012, 2014) has documented movements in excess of hundreds of

kilometers by some of those animals. Schorr et al. (2014) reported the results for an additional eight tagged Cuvier's beaked whales in the same area. Five of these eight whales made journeys of approximately 250 km from their tag deployment location, and one of these five made an extra-regional excursion over 450 km south to Mexico and back again. Given that some beaked whales may routinely move hundreds of kilometers as part of their normal pattern (Schorr et al., 2014), temporarily leaving an area to avoid sonar or other anthropogenic activity may have little cost.

Another approach to investigating long-term consequences of anthropogenic noise exposure has been an attempt to link short-term effects to individuals from anthropogenic stressors with long-term consequences to populations using population models. Population models are well known from many fields in biology including fisheries and wildlife management. These models accept inputs for the population size and changes in vital rates of the population, such as the mean values for survival age, lifetime reproductive success, and recruitment of new individuals into the population. Unfortunately, for acoustic and explosive impacts on marine mammal populations, many of the inputs required by population models are not known. Nowacek et al. (2016) reviewed new technologies, including passive acoustic monitoring, tagging, and the use of unmanned aerial vehicles, which can improve scientists' abilities to study these model inputs and link behavioral changes to individual life functions and ultimately population-level effects. The linkage between immediate behavioral or physiological effects to an individual due to a stressor such as sound, the subsequent effects on that individual's vital rates (growth, survival, and reproduction), and in turn the consequences for the population have been reviewed in National Research Council (2005).

The Population Consequences of Acoustic Disturbance model (National Research Council, 2005) proposes a conceptual model for determining how changes in the vital rates of individuals (i.e., a biologically significant consequence to the individual) translates into biologically significant consequences to the population. In 2009, the U.S. Office of Naval Research set up a working group to transform the Population Consequences of Acoustic Disturbance framework into a mathematical model and include other stressors potentially causing disturbance in addition to noise. The model, now called Population Consequences of Disturbance, has been used for case studies involving bottlenose dolphins, North Atlantic right whales, beaked whales, southern elephant seals, California sea lions, blue whales, humpback whales, and harbor porpoise (Costa et al., 2016a; Costa et al., 2016b; Harwood & King, 2014; Hatch et al., 2012; King et al., 2015; McHuron et al., 2018; New et al., 2013a; New et al., 2013b; New et al., 2014; Pirotta et al., 2018). Currently, the Population Consequences of Disturbance model provides a theoretical framework and identifies types of data that would be needed to assess population-level impacts using this process. The process is complicated and provides a foundation for the type of data that is needed, which is currently lacking for many marine mammal species. Relevant data needed for improving these analytical approaches for population-level consequences resulting from disturbances will continue to be collected during projects funded by the U.S. Navy Marine Species Monitoring Program.

Costa et al. (2016a) emphasized taking into account the size of an animal's home range, whether populations are resident and non-migratory, or if they migrate over long areas and share their feeding or breeding areas with other populations. These factors, coupled with the extent, location, and duration of a disturbance can lead to markedly different impact results. For example, Costa et al. (2016a) modeled seismic surveys with different radii of impacts on the foraging grounds of Bering Sea humpback whales, West Antarctic Peninsula humpback whales, and California Current blue whales, and used data from tagged whales to determine foraging locations and effort on those grounds. They found that for

the blue whales and the West Antarctic humpback whales, less than 19 percent and 16 percent (respectively) of each population would be exposed, and less than 19 percent and 6 percent (respectively) of foraging behavior would be disturbed. This was likely due to the fact that these populations forage for krill over large areas. In contrast, the Bering Sea population of humpback whales had over 90 percent of the population exposed when the disturbance zones extended beyond 50 km, but 100 percent of their foraging time would occur during an exposure when the zone was 25 km or more. These animals forage for fish over a much smaller area, thereby having a limited range for foraging that can be disturbed. Energetic costs were estimated for western gray whales that migrated to possible wintering grounds near China or to the Baja California wintering grounds of eastern gray whales versus the energetic costs of the shorter migration of eastern gray whales (Villegas-Amtmann et al., 2017). Researchers found that when the time spent on the breeding grounds was held constant for both populations, the energetic requirements for the western gray whales were estimated to be 11 percent and 15 percent greater during the migration to Baja California and China, respectively, than for the migration of eastern gray whales, and therefore this population would be more sensitive to energy lost through disturbance.

Pirotta et al. (2018) modeled one reproductive cycle of a female North Pacific blue whale, starting with leaving the breeding grounds off Baja California to begin migrating north to feeding grounds off California, and ending with her return to the breeding grounds, giving birth, and lactating. They modeled this scenario with no disturbance and found 95 percent calf recruitment, under a “normal” environmental perturbation (El Niño-Southern Oscillation) there was a very small reduction in recruitment, and under an “unprecedented” environmental change, recruitment was reduced to 69 percent. An intense, localized anthropogenic disturbance was modeled (although the duration of the event was not provided); if the animals were not allowed to leave the area they did not forage and recruitment dropped to 63 percent. However, if animals could leave the area of the disturbance then there was almost no change to the recruitment rate. Finally, a weak but broader spatial disturbance, where foraging was reduced by 50 percent, caused only a small decrease in calf recruitment to 94 percent.

Using the Population Consequences of Disturbance framework, modeling of the long-term consequences of exposure has been conducted for a variety of marine mammal species and stressors. Even when high and frequent exposure levels are included, few long-term consequences have been predicted. For example, De Silva et al. (2014) conducted a population viability analysis on the long-term impacts of pile driving and construction noise on harbor porpoises and bottlenose dolphins. Despite including the extreme and unlikely assumptions that 25 percent of animals that received PTS would die, and that behavioral displacement from an area would lead to breeding failure, the model only found short-term impacts on the population size and no long-term effects on population viability. Similarly, King et al. (2015) developed a Population Consequences of Disturbance framework using expert elicitation data on impacts from wind farms on harbor porpoises and, even under the worst-case scenarios, predicted less than a 0.5 percent decline in harbor porpoise populations. Nabe-Nelson et al. (2014) also modeled the impact of noise from wind farms on harbor porpoises and predicted that even when assuming a 10 percent reduction in population size if prey is impacted up to two days, the presence of ships and wind turbines did not deplete the population. In contrast, Heinis and De Jong (2015) used the Population Consequences of Disturbance framework to estimate impacts from both pile driving and seismic exploration on harbor porpoises and found a 23 percent decrease in population size over six years, with an increased risk for further reduction with additional disturbance days. These

seemingly contradictory results demonstrate that refinements to models need to be investigated to improve consistency and interpretation of model results.

The Population Consequences of Disturbance model developed by New et al. (2013b) predicted that beaked whales require energy dense prey and high quality habitat, and that non-lethal disturbances that displace whales from that habitat could lead to long-term impacts on fecundity and survival; however, the authors were forced to use many conservative assumptions within their model since many parameters are unknown for beaked whales. As discussed above in Schorr et al. (2014), beaked whales have been tracked roaming over distances of 250 km or more, indicating that temporary displacement from a small area may not preclude finding energy dense prey or high quality habitat. Farmer et al. (2018) developed a bioenergetics framework to examine the impact of foraging disruption on body reserves of individual sperm whales. The authors examined rates of daily foraging disruption to predict the number of days to terminal starvation for various life stages, assuming exposure to seismic surveys. Mothers with calves were found to be most vulnerable to disruptions.

Another Population Consequences of Disturbance model developed in New et al. (2014) predicted elephant seal populations to be relatively robust even with a greater than 50 percent reduction in foraging trips (only a 0.4 percent population decline in the following year). McHuron et al. (2018) modeled the introduction of a generalized disturbance at different times throughout the breeding cycle of California sea lions, with the behavior response being an increase in the duration of a foraging trip by the female. Very short duration disturbances or responses led to little change, particularly if the disturbance was a single event, and changes in the timing of the event in the year had little effect. However, with even relatively short disturbances or mild responses, when a disturbance was modeled as recurring there were resulting reductions in population size and pup recruitment. Often, the effects weren't noticeable for several years, as the impacts on pup recruitment didn't affect the population until those pups were mature.

It should be noted that, in all of these models, assumptions were made and many input variables were unknown and so were estimated using available data. It is still not possible to utilize individual short-term behavioral responses to estimate long-term or population level effects.

The best assessment of long-term consequences from Navy training and testing activities will be to monitor the populations over time within the Study Area. A U.S. workshop on Marine Mammals and Sound (Fitch et al., 2011) indicated a critical need for baseline biological data on marine mammal abundance, distribution, habitat, and behavior over sufficient time and space to evaluate impacts from human-generated activities on long-term population survival. The Navy has developed and implemented comprehensive monitoring plans since 2009 for protected marine mammals occurring on Navy ranges with the goal of assessing the impacts of training and testing activities on marine species and the effectiveness of the Navy's mitigation measures. The results of this long-term monitoring are now being compiled and analyzed for trends in occurrence or abundance over time (e.g., Martin et al., 2017); preliminary results of this analysis at Pacific Missile Range Facility off Kauai, Hawaii indicate no changes in detection rates for several species over the past decade, demonstrating that Navy activities may not be having long-term population-level impacts. This type of analysis can be expanded to the other Navy ranges, such as the Mariana Islands Range Complex. Continued analysis of this 15-year dataset and additional monitoring efforts over time are necessary to fully understand the long-term consequences of exposure to military readiness activities.

### 3.4.2.1.2 Impacts from Sonar and Other Transducer Stressors

Sonar and other transducers proposed for use could be used throughout the Study Area. Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. General categories of these systems are described in Section 3.0.4.1 (Acoustic Stressors). The overall use of sonar and other transducers for training and testing activities would be similar to what is currently conducted (see Tables 2.5-1 and 2.5-2 for details). Although individual activities may vary some from those previously analyzed, the overall determinations presented in the 2015 MITT Final EIS/OEIS remain valid. The quantitative analysis has been improved upon and updated since the 2015 MITT Final EIS/OEIS; therefore, the new analysis is fully presented and described in further detail in the technical report *Quantitative Analysis for Estimating Acoustic and Explosive Impacts to Marine Mammals and Sea Turtles* (U.S. Department of the Navy, 2018a).

Sonar-induced acoustic resonance and bubble formation phenomena are very unlikely to occur under realistic conditions, as discussed in Section 3.4.2.1.1.1 (Injury). Non-auditory injury (i.e., other than PTS) and mortality from sonar and other transducers is so unlikely as to be discountable under normal conditions and is therefore not considered further in this analysis.

The most probable impacts from exposure to sonar and other transducers are PTS, TTS, behavioral reactions, masking, and physiological stress (Section 3.4.2.1.1.2, Hearing Loss; Section 3.4.2.1.1.3, Physiological Stress; Section 3.4.2.1.1.4, Masking; and Section 3.4.2.1.1.5, Behavioral Reactions).

#### 3.4.2.1.2.1 Methods for Analyzing Impacts from Sonar and Other Transducers

The Navy performed a quantitative analysis to estimate the number of times that marine mammals could be affected by sonars and other transducers used during Navy training and testing activities. The Navy's quantitative analysis to determine impacts on marine mammals uses the Navy Acoustic Effects Model to produce initial estimates of the number of times that animals may experience these effects; these estimates are further refined by considering animal avoidance of sound-producing activities and implementation of procedural mitigation measures. The steps of this quantitative analysis are described in Section 3.0.1.2 (Navy's Quantitative Analysis to Determine Impacts on Sea Turtles and Marine Mammals), which takes into account:

- criteria and thresholds used to predict impacts from sonar and other transducers (see below)
- the density and spatial distribution of marine mammals
- the influence of environmental parameters (e.g., temperature, depth, salinity) on sound propagation when estimating the received sound level on the animals


A detailed explanation of this analysis is provided in the technical report titled *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

### Criteria and Thresholds Used to Estimate Impacts from Sonar and Other Transducers

#### *Auditory Weighting Functions*

Animals are not equally sensitive to noise at all frequencies. To capture the frequency-dependent nature of the effects of noise, auditory weighting functions are used (Figure 3.4-5). Auditory weighting functions are mathematical functions that adjust received sound levels to emphasize ranges of best hearing and de-emphasize ranges with less or no auditory sensitivity. They are based on a generic band pass filter and incorporate species-specific hearing abilities to calculate a weighted received sound level in units SPL or SEL. Due to the band pass nature of auditory weighting functions, they resemble an

inverted “U” shape with amplitude plotted as a function of frequency. The flatter portion of the plotted function, where the amplitude is closest to zero, is the emphasized frequency range (i.e., the pass-band), while the frequencies below and above this range (where amplitude declines) are de-emphasized.




Source: For parameters used to generate the functions and more information on weighting function derivation see U.S. Department of the Navy (2017a)

Notes: HF = High-Frequency Cetacean, LF = Low-Frequency Cetacean, and MF = Mid-Frequency Cetacean

**Figure 3.4-5: Navy Auditory Weighting Functions for all Species Groups**

***Hearing Loss from Sonar and Other Transducers***

Defining the TTS and PTS exposure functions (Figure 3.4-6) requires identifying the weighted exposures necessary for TTS and PTS onset from sounds produced by sonar and other transducers. The criteria used to define threshold shifts from non-impulsive sources (e.g., sonar) determines TTS onset as the SEL necessary to induce 6 dB of threshold shift. An SEL 20 dB above the onset of TTS is used in all hearing groups of marine mammals underwater to define the PTS threshold (Southall et al., 2007).



Notes: The solid curve is the exposure function for TTS onset and the large dashed curve is the exposure function for PTS onset. Small dashed lines and asterisks indicate the SEL threshold for TTS and PTS onset in the frequency range of best hearing.

**Figure 3.4-6: TTS and PTS Exposure Functions for Sonar and Other Transducers**

***Behavioral Responses from Sonar and Other Transducers***

Behavioral response criteria are used to estimate the number of animals that may exhibit a behavioral response to sonar and other transducers. See the *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* technical report for detailed information on how the Behavioral Response Functions were derived (U.S. Department of the Navy, 2017a). Developing the new behavioral criteria involved multiple steps. All peer-reviewed published behavioral response studies conducted both in the field and on captive animals were examined in order to understand the breadth of behavioral responses of marine mammals to sonar and other transducers.

The data from the behavioral studies were analyzed by looking for significant responses, or lack thereof, for each experimental session. The terms “significant response” or “significant behavioral response” are used in describing behavioral observations from field or captive animal research that may rise to the

level of “harassment” for military readiness activities. Under the MMPA, for military readiness activities, such as Navy training and testing, behavioral “harassment” is: “any act that *disturbs* or is likely to *disturb* a marine mammal or marine mammal stock in the wild by causing disruption of natural behavioral patterns, including, but not limited to, migration, surfacing, nursing, breeding, feeding, or sheltering, *to a point where such behavioral patterns are abandoned or significantly altered*” (16 U.S.C. section 1362(3)(18)(B)).

The likelihood of injury due to disruption of normal behaviors would depend on many factors, such as the duration of the response, from what the animal is being diverted, and the life history of the animal. Due to the nature of behavioral response research to date, it is not currently possible to ascertain the types of observed reactions that would lead to an abandonment or significant alteration of a natural behavior pattern. Therefore, the Navy has developed a methodology to estimate the possible significance of behavioral reactions and impacts on natural behavior patterns.

Behavioral response severity is described herein as “low,” “moderate,” or “high.” These are derived from the Southall et al. (2007) severity scale. Low severity responses are those behavioral responses that fall within an animal’s range of typical (baseline) behaviors and are unlikely to disrupt an individual to a point where natural behavior patterns are significantly altered or abandoned. Low severity responses include an orientation or startle response, change in respiration, change in heart rate, and change in group spacing or synchrony.

Moderate severity responses could become significant if sustained over a longer duration. What constitutes a long-duration response is different for each situation and species, although it is likely dependent upon the magnitude of the response and species characteristics such as age, body size, feeding strategy, and behavioral state at the time of the exposure. In general, a response could be considered “long-duration” if it lasted for tens of minutes to a few hours, or enough time to significantly disrupt an animal’s daily routine.

Moderate severity responses included

- alter migration path,
- alter locomotion (speed, heading),
- alter dive profiles,
- stop/alter nursing,
- stop/alter breeding,
- stop/alter feeding/foraging,
- stop/alter sheltering/resting,
- stop/alter vocal behavior if tied to foraging or social cohesion, and
- avoid area near sound source.

For the derivation of behavioral criteria, a significant duration was defined as a response that lasted for the duration of exposure or longer, regardless of how long the exposure session may have been. This assumption was made because it was not possible to tell if the behavioral responses would have continued if the exposure had continued. The costs associated with these observed behavioral reactions were not measured, so it is not possible to judge whether reactions would have risen to the level of significance as defined above, although it was conservatively assumed the case. High severity responses

include those responses with immediate consequences (e.g., stranding, mother-calf separation), and were always considered significant behavioral reactions regardless of duration.

Marine mammal species were placed into behavioral criteria groups based on their known or suspected behavioral sensitivities to sound (Figure 3.4-7 through Figure 3.4-9). In most cases, these divisions are driven by taxonomic classifications (e.g., mysticetes, odontocetes). The Odontocete group combines most of the mid- and high-frequency cetaceans, without the beaked whales. These groups are combined as there are not enough data to separate them for behavioral responses.




Figure 3.4-7: Behavioral Response Function for Odontocetes





Figure 3.4-8: Behavioral Response Function for Mysticetes



**Figure 3.4-9: Behavioral Response Function for Beaked Whales**

For all taxa, distances beyond which significant behavioral responses to sonar and other transducers are unlikely to occur, denoted as “cutoff distances,” were defined based on existing data (Table 3.4-3). The distance between the animal and the sound source is a strong factor in determining that animal’s potential reaction (e.g., DeRuiter et al., 2013b). For training and testing events that contain multiple platforms or tactical sonar sources that exceed 215 dB re 1 μPa at 1 m, this cutoff distance is substantially increased (i.e., doubled) from values derived from the literature. The use of multiple platforms and intense sound sources are factors that probably increase responsiveness in marine mammals overall. There are currently few behavioral observations under these circumstances; therefore, the Navy will conservatively predict significant behavioral responses at further ranges for these more intense activities.

**Table 3.4-3: Cutoff Distances for Moderate Source Level, Single Platform Training and Testing Events and for All Other Events with Multiple Platforms or Sonar with Source Levels at or Exceeding 215 dB re 1 μPa at 1 m**

| <i>Criteria Group</i> | <i>Moderate SL/Single Platform Cutoff Distance</i> | <i>High SL/Multi-Platform Cutoff Distance</i> |
|-----------------------|--|---|
| Odontocetes           | 10 km  | 20 km   |
| Mysticetes            | 10 km  | 20 km   |
| Beaked Whales         | 25 km  | 50 km   |

Notes: km = kilometer(s), SL = source level


**Assessing the Severity of Behavioral Responses from Sonar Under Military Readiness**

As discussed above, the terms “significant response” or “significant behavioral response” are used in describing behavioral reactions that may lead to an abandonment or significant alteration of a natural behavior pattern. Due to the limited amount of behavioral response research to date and relatively short durations of observation, it is not possible to ascertain the true significance of the majority of the

observed reactions. When deriving the behavioral criteria, it was assumed that most reactions that lasted for the duration of the sound exposure or longer were significant, even though many of the exposures lasted for 30 minutes or less. Furthermore, the experimental designs used during many of the behavioral response studies were unlike Navy activities in many important ways. These differences include tagging subject animals, following subjects for sometimes hours before the exposure, vectoring towards the subjects after animals began to avoid the sound source, and making multiple close passes on focal groups. This makes the estimated behavioral impacts from Navy activities using the criteria derived from these experiments difficult to interpret. While the state of science does not currently support definitively distinguishing between significant and insignificant behavioral reactions, as described in the technical report titled *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a), the Navy’s analysis incorporates conservative assumptions to account for this uncertainty and therefore likely overestimates the potential impacts.

The estimated behavioral reactions from the Navy’s quantitative analysis are grouped into several categories based on the most powerful sonar source, the number of platforms, the duration, and geographic extent of each Navy activity attributed to the predicted impact. Activities that occur on Navy instrumented ranges or within Navy homeports require special consideration due to the repeated nature of activities in these areas.

Low severity responses are within an animal’s range of typical (baseline) behaviors and are unlikely to disrupt an individual to a point where natural behavior patterns are significantly altered or abandoned. Although the derivation of the Navy’s behavioral criteria did not count low severity responses as significant behavioral responses, in practice, some reactions estimated using the behavioral criteria are likely to be low severity (Figure 3.4-10).



**Figure 3.4-10: Relative Likelihood of a Response Being Significant Based on the Duration and Severity of Behavioral Reactions**

High severity responses are those with a higher potential for direct consequences to growth, survivability, or reproduction. Examples include prolonged separation of females and dependent offspring, panic, flight, stampede, or stranding. High severity reactions would always be considered significant; however, these types of reactions are probably rare under most conditions and may still not lead to direct consequences on survivability. For example, a separation of a killer whale mother-calf pair was observed once during a behavioral response study to an active sonar source (Miller et al., 2014), but the animals were rejoined as soon as the ship had passed. Therefore, although this was a severe response, it did not lead to a negative outcome. Five beaked whale strandings have also occurred associated with U.S. Navy active sonar use as discussed above (see Section 3.4.2.1.1.6, Stranding), but the confluence of factors that contributed to those strandings is now better understood, and the avoidance of those factors has resulted in no known marine mammal strandings associated with U.S. Navy sonar activities for over a decade. The Navy is unable to predict these high severity responses for any activities since the probability of occurrence is apparently very low, although the Navy acknowledges that severe reactions could occasionally occur. In fact, no significant behavioral responses such as panic, stranding, or other severe reactions have been observed during monitoring of actual training or testing activities.

The responses estimated using the Navy's quantitative analysis are most likely to be moderate severity. Moderate severity responses would be considered significant if they were sustained for a duration long enough that it caused an animal to be outside of normal daily variations in feeding, reproduction, resting, migration/movement, or social cohesion. As mentioned previously, the behavioral response functions used within the Navy's quantitative analysis were primarily derived from experiments using short-duration sound exposures that in many cases lasted for less than 30 minutes. If animals exhibited moderate severity reactions for the duration of the exposure or longer, then it was conservatively assumed that the animal experienced a significant behavioral reaction. However, the experiments did not include measurements of costs to animals beyond the immediately observed reactions, and no direct correlations exist between an observed behavioral response and a cost that may result in long-term consequences. Within the Navy's quantitative analysis, many behavioral reactions are estimated from exposure to sonar that may exceed an animal's behavioral threshold for only a single ping to several minutes. While the state of science does not currently support definitively distinguishing between significant and insignificant behavioral reactions, as described in the technical report titled *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a), the Navy's analysis incorporates conservative assumptions to account for this uncertainty and therefore likely overestimates the potential impacts.

### **Accounting for Mitigation**

The Navy will implement mitigation measures to avoid or reduce potential impacts from active sonar on marine mammals, as described in Section 5.3.2.1 (Active Sonar). The benefits of mitigation are conservatively factored into the analysis for Alternative 1 and Alternative 2 of the Proposed Action for training and testing. The Navy's mitigation measures are identical for both action alternatives.

Procedural mitigation measures include a power down or shut down (i.e., power off) of applicable active sonar sources when a marine mammal is observed in a mitigation zone. The mitigation zones for active sonar activities were designed to avoid the potential for marine mammals to be exposed to levels of sound that could result in auditory injury (i.e., PTS) from active sonar to the maximum extent practicable. The mitigation zones for active sonar extend beyond the respective average ranges to auditory injury (including PTS). Therefore, the impact analysis quantifies the potential for procedural

mitigation to reduce the risk of PTS. Two factors are considered when quantifying the effectiveness of procedural mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., active sonar) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

In the quantitative analysis, consideration of mitigation measures means that, for activities that implement mitigation, some model-estimated PTS is considered mitigated to the level of TTS. The impact analysis does not analyze the potential for mitigation to reduce TTS or behavioral effects, even though mitigation could also reduce the likelihood of these effects. In practice, mitigation also protects all unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

The ability to observe the ranges to PTS was estimated for each training or testing event. The ability of Navy Lookouts to detect marine mammals within a mitigation zone is dependent on the animal's presence at the surface and the characteristics of the animal that influence its sightability (such as group size or surface active behavior). The behaviors and characteristics of some species may make them easier to detect. Certain behaviors, such as leaping and breaching, are visible from a great distance and likely increase sighting distances and detections of those species. Environmental conditions under which the training or testing activity could take place are also considered, such as sea surface conditions, weather (e.g., fog or rain), and day versus night.

The Navy will also implement mitigation measures for certain active sonar activities within mitigation areas, as described in Appendix I (Geographic Mitigation Assessment). The benefits of mitigation areas are discussed qualitatively and have not been factored into the quantitative analysis process or reductions in take for the MMPA and ESA impact estimates. Mitigation areas are designed to help avoid or reduce impacts during biologically important life processes within particularly important habitat areas. Therefore, mitigation area benefits are discussed in terms of the context of impact avoidance or reduction.

### **Marine Mammal Avoidance of Sonar and other Transducers**

Because a marine mammal is assumed to initiate avoidance behavior after an initial startle reaction when exposed to relatively high received levels of sound, a marine mammal could reduce its cumulative sound energy exposure over a sonar event with multiple pings (i.e., sound exposures). This would reduce risk of both PTS and TTS, although the quantitative analysis conservatively only considers the potential to reduce instances of PTS by accounting for marine mammals swimming away to avoid repeated high-level sound exposures. All reductions in PTS impacts from likely avoidance behaviors are instead considered TTS impacts.

#### **3.4.2.1.2.2 Impact Ranges for Sonar and Other Transducers**

The following section provides range to effects for sonar and other transducers to specific criteria determined using the Navy Acoustic Effects Model. Marine mammals within these ranges would be predicted to receive the associated effect. Range to effects is important information in not only

predicting acoustic impacts, but also in verifying the accuracy of model results against real-world situations and assessing the level of impact that will likely be mitigated within applicable mitigation zones.

The ranges to the PTS threshold for an exposure of 30 seconds are shown in Table 3.4-4 relative to the marine mammal’s functional hearing group. This period (30 seconds) was chosen based on examining the maximum amount of time a marine mammal would realistically be exposed to levels that could cause the onset of PTS based on platform (e.g., ship) speed and a nominal animal swim speed of approximately 1.5 meters per second. The ranges provided in the table include the average range to PTS, as well as the range from the minimum to the maximum distance at which PTS is possible for each hearing group. Since any hull-mounted sonar, such as the SQS-53, engaged in anti-submarine warfare training would be moving at between 10 and 15 knots and nominally pinging every 50 seconds, the vessel will have traveled a minimum distance of approximately 257 m during the time between those pings (note: 10 knots is the speed used in the Navy Acoustic Effects Model). As a result, there is little overlap of PTS footprints from successive pings, indicating that in most cases, an animal predicted to receive PTS would do so from a single exposure (i.e., ping). For all other bins (besides MF1), PTS ranges are short enough that marine mammals (with a nominal swim speed of approximately 1.5 meters per second) should be able to avoid higher sound levels capable of causing onset PTS within this 30-second period.

For all other functional hearing groups (low-frequency cetaceans and mid-frequency cetaceans), 30-second average PTS zones are substantially shorter. A scenario could occur where an animal does not leave the vicinity of a ship or travels a course parallel to the ship, however, the close distances required make PTS exposure unlikely. For a Navy vessel moving at a nominal 10 knots, it is unlikely a marine mammal could maintain the speed to parallel the ship and receive adequate energy over successive pings to suffer PTS.

The tables below illustrate the range to TTS for 1, 30, 60, and 120 seconds from five representative sonar systems (see Table 3.4-5 through Table 3.4-9). Due to the lower acoustic thresholds for TTS versus PTS, ranges to TTS are longer. Therefore, successive pings can be expected to add together, further increasing the range to onset-TTS.

**Table 3.4-4: Range to Permanent Threshold Shift for Five Representative Sonar Systems**

| <i>Hearing Group</i>     | <i>Approximate PTS (30 seconds) Ranges (meters)<sup>1</sup></i> |                      |                      |                      |                      |
|--------------------------|---|----------------------|----------------------|----------------------|----------------------|
|                          | <i>Sonar bin HF4</i>  | <i>Sonar bin LF4</i> | <i>Sonar bin MF1</i> | <i>Sonar bin MF4</i> | <i>Sonar bin MF5</i> |
| High-frequency cetaceans | 29<br>(22–35)   | 0<br>(0–0)           | 181<br>(180–190)     | 30<br>(30–30)        | 9<br>(8–10)          |
| Low-frequency cetaceans  | 0<br>(0–0)  | 0<br>(0–0)           | 65<br>(65–65)        | 15<br>(15–15)        | 0<br>(0–0)           |
| Mid-frequency cetaceans  | 1<br>(0–1)  | 0<br>(0–0)           | 16<br>(16–16)        | 3<br>(3–3)           | 0<br>(0–0)           |

<sup>1</sup> PTS ranges extend from the sonar or other transducers to the indicated distance. The average range to PTS is provided as well as the range from the estimated minimum to the maximum range to PTS in parentheses.

Notes: HF= high-frequency, LF = low-frequency, MF = mid-frequency, PTS = permanent threshold shift

**Table 3.4-5: Ranges to Temporary Threshold Shift for Sonar Bin HF4 over a Representative Range of Environments within the Study Area**

| <i>Hearing Group</i>     | <i>Approximate TTS Ranges (meters)<sup>1</sup></i> |                   |                   |                    |
|--------------------------|--|-------------------|-------------------|--------------------|
|                          | <i>Sonar Bin HF4</i>                               |                   |                   |                    |
|                          | <i>1 second</i>                                    | <i>30 seconds</i> | <i>60 seconds</i> | <i>120 seconds</i> |
| High-frequency cetaceans | 155<br>(110–210)                                   | 259<br>(180–350)  | 344<br>(240–480)  | 445<br>(300–600)   |
| Low-frequency cetaceans  | 1<br>(0–2)   | 2<br>(1–3)        | 4<br>(3–5)        | 7<br>(5–8)         |
| Mid-frequency cetaceans  | 10<br>(7–12)                                       | 17<br>(12–21)     | 24<br>(17–30)     | 33<br>(25–40)      |

<sup>1</sup> Ranges to TTS represent the model predictions in different areas and seasons within the Study Area. The zone in which animals are expected to suffer TTS extend from onset-PTS to the distance indicated. The average range to TTS is provided as well as the range from the estimated minimum to the maximum range to TTS in parenthesis.

Notes: HF = high-frequency, TTS = temporary threshold shift

**Table 3.4-6: Ranges to Temporary Threshold Shift for Sonar Bin LF4 over a Representative Range of Environments within the Study Area**

| <i>Hearing Group</i>     | <i>Approximate TTS Ranges (meters)<sup>1</sup></i> |                   |                   |                    |
|--------------------------|--|-------------------|-------------------|--------------------|
|                          | <i>Sonar Bin LF4</i>                               |                   |                   |                    |
|                          | <i>1 second</i>                                    | <i>30 seconds</i> | <i>60 seconds</i> | <i>120 seconds</i> |
| High-frequency cetaceans | 0<br>(0–0)   | 0<br>(0–0)        | 0<br>(0–0)        | 0<br>(0–0)         |
| Low-frequency cetaceans  | 3<br>(3–3)   | 4<br>(4–4)        | 6<br>(6–6)        | 9<br>(9–9)         |
| Mid-frequency cetaceans  | 0<br>(0–0)   | 0<br>(0–0)        | 0<br>(0–0)        | 0<br>(0–0)         |

<sup>1</sup> Ranges to TTS represent the model predictions in different areas and seasons within the Study Area. The zone in which animals are expected to suffer TTS extend from onset-PTS to the distance indicated. The average range to TTS is provided as well as the range from the estimated minimum to the maximum range to TTS in parentheses.

Notes: LF = low-frequency, TTS = temporary threshold shift

**Table 3.4-7: Ranges to Temporary Threshold Shift for Sonar Bin MF1 over a Representative Range of Environments within the Study Area**

| <i>Hearing Group</i>     | <i>Approximate TTS Ranges (meters)<sup>1</sup></i> |                        |                        |                        |
|--------------------------|--|------------------------|------------------------|------------------------|
|                          | <i>Sonar Bin MF1</i>                               |                        |                        |                        |
|                          | <i>1 second</i>                                    | <i>30 seconds</i>      | <i>60 seconds</i>      | <i>120 seconds</i>     |
| High-frequency cetaceans | 3,181<br>(2,025–5,025)                             | 3,181<br>(2,025–5,025) | 5,298<br>(2,275–7,775) | 6,436<br>(2,525–9,775) |
| Low-frequency cetaceans  | 898<br>(850–1,025)                                 | 898<br>(850–1,025)     | 1,271<br>(1,025–1,525) | 1,867<br>(1,275–3,025) |
| Mid-frequency cetaceans  | 210<br>(200–210)                                   | 210<br>(200–210)       | 302<br>(300–310)       | 377<br>(370–390)       |

<sup>1</sup> Ranges to TTS represent the model predictions in different areas and seasons within the Study Area. The zone in which animals are expected to suffer TTS extend from onset-PTS to the distance indicated. The average range to TTS is provided as well as the range from the estimated minimum to the maximum range to TTS in parentheses.

Notes: Ranges for 1-sec and 30-sec periods are identical for Bin MF1 because this system nominally pings every 50 seconds; therefore, these periods encompass only a single ping.

Notes: MF = mid-frequency, TTS = temporary threshold shift

**Table 3.4-8: Ranges to Temporary Threshold Shift for Sonar Bin MF4 over a Representative Range of Environments within the Study Area**

| <i>Hearing Group</i>     | <i>Approximate TTS Ranges (meters)<sup>1</sup></i> |                   |                   |                    |
|--------------------------|--|-------------------|-------------------|--------------------|
|                          | <i>Sonar Bin MF4</i>                               |                   |                   |                    |
|                          | <i>1 second</i>                                    | <i>30 seconds</i> | <i>60 seconds</i> | <i>120 seconds</i> |
| High-frequency cetaceans | 232<br>(220–260)                                   | 454<br>(420–600)  | 601<br>(575–875)  | 878<br>(800–1,525) |
| Low-frequency cetaceans  | 85<br>(85–90)                                      | 161<br>(160–170)  | 229<br>(220–250)  | 352<br>(330–410)   |
| Mid-frequency cetaceans  | 22<br>(22–22)                                      | 35<br>(35–35)     | 50<br>(45–50)     | 70<br>(70–70)      |

<sup>1</sup> Ranges to TTS represent the model predictions in different areas and seasons within the Study Area. The zone in which animals are expected to suffer TTS extend from onset-PTS to the distance indicated. The average range to TTS is provided as well as the range from the estimated minimum to the maximum range to TTS in parentheses.

Notes: MF = mid-frequency, TTS = temporary threshold shift

**Table 3.4-9: Ranges to Temporary Threshold Shift for Sonar Bin MF5 over a Representative Range of Environments within the Study Area**

| <i>Hearing Group</i>     | <i>Approximate TTS Ranges (meters)<sup>1</sup></i> |                   |                   |                    |
|--------------------------|--|-------------------|-------------------|--------------------|
|                          | <i>Sonar Bin MF5</i>                               |                   |                   |                    |
|                          | <i>1 second</i>                                    | <i>30 seconds</i> | <i>60 seconds</i> | <i>120 seconds</i> |
| High-frequency cetaceans | 114<br>(110–130)                                   | 114<br>(110–130)  | 168<br>(150–200)  | 249<br>(210–290)   |
| Low-frequency cetaceans  | 11<br>(10–12)                                      | 11<br>(10–12)     | 16<br>(16–17)     | 23<br>(23–24)      |
| Mid-frequency cetaceans  | 5<br>(0–9)   | 5<br>(0–9)        | 12<br>(11–13)     | 18<br>(17–18)      |

<sup>1</sup> Ranges to TTS represent the model predictions in different areas and seasons within the Study Area. The zone in which animals are expected to suffer TTS extend from onset-PTS to the distance indicated. The average range to TTS is provided as well as the range from the estimated minimum to the maximum range to TTS in parentheses.

Notes: MF = mid-frequency, TTS = temporary threshold shift

The range to received sound levels in 6-dB steps from five representative sonar bins and the percentage of animals that may exhibit a significant behavioral response under each behavioral response function are shown in Table 3.4-10 through Table 3.4-14, respectively. See Section 3.4.2.1.2.1 (Methods for Analyzing Impacts from Sonar and Other Transducers) for details on the derivation and use of the behavioral response functions, thresholds, and the cutoff distances.

**Table 3.4-10: Ranges to a Potentially Significant Behavioral Response for Sonar Bin HF4 over a Representative Range of Environments within the Study Area**

| <i>Received Level<br/>(dB re 1 µPa)</i> | <i>Mean Range (meters) with<br/>Minimum and Maximum<br/>Values in Parentheses</i> | <i>Probability of Behavioral Response for Sonar Bin HF4</i> |                  |                     |
|---|---|---|------------------|---------------------|
|   |   | <i>Odontocete</i>   | <i>Mysticete</i> | <i>Beaked Whale</i> |
| 196                                     | 3 (2–4)   | 100%  | 100%             | 100%                |
| 190                                     | 8 (6–10)  | 100%  | 98%              | 100%                |
| 184                                     | 16 (12–20)  | 99%   | 88%              | 100%                |
| 178                                     | 32 (24–40)  | 97%   | 59%              | 100%                |
| 172                                     | 63 (45–80)  | 91%   | 30%              | 99%                 |
| 166                                     | 120 (75–160)  | 78%   | 20%              | 97%                 |
| 160                                     | 225 (120–310)   | 58%   | 18%              | 93%                 |
| 154                                     | 392 (180–550)   | 40%   | 17%              | 83%                 |
| 148                                     | 642 (280–1,275)   | 29%   | 16%              | 66%                 |
| 142                                     | 916 (420–1,775)   | 25%   | 13%              | 45%                 |
| 136                                     | 1,359 (625–2,525)   | 23%   | 9%               | 28%                 |
| 130                                     | 1,821 (950–3,275)   | 20%   | 5%               | 18%                 |
| 124                                     | 2,567 (1,275–5,025)   | 17%   | 2%               | 14%                 |
| 118                                     | 3,457 (1,775–6,025)   | 12%   | 1%               | 12%                 |
| 112                                     | 4,269 (2,275–7,025)   | 6%  | 0%               | 11%                 |
| 106                                     | 5,300 (3,025–8,025)   | 3%  | 0%               | 11%                 |
| 100                                     | 6,254 (3,775–9,275)   | 1%  | 0%               | 8%                  |

Notes: dB re 1 µPa = decibels referenced to 1 micropascal, HF = high-frequency

**Table 3.4-11: Ranges to a Potentially Significant Behavioral Response for Sonar Bin LF4 over a Representative Range of Environments within the Study Area**

| Received Level<br>(dB re 1 $\mu$ Pa) | Mean Range (meters) with<br>Minimum and Maximum<br>Values in Parentheses | Probability of Behavioral Response for Sonar Bin LF4 |           |              |
|--------------------------------------|--|--|-----------|--------------|
|                                      |  | Odontocete   | Mysticete | Beaked Whale |
| 196                                  | 1 (1-1)  | 100%   | 100%      | 100%         |
| 190                                  | 3 (3-3)  | 100%   | 98%       | 100%         |
| 184                                  | 6 (6-6)  | 99%  | 88%       | 100%         |
| 178                                  | 12 (12-12)   | 97%  | 59%       | 100%         |
| 172                                  | 25 (25-25)   | 91%  | 30%       | 99%          |
| 166                                  | 51 (50-55)   | 78%  | 20%       | 97%          |
| 160                                  | 130 (130-160)  | 58%  | 18%       | 93%          |
| 154                                  | 272 (270-300)  | 40%  | 17%       | 83%          |
| 148                                  | 560 (550-675)  | 29%  | 16%       | 66%          |
| 142                                  | 1,048 (1,025-1,525)  | 25%  | 13%       | 45%          |
| 136                                  | 2,213 (1,525-4,525)  | 23%  | 9%        | 28%          |
| 130                                  | 4,550 (2,275-24,025)   | 20%  | 5%        | 18%          |
| 124                                  | 16,903 (4,025-66,275)  | 17%  | 2%        | 14%          |
| 118                                  | 43,256 (7,025-87,775)  | 12%  | 1%        | 12%          |
| 112                                  | 60,155 (7,775-100,000*)  | 6%   | 0%        | 11%          |
| 106                                  | 80,689 (8,775-100,000*)  | 3%   | 0%        | 11%          |
| 100                                  | 92,352 (9,025-100,000*)  | 1%   | 0%        | 8%           |

\* Indicates maximum range to which acoustic model was run, a distance of approximately 100 km from the sound source.

Notes: Cells are shaded if the mean range value for the specified received level exceeds the distance cutoff range for a particular hearing group. Any impacts within the cutoff range for a criteria group are included in the estimated impacts. Cut-off ranges in this table are for activities with high source levels and/or multiple platforms (see Table 3.4-3 for behavioral cut-off distances).

dB re 1  $\mu$ Pa = decibels referenced to 1 micropascal, LF = low-frequency

**Table 3.4-12: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF1 over a Representative Range of Environments within the Study Area**

| <i>Received Level<br/>(dB re 1 <math>\mu</math>Pa)</i> | <i>Mean Range (meters) with<br/>Minimum and Maximum<br/>Values in Parentheses</i> | <i>Probability of Behavioral Response for Sonar Bin MF1</i> |                  |                     |
|--|---|---|------------------|---------------------|
|  |   | <i>Odontocete</i>   | <i>Mysticete</i> | <i>Beaked Whale</i> |
| 196  | 106 (100–110)   | 100%  | 100%             | 100%                |
| 190  | 240 (240–250)   | 100%  | 98%              | 100%                |
| 184  | 501 (490–525)   | 99%   | 88%              | 100%                |
| 178  | 1,019 (975–1,025)   | 97%   | 59%              | 100%                |
| 172  | 3,275 (2,025–5,275)   | 91%   | 30%              | 99%                 |
| 166  | 7,506 (2,525–11,025)  | 78%   | 20%              | 97%                 |
| 160  | 15,261 (4,775–20,775)   | 58%   | 18%              | 93%                 |
| 154  | 27,759 (5,525–36,525)   | 40%   | 17%              | 83%                 |
| 148  | 43,166 (7,525–65,275)   | 29%   | 16%              | 66%                 |
| 142  | 58,781 (8,525–73,525)   | 25%   | 13%              | 45%                 |
| 136  | 71,561 (11,275–90,775)  | 23%   | 9%               | 28%                 |
| 130  | 83,711 (13,025–100,000*)  | 20%   | 5%               | 18%                 |
| 124  | 88,500 (23,525–100,000*)  | 17%   | 2%               | 14%                 |
| 118  | 90,601 (27,025–100,000*)  | 12%   | 1%               | 12%                 |
| 112  | 92,750 (27,025–100,000*)  | 6%  | 0%               | 11%                 |
| 106  | 94,469 (27,025–100,000*)  | 3%  | 0%               | 11%                 |
| 100  | 95,838 (27,025–100,000*)  | 1%  | 0%               | 8%                  |

\* Indicates maximum range to which acoustic model was run, a distance of approximately 100 km from the sound source.

Notes: Cells are shaded if the mean range value for the specified received level exceeds the distance cutoff range for a particular hearing group. Any impacts within the cutoff range for a criteria group are included in the estimated impacts. Cut-off ranges in this table are for activities with high source levels and/or multiple platforms (see Table 3.4-3 for behavioral cut-off distances).

dB re 1  $\mu$ Pa = decibels referenced to 1 micropascal, MF = mid-frequency

**Table 3.4-13: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF4 over a Representative Range of Environments within the Study Area**

| <i>Received Level<br/>(dB re 1 <math>\mu</math>Pa)</i> | <i>Mean Range (meters) with<br/>Minimum and Maximum<br/>Values in Parentheses</i> | <i>Probability of Behavioral Response for Sonar Bin MF4</i> |                  |                     |
|--|---|---|------------------|---------------------|
|  |   | <i>Odontocete</i>   | <i>Mysticete</i> | <i>Beaked Whale</i> |
| 196  | 8 (8–8)   | 100%  | 100%             | 100%                |
| 190  | 17 (17–17)  | 100%  | 98%              | 100%                |
| 184  | 35 (35–35)  | 99%   | 88%              | 100%                |
| 178  | 70 (65–70)  | 97%   | 59%              | 100%                |
| 172  | 141 (140–150)   | 91%   | 30%              | 99%                 |
| 166  | 354 (330–420)   | 78%   | 20%              | 97%                 |
| 160  | 773 (725–1,275)   | 58%   | 18%              | 93%                 |
| 154  | 1,489 (1,025–3,275)   | 40%   | 17%              | 83%                 |
| 148  | 3,106 (1,775–6,775)   | 29%   | 16%              | 66%                 |
| 142  | 8,982 (3,025–18,775)  | 25%   | 13%              | 45%                 |
| 136  | 15,659 (3,775–31,025)   | 23%   | 9%               | 28%                 |
| 130  | 25,228 (4,775–65,775)   | 20%   | 5%               | 18%                 |
| 124  | 41,778 (5,525–73,275)   | 17%   | 2%               | 14%                 |
| 118  | 51,832 (6,025–89,775)   | 12%   | 1%               | 12%                 |
| 112  | 62,390 (6,025–100,000*)   | 6%  | 0%               | 11%                 |
| 106  | 69,235 (6,775–100,000*)   | 3%  | 0%               | 11%                 |
| 100  | 73,656 (7,025–100,000*)   | 1%  | 0%               | 8%                  |

\* Indicates maximum range to which acoustic model was run, a distance of approximately 100 km from the sound source.

Notes: Cells are shaded if the mean range value for the specified received level exceeds the distance cutoff range for a particular hearing group. Any impacts within the cutoff range for a criteria group are included in the estimated impacts. Cut-off ranges in this table are for activities with high source levels and/or multiple platforms (see Table 3.4-3 for behavioral cut-off distances).

dB re 1  $\mu$ Pa = decibels referenced to 1 micropascal, MF = mid-frequency

**Table 3.4-14: Ranges to a Potentially Significant Behavioral Response for Sonar Bin MF5 over a Representative Range of Environments within the Study Area**

| Received Level<br>(dB re 1 $\mu$ Pa) | Mean Range (meters) with<br>Minimum and Maximum<br>Values in Parentheses | Probability of Behavioral Response for Sonar Bin MF5 |           |              |
|--------------------------------------|--|--|-----------|--------------|
|                                      |  | Odontocete   | Mysticete | Beaked Whale |
| 196                                  | 0 (0–0)  | 100%   | 100%      | 100%         |
| 190                                  | 1 (0–3)  | 100%   | 98%       | 100%         |
| 184                                  | 4 (0–7)  | 99%  | 88%       | 100%         |
| 178                                  | 14 (0–15)  | 97%  | 59%       | 100%         |
| 172                                  | 29 (0–30)  | 91%  | 30%       | 99%          |
| 166                                  | 58 (0–60)  | 78%  | 20%       | 97%          |
| 160                                  | 125 (0–150)  | 58%  | 18%       | 93%          |
| 154                                  | 284 (160–525)  | 40%  | 17%       | 83%          |
| 148                                  | 607 (450–1,025)  | 29%  | 16%       | 66%          |
| 142                                  | 1,213 (875–4,025)  | 25%  | 13%       | 45%          |
| 136                                  | 2,695 (1,275–7,025)  | 23%  | 9%        | 28%          |
| 130                                  | 6,301 (2,025–12,525)   | 20%  | 5%        | 18%          |
| 124                                  | 10,145 (3,025–19,525)  | 17%  | 2%        | 14%          |
| 118                                  | 14,359 (3,525–27,025)  | 12%  | 1%        | 12%          |
| 112                                  | 19,194 (3,525–37,275)  | 6%   | 0%        | 11%          |
| 106                                  | 24,153 (4,025–48,025)  | 3%   | 0%        | 11%          |
| 100                                  | 29,325 (5,025–57,775)  | 1%   | 0%        | 8%           |

Notes: Cells are shaded if the mean range value for the specified received level exceeds the distance cutoff range for a particular hearing group. Any impacts within the cutoff range for a criteria group are included in the estimated impacts. Cut-off ranges in this table are for activities with high source levels and/or multiple platforms (see Table 3.4-3 for behavioral cut-off distances).

dB re 1  $\mu$ Pa = decibels referenced to 1 micropascal, MF = mid-frequency

### 3.4.2.1.2.3 Impacts from Sonar and Other Transducers Under the Action Alternatives

Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. Use of sonar and other transducers would typically be transient and temporary. General categories and characteristics of sonar systems and the number of hours these sonars would be operated during training and testing under Alternative 1 and 2 are described in Section 3.0.4.1 (Acoustic Stressors). Activities using sonars and other transducers would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). The major differences between the action alternatives for the purposes of analyzing impacts on marine mammals are:

- Under Alternative 1, training and testing activities would fluctuate each year to account for the natural variation of training cycles and deployment schedules.
- Under Alternative 2, the same type and tempo of military training and testing activities would occur as Alternative 1, but there would be five Joint Multi-Strike Group Exercises (e.g., Valiant Shield) over any five-year period as compared to three under Alternative 1. Additionally, Alternative 2 contemplates three (vice two) Small Joint Coordinated ASW exercises (Multi-Sail/GUAMEX) per year with a 50 percent increase in associated unit-level events (e.g., Missile

Exercise (Surface-to-Air). This would result in an increase of active sonar training compared to Alternative 1. There would also be an increase in the use of active sonar during certain testing events. Alternative 2 reflects the maximum number of training and testing activities that could occur within a given year, and assumes that the maximum number of Fleet exercises would occur every year.

Compared to training and testing activities that use sonar and other transducers that were previously analyzed in the 2015 MITT Final EIS/OEIS under Alternatives 1 and 2, some training and testing activities would increase, decrease, or stay the same from those currently conducted (see Table 2.5-1 and Table 2.5-2 for details). In addition, some new systems using new technologies will be tested under the action alternatives.

Major training exercises are multi-day exercises that transition across large areas and involve multiple anti-submarine warfare assets. It is important to note that, while major training exercises focus on anti-submarine warfare, there are significant periods when active anti-submarine warfare sonars are not in use. Nevertheless, behavioral reactions are assumed more likely to be significant than during other anti-submarine warfare activities due to the duration (i.e., multiple days) and scale (i.e., multiple sonar platforms) of the major training exercises. Although major training exercises tend to move to different locations as the event unfolds, some animals could be exposed multiple times over the course of a few days.

Anti-submarine warfare activities also include unit-level training and coordinated/integrated training, and anti-submarine warfare sonar systems would be active when conducting surface ship and submarine sonar maintenance. Submarine and surface ship sonar maintenance activities involve the use of a single system in a limited manner; therefore, significant reactions to maintenance are less likely than with most other anti-submarine warfare activities. Furthermore, sonar maintenance activities typically occur either pierside or within entrances to harbors where higher levels of anthropogenic activity, including elevated noise levels, already exist. Unit-level training activities typically involve the use of a single vessel or aircraft and last for only a few hours over a small area of ocean. These unit-level training and sonar maintenance activities are limited in scope and duration; therefore, significant behavioral reactions are less likely than with other anti-submarine warfare activities with greater intensity and duration. Unit-level training activities are more likely to occur close to homeports and in the same general locations each time, so resident animals could be more frequently exposed to these types of activities. Coordinated/integrated exercises involve multiple assets and can last for several days transiting across large areas of a range complex. Repeated exposures to some individual marine mammals are likely during coordinated/integrated exercises. However, due to the shorter duration and smaller footprint compared to major training exercises, impacts from these activities are less likely to be significant with the possible exception of resident animals near homeports or Navy instrumented ranges that may incur some repeated exposures.

Anti-submarine warfare testing activities are typically similar to unit-level training. Vessel evaluation testing activities also use the same anti-submarine warfare sonars on ships and submarines. Testing activities that use anti-submarine warfare sonars typically occur in water deeper than approximately 200 m and therefore out of most nearshore habitats where productivity is typically higher (i.e., more food) and many marine mammals have higher abundances. Therefore, significant reactions to anti-submarine warfare and vessel evaluation testing activities are less likely than with larger anti-submarine warfare training activities discussed. Anti-submarine warfare and vessel evaluation testing activities are

more likely to occur close to homeports and testing facilities and in the same general locations each time, so resident animals could be more frequently exposed to these types of activities. These testing activities are limited in scope and duration; therefore, many of the impacts estimated by the quantitative analysis are unlikely to rise to the level of a significant behavioral response.

Mine warfare training activities typically involve a ship, helicopter, or unmanned vehicle using a mine-hunting sonar to locate mines. Most mine warfare sonar systems have a lower source level, higher frequency, and narrower, often downward facing beam pattern as compared to most anti-submarine warfare sonars. Significant reactions in marine mammals have not been reported due to exposure to mine warfare sonars. While individual animals could show short-term and minor responses to mine warfare sonar training activities, these reactions are very unlikely to lead to any costs or long-term consequences for individuals or populations.

Mine warfare testing activities typically involve a ship, helicopter, or unmanned vehicle testing a mine-hunting sonar system. Unmanned underwater vehicle testing also employs many of the same sonar systems as mine warfare testing and usually involves only a single sonar platform (i.e., unmanned underwater vehicle). Most of the sonar systems and other transducers used during these testing activities typically have a lower source level, higher frequency, and narrower, often downward facing beam pattern as compared to most anti-submarine warfare sonars. Significant reactions in marine mammals have not been reported due to exposure to these types of systems sonars. Animals are most likely to show short-term and minor to moderate responses to these testing activities; therefore, many of the impacts estimated by the quantitative analysis are unlikely to rise to the level of a significant behavioral response.

Navigation and object detection activities typically employ ship and submarine-based sonar systems and other transducers to navigate and avoid underwater objects. Significant reactions in marine mammals have not been reported due to exposure to most of the sonars and other transducers typically used in these activities. Some hull-mounted anti-submarine warfare sonars (e.g., bin MF1) have a mode to look for objects in the water such as mines, but this mode uses different source characteristics as compared to the anti-submarine warfare mode. Significant behavioral reactions have not been observed in relation to hull-mounted sonars using object-detection mode; however, significant reactions may be more likely than for all other sonar systems and transducers used within these activities due to the additional presence of a moving vessel and higher source levels. Individual animals could show short-term and minor to moderate responses to these systems, although these reactions are very unlikely to lead to any costs or long-term consequences for individuals or populations.

Other testing activities include testing of individual sonar systems and other transducers for performance and acoustic signature. Most sources used during these exercises have moderate source levels between 160 and 200 dB re 1  $\mu$ Pa @ 1m and are used for a limited duration, up to a few hours in most cases. Significant reactions in marine mammals have not been reported due to exposure to the sonars and other transducers typically used in these activities. Animals are most likely to show short-term and minor to moderate responses to these testing activities; therefore, many of the impacts estimated by the quantitative analysis are unlikely to rise to the level of a significant behavioral response.

Surface warfare activities require limited use of sonar or other transducers as compared to other types of activities discussed above, typically limited to the sonar targeting system of a few torpedoes. The limited scope and duration of sonar use in these activities makes significant behavioral reactions less

likely than with other activities that use anti-submarine warfare sonar systems and other transducers, which are discussed above.

### **Presentation of Estimated Impacts from the Quantitative Analysis**

The results of the analysis of potential impacts on marine mammals from sonars and other transducers (Section 3.4.2.1.2.1, Methods for Analyzing Impacts from Sonar and Other Transducers) are discussed below. The numbers of potential impacts estimated for individual species and stocks of marine mammals from exposure to sonar for training and testing activities under each action alternative are shown in Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) and presented below in figures for each species of marine mammal with any estimated effects (e.g., Figure 3.4-11). The activity categories that are most likely to cause impacts and the most likely region in which impacts could occur are represented in the impact graphics for each species. There is a potential for impacts to occur anywhere within the Study Area where sound from sonar and the species overlap, although only regions or activity categories where 0.5 percent of the impacts or greater are estimated to occur are graphically represented below. All (i.e., grand total) estimated impacts for that species are included, regardless of region or category.

It is important to note when examining the results of the quantitative analysis that the behavioral response functions used to predict the numbers of reactions in this analysis are largely derived from several studies (see Section 3.4.2.1.1.5, Behavioral Reactions). The best available science, including behavioral response studies, was used for deriving these criteria; however, many of the factors inherent in these studies that potentially increased the likelihood and severity of observed responses (e.g., close approaches by multiple vessels, tagging animals, and vectoring towards animals that have already begun avoiding the sound source) would not occur during Navy activities. Because the Navy purposely avoids approaching marine mammals, many of the behavioral responses estimated by the quantitative analysis are unlikely to occur or unlikely to rise to the severity observed during many of the behavioral response studies.

Although the statutory definition of Level B harassment for military readiness activities under the MMPA requires that the natural behavior patterns of a marine mammal be significantly altered or abandoned, the current state of science for determining those thresholds is somewhat unsettled. Therefore, in its analysis of impacts associated with acoustic sources, the Navy is adopting a conservative approach that overestimates the number of takes by Level B harassment. The responses estimated using the Navy's quantitative analysis are most likely to be moderate severity. Moderate severity responses would be considered significant if they were sustained for a duration long enough that it caused an animal to be outside of normal daily variations in feeding, reproduction, resting, migration/movement, or social cohesion. As discussed in Section 3.4.2.1.2.1 (Methods for Analyzing Impacts from Sonar and Other Transducers), the behavioral response functions used within the Navy's quantitative analysis were primarily derived from experiments using short-duration sound exposures lasting, in many cases, for less than 30 minutes. If animals exhibited moderate severity reactions for the duration of the exposure or longer, then it was conservatively assumed that the animal experienced a significant behavioral reaction. However, the experiments did not include measurements of costs to animals beyond the immediately observed reactions, and no direct correlations exist between an observed behavioral response and a cost that may result in long-term consequences. Within the Navy's quantitative analysis, many behavioral reactions are estimated from exposure to sound that may exceed an animal's behavioral threshold for only a single exposure to several minutes. It is likely that many of the estimated behavioral reactions within the Navy's quantitative analysis would not constitute significant behavioral

reactions; however, the numbers of significant versus non-significant behavioral reactions are currently impossible to predict. Consequently, there is a high likelihood that significant numbers of marine mammals exposed to acoustic sources are not significantly altering or abandoning their natural behavior patterns. As such, the overall impact of acoustic sources from military readiness activities on marine mammal species and stocks is negligible (i.e., cannot be reasonably expected to, and is not reasonably likely to, adversely affect the species or stocks through effects on annual rates of recruitment or survival).

### **Mysticetes**

Mysticetes may be exposed to sound from sonar and other transducers associated with training and testing activities. Most low- (less than 1 kHz) and mid- (1–10 kHz) frequency sonars and other transducers produce sounds that are likely to be within the hearing range of mysticetes (Section 3.4.1.6, Hearing and Vocalization). Some high-frequency sonars (greater than 10 kHz) also produce sounds that should be audible to mysticetes, although only smaller species of mysticetes such as minke whales are likely to be able to hear higher frequencies, presumably up to 30 kHz. Therefore, some high-frequency sonars and other transducers with frequency ranges between 10 and 30 kHz may also be audible to some mysticetes. If a sound is within an animal's hearing range then behavioral reactions, physiological stress, masking and hearing loss are potential impacts that must be analyzed. If a marine mammal cannot hear a sound, then behavioral reactions, physiological stress, masking, or hearing loss is not likely to occur. Impact ranges for mysticetes are discussed under low-frequency cetaceans in Section 3.4.2.1.2 (Impacts from Sonar and Other Transducer Stressors).

A few behavioral reactions in mysticetes resulting from exposure to sonar could take place at distances of up to 20 km. Behavioral reactions, however, are much more likely within a few kilometers of the sound source. As discussed above in *Assessing the Severity of Behavioral Responses from Sonar and other Transducers*, the quantitative analysis very likely overestimated the numbers of behavioral reactions due to the underlying nature of the data used to derive the behavioral response functions. Research shows that if mysticetes do respond they may react in a number of ways, depending on the characteristics of the sound source, their experience with the sound source, and whether they are migrating or on seasonal grounds (i.e., breeding or feeding). Behavioral reactions may include alerting, breaking off feeding dives and surfacing, or diving or swimming away. Overall, mysticetes have been observed to be more reactive to acoustic disturbance when a noise source is located directly on their migration route. Mysticetes disturbed while migrating could pause their migration or route around the disturbance. Animals disturbed while engaged in other activities such as feeding or reproductive behaviors may be more likely to ignore or tolerate the disturbance and continue their natural behavior patterns. Therefore, behavioral reactions from mysticetes are likely to be short-term and low to moderate severity.

Some mysticetes may avoid larger activities such as a major training exercise as it moves through an area. Vessels and aircraft associated with training or testing activities are typically in transit during an event (they are not stationary) and activities typically do not use the same training locations day after day during multi-day activities. If an event otherwise focuses on a fixed location, mysticetes may avoid the location of the activity for the duration of the event. If animals are displaced, they would likely return quickly after the event subsides. It is unlikely that most mysticetes would encounter a major training exercise more than once per year. In the ocean, the use of sonar and other transducers is transient and is unlikely to expose the same population of animals repeatedly over a short period except

around homeports and fixed instrumented ranges. Overall, a few behavioral reactions per year by a single individual are unlikely to produce long-term consequences for that individual.

Behavioral research indicates that mysticetes most likely avoid sound sources at levels that would cause any hearing loss (i.e., TTS) (Section 3.4.2.1.1.5, Behavioral Reactions). Therefore, it is likely that the quantitative analysis overestimates TTS in marine mammals because it does not account for animals avoiding sound sources at closer ranges. Mysticetes that do experience PTS or TTS from sonar sounds may have reduced ability to detect biologically important sounds around the frequency band of the sonar until their hearing recovers. Recovery from hearing loss begins almost immediately after the noise exposure ceases and can take a few minutes to a few days to fully recover, depending on the magnitude of the initial threshold shift. TTS would be recoverable and PTS would leave some residual hearing loss. Most TTS, if it does actually occur, would be more likely to be minor to moderate (i.e., less than 20 dB of TTS directly after the exposure) and would recover within a matter of minutes to hours (Section 3.4.2.1.1.2, Hearing Loss). Threshold shifts do not necessarily affect all hearing frequencies equally, and typically manifest themselves at the exposure frequency or within an octave above the exposure frequency. During the period that a mysticete had hearing loss, social calls from conspecifics could be more difficult to detect or interpret if they fell in the octave band of the sonar frequency. Killer whales are a primary predator of mysticetes. Some hearing loss could make killer whale calls more difficult to detect at farther ranges until hearing recovers. It is unclear how or if mysticetes use sound for finding prey or feeding; therefore, it is unknown whether hearing loss would affect a mysticete's ability to locate prey or rate of feeding. A single or even a few minor TTS (less than 20 dB of TTS) to an individual mysticete per year are unlikely to have any long-term consequences for that individual.

Research and observations of masking in marine mammals are discussed in Section 3.4.2.1.1.4 (Masking). Most anti-submarine warfare sonars and countermeasures use mid-frequency ranges and a few use low-frequency ranges. Most of these sonar signals are limited in the temporal, frequency, and spatial domains. The duration of most individual sounds is short, lasting up to a few seconds each. Some systems operate with higher duty cycles or nearly continuously, but typically use lower power. Nevertheless, masking may be more prevalent at closer ranges to these high-duty cycle and continuous active sonar systems. Most anti-submarine warfare activities are geographically dispersed and last for only a few hours, often with intermittent sonar use even within this period. Most anti-submarine warfare sonars also have a narrow frequency band (typically less than one-third octave). These factors reduce the likelihood of sources causing significant masking in mysticetes. High-frequency sonars are typically used for mine hunting, navigation, and object detection (avoidance). High-frequency (greater than 10 kHz) sonars fall outside of the best hearing and vocalization ranges of mysticetes (see Section 3.4.1.6, Hearing and Vocalization). Furthermore, high frequencies (above 10 kHz) attenuate more rapidly in the water due to absorption than do lower frequency signals, thus producing only a small zone of potential masking. Masking in mysticetes due to exposure to high-frequency sonar is unlikely. Potential costs to mysticetes from masking are similar to those discussed above for mild to moderate levels of TTS, with the primary difference being that the effects of masking are only present when the sound source (i.e., sonar) is actively pinging and the effect is over the moment the sound has ceased. By contrast, hearing loss lasts beyond the exposure for a period. Nevertheless, mysticetes that do experience some masking for a short period from low- or mid-frequency sonar may have their ability to communicate with conspecifics reduced, especially at further ranges. However, larger mysticetes (e.g., blue whale, fin whale, sei whale) communicate at frequencies below those of mid-frequency sonar and even most low-frequency sonars. Mysticetes that communicate at higher frequencies (e.g., minke whale) may be affected by some short-term and intermittent masking. Sounds from mid-frequency

sonar could mask killer whale vocalizations making them more difficult to detect, especially at further ranges. It is unknown whether masking would affect a mysticete's ability to feed since it is unclear how or if mysticetes use sound for finding prey or feeding. A single or even a few short periods of masking, if it were to occur, to an individual mysticete per year are unlikely to have any long-term consequences for that individual.

Many activities such as submarine under ice certification and most mine hunting exercises use only high-frequency sonars that are not within mysticetes' hearing range; therefore, there were no predicted effects. Section 3.4.1.6 (Hearing and Vocalization) discusses low-frequency cetacean (i.e., mysticetes) hearing abilities.


***Blue Whale (Endangered Species Act-Listed)***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Blue whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-11 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). For mysticetes, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of blue whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed blue whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-11: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Blue whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-12 or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as Alternative 1, although the numbers of impacts would increase slightly based on the increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of blue whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed blue whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-12: Blue Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Bryde's Whale***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Bryde's whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-13 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Bryde's whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-13: Bryde’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

***Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities***

Bryde’s whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-14 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Bryde’s whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-14: Bryde's Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**


***Fin Whale (Endangered Species Act-Listed)***

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Fin whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-15 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of fin whales incidental to those activities.

Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed fin whales.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-15: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**


**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Fin whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-16 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly

based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of fin whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed fin whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-16: Fin Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Humpback Whale (Endangered Species Act-Listed)***


**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Humpback whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-17 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) or tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact

Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of humpback whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed humpback whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-17: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**


**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Humpback whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-18 below or Appendix E (Estimated Marine Mammal and Sea

Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of humpback whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed humpback whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-18: Humpback Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Minke Whale**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Minke whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-19 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of minke whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-19: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Minke whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-20 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of minke whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-20: Minke Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Omura's Whale**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and testing Activities**

Omura's whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-21 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Omura's whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-21: Omura's Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Omura’s whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-22 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Omura’s whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-22: Omura’s Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**


***Sei Whale (Endangered Species Act-Listed)***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Sei whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-23 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of sei whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed sei whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-23: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Sei whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-24 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of sei whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed sei whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-24: Sei Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Odontocetes**

Odontocetes may be exposed to sound from sonar and other transducers associated with training and testing activities throughout the year. Low- (less than 1 kHz), mid- (1–10 kHz), high-frequency (10–100 kHz), and very high-frequency (100–200 kHz) sonars produce sounds that are likely to be within the audible range of odontocetes (see Section 3.4.1.6, Hearing and Vocalization). If a sound is within an animal’s hearing range then behavioral reactions, physiological stress, masking and hearing loss are potential impacts that must be analyzed. If a marine mammal cannot hear a sound, then behavioral reactions, physiological stress, masking, or hearing loss could not occur. Impact ranges for odontocetes are discussed under mid-frequency cetaceans in Section 3.4.2.1.2 (Impacts from Sonar and Other Transducer Stressors).

A few behavioral reactions in odontocetes (except beaked whales) resulting from exposure to sonar could take place at distances of up to 20 km. Beaked whales have demonstrated a high level of sensitivity to human-made noise and activity; therefore, the quantitative analysis assumes that some beaked whales could experience significant behavioral reactions at distance of up to 50 km from the sound source. Behavioral reactions, however, are much more likely within a few kilometers of the sound source for most species of odontocetes such as delphinids and sperm whales. Even for beaked whales,

as discussed above in *Assessing the Severity of Behavioral Responses from Sonar*, the quantitative analysis very likely overestimated the numbers of behavioral reactions due to the underlying nature of the data used to derive the behavioral response functions.

Research shows that if odontocetes do respond they may react in a number of ways, depending on the characteristics of the sound source and their experience with the sound source. Behavioral reactions may include alerting; breaking off feeding dives and surfacing; or diving or swimming away. Animals disturbed while engaged in other activities such as feeding or reproductive behaviors may be more likely to ignore or tolerate the disturbance and continue their natural behavior patterns. Therefore, most behavioral reactions from odontocetes are likely to be short-term and low to moderate severity.

Large odontocetes such as killer whales and pilot whales have been the subject of behavioral response studies (see Section 3.4.2.1.1.5, Behavioral Reactions). Based on these studies, a number of reactions could occur such as a short-term cessation of natural behavior such as feeding, avoidance of the sound source, or even attraction towards the sound source as seen in pilot whales. Due to the factors involved in Navy training exercises versus the conditions under which pilot whales and killer whales were exposed during behavioral response studies, large odontocetes are unlikely to have more than short-term and moderate severity reactions to sounds from sonar or other human disturbance, and typically only at ranges within a few kilometers. Most estimated impacts are due to anti-submarine warfare activities, which could vary in duration and intensity. Anti-submarine warfare unit-level exercises and maintenance typically last for a matter of a few hours and involves a limited amount of sonar use so significant responses would be less likely than with longer and more intense exercises (more sonar systems and vessels). Coordinated/integrated anti-submarine warfare exercises involve multiple sonar systems and can last for a period of days, making significant response more likely. A single or few short-lived TTS or behavioral reactions per year are unlikely to have any significant costs or long-term consequences for individuals.

Small odontocetes have been the subject of behavioral response studies and observations in the field (see Section 3.4.2.1.1.5, Behavioral Reactions). Based on these studies, small odontocetes (dolphins) appear to be less sensitive to sound and human disturbance than other cetacean species. If reactions did occur, they could consist of a short-term behavior response such as cessation of feeding, avoidance of the sound source, or even attraction towards the sound source. Small odontocetes are unlikely to have more than short-term and moderate severity reactions to sounds from sonar or other human disturbance, and typically only at ranges within a few kilometers. Most estimated impacts are due to anti-submarine warfare activities, which could vary in duration and intensity. Anti-submarine warfare unit-level exercises and maintenance typically last for a matter of a few hours and involve a limited amount of sonar use so significant responses would be less likely than with longer and more intense exercises (more sonar systems and vessels). Coordinated/integrated anti-submarine warfare exercises involve multiple sonar systems and can last for a period of days, making significant response more likely. Some bottlenose dolphin estimated impacts could also occur due to navigation and object avoidance (detection) since these activities typically occur entering and leaving Navy homeports that overlap the distribution of coastal populations of this species. Navigation and object avoidance (detection) activities normally involve a single ship or submarine using a limited amount of sonar, therefore significant reactions are unlikely. A single or few short-lived TTS or behavioral reactions per year are unlikely to have any significant costs or long-term consequences for individuals.

Some odontocetes may avoid larger activities such as a major training exercise as it moves through an area. Vessels and aircraft associated with training or testing activities are typically in transit during an

event (they are not stationary) and activities typically do not use the same training locations day-after-day during multi-day activities. If an event otherwise focuses on a fixed location, sensitive species of odontocetes, such as beaked whales, may avoid the location of the activity for the duration of the event. Section 3.4.2.1.1.5 (Behavioral Reactions) discusses these species' observed reactions to sonar and other transducers. If animals are displaced, they would likely return after the sonar activity subsides within an area, as seen in Blainville's beaked whales in the Bahamas (Tyack et al., 2011) and Hawaii (Henderson et al., 2015b; Henderson et al., 2016; Manzano-Roth et al., 2016). This would allow the animal to recover from any energy expenditure or missed resources, reducing the likelihood of long-term consequences for the individual. It is unlikely that most individuals would encounter a major training exercise more than once per year due to where these activities are typically conducted. Outside of Navy instrumented ranges and homeports, the use of sonar and other transducers is transient and is unlikely to expose the same population of animals repeatedly over a short period. However, a few behavioral reactions per year from a single individual are unlikely to produce long-term consequences for that individual.

Behavioral research indicates that most odontocetes avoid sound sources at levels that would cause any temporary hearing loss (i.e., TTS) (see Section 3.4.2.1.1.5, Behavioral Reactions). TTS and even PTS is more likely for high-frequency cetaceans, such as Kogia whales, because hearing loss thresholds for these animals are lower than for all other marine mammals. These species have demonstrated a high level of sensitivity to human-made sound and activities and may avoid at further distances. This increased distance could avoid or minimize hearing loss for these species as well, especially as compared to the estimates from the quantitative analysis. Therefore, it is likely that the quantitative analysis overestimates TTS and PTS in marine mammals because it does not account for animals avoiding sound sources at closer ranges. Recovery from hearing loss begins almost immediately after the noise exposure ceases and can take a few minutes to a few days to fully recover, depending on the magnitude of the initial threshold shift. TTS would be recoverable and PTS would leave some residual hearing loss. Most TTS, if it does actually occur, would be more likely to be minor to moderate (i.e., less than 20 dB of TTS directly after the exposure) and would recover within a matter of minutes to hours. Threshold shifts do not necessarily affect all hearing frequencies equally, and typically manifest themselves at the exposure frequency or within an octave above the exposure frequency. During the period that an odontocete had hearing loss, social calls from conspecifics could be more difficult to detect or interpret. Killer whales are a primary predator of odontocetes. Some hearing loss could make killer whale calls more difficult to detect at further ranges until hearing recovers. Odontocetes use echolocation clicks to find and capture prey. These echolocation clicks and vocalizations are at frequencies above a few tens of kHz for delphinids, beaked whales, and sperm whales, and above 100 kHz for Kogia whales. Therefore, echolocation associated with feeding and navigation in odontocetes is unlikely to be affected by threshold shift at lower frequencies and should not have any significant effect on an odontocete's ability to locate prey or navigate, even in the short-term. Therefore, a single or even a few minor TTS (less than 20 dB of TTS) to an individual odontocete per year are unlikely to have any long-term consequences for that individual. Minor PTS (a few dB or less) in an individual could have no to minor long-term consequences for individuals.

Research and observations of masking in marine mammals are discussed in Section 3.4.2.1.1.4 (Masking). Many anti-submarine warfare sonars and countermeasures use low- and mid-frequency sonar. Most low- and mid-frequency sonar signals (i.e., sounds) are limited in their temporal, frequency, and spatial domains. The duration of most individual sounds is short, lasting up to a few seconds each. Some systems operate with higher duty cycles or nearly continuously, but typically use lower power. Nevertheless, masking may be more prevalent at closer ranges to these high-duty cycle and continuous

active sonar systems. Most anti-submarine warfare activities are geographically dispersed and last for only a few hours, often with intermittent sonar use even within this period. Most anti-submarine warfare sonars also have a narrow frequency band (typically much less than one-third octave). These factors reduce the likelihood of sources causing significant masking in odontocetes due to exposure to sonar used during anti-submarine warfare activities. Odontocetes may experience some limited masking at closer ranges from high-frequency sonars and other transducers; however, the frequency band of the sonar is narrow, limiting the likelihood of masking. High-frequency sonars are typically used for mine hunting, navigation, and object detection (avoidance). Potential costs to odontocetes from masking are similar to those discussed above for mild to moderate levels of TTS, with the primary difference being that the effects of masking are only present when the sound source (i.e., sonar) is actively pinging and the effect is over the moment the sound has ceased.

Nevertheless, odontocetes that do experience some masking from sonar or other transducers may have their ability to communicate with conspecifics reduced, especially at further ranges. Sounds from mid-frequency sonar could mask killer whale vocalizations making them more difficult to detect, especially at further ranges. As discussed above for TTS, odontocetes use echolocation to find prey and navigate. The echolocation clicks of odontocetes are above the frequencies of most sonar systems, especially those used during anti-submarine warfare. Therefore, echolocation associated with feeding and navigation in odontocetes is unlikely to be masked by sounds from sonars or other transducers. A single or even a few short periods of masking, if it were to occur, to an individual odontocete per year are unlikely to have any long-term consequences for that individual.

#### ***Beaked Whales***

Beaked whales within the Study Area include: Blainville's beaked whale, Cuvier's beaked whale, ginkgo-toothed beaked whale, and Longman's beaked whale. As discussed above for odontocetes overall, the quantitative analysis overestimates hearing loss in marine mammals because behavioral response research has shown that most marine mammals are likely to avoid sound levels that could cause more than minor to moderate TTS (6–20 dB). Specifically for beaked whales, behavioral response research discussed below and in Section 3.4.2.1.1.5 (Behavioral Reactions) has demonstrated that beaked whales are sensitive to sound from sonars and usually avoid sound sources by 10 or more kilometers. These are well beyond the ranges to TTS for mid-frequency cetaceans such as beaked whales. Therefore, any TTS predicted by the quantitative analysis is unlikely to occur in beaked whales.

Research and observations (Section 3.4.2.1.1.5, Behavioral Reactions) show that if beaked whales are exposed to sonar or other transducers they may startle, break off feeding dives, and avoid the area of the sound source at levels ranging between 95 and 157 dB re 1  $\mu$ Pa (McCarthy et al., 2011). Furthermore, in research done at the Navy's fixed tracking range in the Bahamas and Hawaii, animals leave the immediate area of the anti-submarine warfare training exercise but return within a few days after the event ends (Henderson et al., 2015b; Henderson et al., 2016; Manzano-Roth et al., 2016; Tyack et al., 2011). Populations of beaked whales and other odontocetes on Navy fixed ranges that have been operating for decades appear to be stable, and analysis is ongoing. Significant behavioral reactions seem likely in most cases if beaked whales are exposed to anti-submarine sonar within a few tens of kilometers, especially for prolonged periods (a few hours or more) since this is one of the most sensitive marine mammal groups to human-made sound of any species or group studied to date.


Based on the best available science, the Navy believes that beaked whales that exhibit a significant behavioral reaction due to sonar and other transducers would generally not have long-term consequences for individuals or populations. However, because of a lack of scientific consensus

regarding the causal link between sonar and stranding events, NMFS has stated in a letter to the Navy dated October 2006 that it “cannot conclude with certainty the degree to which mitigation measures would eliminate or reduce the potential for serious injury or mortality.” The Navy does not anticipate that marine mammal strandings or mortality would result from the operation of sonar during Navy exercises within the Study Area. Additionally, through the MMPA process (which allows for adaptive management), NMFS and the Navy will determine the appropriate way to proceed in the event that a causal relationship were to be found between Navy activities and a future stranding.

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**


Beaked whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-25 through Figure 3.4-28 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Blainville’s, Cuvier’s, ginkgo-toothed, and Longman’s beaked whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-25: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-26: Cuvier's Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-27: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-28: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

***Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities***


Beaked whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-29 through Figure 3.4-32 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Blainville’s, Cuvier’s, ginkgo-toothed, and Longman’s beaked whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-29: Blainville’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-30: Cuvier's Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-31: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-32: Longman’s Beaked Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Common Bottlenose Dolphin***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Bottlenose dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-33 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of bottlenose dolphins incidental to those activities.*




Note: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-33: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

***Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities***

Bottlenose dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-34 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of bottlenose dolphins incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-34: Bottlenose Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Kogia Whales***


Kogia whales include two species that are often difficult to distinguish from one another: dwarf sperm whales and pygmy sperm whales; however, impacts to the populations of dwarf and pygmy sperm whales are modeled separately. TTS and PTS thresholds for high-frequency cetaceans, such as Kogia whales are lower than for all other marine mammals, which leads to a higher number of estimated hearing loss impacts relative to the number of animals exposed to the sound as compared to other hearing groups (e.g., mid-frequency cetaceans).

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Kogia whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions, TTS, and


PTS under Alternative 1. See Figure 3.4-35 and Figure 3.4-36 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. PTS could reduce an animal’s ability to detect biologically important sounds; however, as discussed above, a small threshold shift due to exposure to sonar is unlikely to affect the hearing range that Kogia whales rely upon. Nevertheless, PTS could have minor long-term consequences for individuals. This minor consequence for an individual is unlikely to have any long-term consequences for the species. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Kogia whales (i.e., dwarf and pygmy sperm whales) incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-35: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-36: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**


Kogia whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions, TTS, and PTS under Alternative 2. See Figure 3.4-37 and Figure 3.4-38 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Kogia whales (i.e., dwarf and pygmy sperm whales) incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-37: Dwarf Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts.  
(2) MIRC = Mariana Islands Range Complex


**Figure 3.4-38: Pygmy Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***False Killer Whale***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

False killer whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-39 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of false killer whales incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-39: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

False killer whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-40 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of false killer whales incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-40: False Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Fraser’s Dolphin**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Fraser’s dolphin may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-41 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Fraser’s dolphin incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-41: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Fraser’s dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-42 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Fraser’s dolphin incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-42: Fraser’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

*Killer Whale*

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Killer whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-43 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of killer whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-43: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Killer whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-44 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of killer whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-44: Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Melon-Headed Whale**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Melon-headed whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-45 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of melon-headed whales incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-45: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Melon-headed whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-46 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of melon-headed whales incidental to those activities.



Note: Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-46: Melon-Headed Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Pantropical Spotted Dolphin***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Pantropical spotted dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-47 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of pantropical spotted dolphins incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-47: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Pantropical spotted dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-48 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of pantropical spotted dolphins incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-48: Pantropical Spotted Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Pygmy Killer Whale***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Pygmy killer whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-49 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of pygmy killer whales incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-49: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Pygmy killer whale may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-50 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of pygmy killer whales incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-50: Pygmy Killer Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Risso's Dolphin***

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Risso's dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-51 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of Risso’s dolphins incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-51: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Risso’s dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-52 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2 versus Alternative 1.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of Risso’s dolphins incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-52: Risso’s Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

***Rough-Toothed Dolphin***

***Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities***

Rough-toothed dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-53 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of rough-toothed dolphins incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-53: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Rough-toothed dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-54 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of rough-toothed dolphins incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-54: Rough-Toothed Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**Short-Finned Pilot Whale**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Short-finned pilot whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-55 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of short-finned pilot whales incidental to those activities.




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-55: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Short-finned pilot whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-56 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of short-finned pilot whales incidental to those activities.



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-56: Short-Finned Pilot Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**


***Sperm Whale (Endangered Species Act-Listed)***

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Sperm whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-57 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of sperm whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed sperm whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-57: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**


**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Sperm whales may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-58 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly

based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of sperm whales incidental to those activities.*

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed sperm whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-58: Sperm Whale Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**


**Spinner Dolphin**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Spinner dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-59 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact

Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of spinner dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-59: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Spinner dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-60 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact

Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of spinner dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-60: Spinner Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**


**Striped Dolphin**

**Impacts from Sonar and Other Transducers Under Alternative 1 for Training and Testing Activities**

Striped dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 1. See Figure 3.4-61 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact

Ranges for Sonar and Other Transducers). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 would result in the unintentional taking of striped dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-61: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 1**

**Impacts from Sonar and Other Transducers Under Alternative 2 for Training and Testing Activities**

Striped dolphins may be exposed to sounds from sonar and other transducers associated with training and testing activities throughout the year. The quantitative analysis estimates behavioral reactions and TTS under Alternative 2. See Figure 3.4-62 below or Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) for tabular results. Impact ranges for this species are discussed in Section 3.4.2.1.2.2 (Impact

Ranges for Sonar and Other Transducers). Potential impacts under Alternative 2 from sonar and other transducers would be similar in type as for Alternative 1, although the numbers of impacts would increase slightly based on the slight increase in sonar use associated with training and testing activities under Alternative 2.

*Pursuant to the MMPA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 would result in the unintentional taking of striped dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. No PTS is estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-62: Striped Dolphin Impacts Estimated per Year from Sonar and Other Transducers Used During Training and Testing Under Alternative 2**

**3.4.2.1.2.4 Impacts from Sonar and Other Transducers Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors from the use of sonar and other transducers, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that use sonar and other

transducers within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from sonar and other transducers on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

#### **3.4.2.1.3 Impacts from Vessel Noise**

Marine mammals may be exposed to noise from vessel movement. A detailed description of the acoustic characteristics and typical sound levels of vessel noise are in Section 3.0.4.1.2 (Vessel Noise). Vessel movements involve transits to and from ports to various locations within the Study Area, including commercial ship traffic as well as recreational vessels in addition to U.S. Navy vessels. Many ongoing and proposed training and testing activities within the Study Area involve maneuvers by various types of surface ships, boats, and submarines (collectively referred to as vessels). Section 3.4.2.1.1 (Background) summarizes and synthesizes available information on behavioral reactions, masking, and physiological stress due to noise exposure, including vessel noise (Section 3.4.2.1.1.2, Hearing Loss; Section 3.4.2.1.1.3, Physiological Stress; Section 3.4.2.1.1.4, Masking; and Section 3.4.2.1.1.5, Behavioral Reactions).

Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Table 2.5-1 and Table 2.5-2 for proposed activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The Navy will implement mitigation measures for vessel movement to avoid the potential for marine mammal vessel strikes, as discussed in Section 5.3.4.1 (Vessel Movement). The mitigation for vessel movement (i.e., maneuvering to maintain a specified distance from a marine mammal) will also help the Navy avoid or reduce potential impacts from vessel noise on marine mammals.

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors (i.e., vessel noise) from the use of vessels, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that produce vessel noise within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from vessel noise on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

*Pursuant to the MMPA, vessel noise during training and testing activities as described under Alternative 1 and Alternative 2 would not result in the unintentional taking of marine mammals.*

*Pursuant to the ESA, vessel noise during training and testing activities as described under Alternative 1 and Alternative 2 may affect ESA-listed marine mammals.*

#### **3.4.2.1.4 Impacts from Aircraft Noise**

Marine mammals may be exposed to aircraft-generated noise throughout the Study Area. Fixed- and rotary-wing aircraft are used for a variety of training and testing activities throughout the Study Area. Tilt-rotor impacts would be similar to fixed-wing or helicopter impacts depending which mode the aircraft is in. Most of these sounds would be concentrated around airbases and fixed ranges within the

range complex. Aircraft produce extensive airborne noise from either turbofan or turbojet engines. An infrequent type of aircraft noise is the sonic boom, produced when the aircraft exceeds the speed of sound. Rotary-wing aircraft produce low-frequency sound and vibration (Pepper et al., 2003). Section 3.4.2.1.1 (Background) summarizes and synthesizes available information on behavioral reactions, masking, and physiological stress due to noise exposure, including aircraft noise (Section 3.4.2.1.1.2, Hearing Loss; Section 3.4.2.1.1.3, Physiological Stress; Section 3.4.2.1.1.4, Masking; and Section 3.4.2.1.1.5, Behavioral Reactions).

A detailed description of aircraft noise as a stressor is in Section 3.0.4.1.3 (Aircraft Noise). Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Table 2.5-1 and Table 2.5-2 for proposed training and testing activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors (i.e., aircraft noise) from the use of aircraft, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that produce aircraft noise within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from aircraft noise on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

*Pursuant to the MMPA, aircraft noise during training and testing activities as described under Alternative 1 and Alternative 2 would not result in the unintentional taking of marine mammals.*

*Pursuant to the ESA, aircraft noise during training and testing activities as described under Alternative 1 and Alternative 2 may affect ESA-listed marine mammals.*

#### **3.4.2.1.5 Impacts from Weapon Noise**

Marine mammals may be exposed to sounds caused by the firing of weapons, objects in flight, and inert impact of non-explosive munitions on the water's surface, which are described in Section 3.0.4.1.4 (Weapon Noise). In general, these are impulsive sounds generated in close vicinity to or at the water surface, with the exception of items that are launched underwater. The firing of a weapon may have several components of associated noise. Firing of guns could include sound generated in air by firing a gun (muzzle blast) and a crack sound due to a low amplitude shock wave generated by a supersonic projectile flying through the air. Most in-air sound would be reflected at the air-water interface.

Underwater sounds would be strongest just below the surface and directly under the firing point. Any sound that enters the water only does so within a narrow cone below the firing point or path of the projectile. Vibration from the blast propagating through a ship's hull, the sound generated by the impact of an object with the water surface, and the sound generated by launching an object underwater are other sources of impulsive sound in the water. Sound due to missile and target launches is typically at a maximum at initiation of the booster rocket and rapidly fades as the missile or target travels downrange.

Section 3.4.2.1.1 (Background) summarizes and synthesizes available information on behavioral reactions, masking, and physiological stress due to noise exposure, including aircraft noise (Section

3.4.2.1.1.2, Hearing Loss; Section 3.4.2.1.1.3, Physiological Stress; Section 3.4.2.1.1.4, Masking; and Section 3.4.2.1.1.5, Behavioral Reactions).

Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Table 2.5-1 and Table 2.5-2 for proposed training and testing activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The Navy will implement mitigation measures to avoid or reduce potential impacts from weapon noise during large-caliber gunnery activities, as discussed in Section 5.3.2.2 (Weapons Firing Noise).

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur; however, weapon noise would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Under the No Action Alternative, discontinuing training and testing activities that produce weapon noise within the marine environment where training and testing activities have historically been conducted would reduce the potential for impacts from weapon noise on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

*Pursuant to the MMPA, weapon noise during training and testing activities as described under Alternative 1 and Alternative 2 would not result in the unintentional taking of marine mammals.*

*Pursuant to the ESA, weapon noise during training and testing activities as described under Alternative 1 and Alternative 2 may affect ESA-listed marine mammals.*

### **3.4.2.2 Explosive Stressors**

Assessing whether an explosive detonation may disturb or injure a marine mammal involves understanding the characteristics of the explosive sources, the marine mammals that may be present near the sources, the physiological effects of a close explosive exposure, and the effects of impulsive sound on marine mammal hearing and behavior. Many other factors besides the received level or pressure wave of an explosion, such as the animal's physical condition and size, prior experience with the explosive sound, and proximity to the explosion, may influence physiological effects and behavioral reactions.

The ways in which an explosive exposure could result in immediate effects or lead to long-term consequences for an animal are explained in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities). The following background section discusses what is currently known about explosive effects to marine mammals.

Due to new acoustic impact criteria, marine mammal densities, and revisions to the acoustics effects model, the analysis provided in Section 3.4.2.2.2 (Impacts from Explosive Stressors) of this SEIS/OEIS supplants the 2015 MITT Final EIS/OEIS for marine mammals and changes estimated impacts for some species since the 2015 MITT Final EIS/OEIS.

#### **3.4.2.2.1 Background**

##### **3.4.2.2.1.1 Injury**

Injury refers to the direct effects on the tissues or organs of an animal due to exposure to pressure waves. Injury in marine mammals can be caused directly by exposure to explosions. Section 3.0.4.7

(Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) provides additional information on injury and the framework used to analyze this potential impact.

### **Injury due to Explosives**

Explosive injury to marine mammals would consist of primary blast injury, which refers to those injuries that result from the compression of a body exposed to a blast wave and is usually observed as barotrauma of gas-containing structures (e.g., lung and gut) and structural damage to the auditory system (Greaves et al., 1943; Office of the Surgeon General, 1991; Richmond et al., 1973). The near instantaneous high magnitude pressure change near an explosion can injure an animal where tissue material properties significantly differ from the surrounding environment, such as around air-filled cavities such as in the lungs or gastrointestinal tract. Large pressure changes at tissue-air interfaces in the lungs and gastrointestinal tract may cause tissue rupture, resulting in a range of injuries depending on degree of exposure. The lungs are typically the first site to show any damage, while the solid organs (e.g., liver, spleen, and kidney) are more resistant to blast injury (Clark & Ward, 1943). Recoverable injuries would include slight lung injury, such as capillary interstitial bleeding, and contusions to the gastrointestinal tract. More severe injuries, such as tissue lacerations, major hemorrhage, organ rupture, or air in the chest cavity (pneumothorax), would significantly reduce fitness and likely cause death in the wild. Rupture of the lung may also introduce air into the vascular system, producing air emboli that can cause a stroke or heart attack by restricting oxygen delivery to critical organs.

If an animal is exposed to an explosive blast underwater, the likelihood of injury depends on the charge size, the geometry of the exposure (distance to the charge, depth of the animal and the charge), and the size of the animal. In general, an animal would be less susceptible to injury near the water surface because the pressure wave reflected from the water surface would interfere with the direct path pressure wave, reducing positive pressure exposure. Susceptibility would increase with depth, until normal lung collapse (due to increasing hydrostatic pressure) and increasing ambient pressures again reduce susceptibility. See Appendix H (Acoustic and Explosive Concepts) for an overview of explosive propagation and an explanation of explosive effects on gas cavities.

The only known occurrence of mortality or injury to a marine mammal due to a Navy training or testing event involving explosives occurred in March 2011 in nearshore waters off San Diego, California, at the Silver Strand Training Complex. This area had been used for underwater demolitions training for at least three decades without prior known incident. On this occasion, however, a group of approximately 100–150 long-beaked common dolphins entered the mitigation zone surrounding an area where a time-delayed firing device had been initiated on an explosive with a net explosive weight of 8.76 pounds (lb.) (3.97 kilograms [kg]) placed at a depth of 48 ft. (14.6 m). Approximately one minute after detonation, three animals were observed dead at the surface. The Navy recovered those animals and transferred them to the local stranding network for necropsy. A fourth animal was discovered stranded and dead 42 NM to the north of the detonation three days later. It is unknown exactly how close those four animals were to the detonation. Upon necropsy, all four animals were found to have sustained typical mammalian primary blast injuries (Danil & St Leger, 2011).

Relatively little is known about auditory system trauma in marine mammals resulting from explosive exposure, although it is assumed that auditory structures would be vulnerable to blast injuries. Auditory trauma was found in two humpback whales that died following the detonation of a 5,000 kg explosive used off Newfoundland during demolition of an offshore oil rig platform (Ketten et al., 1993), but the proximity of the whales to the detonation was unknown. Eardrum rupture was examined in submerged

terrestrial mammals exposed to underwater explosions (Richmond et al., 1973; Yelverton et al., 1973); however, results may not be applicable to the anatomical adaptations for underwater hearing in marine mammals. In this discussion, primary blast injury to auditory tissues is considered gross structural tissue damage distinct from threshold shift or other auditory effects (see Section 3.4.2.2.1.2, Hearing Loss).

Controlled tests with a variety of lab animals (mice, rats, dogs, pigs, sheep, and other species) are the best data sources on actual injury to mammals due to underwater exposure to explosions. In the early 1970s, the Lovelace Foundation for Medical Education and Research conducted a series of tests in an artificial pond at Kirtland Air Force Base, New Mexico, to determine the effects of underwater explosions on mammals, with the goal of determining safe ranges for human divers. The resulting data were summarized in two reports (Richmond et al., 1973; Yelverton et al., 1973). Specific physiological observations for each test animal are documented in Richmond et al. (1973). Gas-containing internal organs, such as lungs and intestines, were the principal damage sites in submerged terrestrial mammals; this is consistent with earlier studies of mammal exposures to underwater explosions in which lungs were consistently the first areas to show damage, with less consistent damage observed in the gastrointestinal tract (Clark & Ward, 1943; Greaves et al., 1943). Results from all of these tests suggest two explosive metrics are predictive of explosive injury: peak pressure and impulse.

#### ***Impulse as a Predictor of Explosive Injury***

In the Lovelace studies, acoustic impulse was found to be the metric most related to degree of injury, and the size of an animal's gas-containing cavities was thought to play a role in blast injury susceptibility. The lungs of most marine mammals are similar in proportion to overall body size as those of terrestrial mammals, so the magnitude of lung damage in the tests may approximate the magnitude of injury to marine mammals when scaled for body size. Within the marine mammals, mysticetes and deeper divers (e.g., Kogiidae, Physeteridae, Ziphiidae) tend to have lung-to-body size ratios that are smaller and more similar to terrestrial animal ratios than the shallow diving odontocetes (e.g., Phocoenidae, Delphinidae) and pinnipeds (Fahlman et al., 2014a; Piscitelli et al., 2010). The use of test data with smaller lung-to-body ratios to set injury thresholds may result in a more conservative estimate of potential for damaging effects (i.e., lower thresholds) for animals with larger lung-to-body ratios.

For these shallow exposures of small terrestrial mammals (masses ranging from 3.4 to 50 kg) to underwater detonations, Richmond et al. (1973) reported that no blast injuries were observed when exposures were less than 6 pounds per square inch (psi) per millisecond (psi-ms) (40 pascals second [Pa-s]), no instances of slight lung hemorrhage occurred below 20 psi-ms (140 Pa-s), and instances of no lung damage were observed in some exposures at higher levels up to 40 psi-ms (280 Pa-s). An impulse of 34 psi-ms (230 Pa-s) resulted in about 50 percent incidence of slight lung hemorrhage. About half of the animals had gastrointestinal tract contusions (with slight ulceration, i.e., some perforation of the mucosal layer) at exposures of 25–27 psi-ms (170–190 Pa-s). Lung injuries were found to be slightly more prevalent than gastrointestinal tract injuries for the same exposure.

The Lovelace subject animals were exposed near the water surface; therefore, depth effects were not discernible in this data set. In addition, this data set included only small terrestrial animals, whereas marine mammals may be several orders of magnitude larger and have respiratory structures adapted for the high pressures experienced at depth. Goertner (1982) examined how lung cavity size would affect susceptibility to blast injury by considering both marine mammal size and depth in a bubble oscillation model of the lung. Animal depth relates to injury susceptibility in two ways: injury is related to the relative increase in explosive pressure over hydrostatic pressure, and lung collapse with depth reduces the potential for air cavity oscillatory damage. The period over which an impulse must be delivered to

cause damage is assumed to be related to the natural oscillation period of an animal's lung, which depends on lung size.

Because gas-containing organs are more vulnerable to primary blast injury, adaptations for diving that allow for collapse of lung tissues with depth may make animals less vulnerable to lung injury with depth. Adaptations for diving include a flexible thoracic cavity, distensible veins that can fill space as air compresses, elastic lung tissue, and resilient tracheas with interlocking cartilaginous rings that provide strength and flexibility (Ridgway, 1972). Older literature suggested complete lung collapse depths at approximately 70 m for dolphins (Ridgway & Howard, 1979) and 20–50 m for phocid seals (Falke et al., 1985; Kooyman et al., 1972). Follow-on work by Kooyman and Sinnott (1982), in which pulmonary shunting was studied in harbor seals and sea lions, suggested that complete lung collapse for these species would be about 170 m and about 180 m, respectively. More recently, evidence in sea lions suggests that complete collapse might not occur until depths as great as 225 m; although the depth of collapse and depth of the dive are related, sea lions can affect the depth of lung collapse by varying the amount of air inhaled on a dive (McDonald & Ponganis, 2012). This is an important consideration for all divers who can modulate lung volume and gas exchange prior to diving via the degree of inhalation and during diving via exhalation (Fahlman et al., 2009); indeed, there are noted differences in pre-dive respiratory behavior, with some marine mammals exhibiting pre-dive exhalation to reduce the lung volume [e.g., phocid seals (Kooyman et al., 1973)].

#### ***Peak Pressure as a Predictor of Explosive Injury***

High instantaneous peak pressures can cause damaging tissue distortion. Goertner (1982) suggested a peak overpressure gastrointestinal tract injury criterion because the size of gas bubbles in the gastrointestinal tract are variable, and their oscillation period could be short relative to primary blast wave exposure duration. The potential for gastrointestinal tract injury, therefore, may not be adequately modeled by the single oscillation bubble methodology used to estimate lung injury due to impulse. Like impulse, however, high instantaneous pressures may damage many parts of the body, but damage to the gastrointestinal tract is used as an indicator of any peak pressure-induced injury due to its vulnerability.

Older military reports documenting exposure of human divers to blast exposure generally describe peak pressure exposures around 100 psi (237 dB re 1  $\mu$ Pa peak) to feel like slight pressure or stinging sensation on skin, with no enduring effects (Christian & Gaspin, 1974). Around 200 psi, the shock wave felt like a blow to the head and chest. Data from the Lovelace Foundation experiments show instances of gastrointestinal tract contusions after exposures up to 1147 psi peak pressure, while exposures of up to 588 psi peak pressure resulted in many instances of no observed gastrointestinal tract effects. The lowest exposure for which slight contusions to the gastrointestinal tract were reported was 237 dB re 1  $\mu$ Pa peak. As a vulnerable gas-containing organ, the gastrointestinal tract is vulnerable to both high peak pressure and high impulse, which may vary to differing extents due to blast exposure conditions (i.e., animal depth, distance from the charge). This likely explains the range of effects seen at similar peak pressure exposure levels and shows the utility of considering both peak pressure and impulse when analyzing the potential for injury due to explosives.

#### **3.4.2.2.1.2 Hearing Loss**

Exposure to intense sound may result in noise-induced hearing loss that persists after cessation of the noise exposure. Hearing loss may be temporary or permanent, depending on factors such as the exposure frequency, received SPL, temporal pattern, and duration. The frequencies affected by hearing

loss may vary depending on the exposure frequency, with frequencies at and above the exposure frequency most strongly affected. The amount of hearing loss may range from slight to profound, depending on the ability of the individual to hear at the affected frequencies. Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) provides additional information on hearing loss and the framework used to analyze this potential impact.

Hearing loss has only been studied in a few species of marine mammals, although hearing studies with terrestrial mammals are also informative. There are no direct measurements of hearing loss in marine mammals due to exposure to explosive sources. The sound resulting from an explosive detonation is considered an impulsive sound and shares important qualities (e.g., short duration and fast rise time) with other impulsive sounds such as those produced by air guns. General research findings regarding TTS and PTS in marine mammals, as well as findings specific to exposure to other impulsive sound sources, are discussed in Section 3.4.2.1.1.2 (Hearing Loss).

#### **Threshold Shift due to Impulsive Sound Sources**

Marine mammal TTS data from impulsive sources are limited to two studies with measured TTS of 6 dB or more: Finneran et al. (2002) reported behaviorally measured TTSs of 6 and 7 dB in a beluga exposed to single impulses from a seismic water gun, and Lucke et al. (2009) reported auditory evoked potential-measured TTS of 7–20 dB in a harbor porpoise exposed to single impulses from a seismic air gun.

In addition to these data, Kastelein et al. (2015a) reported behaviorally measured mean TTS of 4 dB at 8 kHz and 2 dB at 4 kHz after a harbor porpoise was exposed to a series of impulsive sounds produced by broadcasting underwater recordings of impact pile driving strikes through underwater sound projectors. The cumulative SEL was approximately 180 dB re 1  $\mu\text{Pa}^2\text{s}$ . The pressure waveforms for the simulated pile strikes exhibited significant “ringing” not present in the original recordings, and most of the energy in the broadcasts was between 500 and 800 Hz. As a result, some questions exist regarding whether the fatiguing signals were representative of underwater pressure signatures from impact pile driving.

Several impulsive noise exposure studies have also been conducted without behaviorally measurable TTS. Finneran et al. (2000) exposed dolphins and belugas to single impulses from an “explosion simulator,” and Finneran et al. (2015) exposed three dolphins to sequences of 10 impulses from a seismic air gun (maximum cumulative SEL = 193 to 195 dB re 1  $\mu\text{Pa}^2\text{s}$ , peak SPL = 196 to 210 dB re 1  $\mu\text{Pa}$ ) without measurable TTS. Finneran et al. (2003b) exposed two sea lions to single impulses from an arc-gap transducer with no measurable TTS (maximum unweighted SEL = 163 dB re 1  $\mu\text{Pa}^2\text{s}$ , peak SPL = 183 dB re 1  $\mu\text{Pa}$ ).

#### **3.4.2.2.1.3 Physiological Stress**

Marine mammals naturally experience stress within their environment and as part of their life histories. The stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor. However, if the magnitude and duration of the stress response is too great or too long, then it can have negative consequences to the organism (e.g., decreased immune function, decreased reproduction). Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities) provides additional information on physiological stress and the framework used to analyze this potential impact.

There are no direct measurements of physiological stress in marine mammals due to exposure to explosive sources. General research findings regarding physiological stress in marine mammals due to exposure to sound and other stressors are discussed in detail in Section 3.4.2.1.1.3 (Physiological Stress). Because there are many unknowns regarding the occurrence of acoustically induced stress responses in marine mammals, it is assumed that any physiological response (e.g., hearing loss or injury) or significant behavioral response is also associated with a stress response.

#### **3.4.2.2.1.4 Masking**

Masking occurs when one sound, distinguished as the “noise,” interferes with the detection, discrimination, or recognition of another sound. The quantitative definition of masking is the amount in decibels an auditory detection, discrimination, or recognition threshold is raised in the presence of a masker (Erbe et al., 2015). As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), masking can effectively limit the distance over which a marine mammal can communicate, detect biologically relevant sounds, and echolocate (odontocetes). Masking only occurs in the presence of the masking noise and does not persist after the cessation of the noise. Masking can lead to vocal changes (e.g., Lombard effect, increasing amplitude, or changing frequency) and behavior changes (e.g., cessation of foraging, leaving an area) to both signalers and receivers, in an attempt to compensate for noise levels (Erbe et al., 2016).

There are no direct observations of masking in marine mammals due to exposure to explosive sources. Potential masking from explosive sounds is likely to be similar to masking studied for other impulsive sounds such as air guns. Masking could occur in mysticetes due to the overlap between their low-frequency vocalizations and the dominant frequencies of air gun pulses, however, masking in odontocetes is less likely unless the seismic survey activity is in close range when the pulses are more broadband. For example, differential vocal responses in marine mammals were documented in the presence of seismic survey noise. An overall decrease in vocalizations during active surveying was noted in large marine mammal groups (Potter et al., 2007), while blue whale feeding/social calls increased when seismic exploration was underway (Di Lorio & Clark, 2010), indicative of a possible compensatory response to the increased noise level. Bowhead whales were found to increase call rates in the presence of seismic air gun noise at lower received levels (below 100 dB re 1  $\mu\text{Pa}^2\text{s}$  cumulative SEL), but once the received level rose above 127 dB re 1  $\mu\text{Pa}^2\text{s}$  cumulative SEL the call rate began decreasing, and stopped altogether once received levels reached 170 dB re 1  $\mu\text{Pa}^2\text{s}$  cumulative SEL (Blackwell et al., 2015). Nieukirk et al. (2012) recorded both seismic surveys and fin whale 20 Hz calls at various locations around the mid-Atlantic Ocean and hypothesized that distant seismic noise could mask those calls, thereby decreasing the communication range of fin whales, whose vocalizations may propagate over 400 km to reach conspecifics (Spiesberger & Fristrup, 1990). A spotted and ringed seal in captivity were exposed to seismic air gun sounds recorded within 1 km and 30 km of an air gun survey conducted in shallow (<40 m.) water. They were then tested on their ability to detect a 500 ms upswEEP centered at 100 Hz at different points in the air gun pulse (start, middle, and end). Based on these results, a 100 Hz vocalization with a source level of 130 dB re 1  $\mu\text{Pa}$  would not be detected above a seismic survey 1 km away unless the animal was within 1–5 m, and would not be detected above a survey 30 km away beyond 46 m (Sills et al., 2017).

#### **3.4.2.2.1.5 Behavioral Reactions**

As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), any stimuli in the environment can cause a behavioral response in marine mammals, including noise from explosions. There are few direct observations of behavioral reactions from marine

mammals due to exposure to explosive sounds. Lammers et al. (2017) recorded dolphin detections near naval mine neutralization exercises and found that although the immediate response (within 30 seconds of the explosion) was an increase in whistles relative to the 30 seconds before the explosion, there was a reduction in daytime acoustic activity during the day of and the day after the exercise within 6 km. However, the nighttime activity did not seem to be different than that prior to the exercise, and two days after there appeared to be an increase in daytime acoustic activity, indicating a rapid return to the area by the dolphins (Lammers et al., 2017). Vallejo et al. (2017) report on boat-based line-transect surveys which were run over 10 years in an area where an offshore wind farm was built; these surveys included the periods of preconstruction, construction, and post-construction. Harbor porpoise were observed throughout the area during all three phases, but were not detected within the footprint of the windfarm during the construction phase, and were overall less frequent throughout the study area. However, they returned after the construction was completed at a slightly higher level than in the preconstruction phase. Furthermore, there was no large-scale displacement of harbor porpoises during construction, and in fact their avoidance behavior only occurred out to about 18 km, in contrast to the approximately 25 km avoidance distance found in other windfarm construction and pile driving monitoring efforts.

Impulsive signals, particularly at close range, have a rapid rise time and higher instantaneous peak pressure than other signal types, making them more likely to cause startle responses or avoidance responses. However, at long distances the rise time increases as the signal duration lengthens (similar to a “ringing” sound), making the impulsive signal more similar to a non-impulsive signal. Behavioral reactions from explosive sounds are likely to be similar to reactions studied for other impulsive sounds, such as those produced by air guns and impact pile driving. Data on behavioral responses to impulsive sound sources are limited across all marine mammal groups, with only a few studies available for mysticetes and odontocetes. Most data have come from seismic surveys that occur over long durations (e.g., on the order of days to weeks), and typically utilize large multi-air gun arrays that fire repeatedly. While seismic data provide the best available science for assessing behavioral responses to impulsive sounds by marine mammals, it is likely that these responses represent a worst-case scenario compared to responses to explosives used in Navy activities, which would typically consist of single impulses or a cluster of impulses, rather than long-duration, repeated impulses.

### **Mysticetes**

Baleen whales have shown a variety of responses to impulsive sound sources, including avoidance, attraction to the source, reduced surface intervals, altered swimming behavior, and changes in vocalization rates (Gordon et al., 2003; McCauley et al., 2000; Richardson et al., 1985; Southall et al., 2007). Studies have been conducted on many baleen whale species, including gray, humpback, blue, fin and bowhead whales; it is assumed that these responses are representative of all baleen whale species. The behavioral state of the whale seems to be an integral part of whether or not the animal responds and how they respond, as does the location and movement of the sound source, more than the received level of the sound.

Migratory behavior seems to lead to a higher likelihood of response, with some species demonstrating more sensitivity than others. For example, migrating gray whales showed avoidance responses to seismic vessels at received levels between 164 and 190 dB re 1  $\mu$ Pa (Malme et al., 1986, 1988). Similarly, migrating humpback whales showed avoidance behavior at ranges of 5–8 km from a seismic array during observational studies and controlled exposure experiments in one Australian study (McCauley et al., 1998), and in another Australian study decreased their dive times and reduced their swimming

speeds (Dunlop et al., 2015). However, when comparing received levels and behavioral responses when using ramp-up versus a constant noise level of air guns, humpback whales did not change their dive behavior but did deviate from their predicted heading and decrease their swim speeds (Dunlop et al., 2016). In addition, the whales demonstrated more course deviation during the constant source trials but reduced travel speeds more in the ramp-up trials; in either case there was no dose-response relationship with the received level of the air gun noise, and similar responses were observed in control trials with vessel movement but no air guns, so some of the response was likely due to the presence of the vessel and not the received level of the air guns. When looking at the relationships between proximity, received level, and behavioral response, Dunlop et al. (2017) used responses to two different air guns and found responses occurred more towards the smaller, closer source than to the larger source at the same received level, demonstrating the importance of proximity. Responses were found to be more likely when the source was within 3 km or above 140 dB re 1  $\mu$ Pa, although responses were variable and some animals did not respond at those values while others responded below them. In addition, responses were generally small, with course deviations of only around 500 m, and short-term (Dunlop et al., 2017). McDonald et al. (1995) tracked a blue whale with seafloor seismometers and reported that it stopped vocalizing and changed its travel direction at a range of 10 km from the seismic vessel (estimated received level 143 dB re 1  $\mu$ Pa peak-to-peak). Bowhead whales seem to be the most sensitive species, perhaps due to a higher overlap between bowhead whale distribution and seismic surveys in Arctic and sub-Arctic waters, as well as a recent history of being hunted. While most bowhead whales did not show active avoidance until within 8 km of seismic vessels (Richardson et al., 1995b), some whales avoided vessels by more than 20 km at received levels as low as 120 dB re 1  $\mu$ Pa. Additionally, Malme et al. (1988) observed clear changes in diving and breathing patterns in bowheads at ranges up to 73 km from seismic vessels, with received levels as low as 125 dB re 1  $\mu$ Pa. Bowhead whales may also avoid the area around seismic surveys, from 6 to 8 km (Koski and Johnson 1987, as cited in Gordon et al., 2003) out to 20 or 30 km (Richardson et al., 1999). However, work by Robertson (2014) supports the idea that behavioral responses are contextually dependent, and that during seismic operations bowhead whales may be less “available” for counting due to alterations in dive behavior but may not have left the area after all.

In contrast, noise from seismic surveys was not found to impact feeding behavior or exhalation rates in western gray whales while resting or diving off the coast of Russia (Gailey et al., 2007; Yazvenko et al., 2007); however, the increase in vessel traffic associated with the surveys and the proximity of the vessels to the whales did affect the orientation of the whales relative to the vessels and shortened their dive-surface intervals (Gailey et al., 2016). Todd et al. (1996) found no clear short-term behavioral responses by foraging humpbacks to explosions associated with construction operations in Newfoundland but did see a trend of increased rates of net entanglement closer to the noise source, possibly indicating a reduction in net detection associated with the noise through masking or TTS. Distributions of fin and minke whales were modeled with a suite of environmental variables along with the occurrence or absence of seismic surveys, and no evidence of a decrease in sighting rates relative to seismic activity was found for either species (Vilela et al., 2016). Their distributions were driven entirely by environmental variables, particularly those linked to prey, including warmer sea surface temperatures, higher chlorophyll-a values, and higher photosynthetically available radiation (a measure of primary productivity).

Vocal responses to seismic surveys have been observed in a number of baleen whale species, including a cessation of calling, a shift in frequency, increases in amplitude or call rate, or a combination of these strategies. Blue whale feeding/social calls were found to increase when seismic exploration was

underway, with seismic pulses at average received SELs of 131 dB re 1  $\mu\text{Pa}^2\text{s}$  (Di Lorio & Clark, 2010), a potentially compensatory response to increased noise level. Responses by fin whales to a 10-day seismic survey in the Mediterranean Sea included possible decreased 20-Hz call production and movement of animals from the area based on lower received levels and changes in bearings (Castellote et al., 2012). However, similarly distant seismic surveys elicited no apparent vocal response from fin whales in the mid-Atlantic Ocean; instead, Nieukirk et al. (2012) hypothesized that 20-Hz calls may have been masked from the receiver by distant seismic noise. Models of humpback whale songs off Angola showed significant seasonal and diel variation, but also showed a decrease in the number of singers with increasing received levels of air gun pulses (Cerchio et al., 2014). Bowhead whale calling rates decreased significantly at sites near seismic surveys (41–45 km) where median received levels were between 116–129 dB re 1  $\mu\text{Pa}$ , and did not decrease at sites further from the seismic surveys (greater than 104 km) where median received levels were 99–108 dB re 1  $\mu\text{Pa}$  (Blackwell et al., 2013). In fact, bowhead whale calling rates increased at the lower received levels, began decreasing at around 127 dB re 1  $\mu\text{Pa}^2\text{s}$  cumulative SEL, and ceased altogether at received levels over 170 dB re 1  $\mu\text{Pa}^2\text{s}$  cumulative SEL (Blackwell et al., 2015). Similar patterns were observed for bowhead vocalizations in the presence of tonal sounds associated with drilling activities, and were amplified in the presence of both the tonal sounds and air gun pulses (Blackwell et al., 2017).

Mysticetes seem to be the most sensitive taxonomic group of marine mammals to impulsive sound sources, with possible avoidance responses occurring out to 30 km and vocal changes occurring in response to sounds over 100 km away. However, responses appear to be behaviorally mediated, with most avoidance responses occurring during migration behavior and little observed response during feeding behavior. These response patterns are likely to hold true for Navy impulsive sources; however, Navy impulsive sources would largely be stationary (e.g., explosives fired at a fixed target), and short-term (on the order of hours rather than days or weeks) than were found in these studies, and so responses would likely occur in closer proximity or not at all.

### **Odontocetes**

Few data are available on odontocete responses to impulsive sound sources, with only a few studies on responses to seismic surveys, pile driving, and construction activity available. However, odontocetes appear to be less sensitive to impulsive sound than mysticetes, with responses occurring at much closer distances. This may be due to the predominance of low-frequency sound associated with these sources that propagates long distances and overlaps with the range of best hearing for mysticetes but is below that range for odontocetes. The exception to this is the harbor porpoise, which has been shown to be highly sensitive to most sound sources, avoiding both stationary (e.g., pile driving) and moving (e.g., seismic survey vessels) impulsive sound sources out to approximately 20 km (e.g., Haelters et al., 2014; Pirodda et al., 2014). However, even this response is short-term, with porpoises returning to the area within hours after the cessation of the noise.

Madsen et al. (2006a) and Miller et al. (2009) tagged and monitored eight sperm whales in the Gulf of Mexico exposed to seismic air gun surveys. Sound sources were from approximately 2 to 7 NM away from the whales, and received levels were as high as 162 dB SPL re 1  $\mu\text{Pa}$  (Madsen et al., 2006). The whales showed no horizontal avoidance, although one whale rested at the water's surface for an extended period of time until air guns ceased firing (Miller et al., 2009). While the remaining whales continued to execute foraging dives throughout exposure, tag data suggested there may have been subtle effects of noise on foraging behavior (Miller et al., 2009). Similarly, Weir (2008) observed that seismic air gun surveys along the Angolan coast did not significantly reduce the encounter rate of sperm

whales during the 10-month survey period, nor were avoidance behaviors to air gun impulsive sounds observed. In contrast, Atlantic spotted dolphins did show a significant, short-term avoidance response to air gun impulses within approximately 1 km of the source (Weir, 2008). The dolphins were observed at greater distances from the vessel when the air gun was in use, and when the air gun was not in use they readily approached the vessel to bow ride.

Captive bottlenose dolphins sometimes vocalized or were reluctant to return to the test station after exposure to single impulses from a seismic water gun (Finneran et al., 2002). When exposed to multiple impulses from a seismic air gun, some dolphins turned their heads away from the sound source just before the impulse, showing that they could anticipate the timing of the impulses and perhaps reduce the received level (Finneran et al., 2015). During construction (including the blasting of old bastions) of a bridge over a waterway commonly used by the Tampa Bay, FL, stock of bottlenose dolphins, the use of the area by females decreased while males displayed high site fidelity and continued using the area, perhaps indicating differential habitat uses between the sexes (Weaver, 2015).

A study was conducted on the response of harbor porpoises to a seismic survey using aerial surveys and C-PODs (an autonomous recording device that counts odontocete clicks); the animals appeared to have left the area of the survey and decreased their foraging activity within 5–10 km, as evidenced by both a decrease in vocalizations near the survey and an increase in vocalizations at a distance (Pirota et al., 2014; Thompson et al., 2013). However, the animals returned within a day after the air gun operation ceased, and the decrease in occurrence over the survey period was small relative to the observed natural seasonal decrease compared to the previous year. A number of studies (Brandt et al., 2011; Dähne et al., 2014; Haelters et al., 2014; Thompson et al., 2010; Tougaard et al., 2005; Tougaard et al., 2009) also found strong avoidance responses by harbor porpoises out to 20 km during pile driving; however, all studies found that the animals returned to the area after the cessation of pile driving. When bubble curtains were deployed around pile driving, the avoidance distance appeared to be reduced to half that distance (12 km), and the response only lasted about five hours rather than a day before the animals returned to the area (Dähne et al., 2017). Kastelein et al. (2013b) exposed a captive harbor porpoise to impact pile driving sounds, and found that above 136 dB re 1  $\mu$ Pa (zero-to-peak) the animal's respiration rates increased, and at higher levels it jumped more frequently. Bergstrom et al. (2014) found that although there was a high likelihood of acoustic disturbance during wind farm construction (including pile driving), the impact was short-term. Graham et al. (2017) assessed the occurrence of bottlenose dolphins and harbor porpoises over different area and time scales with and without impact and vibratory pile driving. While there were fewer hours with bottlenose dolphin detections and reduced detection durations within the pile driving area and increased detection durations outside the area, the effects sizes were small, and the reduced harbor porpoise encounter duration was attributed to seasonal changes outside the influence of the pile driving. However, received levels in this area were lower due to propagation effects than in the other areas described above, which may have led to the lack of or reduced response.

Odontocete behavioral responses to impulsive sound sources are likely species- and context-dependent, with most species demonstrating little to no apparent response. Responses might be expected within close proximity to a noise source, under specific behavioral conditions such as females with offspring, or for sensitive species such as harbor porpoises.

#### **3.4.2.2.1.6 Stranding**

When a marine mammal (alive or dead) swims or floats onto shore and becomes beached or incapable of returning to sea, the event is termed a "stranding" (Geraci et al., 1999; Geraci & Lounsbury, 2005;

Perrin & Geraci, 2002). Specifically, under U.S. law, a stranding is an event in the wild where: “(A) a marine mammal is dead and is (i) on a beach or shore of the United States; or (ii) in waters under the jurisdiction of the United States (including any navigable waters); or (B) a marine mammal is alive and is (i) on a beach or shore of the United States and is unable to return to the water; (ii) on a beach or shore of the United States and, although able to return to the water, is in need of medical attention; or (iii) in the waters under the jurisdiction of the United States (including any navigable waters), but is unable to return to its natural habitat under its own power or without assistance” (16 U.S.C. section 1421h).

Impulsive sources (e.g., explosions) also have the potential to contribute to strandings, but such occurrences are even less common than those that have been related to certain sonar activities. During a Navy training event on March 4, 2011, at the Silver Strand Training Complex in San Diego, California, three long-beaked common dolphins were killed by an underwater detonation. Further details are provided above. Discussions of mitigation measures associated with these and other training and testing events are presented in Chapter 5 (Mitigation).

#### **3.4.2.2.1.7 Long-Term Consequences**

Long-term consequences to a population are determined by examining changes in the population growth rate. For additional information on the determination of long-term consequences, see Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities). Physical effects from explosive sources that could lead to a reduction in the population growth rate include mortality or injury, which could remove animals from the reproductive pool, and permanent hearing impairment or chronic masking, which could impact navigation, foraging, predator avoidance, or communication. The long-term consequences due to individual behavioral reactions, masking, and short-term instances of physiological stress are especially difficult to predict because individual experience over time can create complex contingencies, especially for long-lived animals like marine mammals. For example, a lost reproductive opportunity could be a measurable cost to the individual; however, short-term costs may be recouped during the life of an otherwise healthy individual. These factors are taken into consideration when assessing risk of long-term consequences.

#### **3.4.2.2.2 Impacts from Explosive Stressors**

Marine mammals could be exposed to energy, sound, and fragments from explosions in the water and near the water surface associated with the proposed activities. Energy from an explosion is capable of causing mortality, injury, hearing loss, a behavioral response, masking, or physiological stress, depending on the level and duration of exposure.

The death of an animal would eliminate future reproductive potential, which is considered in the analysis of potential long-term consequences to the population. Exposures that result in non-auditory injuries or PTS may limit an animal’s ability to find food, communicate with other animals, or interpret the surrounding environment. Impairment of these abilities can decrease an individual’s chance of survival or impact its ability to successfully reproduce. TTS can also impair an animal’s abilities, but the individual is likely to recover quickly with little significant effect.

Explosions in the ocean or near the water surface can introduce loud, impulsive, broadband sounds into the marine environment. These sounds, which are within the audible range of most marine mammals, could cause behavioral reactions, masking, and elevated physiological stress. Behavioral responses can include shorter surfacings, shorter dives, fewer blows (breaths) per surfacing, longer intervals between blows, ceasing or increasing vocalizations, shortening or lengthening vocalizations, and changing frequency or intensity of vocalizations (National Research Council 2005). Sounds from explosives could

also mask biologically important sounds; however, the duration of individual sounds is very short, reducing the likelihood of substantial auditory masking.

#### **3.4.2.2.2.1 Methods for Analyzing Impacts from Explosives**

The Navy performed a quantitative analysis to estimate the number of times that marine mammals could be impacted by explosions used during Navy training and testing activities. The Navy's quantitative analysis to determine impacts on marine mammals uses the Navy Acoustic Effects Model to produce initial estimates of the number of instances that animals may experience these effects; these estimates are further refined by considering animal avoidance of sound-producing activities and implementation of procedural mitigation measures. The steps of this quantitative analysis are described in Section 3.0.1.2 (Navy's Quantitative Analysis to Determine Impacts on Sea Turtles and Marine Mammals), which takes into account:

- criteria and thresholds used to predict impacts from explosives (see below)
- the density and spatial distribution of marine mammals
- the influence of environmental parameters (e.g., temperature, depth, salinity) on sound propagation and explosive energy when estimating the received sound level and pressure on the animals

A detailed explanation of this analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

#### **Criteria and Thresholds used to Estimate Impacts to Marine Mammals from Explosives**

##### ***Mortality and Injury from Explosives***

As discussed above in Section 3.4.2.2.1.1 (Injury), two metrics have been identified as predictive of injury: impulse and peak pressure. Peak pressure contributes to the "crack" or "stinging" sensation of a blast wave, compared to the "thump" associated with received impulse. Older military reports documenting exposure of human divers to blast exposure generally describe peak pressure exposures around 100 psi (237 dB re 1  $\mu$ Pa SPL peak) to feel like slight pressure or stinging sensation on skin, with no enduring effects (Christian & Gaspin, 1974).

Because data on explosive injury do not indicate a set threshold for injury, rather a range of risk for explosive exposures, two sets of criteria are provided for use in non-auditory injury assessment. The exposure thresholds are used to estimate the number of animals that may be affected during Navy training and testing activities (Table 3.4-15). The thresholds for the farthest range to effect are based on the received level at which one percent risk is predicted and are useful for assessing potential effects to marine mammals and the level of potential impacts covered by the mitigation zones. Increasing animal mass and increasing animal depth both increase the impulse thresholds (i.e., decrease susceptibility), whereas smaller mass and decreased animal depth reduce the impulse thresholds (i.e., increase susceptibility). For impact assessment, marine mammal populations are assumed to be 70 percent adult and 30 percent calf/pup. Sub-adult masses are used to determine onset of effect, in order to estimate the farthest range at which an effect may first be observable. The derivation of these injury criteria and the species mass estimates are provided in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).

**Table 3.4-15: Criteria to Quantitatively Assess Non-Auditory Injury Due to Underwater Explosions**

| <i>Impact Category</i> | <i>Impact Threshold</i>                                  | <i>Threshold for Farthest Range to Effect<sup>2</sup></i> |
|------------------------|--|---|
| Mortality <sup>1</sup> | $144M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s  | $103 \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s          |
| Injury <sup>1</sup>    | $65.8M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s | $47.5M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s  |
|                        | 243 dB re 1 μPa SPL peak                                 | 237 dB re 1 μPa SPL peak                                  |

<sup>1</sup> Impulse delivered over 20 percent of the estimated lung resonance period. See U.S. Department of the Navy (2017a).


<sup>2</sup> Threshold for one percent risk used to assess mitigation effectiveness.

Notes: D = animal depth (m), dB re 1 μPa = decibels referenced to 1 micropascal, M = animal mass (kg), Pa-s = Pascal-second, SPL = sound pressure level

When explosive ordnance (e.g., bomb or missile) detonates, fragments of the weapon are thrown at high-velocity from the detonation point, which can injure or kill marine mammals if they are struck. Risk of fragment injury reduces exponentially with distance as the fragment density is reduced. Fragments underwater tend to be larger than fragments produced by in-air explosions (Swisdak & Montanaro, 1992). Underwater, the friction of the water would quickly slow these fragments to a point where they no longer pose a threat. On the other hand, the blast wave from an explosive detonation moves efficiently through the seawater. Because the ranges to mortality and injury due to exposure to the blast wave are likely to far exceed the zone where fragments could injure or kill an animal, the above thresholds are assumed to encompass risk due to fragmentation.

**Auditory Weighting Functions**

Animals are not equally sensitive to noise at all frequencies. To capture the frequency-dependent nature of the effects of noise, auditory weighting functions are used (Figure 3.4-63). Auditory weighting functions are mathematical functions based on a generic band-pass filter and incorporate species-specific hearing abilities to calculate a weighted received sound level in units SPL or SEL. Due to the band pass nature of auditory weighting functions, they resemble an inverted “U” shape with amplitude plotted as a function of frequency. The flatter portion of the plotted function, where the amplitude is closest to zero, is the emphasized frequency range (i.e., the pass-band), while the frequencies below and above this range (where amplitude declines) are de-emphasized.




Source: See Finneran (2015) for parameters used to generate the functions and more information on weighting function derivation.

Notes: MF = mid-frequency cetacean, HF = high-frequency cetacean, LF = low-frequency cetacean

**Figure 3.4-63: Navy Phase III Weighting Functions for All Species Groups**

**Hearing Loss from Explosives**

Criteria used to define threshold shifts from explosions are derived from the two known studies designed to induce TTS in marine mammals from impulsive sources. Finneran et al. (2002) reported behaviorally measured TTS of 6 and 7 dB in a beluga exposed to single impulses from a seismic water gun, and Lucke et al. (2009) reported auditory evoked potential-measured TTS of 7–20 dB in a harbor porpoise exposed to single impulses from a seismic air gun. Since marine mammal PTS data from impulsive noise exposures do not exist, onset-PTS levels for all groups were estimated by adding 15 dB to the threshold for non-impulsive sources. This relationship was derived by Southall et al. (2007) from impulsive noise TTS growth rates in chinchillas. These frequency-dependent thresholds are depicted by the exposure functions for each group’s range of best hearing (Figure 3.4-64). Weighted sound exposure thresholds for underwater explosive sounds used in the analysis are shown in Table 3.4-16.



Notes: The dark dashed curve is the exposure function for PTS onset, the solid black curve is the exposure function for TTS onset, and the light grey curve is the exposure function for behavioral response. Small dashed lines indicate the SEL threshold for behavioral response, TTS, and PTS onset at each group's most sensitive frequency (i.e., the weighted SEL threshold).

**Figure 3.4-64: Navy Phase III Behavioral, TTS and PTS Exposure Functions for Explosives**

**Table 3.4-16: Navy Phase III Weighted Sound Exposure Thresholds for Underwater Explosive Sounds**

| <i>Hearing Group</i>    | <i>Explosive Sound Source</i>       |                                |                                       |                                |                                       |
|-------------------------|-------------------------------------|--------------------------------|---------------------------------------|--------------------------------|---------------------------------------|
|                         | <i>Behavior (SEL) weighted (dB)</i> | <i>TTS (SEL) weighted (dB)</i> | <i>TTS (Peak SPL) unweighted (dB)</i> | <i>PTS (SEL) weighted (dB)</i> | <i>PTS (Peak SPL) unweighted (dB)</i> |
| Low-frequency Cetacean  | 163                                 | 168                            | 213                                   | 183                            | 219                                   |
| Mid-frequency Cetacean  | 165                                 | 170                            | 224                                   | 185                            | 230                                   |
| High-frequency Cetacean | 135                                 | 140                            | 196                                   | 155                            | 202                                   |

Notes: dB = decibels, PTS = permanent threshold shift, SEL = sound exposure level, SPL = sound pressure level, TTS = temporary threshold shift

***Behavioral Responses from Explosives***

If more than one explosive event occurs within any given 24-hour period within a training or testing activity, criteria are applied to predict the number of animals that may have a behavioral reaction. For exercises with multiple explosions, the behavioral threshold used in this analysis is 5 dB less than the TTS onset threshold (in SEL). This value is derived from observed onsets of behavioral response by test subjects (bottlenose dolphins) during non-impulsive TTS testing (Schlundt et al., 2000).

Some multiple explosive exercises, such as certain naval gunnery exercises, may be treated as a single event because a few explosions occur closely spaced within a very short time (a few seconds). For single explosions at received sound levels below hearing loss thresholds, the most likely behavioral response is a brief alerting or orienting response. Since no further sounds follow the initial brief impulses, significant behavioral reactions would not be expected to occur. This reasoning was applied to previous shock trials (63 FR 230; 66 FR 87; 73 FR 143) and is extended to the criteria used in this analysis.

**Accounting for Mitigation**

The Navy will implement mitigation measures to avoid or reduce potential impacts from explosives on marine mammals, as described in Section 5.3.3 (Explosive Stressors). The benefits of mitigation are conservatively factored into the analysis for Alternative 1 and Alternative 2 of the Proposed Action for training and testing. The Navy’s mitigation measures are identical for both action alternatives.

Procedural mitigation measures include delaying or ceasing applicable detonations when a marine mammal is observed in a mitigation zone. The mitigation zones for explosives extend beyond the respective average ranges to mortality. Therefore, the impact analysis quantifies the potential for procedural mitigation to reduce the risk of mortality due to exposure to explosives. Two factors are considered when quantifying the effectiveness of mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., an explosive activity) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

In the quantitative analysis, consideration of mitigation measures means that, for activities that implement mitigation, model-estimated mortality is considered mitigated to the level of injury. The impact analysis does not analyze the potential for mitigation to reduce non-auditory injury, PTS, TTS, or behavioral effects, even though mitigation would also reduce the likelihood of these effects. In practice, mitigation also protects all unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

The ability to observe the ranges to mortality was estimated for each training or testing event. The ability of Navy Lookouts to detect marine mammals within a mitigation zone is dependent on the animal's presence at the surface and the characteristics of the animal that influence its sightability (such as group size or surface active behavior). The behaviors and characteristics of some species may make them easier to detect. Certain behaviors, such as leaping and breaching, are visible from a great distance and likely increase sighting distances and detections of those species. Environmental conditions under which the training or testing activity could take place are also considered, such as sea surface conditions, weather (e.g., fog or rain), and day versus night.

The Navy will also implement mitigation measures for certain explosive activities within mitigation areas, as described in Appendix I (Geographic Mitigation Assessment). The benefits of mitigation areas are discussed qualitatively and have not been factored into the quantitative analysis process or reductions in take for the MMPA and ESA impact estimates. Mitigation areas are designed to help avoid or reduce impacts during biologically important life processes within particularly important habitat areas. Therefore, mitigation area benefits are discussed in terms of the context of impact avoidance or reduction.

#### **3.4.2.2.2.2 Impact Ranges for Explosives**

The following section provides the range (distance) over which specific physiological or behavioral effects are expected to occur based on the explosive criteria and the explosive propagation calculations from the Navy Acoustic Effects Model (Section 3.4.2.2.2.1, Methods for Analyzing Impacts from Explosives). The range to effects are shown for a range of explosive bins, from E1 (up to 0.25 lb. net explosive weight) to E12 (up to 1,000 lb. net explosive weight). Ranges are determined by modeling the distance that noise from an explosion will need to propagate to reach exposure level thresholds specific to a hearing group that would cause behavioral response, TTS, PTS, and non-auditory injury. Range to effects is important information in not only predicting impacts from explosives, but also in verifying the accuracy of model results against real-world situations and assessing the level of impact that would likely be mitigated within applicable mitigation zones.

Table 3.4-17 shows the minimum, average, and maximum ranges due to varying propagation conditions to non-auditory injury as a function of animal mass and explosive bin. Ranges to gastrointestinal tract injury typically exceed ranges to slight lung injury; therefore, the maximum range to effect is not mass-dependent. Animals within these water volumes would be expected to receive minor injuries at the outer ranges, increasing to more substantial injuries, and finally mortality as an animal approaches the detonation point. Ranges to mortality, based on animal mass, are shown in Table 3.4-18.

The following tables (Table 3.4-19 through Table 3.4-24) show the minimum, average, and maximum ranges to onset of auditory and behavioral effects based on the thresholds described in Section 3.4.2.2.2.1 (Methods for Analyzing Impacts from Explosives). Ranges are provided for a representative

source depth and cluster size (the number of rounds fired [or buoys dropped] within a very short duration) for each bin. For events with multiple explosions, sound from successive explosions can be expected to accumulate and increase the range to the onset of an impact based on SEL thresholds. Modeled ranges to TTS and PTS based on peak pressure for a single explosion generally exceed the modeled ranges based on SEL even when accumulated for multiple explosions. Peak pressure based ranges are estimated using the best available science; however, data on peak pressure at far distances from explosions are very limited. For additional information on how ranges to impacts from explosions were estimated, see the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

**Table 3.4-17: Ranges to Non-Auditory Injury (in meters) for All Marine Mammal Hearing Groups**

| <i>Bin</i> | <i>Range to Non-Auditory Injury (meters)<sup>1</sup></i> |
|------------|--|
| E1         | 12<br>(11–13)  |
| E2         | 16<br>(15–16)  |
| E3         | 25<br>(25–25)  |
| E4         | 30<br>(30–35)  |
| E5         | 40<br>(40–65)  |
| E6         | 52<br>(50–60)  |
| E7         | 120<br>(120–120)   |
| E8         | 98<br>(90–150)   |
| E9         | 123<br>(120–270)   |
| E10        | 155<br>(150–430)   |
| E11        | 418<br>(410–420)   |
| E12        | 195<br>(180–675)   |

<sup>1</sup> Average distance is shown with the minimum and maximum distances due to varying propagation environments in parentheses.

Notes: All ranges to non-auditory injury within this table are driven by gastrointestinal tract injury thresholds regardless of animal mass.

**Table 3.4-18: Ranges to Mortality (in meters) for All Marine Mammal Hearing Groups as a Function of Animal Mass**

| <i>Bin</i> | <i>Range to Mortality (meters) for Various Animal Mass Intervals (kg)<sup>1</sup></i> |                |                 |                 |                  |                  |
|------------|---|----------------|-----------------|-----------------|------------------|------------------|
|            | <i>10 kg</i>  | <i>250 kg</i>  | <i>1,000 kg</i> | <i>5,000 kg</i> | <i>25,000 kg</i> | <i>72,000 kg</i> |
| E1         | 3<br>(3-3)  | 1<br>(0-2)     | 0<br>(0-0)      | 0<br>(0-0)      | 0<br>(0-0)       | 0<br>(0-0)       |
| E2         | 4<br>(3-4)  | 2<br>(1-3)     | 1<br>(0-1)      | 0<br>(0-0)      | 0<br>(0-0)       | 0<br>(0-0)       |
| E3         | 9<br>(7-10)   | 4<br>(2-8)     | 2<br>(1-2)      | 1<br>(0-1)      | 0<br>(0-0)       | 0<br>(0-0)       |
| E4         | 13<br>(12-15)   | 7<br>(4-12)    | 3<br>(3-4)      | 2<br>(1-3)      | 1<br>(1-1)       | 1<br>(0-1)       |
| E5         | 13<br>(12-30)   | 7<br>(4-25)    | 3<br>(2-7)      | 2<br>(1-5)      | 1<br>(1-2)       | 1<br>(0-2)       |
| E6         | 16<br>(15-25)   | 9<br>(5-23)    | 4<br>(3-8)      | 3<br>(2-6)      | 1<br>(1-2)       | 1<br>(1-2)       |
| E7         | 55<br>(55-55)   | 26<br>(18-40)  | 13<br>(11-15)   | 9<br>(7-10)     | 4<br>(4-4)       | 3<br>(2-3)       |
| E8         | 42<br>(25-65)   | 22<br>(9-50)   | 11<br>(6-19)    | 8<br>(4-13)     | 4<br>(2-6)       | 3<br>(1-5)       |
| E9         | 33<br>(30-35)   | 20<br>(13-30)  | 10<br>(9-12)    | 7<br>(5-9)      | 4<br>(3-4)       | 3<br>(2-3)       |
| E10        | 55<br>(40-170)  | 24<br>(16-35)  | 13<br>(11-15)   | 9<br>(7-11)     | 5<br>(4-5)       | 4<br>(3-4)       |
| E11        | 206<br>(200-210)  | 98<br>(55-170) | 44<br>(35-50)   | 30<br>(25-35)   | 16<br>(14-18)    | 12<br>(10-15)    |
| E12        | 86<br>(50-270)  | 35<br>(20-210) | 16<br>(13-19)   | 11<br>(9-13)    | 6<br>(5-6)       | 5<br>(4-5)       |

<sup>1</sup>Average distance to mortality (meters) is depicted above the minimum and maximum distances, which are in parentheses for each animal mass interval.

Note: Kg = kilogram

**Table 3.4-19: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for High-Frequency Cetaceans**

| <i>Range to Effects for Explosives: High-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|--|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>   | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E1   | 0.1                          | 1                   | 353<br>(340–370)             | 1,303<br>(1,275–1,775)       | 2,139<br>(2,025–4,275)              |
|  |                              | 18                  | 1,031<br>(1,025–1,275)       | 3,409<br>(2,525–8,025)       | 4,208<br>(3,025–11,525)             |
| E2   | 0.1                          | 1                   | 431<br>(410–700)             | 1,691<br>(1,525–2,775)       | 2,550<br>(2,025–4,525)              |
|  |                              | 5                   | 819<br>(775–1,275)           | 2,896<br>(2,275–6,775)       | 3,627<br>(2,525–10,275)             |
| E3   | 0.1                          | 1                   | 649<br>(625–700)             | 2,439<br>(2,025–4,525)       | 3,329<br>(2,525–7,525)              |
|  |                              | 12                  | 1,682<br>(1,525–2,275)       | 4,196<br>(3,025–11,525)      | 5,388<br>(4,525–16,275)             |
|  | 18.25                        | 1                   | 720<br>(675–775)             | 4,214<br>(2,275–6,275)       | 7,126<br>(3,525–8,775)              |
|  |                              | 12                  | 1,798<br>(1,525–2,775)       | 10,872<br>(4,525–13,775)     | 14,553<br>(5,525–17,775)            |
| E4   | 10                           | 2                   | 1,365<br>(1,025–2,775)       | 7,097<br>(4,275–10,025)      | 9,939<br>(5,025–15,275)             |
|  | 60                           | 2                   | 1,056<br>(875–2,275)         | 3,746<br>(2,775–5,775)       | 5,262<br>(3,025–7,775)              |
| E5   | 0.1                          | 20                  | 2,926<br>(1,525–6,275)       | 6,741<br>(4,525–16,025)      | 9,161<br>(4,775–20,025)             |
|  | 30                           | 20                  | 4,199<br>(3,025–6,275)       | 13,783<br>(8,775–17,775)     | 17,360<br>(10,525–22,775)           |
| E6   | 0.1                          | 1                   | 1,031<br>(1,025–1,275)       | 3,693<br>(2,025–8,025)       | 4,659<br>(3,025–12,775)             |
|  | 30                           | 1                   | 1,268<br>(1,025–1,275)       | 7,277<br>(3,775–8,775)       | 10,688<br>(5,275–12,525)            |
| E7   | 28                           | 1                   | 1,711<br>(1,525–2,025)       | 8,732<br>(4,275–11,775)      | 12,575<br>(4,275–16,025)            |
| E8   | 0.1                          | 1                   | 1,790<br>(1,775–3,025)       | 4,581<br>(4,025–10,775)      | 6,028<br>(4,525–15,775)             |
|  | 45.75                        | 1                   | 1,842<br>(1,525–2,025)       | 9,040<br>(4,525–12,775)      | 12,729<br>(5,025–18,525)            |
| E9   | 0.1                          | 1                   | 2,343<br>(2,275–4,525)       | 5,212<br>(4,025–13,275)      | 7,573<br>(5,025–17,025)             |

**Table 3.4-19: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for High-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: High-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|--|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>   | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E10  | 0.1                          | 1                   | 2,758<br>(2,275–5,025)       | 6,209<br>(4,275–16,525)      | 8,578<br>(5,275–19,775)             |
| E11  | 45.75                        | 1                   | 3,005<br>(2,525–3,775)       | 11,648<br>(5,025–18,775)     | 14,912<br>(6,525–24,775)            |
|  | 91.4                         | 1                   | 3,234<br>(2,525–4,525)       | 5,772<br>(4,775–11,775)      | 7,197<br>(5,775–14,025)             |
| E12  | 0.1                          | 1                   | 3,172<br>(3,025–6,525)       | 7,058<br>(5,025–17,025)      | 9,262<br>(6,025–21,775)             |
|  |                              | 4                   | 4,209<br>(3,775–10,025)      | 9,817<br>(6,275–22,025)      | 12,432<br>(7,525–27,775)            |

<sup>1</sup>Average distance (meters) to PTS, TTS, and behavioral thresholds are depicted above the minimum and maximum distances which are in parentheses. Values depict the range produced by SEL hearing threshold criteria levels. PTS = permanent threshold shift, SEL = sound exposure level, TTS = temporary threshold shift

**Table 3.4-20: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for High-Frequency Cetaceans**

| <i>Range to Effects for Explosives: High-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|--|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>   | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E1   | 0.1                          | 745<br>(700–775)             | 1,275<br>(1,275–1,275)       |
| E2   | 0.1                          | 912<br>(380–975)             | 1,498<br>(725–1,525)         |
| E3   | 0.1                          | 1,525<br>(1,525–1,525)       | 2,397<br>(2,025–2,525)       |
|  | 18.25                        | 1,561<br>(1,525–2,775)       | 2,919<br>(2,775–3,525)       |
| E4   | 10                           | 2,076<br>(1,775–2,525)       | 5,565<br>(3,525–7,775)       |
|  | 60                           | 2,364<br>(1,775–4,775)       | 4,044<br>(2,025–5,275)       |
| E5   | 0.1                          | 2,267<br>(1,025–3,275)       | 3,093<br>(1,275–5,775)       |
|  | 30                           | 2,567<br>(2,275–2,775)       | 3,747<br>(3,025–5,275)       |

**Table 3.4-20: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for High-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: High-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|--|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>   | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E6   | 0.1                          | 2,546<br>(1,275–4,525)       | 3,356<br>(1,525–6,525)       |
|  | 30                           | 3,242<br>(2,775–3,525)       | 4,598<br>(3,525–5,275)       |
| E7   | 28                           | 4,261<br>(3,025–5,025)       | 7,782<br>(3,775–12,525)      |
| E8   | 0.1                          | 3,458<br>(3,025–6,525)       | 4,324<br>(3,775–8,275)       |
|  | 45.75                        | 4,790<br>(4,275–6,525)       | 11,013<br>(4,775–23,775)     |
| E9   | 0.1                          | 3,870<br>(3,275–8,025)       | 4,620<br>(3,775–10,275)      |
| E10  | 0.1                          | 3,993<br>(2,525–9,275)       | 5,076<br>(2,775–16,025)      |
| E11  | 45.75                        | 8,388<br>(4,775–24,275)      | 17,386<br>(5,025–33,275)     |
|  | 91.4                         | 5,051<br>(4,025–7,525)       | 7,065<br>(4,275–26,525)      |
| E12  | 0.1                          | 4,519<br>(3,775–9,775)       | 5,678<br>(4,275–13,025)      |

<sup>1</sup> Average distance (meters) is shown with the minimum and maximum distances due to varying propagation environments in parentheses. PTS = permanent threshold shift, TTS = temporary threshold shift

**Table 3.4-21: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Low-Frequency Cetaceans**

| <i>Range to Effects for Explosives: Low-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|---|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E1  | 0.1                          | 1                   | 51<br>(50–55)                | 231<br>(200–250)             | 378<br>(280–410)                    |
|   |                              | 18                  | 183<br>(170–190)             | 691<br>(450–775)             | 934<br>(575–1,275)                  |
| E2  | 0.1                          | 1                   | 66<br>(65–70)                | 291<br>(220–320)             | 463<br>(330–500)                    |
|   |                              | 5                   | 134<br>(110–140)             | 543<br>(370–600)             | 769<br>(490–950)                    |
| E3  | 0.1                          | 1                   | 113<br>(110–120)             | 477<br>(330–525)             | 689<br>(440–825)                    |
|   |                              | 12                  | 327<br>(250–370)             | 952<br>(600–1,525)           | 1,240<br>(775–4,025)                |
|   | 18.25                        | 1                   | 200<br>(200–200)             | 955<br>(925–1,000)           | 1,534<br>(1,275–1,775)              |
|   |                              | 12                  | 625<br>(600–625)             | 5,517<br>(2,275–7,775)       | 10,299<br>(3,775–13,025)            |
| E4  | 10                           | 2                   | 429<br>(370–600)             | 2,108<br>(1,775–2,775)       | 4,663<br>(3,025–6,025)              |
|   | 60                           | 2                   | 367<br>(340–470)             | 1,595<br>(1,025–2,025)       | 2,468<br>(1,525–4,275)              |
| E5  | 0.1                          | 20                  | 702<br>(380–1,275)           | 1,667<br>(850–11,025)        | 2,998<br>(1,025–19,775)             |
|   | 30                           | 20                  | 1,794<br>(1,275–2,775)       | 8,341<br>(3,775–11,525)      | 13,946<br>(4,025–22,275)            |
| E6  | 0.1                          | 1                   | 250<br>(190–410)             | 882<br>(480–1,775)           | 1,089<br>(625–6,525)                |
|   | 30                           | 1                   | 495<br>(490–500)             | 2,315<br>(2,025–2,525)       | 5,446<br>(3,275–6,025)              |
| E7  | 28                           | 1                   | 794<br>(775–900)             | 4,892<br>(2,775–6,275)       | 9,008<br>(3,775–12,525)             |
| E8  | 0.1                          | 1                   | 415<br>(270–725)             | 1,193<br>(625–4,275)         | 1,818<br>(825–8,525)                |
|   | 45.75                        | 1                   | 952<br>(900–975)             | 6,294<br>(3,025–9,525)       | 12,263<br>(4,275–20,025)            |
| E9  | 0.1                          | 1                   | 573<br>(320–1,025)           | 1,516<br>(725–7,275)         | 2,411<br>(950–14,275)               |

**Table 3.4-21: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Low-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: Low-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|---|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E10   | 0.1                          | 1                   | 715<br>(370–1,525)           | 2,088<br>(825–28,275)        | 4,378<br>(1,025–32,275)             |
| E11   | 45.75                        | 1                   | 1,881<br>(1,525–2,275)       | 12,425<br>(4,275–27,275)     | 23,054<br>(7,025–65,275)            |
|   | 91.4                         | 1                   | 1,634<br>(1,275–2,525)       | 5,686<br>(3,775–11,275)      | 11,618<br>(5,525–64,275)            |
| E12   | 0.1                          | 1                   | 790<br>(420–2,775)           | 2,698<br>(925–25,275)        | 6,032<br>(1,025–31,275)             |
|   |                              | 4                   | 1,196<br>(575–6,025)         | 6,876<br>(1,525–31,275)      | 13,073<br>(3,775–64,275)            |

<sup>1</sup>Average distance (meters) to PTS, TTS, and behavioral thresholds are depicted above the minimum and maximum distances which are in parentheses. Values depict the range produced by SEL hearing threshold criteria levels.

Notes: PTS = permanent threshold shift, SEL = sound exposure level, TTS = temporary threshold shift

**Table 3.4-22: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Low-Frequency Cetaceans**

| <i>Range to Effects for Explosives: Low-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|---|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E1  | 0.1                          | 135<br>(130–140)             | 249<br>(220–270)             |
| E2  | 0.1                          | 173<br>(120–180)             | 305<br>(180–330)             |
| E3  | 0.1                          | 292<br>(240–310)             | 499<br>(330–550)             |
|   | 18.25                        | 310<br>(310–310)             | 583<br>(550–600)             |
| E4  | 10                           | 396<br>(390–420)             | 738<br>(725–750)             |
|   | 60                           | 420<br>(380–775)             | 846<br>(575–2,025)           |
| E5  | 0.1                          | 451<br>(310–525)             | 740<br>(410–1,025)           |
|   | 30                           | 521<br>(490–600)             | 971<br>(925–1,025)           |

**Table 3.4-22: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Low-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: Low-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|---|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E6  | 0.1                          | 547<br>(350–700)             | 842<br>(460–1,275)           |
|   | 30                           | 622<br>(600–650)             | 1,025<br>(1,025–1,025)       |
| E7  | 28                           | 927<br>(900–950)             | 1,524<br>(1,275–1,525)       |
| E8  | 0.1                          | 799<br>(450–925)             | 1,030<br>(575–1,775)         |
|   | 45.75                        | 1,025<br>(1,025–1,025)       | 1,778<br>(1,525–2,025)       |
| E9  | 0.1                          | 947<br>(500–1,275)           | 1,294<br>(675–3,025)         |
| E10   | 0.1                          | 1,032<br>(550–1,775)         | 1,388<br>(800–4,275)         |
| E11   | 45.75                        | 1,778<br>(1,525–2,025)       | 3,067<br>(2,275–11,275)      |
|   | 91.4                         | 1,676<br>(1,275–3,275)       | 2,442<br>(2,025–3,525)       |
| E12   | 0.1                          | 1,151<br>(625–2,525)         | 1,762<br>(900–5,275)         |

<sup>1</sup> Average distance (meters) is shown with the minimum and maximum distances due to varying propagation environments in parentheses.

Notes: PTS = permanent threshold shift, TTS = temporary threshold shift

**Table 3.4-23: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Mid-Frequency Cetaceans**

| <i>Range to Effects for Explosives: Mid-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|---|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E1  | 0.1                          | 1                   | 25<br>(25–25)                | 116<br>(110–120)             | 199<br>(190–210)                    |
|   |                              | 18                  | 94<br>(90–100)               | 415<br>(390–440)             | 646<br>(525–700)                    |
| E2  | 0.1                          | 1                   | 30<br>(30–35)                | 146<br>(140–170)             | 248<br>(230–370)                    |
|   |                              | 5                   | 63<br>(60–70)                | 301<br>(280–410)             | 481<br>(430–675)                    |
| E3  | 0.1                          | 1                   | 50<br>(50–50)                | 233<br>(220–250)             | 381<br>(360–400)                    |
|   |                              | 12                  | 155<br>(150–160)             | 642<br>(525–700)             | 977<br>(700–1,025)                  |
|   | 18.25                        | 1                   | 40<br>(40–40)                | 202<br>(190–220)             | 332<br>(320–350)                    |
|   |                              | 12                  | 126<br>(120–130)             | 729<br>(675–775)             | 1,025<br>(1,025–1,025)              |
| E4  | 10                           | 2                   | 76<br>(70–90)                | 464<br>(410–550)             | 783<br>(650–975)                    |
|   | 60                           | 2                   | 60<br>(60–60)                | 347<br>(310–675)             | 575<br>(525–900)                    |
| E5  | 0.1                          | 20                  | 290<br>(280–300)             | 1,001<br>(750–1,275)         | 1,613<br>(925–3,275)                |
|   | 30                           | 20                  | 297<br>(240–420)             | 1,608<br>(1,275–2,775)       | 2,307<br>(2,025–2,775)              |
| E6  | 0.1                          | 1                   | 98<br>(95–100)               | 430<br>(400–450)             | 669<br>(550–725)                    |
|   | 30                           | 1                   | 78<br>(75–80)                | 389<br>(370–410)             | 619<br>(600–650)                    |
| E7  | 28                           | 1                   | 110<br>(110–110)             | 527<br>(500–575)             | 1,025<br>(1,025–1,025)              |
| E8  | 0.1                          | 1                   | 162<br>(150–170)             | 665<br>(550–700)             | 982<br>(725–1,025)                  |
|   | 45.75                        | 1                   | 127<br>(120–130)             | 611<br>(600–625)             | 985<br>(950–1,025)                  |
| E9  | 0.1                          | 1                   | 215<br>(210–220)             | 866<br>(625–1,000)           | 1,218<br>(800–1,525)                |

**Table 3.4-23: SEL-Based Ranges to Onset PTS, Onset TTS, and Behavioral Reaction (in meters) for Mid-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: Mid-Frequency Cetaceans<sup>1</sup></i> |                              |                     |                              |                              |                                     |
|---|------------------------------|---------------------|------------------------------|------------------------------|-------------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> | <i>Range to Behavioral (meters)</i> |
| E10   | 0.1                          | 1                   | 270<br>(250–280)             | 985<br>(700–1,275)           | 1,506<br>(875–2,525)                |
| E11   | 45.75                        | 1                   | 241<br>(230–250)             | 1,059<br>(1,000–1,275)       | 1,874<br>(1,525–2,025)              |
|   | 91.4                         | 1                   | 237<br>(230–270)             | 1,123<br>(900–2,025)         | 1,731<br>(1,275–2,775)              |
| E12   | 0.1                          | 1                   | 332<br>(320–370)             | 1,196<br>(825–1,525)         | 1,766<br>(1,025–3,525)              |
|   |                              | 4                   | 572<br>(500–600)             | 1,932<br>(1,025–4,025)       | 2,708<br>(1,275–6,775)              |

<sup>1</sup>Average distance (meters) to PTS, TTS, and behavioral thresholds are depicted above the minimum and maximum distances which are in parentheses. Values depict the range produced by SEL hearing threshold criteria levels.

Notes: PTS = permanent threshold shift, SEL = sound exposure level, TTS = temporary threshold shift

**Table 3.4-24: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Mid-Frequency Cetaceans**

| <i>Range to Effects for Explosives: Mid-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|---|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E1  | 0.1                          | 43<br>(40–45)                | 84<br>(80–90)                |
| E2  | 0.1                          | 58<br>(55–60)                | 105<br>(95–110)              |
| E3  | 0.1                          | 98<br>(95–100)               | 183<br>(170–190)             |
|   | 18.25                        | 100<br>(100–100)             | 180<br>(180–180)             |
| E4  | 10                           | 120<br>(120–120)             | 255<br>(250–260)             |
|   | 60                           | 123<br>(120–130)             | 239<br>(230–340)             |
| E5  | 0.1                          | 155<br>(150–160)             | 288<br>(270–300)             |
|   | 30                           | 168<br>(160–190)             | 310<br>(290–350)             |

**Table 3.4-24: Peak Pressure Based Ranges to Onset PTS and Onset TTS (in meters) for Mid-Frequency Cetaceans (continued)**

| <i>Range to Effects for Explosives: Mid-Frequency Cetaceans<sup>1</sup></i> |                              |                              |                              |
|---|------------------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E6  | 0.1                          | 197<br>(190–210)             | 359<br>(320–400)             |
|   | 30                           | 200<br>(200–200)             | 380<br>(380–380)             |
| E7  | 28                           | 296<br>(290–300)             | 525<br>(525–525)             |
| E8  | 0.1                          | 333<br>(310–340)             | 574<br>(440–625)             |
|   | 45.75                        | 351<br>(350–370)             | 629<br>(625–725)             |
| E9  | 0.1                          | 442<br>(370–460)             | 757<br>(500–850)             |
| E10   | 0.1                          | 546<br>(420–700)             | 939<br>(550–1,275)           |
| E11   | 45.75                        | 662<br>(650–800)             | 1,104<br>(1,025–1,275)       |
|   | 91.4                         | 748<br>(600–1,525)           | 1,353<br>(1,000–2,525)       |
| E12   | 0.1                          | 663<br>(470–725)             | 1,064<br>(625–1,275)         |

<sup>1</sup> Average distance (meters) is shown with the minimum and maximum distances due to varying propagation environments in parentheses.

Notes: PTS = permanent threshold shift, TTS = temporary threshold shift

#### **3.4.2.2.3 Impacts from Explosive Stressors Under the Action Alternatives**

Under Alternative 1, there could be fluctuation in the amount of explosives use that could occur annually, although potential impacts would be similar from year to year. The number of impulsive sources in this SEIS/OEIS compared with the totals analyzed in the 2015 MITT Final EIS/OEIS are described in Table 2.5-1 and Table 2.5-2. This comparison applies to both Alternative 1 and Alternative 2, because the number of explosives used would be almost identical under each alternative.

The number of torpedo testing events (both explosive and non-explosive) planned under Alternative 1 testing can vary slightly from year to year; however, all other training and testing activities that involve the use of explosives would remain consistent from year to year. Alternative 1 results are presented for a maximum explosive use year; however, during most years, explosive use would be less, resulting in fewer potential impacts. The numbers of activities planned under Alternative 2 are consistent from year to year and would increase slightly compared to activities planned under Alternative 1. The numbers of explosives used under each alternative are described in Section 3.0.4.2 (Explosive Stressors).

## Presentation of Estimated Impacts from the Quantitative Analysis

The results of the analysis of potential impacts on marine mammals from explosives (see above Section 3.4.2.2.2.1, Methods for Analyzing Impacts from Explosives) are discussed below. The numbers of potential impacts estimated for individual species of marine mammals from exposure to explosive energy and sound for training activities under Alternative 1 and 2 are shown in Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Additionally, estimated numbers of potential impacts from the quantitative analysis for each species are presented below (e.g., Figure 3.4-65). The most likely regions and activity categories from which the impacts could occur are displayed in the impact graphics for each species. There is a potential for impacts to occur anywhere within the Study Area where sound and energy from explosions and the species overlap, although only regions or activity categories where 0.5 percent of the impacts, or greater, are estimated to occur are graphically represented on the impact graphics below. All (i.e., grand total) estimated impacts are also included, regardless of region or category.

### Mysticetes

Mysticetes may be exposed to sound and energy from explosions associated with training and testing activities throughout the year. Explosions produce sounds that are within the hearing range of mysticetes (see Section 3.4.1.6, Hearing and Vocalization). Potential impacts from explosive energy and sound include non-auditory injury, behavioral reactions, physiological stress, masking, and hearing loss. The quantitative analysis estimates the number of behavioral reactions and TTS in mysticetes. Impact ranges for mysticetes exposed to explosive sound and energy are discussed under low-frequency cetaceans in Section 3.4.2.2.2.2 (Impact Ranges for Explosives).

Mysticetes that do experience threshold shift from explosive sounds may have reduced ability to detect biologically important sounds (e.g., social vocalizations) until their hearing recovers. Recovery from threshold shift begins almost immediately after the noise exposure ceases and can take a few minutes to a few days, depending on the severity of the initial shift, to recover. TTS would recover fully and PTS would leave some residual hearing loss. Threshold shifts do not necessarily affect all hearing frequencies equally, and typically manifest themselves at the exposure frequency or within an octave above the exposure frequency. Noise from explosions is broadband with most energy below a few hundred Hertz; therefore, any hearing loss from exposure to explosive sounds is likely to be broadband with effects predominantly at lower frequencies. During the short period that a mysticete had TTS, or permanently for PTS, social calls from conspecifics could be more difficult to detect or interpret, the ability to detect predators may be reduced, and the ability to detect and avoid sounds from approaching vessels or other stressors might be reduced. It is unclear how or if mysticetes use sound for finding prey or feeding; therefore, it is unknown whether a TTS would affect a mysticete's ability to locate prey or rate of feeding.

Research and observations of auditory masking in marine mammals due to impulsive sounds are discussed in Section 3.4.2.2.1.4 (Masking). Explosions introduce low-frequency, broadband sounds into the environment, which could mask hearing thresholds in mysticetes that are nearby, although sounds from explosions last for only a few seconds at most. Masking due to time-isolated detonations would not be significant. Activities that have multiple detonations such as some naval gunfire exercises could create some masking for mysticetes in the area over the short duration of the event. Potential costs to mysticetes from masking are similar to those discussed above for TTS, with the primary difference being

that the effects of masking are only present when the sound from the explosion is present within the water and the effect is over the moment the sound has ceased.

Research and observations (see Section 3.4.2.2.1.5, Behavioral Reactions) show that if mysticetes are exposed to impulsive sounds such as those from explosives, they may react in a variety of ways, which may include alerting, startling, breaking off feeding dives and surfacing, diving or swimming away, changing vocalization, or showing no response at all. Overall, mysticetes have been observed to be more reactive to acoustic disturbance when a noise source is located directly on their migration route. Mysticetes disturbed while migrating could pause their migration or route around the disturbance. Animals disturbed while engaged in other activities such as feeding or reproductive behaviors may be more likely to ignore or tolerate the disturbance and continue their natural behavior patterns. Because noise from most activities using explosives is short-term and intermittent, and because detonations usually occur within a small area, behavioral reactions from mysticetes are likely to be short-term and low to moderate severity.

Physiological stress could be caused by injury or hearing loss and could accompany any behavioral reaction as well. Research and observations of physiological stress in marine mammals are discussed in Section 3.4.2.2.1.3 (Physiological Stress). Due to the short-term and intermittent use of explosives, physiological stress is also likely to be short-term and intermittent. Long-term consequences from physiological stress due to the sound of explosives would not be expected.

***Blue Whale (Endangered Species Act-Listed)***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Blue whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no blue whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of blue whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 1 would not affect ESA-listed blue whales.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Blue whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no blue whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of blue whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 2 would not affect ESA-listed blue whales.*


***Bryde's Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Bryde's whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS (see Figure 3.4-65 and tabular results

in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for mysticetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of Bryde’s whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-65: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly in location and number (see Figure 3.4-66) compared to the impacts shown and discussed above

in Impacts from Explosives Under Alternative 1 for Training and Testing Activities. The primary distinction is that explosive use would be consistent year to year under Alternative 2 and the total number of activities would increase slightly compared to Alternative 1.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of Bryde’s whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-66: Bryde’s Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Fin Whale (Endangered Species Act-Listed)***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Fin whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no fin whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of fin whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 1 would not affect ESA-listed fin whales.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Fin whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no fin whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of fin whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 2 would not affect ESA-listed fin whales.*


**Humpback Whale (Endangered Species Act-Listed)**

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Humpback whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS (see Figure 3.4-67 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for mysticetes above, even a few minor to moderate TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of humpback whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 1 may affect ESA-listed humpback whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.

(2) MIRC = Mariana Islands Range Complex


**Figure 3.4-67: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly in location and number (see Figure 3.4-68) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities. The primary distinction is that explosive use would be consistent year to year under Alternative 2 and the total number of activities would increase slightly compared to Alternative 1.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of humpback whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 2 may affect ESA-listed humpback whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-68: Humpback Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Minke Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Minke whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS (see Figure 3.4-69 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for mysticetes above, even a few minor to moderate behavioral reactions or TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of minke whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-69: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training with explosives would differ slightly in activity and location (see Figure 3.4-70) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities; however, the total number of impacts would remain the same.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of minke whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-70: Minke Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Omura's Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Omura's whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no Omura's whales would be impacted. Long-term consequences for individuals or the species would not be expected.


*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of Omura's whales incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Omura's whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 2, estimates behavioral reactions (see Figure 3.4-71 and tabular

results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for mysticetes above, even a few minor to moderate behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of Omura's whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No TTS, PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-71: Omura's Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**


*Sei Whale (Endangered Species Act-Listed)*

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Sei whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS (see Figure 3.4-72 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for mysticetes above, even a few minor to moderate behavioral reactions or TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of sei whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 1 may affect ESA-listed sei whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-72: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training with explosives would differ slightly in location (see Figure 3.4-73) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities; however, the total number of impacts would remain the same.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of sei whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 2 may affect ESA-listed sei whales.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No TTS, or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-73: Sei Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

**Odontocetes**

Odontocetes may be exposed to sound and energy from explosives associated with training and testing activities throughout the year. Explosions produce sounds that are within the hearing range of odontocetes (see Section 3.4.1.6, Hearing and Vocalization). Potential impacts from explosive energy and sound include non-auditory injury, behavioral reactions, physiological stress, masking and hearing loss. Impact ranges for odontocetes exposed to explosive sound and energy are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives) under mid-frequency cetaceans for most species, and under high-frequency cetaceans for Kogia whales.

Non-auditory injuries to odontocetes, if they did occur, could include anything from mild injuries that are recoverable and are unlikely to have long-term consequences, to more serious injuries, including mortality. It is possible for marine mammals to be injured or killed by an explosion in isolated instances. Animals that did sustain injury could have long-term consequences for that individual. Considering that dolphin species for which these impacts are predicted have populations with tens to hundreds of

thousands of animals, removing several animals from the population would be unlikely to have measurable long-term consequences for the species or stocks. As discussed in Section 5.3.3 (Explosive Stressors), the Navy will implement procedural mitigation measures to delay or cease detonations when a marine mammal is sighted in a mitigation zone to avoid or reduce potential explosive impacts.

Odontocetes that do experience a hearing threshold shift from explosive sounds may have reduced ability to detect biologically important sounds (e.g., social vocalizations) until their hearing recovers. Recovery from a hearing threshold shift begins almost immediately after the noise exposure ceases. A threshold shift can take a few minutes to a few days, depending on the severity of the initial shift, to recover. TTS would recover fully and PTS would leave some residual hearing loss. Threshold shifts do not necessarily affect all hearing frequencies equally, and typically manifest themselves at the exposure frequency or within an octave above the exposure frequency. Noise from explosions is broadband with most energy below a few hundred Hertz; therefore, any hearing loss from exposure to explosive sounds is likely to be broadband with effects predominantly at lower frequencies. During the period that an odontocete had hearing loss, social calls from conspecifics and sounds from predators such as killer whale vocalizations could be more difficult to detect or interpret, although many of these sounds may be above the frequencies of the threshold shift. Odontocetes use echolocation clicks to find and capture prey. These echolocation clicks and vocalizations are at frequencies above a few kHz, which are less likely to be affected by threshold shift at lower frequencies, and should not affect odontocete's ability to locate prey or rate of feeding.

Research and observations of masking in marine mammals due to impulsive sounds are discussed in Section 3.4.2.2.1.4 (Masking). Explosions introduce low-frequency, broadband sounds into the environment, which could mask hearing thresholds in odontocetes that are nearby, although sounds from explosions last for only a few seconds at most. Also, odontocetes typically communicate, vocalize, and echolocate at higher frequencies that would be less affected by masking noise at lower frequencies such as those produced by an explosion. Masking due to time-isolated detonations would not be significant. Activities that have multiple detonations such as some naval gunfire exercises could create some masking for odontocetes in the area over the short duration of the event. Potential costs to odontocetes from masking are similar to those discussed above for TTS, with the primary difference being that the effects of masking are only present when the sound from the explosion is present within the water and the effect is over the moment the sound has ceased.

Research and observations (see Section 3.4.2.2.1.5, Behavioral Reactions) show that odontocetes do not typically show strong behavioral reactions to impulsive sounds such as explosions. Reactions, if they did occur, would likely be limited to short ranges, within a few kilometers of multiple explosions. Reactions could include alerting, startling, breaking off feeding dives and surfacing, diving or swimming away, change in vocalization, or showing no response at all. Animals disturbed while engaged in other activities such as feeding or reproductive behaviors may be more likely to ignore or tolerate the disturbance and continue their natural behavior patterns. Because noise from most activities using explosives is short-term and intermittent, and because detonations usually occur within a small area, behavioral reactions from odontocetes are likely to be short-term and low to moderate severity.

Physiological stress could be caused by injury or hearing loss and could accompany any behavioral reaction as well. Research and observations of physiological stress in marine mammals are discussed in Section 3.4.2.2.1.3 (Physiological Stress). Due to the short-term and intermittent use of explosives, physiological stress is also likely to be short-term and intermittent. Long-term consequences from physiological stress due to the sound of explosives would not be expected.

***Beaked Whales***


Beaked whales within the Study Area include: Blainville's beaked whale, Cuvier's beaked whale, ginkgo-toothed beaked whale, and Longman's beaked whale. Research and observations (Section 3.4.2.2.1.5, Behavioral Reactions) show that beaked whales are sensitive to human disturbance including noise from sonars, although no research on specific reactions to impulsive or explosion noise is available.

Odontocetes overall have shown little responsiveness to impulsive sounds although it is likely that beaked whales are more reactive than most other odontocetes. Reactions could include alerting, startling, breaking off feeding dives and surfacing, diving or swimming away, change in vocalization, or showing no response at all. Beaked whales on Navy ranges have been observed leaving the area for a few days during sonar training exercises. It is reasonable to expect that animals may leave an area of more intense explosive activity for a few days, however most explosive use during Navy activities is short-duration consisting of only a single or few closely timed explosions (i.e., detonated within a few minutes) with a limited footprint due to a single detonation point. Because noise from most activities using explosives is short-term and intermittent and because detonations usually occur within a small area, behavioral reactions from beaked whales are likely to be short-term and moderate severity.

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**


Beaked whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS for only ginkgo-toothed and Longman's beaked whales (see Figure 3.4-74 and Figure 3.4-75, and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for these species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). No impacts are estimated for Blainville's beaked whale or Cuvier's beaked whale. As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of ginkgo-toothed and Longman's beaked whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-74: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.


(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-75: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**


The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions and TTS for only ginkgo-toothed and Longman’s beaked whales. Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly in location (see Figure 3.4-76 and Figure 3.4-77) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities; however, the total number of impacts would remain the same.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of ginkgo-toothed and Longman’s beaked whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.  
 (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-76: Ginkgo-Toothed Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species.  
(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-77: Longman’s Beaked Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Common Bottlenose Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Bottlenose dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no bottlenose dolphins would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of bottlenose dolphins incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Bottlenose dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no bottlenose

dolphins would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of bottlenose dolphins incidental to those activities.*


#### ***Kogia Whales***

Kogia whales include two species that are often difficult to distinguish from one another: dwarf sperm whales and pygmy sperm whales; however, impacts to the populations of dwarf and pygmy sperm whales are modeled separately. TTS and PTS thresholds for high-frequency cetaceans, such as Kogia whales are lower than for all other marine mammals, which leads to a higher number of estimated hearing loss impacts relative to the number of animals exposed to the sound as compared to other hearing groups (e.g., mid-frequency cetaceans).

#### **Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Kogia whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions, TTS, and PTS (see Figure 3.4-78 through Figure 3.4-79, and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for these species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. PTS in an individual could have no to minor long-term consequences for individuals although a single minor long-term consequence for an individual is unlikely to lead to long-term consequences for a population. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.


*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of dwarf and pygmy sperm whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species.

(2) MIRC = Mariana Islands Range Complex

**Figure 3.4-78: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-79: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**


Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly (see Figure 3.4-80 and Figure 3.4-81) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of dwarf and pygmy sperm whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-80: Dwarf Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-81: Pygmy Sperm Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***False Killer Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

False killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no false killer whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of false killer whales incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training Activities**

False killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no false killer

whales would be impacted. Long-term consequences for individuals or the species would not be expected.


*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of false killer whales incidental to those activities.*

**Fraser's Dolphin**

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Fraser's dolphin may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 1, estimates behavioral reaction, TTS, and PTS (see Figure 3.4-82 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. PTS in an individual could have no to minor long-term consequences for individuals although a single minor long-term consequence for an individual is unlikely to lead to long-term consequences for a population. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of Fraser's dolphins incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-82: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would increase slightly (see Figure 3.4-83) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of Fraser’s dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-83: Fraser’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Killer Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no killer whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of killer whales incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training Activities**

Killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no killer whales would be impacted. Long-term consequences for individuals or the species would not be expected.


*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of killer whales incidental to those activities.*

**Melon-Headed Whale**

**Impacts from Explosives Under Alternative 1 for Training Activities**

Melon-headed whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 1, estimates behavioral reaction and TTS (see Figure 3.4-84 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of melon-headed whales incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-84: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly by activity and location (see Figure 3.4-85) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities; however, the total number of impacts would remain the same.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of melon-headed whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No impacts are estimated for training activities. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-85: Melon-Headed Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**


***Pantropical Spotted Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Pantropical spotted dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosives per year under Alternative 1, estimates behavioral reactions, TTS and PTS (see Figure 3.4-86 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. PTS could reduce an animal’s ability to detect biologically important sounds; however, as discussed above, a small threshold shift due to exposure to sonar is unlikely to affect the hearing range that pantropical spotted dolphins rely upon. Nevertheless, PTS could have minor long-term consequences for individuals. This minor consequence for

an individual is unlikely to have any long-term consequences for the species. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of pantropical spotted dolphins incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-86: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would increase slightly (see Figure 3.4-87) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of pantropical spotted dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-87: Pantropical Spotted Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Pygmy Killer Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Pygmy killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no pygmy killer whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of pygmy killer whales incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training Activities**

Pygmy killer whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no pygmy killer whales would be impacted. Long-term consequences for individuals or the species would not be expected.


*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of pygmy killer whales incidental to those activities.*

***Risso's Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Risso's dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 1, estimates a behavioral reaction and TTS (see Figure 3.4-88 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS or behavioral reactions to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of Risso's dolphins incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-88: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly (see Figure 3.4-89) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of Risso’s dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-89: Risso’s Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Rough-Toothed Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Rough-toothed dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no rough-toothed dolphins would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of rough-toothed dolphins incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Rough-toothed dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no rough-

toothed dolphins would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of rough-toothed dolphins incidental to those activities.*

***Short-Finned Pilot Whale***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**


Short-finned pilot whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no short-finned pilot whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of short-finned pilot whales incidental to those activities.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Short-finned pilot whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 2, estimates behavioral reaction (see Figure 3.4-90 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of short-finned pilot whales incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No TTS, PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-90: Short-Finned Pilot Whale Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

***Sperm Whale (Endangered Species Act-Listed)***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Sperm whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no sperm whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would not result in the unintentional taking of sperm whales incidental to those activities.*

*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 1 would not affect ESA-listed sperm whales.*

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Sperm whales may be exposed to sound or energy from explosions associated with training and testing activities throughout the year, although the quantitative analysis estimates that no sperm whales would be impacted. Long-term consequences for individuals or the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would not result in the unintentional taking of sperm whales incidental to those activities.*


*Pursuant to the ESA, the use of explosives during training and testing activities as described under Alternative 2 would not affect ESA-listed sperm whales.*

***Spinner Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Spinner dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 1, estimates TTS (see Figure 3.4-91 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to moderate TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of spinner dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No behavioral responses, PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex


**Figure 3.4-91: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Spinner dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 2, estimates TTS and PTS (see Figure 3.4-92 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. As described for odontocetes above, even a few minor to moderate TTS to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. PTS could reduce an animal’s ability to detect biologically important sounds; however, as discussed above, a small threshold shift due to exposure to sonar is unlikely to affect the hearing range that spinner dolphins rely upon. Nevertheless, PTS could have minor long-term consequences for individuals. This minor consequence for an individual is unlikely to have any long-term consequences for the species. Considering these factors and the mitigation measures that will be

implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of spinner dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No behavioral responses or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-92: Spinner Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**


***Striped Dolphin***

**Impacts from Explosives Under Alternative 1 for Training and Testing Activities**

Striped dolphins may be exposed to sound or energy from explosions associated with training and testing activities throughout the year. The quantitative analysis, using the maximum number of explosions per year under Alternative 1, estimates behavioral reaction and TTS (see Figure 3.4-93 and tabular results in Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Estimated impacts most years would be less based on fewer explosions. Impact ranges for this species are discussed in Section 3.4.2.2.2.2 (Impact Ranges for Explosives). As described for odontocetes above, even a few minor to

moderate TTS or behavioral reaction to an individual over the course of a year are unlikely to have any significant costs or long-term consequences for that individual. Considering these factors and the mitigation measures that will be implemented as described in Chapter 5 (Mitigation), long-term consequences for the species would not be expected.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 1 would result in the unintentional taking of striped dolphins incidental to those activities.*




Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-93: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

**Impacts from Explosives Under Alternative 2 for Training and Testing Activities**

Potential annual impacts under Alternative 2 from training and testing with explosives would differ slightly (see Figure 3.4-94) compared to the impacts shown and discussed above in Impacts from Explosives Under Alternative 1 for Training and Testing Activities; however, the total number of impacts would remain the same.

*Pursuant to the MMPA, the use of explosives during training and testing activities as described under Alternative 2 would result in the unintentional taking of striped dolphins incidental to those activities.*



Notes: (1) Region and Activity bar charts show categories +/- 0.5 percent of the estimated impacts. Estimated impacts most years would be less based on fewer explosions. No PTS or injuries (non-auditory) are estimated for this species. (2) MIRC = Mariana Islands Range Complex

**Figure 3.4-94: Striped Dolphin Impacts Estimated per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 2**

**3.4.2.2.2.4 Impacts from Explosive Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosive stressors, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that use explosives within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from explosive stressors on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

### 3.4.2.3 Energy Stressors

Energy stressors are discussed in Section 3.0.4.3 (Energy Stressors) of this SEIS/OEIS. The energy stressors that may impact marine mammals include in-water electromagnetic devices and high-energy lasers. NMFS has previously determined in documents and analyses associated with two prior Navy training and testing EIS/OEISs within the MITT Study Area that in-water electromagnetic devices would not result in harassment or the incidental taking of marine mammals (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). These determinations for this stressor were recently reaffirmed for a third time by NMFS for the same actions elsewhere (83 FR 10954 & 83 FR 29872). Since the 2015 MITT Final EIS/OEIS and with the increased use of undersea power cables associated with offshore energy generation, there has been renewed scientific interest in electromagnetic fields possibly affecting migrating marine mammals (Gill et al., 2014; Kremers et al., 2014; Kremers et al., 2016; Zellar et al., 2017). Horton et al. (2017) have indicated that future experiments involving empirical observation of free-ranging animals are still required for there to be sufficient evidence demonstrating causal relations between marine mammal movement decisions and environmental cues such as the earth's magnetic field. These additional scientific findings do not change in any way the rationale for the dismissal of in-water electromagnetic devices as presented in the 2015 analyses given a negligible or discountable impact on marine mammal populations or species.

The 2015 MITT Final EIS/OEIS covered the use of low-energy lasers in Section 3.0.5.2.2.3 (Lasers), but high-energy laser weapons were not part of the Proposed Action in the 2015 MITT Final EIS/OEIS. The use of high-energy lasers represents a new sub-stressor as part of an existing activity in this SEIS/OEIS. As discussed in this SEIS/OEIS, Section 3.0.4.3.2.2 (High-Energy Lasers), high-energy lasers are designed to disable surface targets, rendering them immobile. The primary concern is the potential for a marine mammal to be struck with the laser beam at or near the water's surface, where extended exposure could result in injury or death.

As described in Section 3.0.4.3.2 (Lasers), high-energy laser weapons testing activities involve evaluating the effectiveness of a high-energy laser deployed from a surface ship to create small but critical failures in potential targets from short ranges. The concern with the proposed use of high-energy lasers is the potential for a marine mammal to be exposed to the laser beam if the laser beam missed the target, if the animal was above the ocean surface, and if the animal was in the direct path of the laser beam in front of or directly behind the target. The Navy conducted statistical modeling to estimate the probability of a marine mammal being struck by a high-energy laser during training and testing activities. The probability was estimated for a location off Southern California where the density of marine mammals is generally higher than in the Study Area and where more training and testing activities using high-energy lasers are conducted. The results of the analysis showed that there is a very low probability of a direct strike by a high-energy laser on a marine mammal, and that the likelihood of a strike occurring is therefore discountable (U.S. Department of the Navy, 2017b). Given that marine mammal densities are lower in the Study Area and fewer activities using high-energy lasers would be conducted in the Study Area, it is reasonable to conclude that the probability of a direct strike is even lower than predicted off Southern California. Therefore, it is reasonable to conclude that marine mammals in the Study Area are not likely to be struck by a high-energy laser. Training and testing activities have the potential to expose marine mammals that occur within the Study Area to this energy stressor. However, given the short ranges involved in the activities involving high-energy lasers, the aim point being a surface target, the inherent precision of the weapon and its targeting system, and the fact that marine

mammals spend up to 90 percent of their time under the water (Costa, 1993; Costa & Block, 2009), indicates that impacts on marine mammals from high-energy lasers should not be expected to occur.

#### **3.4.2.3.1 Impacts from Energy Stressors Under Alternative 1**

Under Alternative 1, the number of proposed training and testing activities involving the use of in-water electromagnetic devices would decrease in comparison to the 2015 MITT Final EIS/OEIS (see Table 3.0-9). These activities would occur in the same locations and in a similar manner as previously analyzed. Therefore, as stated in the 2015 MITT Final EIS/OEIS and based on the new science summarized above, the impact of in-water electromagnetic devices on marine mammals are not expected.

Under Alternative 1, there would be up to 54 activities annually that include the use of high-energy lasers (Table 3.0-10). As discussed above, impacts on marine mammals from high-energy lasers are not expected because of the very low probability of a direct strike by a high-energy laser on a marine mammal.

Impacts on marine mammals from energy stressors, including in-water electromagnetic devices and high-energy lasers, are not expected to occur under Alternative 1.

*The use of in-water electromagnetic devices and high-energy lasers as described under Alternative 1 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of in-water electromagnetic devices and high-energy lasers as described under Alternative 1 may affect ESA-listed marine mammals.*

#### **3.4.2.3.2 Impacts from Energy Stressors Under Alternative 2**

Under Alternative 2, the number of proposed training and testing activities involving energy stressors would be the same as Alternative 1 for in-water electromagnetic devices (Table 3.0-9). Under Alternative 2, the use of high-energy lasers would increase as compared to Alternative 1 (Table 3.0-10). There would be no change regarding the impact conclusions for energy stressors as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, impacts on marine mammals under Alternative 2 from energy stressors, including high-energy lasers, are not expected to occur.

*The use of in-water electromagnetic devices and high-energy lasers as described under Alternative 2 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of in-water electromagnetic devices and high-energy lasers as described under Alternative 2 may affect ESA-listed marine mammals.*

#### **3.4.2.3.3 Impacts from Energy Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with the Proposed Action would continue to occur. Energy stressors from the use of in-water electromagnetic devices and high-energy lasers, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that produce energy stressors within the marine environment where training and testing activities have

historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from energy stressors on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

#### **3.4.2.4 Physical Disturbance and Strike Stressors**

Physical disturbance and strike stressors are discussed in Section 3.0.4.4 (Physical Disturbance and Strike Stressors) of this SEIS/OEIS. The physical disturbance and strike stressors that may impact marine mammals include (1) vessels and in-water devices, (2) military expended materials, and (3) seafloor devices. The annual number of activities including vessels and in-water devices, the annual number of military expended materials, and the annual number of activities including seafloor devices are shown in Table 3.0-12 through Table 3.0-18.

Since 1995, the U.S. Navy has reported all known or suspected vessel collisions with whales to NMFS, and there have been no known collisions between Navy vessels and whales in the MITT Study Area associated with any of the proposed training or testing activities. The Navy has several standard operating procedures and mitigation measures for vessel safety that will benefit marine mammals through a reduction in the potential for vessel strike, as discussed in Section 2.3.3 (Standard Operating Procedures) and Chapter 5 (Mitigation). Based on the absence of any Navy vessel strikes associated with the Proposed Action in the Study Area, the general reduction in strike incidents Navy-wide since introduction of the Marine Species Awareness Training in 2006, and the future reduction in vessel and in-water device use in comparison to the ongoing actions (see Tables 3.0-12 and 3.0-13), the Navy does not anticipate the occurrence of future vessel strikes to marine mammals within the Study Area during training and testing activities. For these reasons, the Navy is not requesting authorization of a take by vessel strike during Navy training and testing activities in the MITT Study Area.

Most in-water devices, such as unmanned underwater vehicles and towed devices, will move slowly through the water and are highly unlikely to strike marine mammals because the mammal could easily avoid the device. In-water devices towed by manned platforms are unlikely to strike a marine mammal because of the observers on the towing platform and other standard safety measures employed when towing in-water devices. In-water devices that could pose a higher risk to marine mammals are those operated at high speeds and unmanned, such as torpedoes. The Navy reviewed torpedo design features and a large number of previous anti-submarine warfare torpedo exercises to assess the potential of torpedo strikes on marine mammals. The tactical software that guides U.S. Navy torpedoes is sophisticated and should not identify a marine mammal as a target. All training and testing torpedoes are recovered after being fired at targets and are reconfigured for re-use. Review of the exercise torpedo records indicates there has never been an impact on a marine mammal. In thousands of exercises in which torpedoes were fired or in-water devices used, there have been no recorded or reported instances of a marine mammal strike.

As part of military expended materials, small-caliber munitions are inert, are meant to be aimed at targets, and are not long-range weapons. As a result, marine mammals are extremely unlikely to be disturbed or struck by expended small-caliber munitions. There have been no known instances of a seafloor device (such as an anchor) striking a marine mammal as it was being deployed or recovered.

In short, there have been no known instances of physical disturbance or strike to any marine mammals as a result of training and testing activities prior to or since the 2015 MITT Final EIS/OEIS. As described in Section 5.3.4 (Physical Disturbance and Strike Stressors), the Navy will continue to implement mitigation measures for applicable vessel movements, towed in-water devices, and military expended materials

during non-explosive activities. The mitigation measures will further avoid or reduce the already low potential for impacts on marine mammals during activities involving physical disturbance or strike stressors.

NMFS has previously determined in documents and analyses associated with two prior Navy training and testing EIS/OEISs within the MITT Study Area that physical disturbance and strike stressors would not result in harassment or the incidental taking of marine mammals (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). There has been no subsequent emergent science that would necessitate changes to these conclusions, reached in association with the 2015 MITT Final EIS/OEIS analyses, regarding physical disturbance and strike stressors being dismissed as having a negligible or discountable impact on marine mammal populations or species.

#### **3.4.2.4.1 Impacts from Physical Disturbance and Strike Stressors Under Alternative 1**

Under Alternative 1, analysis of the individual sub-stressors including the use of vessels and in-water devices, military expended materials, and seafloor devices presented in Section 3.0.4.4 (Physical Disturbance and Strike Stressors) indicates that those items having the most potential to affect marine mammals have decreased in comparison to the 2015 MITT Final EIS/OEIS (Tables 3.0-12 through 3.0-18). This assumes the dismissal of small-caliber munitions for the reasons noted above.

Given the reduction in physical disturbance and strike stressors for this SEIS/OEIS, the findings presented in the 2015 MITT Final EIS/OEIS, Section 3.4.4.4 (Physical Disturbance and Strike Stressors), the MMPA authorization (80 FR 46112), and the NMFS Biological Opinion, the findings associated with the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015b) remain valid.

*The use of vessels and in-water devices, military expended materials, and seafloor devices under Alternative 1 would not result in the unintentional taking of marine mammals incidental to those activities as defined as defined under MMPA.*

*Pursuant to the ESA, the use of vessels and in-water devices, military expended materials, and seafloor devices as summarized above under Alternative 1 may affect ESA-listed marine mammals.*

#### **3.4.2.4.2 Impacts from Physical Disturbance and Strike Stressors Under Alternative 2**

Under Alternative 2, physical disturbance and strike stressors during training and testing activities would decrease in comparison to the 2015 (Tables 3.0-12 through 3.0-18) assuming the dismissal of small-caliber munitions use for the reasons noted above. Under Alternative 2, there would be additional physical disturbance and strike stressors in comparison to Alternative 1, but the conclusions remain the same. Therefore, impacts on marine mammals from physical disturbance and strike stressors are not expected to occur.

*The use of vessels and in-water devices, military expended materials, and seafloor devices under Alternative 2 would not result in the unintentional taking of marine mammals incidental to those activities as defined under MMPA.*

*Pursuant to the ESA, the use of vessels and in-water devices, military expended materials, and seafloor devices may affect ESA-listed marine mammals.*

#### 3.4.2.4.3 Impacts from Physical Disturbance and Strike Stressors Under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with the Proposed Action would continue to occur. Physical disturbance and strike stressors from the use of vessels and in-water devices, military expended materials, and seafloor devices, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing training and testing activities would result in fewer activities that produce physical disturbance and strike stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from physical disturbance and strike stressors on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

#### 3.4.2.5 Entanglement Stressors

Entanglement stressors are discussed in Section 3.0.4.5 (Entanglement Stressors) of this SEIS/OEIS. Entanglement stressors considered for marine mammals include (1) wires and cables, and (2) decelerators/parachutes. The annual numbers of wires and cables and decelerators/parachutes proposed under the alternatives and in comparison to current ongoing activities are presented in Tables 3.0-20 through 3.0-22. There have been no known instances of any marine mammals being entangled in wires and cables, or decelerators/parachutes associated with Navy training and testing activities prior to or since the 2015 MITT Final EIS/OEIS.

NMFS has previously determined in documents and analyses associated with two prior Navy training and testing EIS/OEISs within the MITT Study Area that entanglement stressors would not result in harassment or the incidental taking of marine mammals (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). There has been no subsequent emergent science that would necessitate changes to these conclusions, reached in association with the 2015 MITT Final EIS/OEIS analyses, regarding this stressor being dismissed as having a negligible or discountable impact on marine mammal populations or species. These determinations for this stressor were recently reaffirmed for a third time by NMFS for the same actions elsewhere (83 FR 10954 & 83 FR 29872).

##### 3.4.2.5.1 Impacts from Entanglement Stressors Under Alternative 1

Under Alternative 1, the annual number of entanglement stressors would decrease in comparison to the current ongoing activities (Tables 3.0-20 through 3.0-22). Therefore, the analysis from the 2015 MITT Final EIS/OEIS remains valid. The analysis presented in the 2015 MITT Final EIS/OEIS (Section 3.4.4.5, Entanglement Stressors), the MMPA authorization (80 FR 46112), and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015b) determined that entanglement stressors associated with the Navy's Proposed Action can be dismissed as having a negligible or discountable impact on marine mammal populations or species.

*The use of wires and cables and decelerators/parachutes as described under Alternative 1 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of wires and cables and parachutes/decelerators as described under Alternative 1 may affect ESA-listed marine mammals.*

#### 3.4.2.5.2 Impacts from Entanglement Stressors Under Alternative 2

Under Alternative 2, the number of entanglement stressors would decrease in comparison to current ongoing activities (Tables 3.0-20 through 3.0-22). In comparison to Alternative 1, there would be a slight increase under Alternative 2 for entanglement stressors; however, the combined number of annual entanglement stressors (fiber optic cable, guidance wire, and decelerators/parachutes) decreases when compared to the 2015 MITT Final EIS/OEIS. Therefore, the analysis and conclusions presented in the 2015 MITT Final EIS/OEIS (Section 3.4.4.5, Entanglement Stressors), the MMPA authorization (80 FR 46112), and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015b) remain valid. Impacts on marine mammals from the use of entanglement stressors are not anticipated.

*The use of wires and cables and decelerators/parachutes as described under Alternative 2 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of wires and cables and parachutes/decelerators as described under Alternative 2 may affect ESA-listed marine mammals.*

#### 3.4.2.5.2.1 Impacts from Entanglement Stressors Under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with the Proposed Action would continue to occur. Entanglement stressors from the use of wires and cables and decelerators/parachutes, as described above, would not be introduced into the marine environment from the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer activities that use entanglement stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from entanglement stressors on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

#### 3.4.2.6 Ingestion Stressors

Ingestion stressors are discussed in Section 3.0.4.6 (Ingestion Stressors) of this SEIS/OEIS. Types of materials that could become ingestion stressors during training and testing in the Study Area include military expended materials – munitions and military expended materials – other than munitions. The annual number of activities including military expended materials are shown in Tables 3.0-14 through 3.0-17, 3.0-20 and 3.0-21, and Tables 3.0-23 through 3.0-25. As discussed in Section 3.4.4.6.3 (Impacts from Munitions) of the 2015 MITT Final EIS/OEIS, the number of munitions and explosive munitions fragments that an individual marine mammal could encounter would generally be low, based on the patchy distribution of both the munitions and the habitats where marine mammals forage. For the more numerous small-caliber munitions, these expended material type items are inert, small, do not resemble prey items, and end up as part of the seafloor, where they are unlikely to be encountered by most marine mammals. In addition, it is assumed for marine mammal species that may feed at the seafloor, that they would not ingest every munition or munition's fragment encountered; if a munition or munition's fragment were ingested, an animal may attempt to reject it when it realizes the item is not food. There is evidence indicating that even ingestion of certain metal items (e.g., hooks) may not result in injury or mortality to the individual (Wells et al., 2008; West, 2016).

NMFS has previously determined in documents and analyses associated with two prior Navy training and testing EIS/OEISs within the MITT Study Area that ingestion stressors would not result in harassment or the incidental taking of marine mammals (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). There has been no subsequent emergent science that would necessitate changes to these conclusions reached in association with the 2015 MITT Final EIS/OEIS analyses regarding this stressor being dismissed as having a negligible or discountable impact on marine mammal populations or species. These determinations for this stressor were recently reaffirmed for a third time by NMFS for the same actions elsewhere (83 FR 10954 & 83 FR 29872).

#### **3.4.2.6.1 Impacts from Ingestion Stressors Under Alternative 1**

Under Alternative 1, analysis of the individual sub-stressors presented in Section 3.0.4.6 (Ingestion Stressors) indicates that those items considered ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) having the most potential to affect marine mammals have decreased (Tables 3.0-14 through 3.0.17, Table 3.0-20 and 3.0-21, and Tables 3.0-23 through 3.0-25). For the reasons noted above, the Navy has determined that potential impacts from ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) would not be substantially different from the 2015 MITT Final EIS/OEIS. In the 2015 analysis of training and testing activities within the Study Area, NMFS determined that ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) would not result in harassment or the incidental taking of marine mammals activities (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). The activities expending munitions and other military expended materials analyzed in this SEIS/OEIS under Alternative 1 are not a significant change over what was analyzed in the 2015 MITT Final EIS/OEIS, and there has been no new science necessitating a revision of the 2015 conclusions in that regard. Impacts on marine mammals from ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) associated with Navy activities in the Study Area are not anticipated.

*The use of military expended materials as described under Alternative 1 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of military expended materials as described under Alternative 1 may affect ESA-listed marine mammals.*

#### **3.4.2.6.2 Impacts from Ingestion Stressors Under Alternative 2**

Under Alternative 2, the use of military expended materials would decrease under this SEIS/OEIS in comparison to the ongoing activities, with the exception of increased small-caliber munitions use (Tables 3.0-14 through 3-17 and Tables 3.0-22 through 3.0-24). Under Alternative 2, increases as compared to Alternative 1 do not change the impact conclusions for ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, impacts on marine mammals from ingestion of military expended materials under Alternative 2 are not expected.

*The use of military expended materials as described under Alternative 2 would not result in the unintentional taking of marine mammals incidental to those activities as defined under the MMPA.*

*Pursuant to the ESA, the use of military expended materials as described under Alternative 2 may affect ESA-listed marine mammals.*

#### **3.4.2.6.3 Impacts from Ingestion Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with the Proposed Action would continue to occur. Ingestion stressors from the use of military expended materials – munitions and military expended materials – other than munitions, as described above, would not be introduced into the marine environment under the Proposed Action. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of on-going training and testing activities. Discontinuing the training and testing activities would result in fewer activities that produce ingestion stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would reduce the potential for impacts from ingestion stressors on marine mammals, but would not measurably improve the overall distribution or abundance of marine mammals.

#### **3.4.2.7 Secondary Stressors**

As discussed in Section 3.4.4.7 (Secondary Stressors) of the 2015 MITT Final EIS/OEIS, secondary stressors from training and testing activities were analyzed for potential indirect impacts on marine mammals via habitat degradation or an effect on prey availability. These stressors included (1) explosives, (2) explosive byproducts and unexploded ordnance, (3) metals, (4) chemicals, and (5) transmission of marine mammal diseases and parasites. Analyses of the potential impacts on sediments and water quality are discussed in detail in Section 3.1 (Sediments and Water Quality) of the 2015 MITT Final EIS/OEIS and in Section 3.1 (Sediments and Water Quality) of this SEIS/OEIS. NMFS has previously determined in documents and analyses associated with two prior Navy training and testing EIS/OEISs within the MITT Study Area that secondary stressors would not result in harassment or the incidental taking of marine mammals (80 FR 46112) and would not result in significant adverse impacts or jeopardize the continued existence of any ESA-listed marine mammals (National Oceanic and Atmospheric Administration, 2015b). There has been no subsequent emergent science since 2015 that would necessitate changes to the analysis of secondary stressors being dismissed as having a negligible or discountable impact on marine mammal populations or species. These determinations for secondary stressors were recently reaffirmed for a third time by NMFS for the same actions elsewhere (83 FR 10954 and 83 FR 29872).

The current analysis has concluded that the relatively low solubility of most explosives and their degradation products means that concentrations of these contaminants in the marine environment, from either high-order or low-order detonations, are relatively low and readily diluted. For example, degradation products of Royal Demolition Explosive are not toxic to marine organisms at realistic exposure levels (Rosen & Lotufo, 2010). Any remnant undetonated components from explosives such as TNT, royal demolition explosive, and high melting explosive experience rapid biological and photochemical degradation in marine systems (Cruz-Uribe et al., 2007; Juhasz & Naidu, 2007; Pavlostathis & Jackson, 2002; Singh et al., 2009; Walker et al., 2006).

For undetonated munitions, the concentration of unexploded ordnance, explosion byproducts, metals, and other chemicals would never exceed that of a World War II dump site, where studies found minimal

concentrations were detected only within a few feet of the ordnance (Briggs et al., 2016; Edwards & Beldowski, 2016; Edwards et al., 2016a; Edwards et al., 2016b; Kelley et al., 2016; Koide et al., 2015; University of Hawaii, 2010). Annual surveys conducted in the waters off Farallon de Medinilla from 1999 through 2012 found no evidence that the habitat had changed or been adversely impacted to a significant degree by the training activities that have been conducted there (Smith & Marx, 2016). Therefore, long-term secondary effects on marine mammal habitat or prey would be negligible.

The potential transmission of diseases or parasites from Navy marine mammals to indigenous marine mammals is highly unlikely for the following reasons: the Navy marine mammals spend a very small amount of time in the open ocean; the Navy trainers have excellent control over the animals; the Navy follows procedures for the collection and proper disposal of marine mammal waste; the Navy's marine mammals are screened and receive exceptional veterinarian care; the Navy conducts visual monitoring for indigenous marine mammals to avoid any interactions with Navy marine mammals; and the Navy has a track record of over 40 years, with zero known incidents. As described in detail in Section 3.4.4.7.5 (Transmission of Marine Mammal Diseases and Parasites) in the 2015 MITT Final EIS/OEIS, there is no scientific basis to conclude that the use of Navy marine mammals during training and testing activities would have an indirect impact on wild marine mammals.

*Secondary stressors from training and testing activities as described under Alternative 1, Alternative 2, and the No Action Alternative, would not result in the unintentional taking of marine mammals incidental to those stressors as defined under the MMPA.*

*Pursuant to the ESA, secondary stressors from training and testing activities, as described under Alternative 1, Alternative 2, and the No Action Alternative, may affect ESA-listed marine mammals.*

### **3.4.3 Summary of Potential Impacts on Marine Mammals**

As described in Section 3.0.5.4 (Resource-Specific Impacts Analysis for Multiple Stressors) in the 2015 MITT Final EIS/OEIS, this section evaluates the potential for combined impacts of all the stressors from the Proposed Action. The analysis and conclusions for the potential impacts from each of the individual stressors are discussed in Section 3.4.2.1 (Acoustic Stressors) through Section 3.4.2.6 (Ingestion Stressors) and, for ESA-listed species, summarized in Section 3.4.5 (Endangered Species Act Determinations).

Understanding the combined effects of stressors on marine organisms in general and marine mammal populations in particular is extremely difficult to predict (National Academies of Sciences Engineering and Medicine, 2017). Recognizing the difficulties with measuring trends in marine mammal populations, the focus has been on indicators for adverse impacts, including health and other population metrics (National Academies of Sciences Engineering and Medicine, 2017). This recommended use of population indicators is the approach Navy presented in the 2015 MITT Final EIS/OEIS (Section 3.4.5, Summary of Impacts on Marine Mammals) and formed part of the 2015 analyses by NMFS in their MMPA authorization (80 FR 46112), and the Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015b).

Stressors associated with training and testing activities do not typically occur in isolation but rather occur in some combination. For example, mine neutralization activities include elements of acoustic, physical disturbance and strike, entanglement, ingestion, and secondary stressors that are all coincident in space and time. An analysis of the combined impacts of all stressors considers the potential

consequences of additive stressors and synergistic stressors, as described below. This analysis makes the reasonable assumption, which is supported by the acoustic effects modeling, that the majority of exposures to stressors are non-lethal, and instead focuses on consequences potentially impacting marine mammal fitness (e.g., physiology, behavior, reproductive potential).

There are generally two ways that a marine mammal could be exposed to multiple additive stressors. The first would be if a marine mammal were exposed to multiple sources of stress from a single event or activity within a single event (e.g., a mine warfare event may include the use of a sound source and a vessel). The potential for a combination of these impacts from a single activity would depend on the range to effects of each of the stressors and the response or lack of response to that stressor. Most of the activities proposed under Alternative 1 generally involve the use of moving platforms (e.g., ships, torpedoes, aircraft) that may produce one or more stressors; therefore, it is likely that if a marine mammal were within the potential impact range of those activities, it may be impacted by multiple stressors simultaneously. Individual stressors that would otherwise have minimal to no impact may combine to have a measurable response. However, due to the wide dispersion of stressors, speed of the platforms, general dynamic movement of many training and testing activities, and behavioral avoidance exhibited by many marine mammal species, it is very unlikely that a marine mammal would remain in the potential impact range of multiple sources or sequential events. Exposure to multiple stressors is more likely to occur at an instrumented range where training and testing activities using multiple platforms may be concentrated during a particular event. In such cases involving a relatively small area on an instrumented range, a behavioral reaction resulting in avoidance of the immediate vicinity of the activity would reduce the likelihood of exposure to additional stressors. Nevertheless, the majority of the proposed activities are unit-level training and testing activities, which are conducted in the open ocean. Unit-level events occur over a small spatial scale (one to a few square miles) and with few participants (usually one or two) or short duration (the order of a few hours or less).

Secondly, a marine mammal could be exposed to multiple training and testing activities over the course of its life; however, training and testing activities are generally separated in space and time in such a way that it would be unlikely that any individual marine mammal would be exposed to stressors from multiple activities within a short timeframe. However, animals with a home range intersecting an area of concentrated Navy activity have elevated exposure risks relative to animals that simply transit the area through a migratory corridor.

Multiple stressors may also have synergistic effects. For example, marine mammals that experience temporary hearing loss or injury from acoustic stressors could be more susceptible to physical strike and disturbance stressors via a decreased ability to detect and avoid threats. Marine mammals that experience behavioral and physiological consequences of ingestion stressors could be more susceptible to entanglement and physical strike stressors via malnourishment and disorientation. These interactions are speculative, and without data on the combination of multiple Navy stressors, the synergistic impacts from the combination of Navy stressors are difficult to predict in any meaningful way. Research and monitoring efforts have included before, during, and after-event observations and surveys, data collection through conducting long-term studies in areas of Navy activity, occurrence surveys over large geographic areas, biopsy of animals occurring in areas of Navy activity, and tagging studies where animals are exposed to Navy stressors. These efforts are intended to contribute to the overall understanding of what impacts may be occurring overall to animals in these areas. To date, the findings from the research and monitoring and the regulatory conclusions from previous analyses by NMFS in the MMPA authorization (80 FR 46112) and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS

(National Oceanic and Atmospheric Administration, 2015b) have been that the majority of impacts from training and testing activities are not expected to have deleterious impacts on the fitness of any individuals or long-term consequences to populations of marine mammals.

#### **3.4.3.1 Combined Impacts of All Stressors Under Alternative 1**

Although potential impacts on certain marine mammal species from training and testing activities under Alternative 1 may include injury to individuals, those injuries are not expected to lead to long-term consequences for populations. The potential impacts anticipated from Alternative 1 are summarized in Section 3.4.4 (Marine Mammal Protection Act Determinations) and Section 3.4.5 (Endangered Species Act Determinations) for each regulation applicable to marine mammals. For a discussion of cumulative impacts, see Chapter 4 (Cumulative Impacts). For a discussion of mitigation, see Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment).

#### **3.4.3.2 Combined Impacts of All Stressors Under Alternative 2**

As detailed previously in this section, some training and testing activities proposed under Alternative 2 would be an increase over what is proposed for Alternative 1. However, this increase is not expected to significantly increase the potential for impacts over what is analyzed for Alternative 1. The analysis presented in Section 3.4.3.1 (Combined Impacts of All Stressors Under Alternative 1) would similarly apply to Alternative 2. The combined impacts of all stressors for training and testing activities under Alternative 2 are not expected to have deleterious impacts or long-term consequences to populations of marine mammals.

#### **3.4.3.3 Combined Impacts of All Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with the Proposed Action would continue to occur. All stressors associated with training and testing activities would not be introduced into the marine environment. Therefore, baseline conditions of the existing environment would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

#### **3.4.3.4 Summary of Monitoring and Observations During Navy Activities Since 2015**

As provided in detail in the 2015 MITT Final EIS/OEIS, Section 3.4.5.2 (Summary of Observations During Previous Navy Activities), the results of previous monitoring and research since 2006 taking place in and around Navy ranges and occurring before, during, and after Navy training and testing events, have been included as part of the Navy analyses as well as the analyses by NMFS in their MMPA authorization (National Oceanic and Atmospheric Administration, 2015b), and the Biological Opinion for the 2015 MITT Final EIS/OEIS (National Marine Fisheries Service, 2015b).

Since 2006, the Navy, non-Navy marine mammal scientists, and research groups and institutions have conducted scientific monitoring and research in and around ocean areas in the Atlantic and Pacific where the Navy has been and proposes to continue training and testing. The analysis provided in this SEIS/OEIS will be the third time Navy training and testing activities at sea have been comprehensively analyzed in the Study Area. Data collected from Navy monitoring, scientific research findings, and annual reports have been provided to NMFS; this public record is informative as part of the analysis of impacts on marine mammals in general for a variety of reasons, including species distribution, habitat use, and evaluation of potential responses to Navy activities.

Monitoring across Navy training and testing ranges is performed using a variety of methods, including visual surveys from surface vessels and aircraft; in addition, passive acoustics before, during, and after

Navy activities have been conducted. The Navy also has continued to contribute to funding of basic research, including behavioral response studies specifically designed to determine the effects to marine mammals from the Navy's use of mid-frequency sonar and other transducers.

The majority of the training and testing activities Navy is proposing in the Study Area are similar if not nearly identical to activities that have been occurring in the same locations for decades. For example, the mid-frequency sonar system on the Navy's cruisers and destroyers training in the Study Area have the same sonar system components in the water as those first deployed in the 1970s. While the signal analysis and computing processes onboard these ships have been upgraded with modern technology, the sonar transducers, which put signals into the water, have not changed. For this reason, the history of past marine mammal observations, research, and monitoring remain applicable to the analysis of effects from the proposed future training and testing activities in the Study Area.

It is still the case that in the Pacific, the vast majority of scientific field work, research, and monitoring efforts have been expended in Southern California and Hawaii where Navy training and testing activities have been most concentrated. Since 2006, the Navy has been submitting exercise reports and monitoring reports to NMFS for the Navy's range complexes, including monitoring conducted before, during, and after training and testing activities. These publically available exercise reports, monitoring reports, and the associated scientific research findings have been integrated into decisions regarding the focus for subsequent research and monitoring as determined in collaborations between Navy, NMFS, Marine Mammal Commission, and other marine resource subject matter experts using an adaptive management approach.

In the Study Area, because training and testing events are less frequent and general small in scope by comparison to other Navy areas, the majority of Navy's research effort has been focused elsewhere. Despite this, funding by the Navy has provided nearly the entirety of marine mammal science collected in the Mariana Islands. In fact, prior to Navy funding of marine mammal science, there had not been any dedicated marine mammal surveys performed in the Mariana Islands. The bulk of these Navy-funded research efforts span two primary methodologies: small-vessel surveys and bottom-moored acoustic deployments. These primary data collection methods have been supplemented by additional results from autonomous gliders acoustic survey, acoustic towed-arrays, visual survey from shore-stations, marine mammal observers on large-vessel surveys, and further analysis and collection of incidental and stranding data. Since the 2015 MITT Final EIS/OEIS, new research has continued to be funded by Navy in the Mariana Islands and has included, but is not limited to the following findings:

- The continuation of annual small vessel nearshore surveys, sightings, satellite tagging, biopsy and genetic analysis, photo-identification, and opportunistic acoustic recording off Guam, Saipan, Tinian, Rota, and Aguigan in partnership with NMFS (Hill et al., 2015b; Hill et al., 2016b; Hill et al., 2017a). The satellite tagging and genetic analyses have resulted in the first information discovered on the movement patterns, habitat preference, and population structure of multiple odontocete species in the Study Area.
- Since 2015, the addition of a series of small vessel surveys in the winter season dedicated to humpback whales has provided new information relating to the occurrence, calving behavior, and population identity of this species (Hill et al., 2016a; Hill et al., 2017b), which has not previously been sighted during the previous small vessel surveys in the summer or winter. This work has included sighting data, photo ID matches of individuals to other areas demonstrating migration as well as re-sights within the Mariana Islands across different years, and the collection of biopsy samples for genetic analyses of populations.

- The continued deployment of passive acoustic monitoring devices and analysis of acoustic data obtained using bottom-moored acoustic recording devices deployed by NMFS has provided information on the presence and seasonal occurrence of mysticetes, as well as the occurrence of cryptic odontocetes typically found offshore, including beaked whales and Kogia whales (Hill et al., 2015b; Hill et al., 2016a; Hill et al., 2016b; Hill et al., 2017a; Munger et al., 2015; Norris et al., 2017; Oleson et al., 2015; Yack et al., 2016).
- Acoustic surveys using autonomous gliders were used to characterize the occurrence of odontocetes and mysticetes in abyssal offshore waters near Guam and Commonwealth of the Northern Mariana Islands (CNMI), including species not seen in the small vessel visual survey series such as killer whales and Risso's dolphins. Analysis of collected data also provided new information on the seasonality of baleen whales, patterns of beaked whale occurrence and potential call variability, and identification of new unknown marine mammal calls (Klinck et al., 2016b; Nieu Kirk et al., 2016).
- Visual surveys were conducted from a shore-station at high elevation on the north shore of Guam to document the nearshore occurrence of marine mammals in waters where small vessel visual surveys are challenging due to regularly high sea states (Deakos & Richlen, 2015; Deakos et al., 2016).
- Analysis of archive data, including marine mammal sightings during Guam Department of Agriculture Division of Aquatic and Wildlife Resources aerial surveys undertaken between 1963 and 2012 (Martin et al., 2016).
- Analysis of archived acoustic towed-array data for an assessment of the abundance and density of minke whales (Norris et al., 2017), abundance and density of sperm whales (Yack et al., 2016), and the characterization of sei and humpback whale vocalizations (Norris et al., 2014).

As detailed in the 2015 MITT Final EIS/OEIS, these reporting, monitoring, and research efforts by the Navy have added to the baseline data for marine mammal species inhabiting the Study Area. In addition, subsequent research and monitoring across the Navy has continued to broaden the sample of observations regarding the general health of marine mammal populations in locations where Navy has been conducting training and testing activities for decades, which has been considered in the analysis of marine mammal impacts presented in this SEIS/OEIS in the same manner that the previous findings were used in the 2015 MITT Final EIS/OEIS, the NMFS authorization of takes under MMPA (National Oceanic and Atmospheric Administration, 2015b), and the NMFS Biological Opinion pursuant to the ESA (National Marine Fisheries Service, 2015b).

This public record of training and testing activities, monitoring, and research from across the Navy range complexes in the Pacific and Atlantic now spans more than 13 years. Given that this record involves many of the same Navy training and testing activities being considered for the Study Area, and includes all the marine mammal taxonomic families present in the Study Area, many of the same species, and perhaps some of the same populations as they seasonally migrate from other range complexes, this compendium of Navy reporting is directly applicable to the Study Area. It was the Navy's assessment in the 2015 MITT Final EIS/OEIS and that of NMFS, as reflected in their analysis of previous Navy training and testing in the Study Area (National Marine Fisheries Service, 2015b; National Oceanic and Atmospheric Administration, 2015b), that it was unlikely there would be impacts on populations of marine mammals (such as whales, dolphins, and pinnipeds) having any long-term consequences as a result of the proposed continuation of training and testing in the Study Area. This assessment of likelihood is based on four indicators from areas in the eastern Pacific where Navy training and testing

has been ongoing for decades: (1) evidence suggesting or documenting increases in the numbers of marine mammals present, (2) examples of documented presence and site fidelity of species and long-term residence by individual animals of some species, (3) use of training and testing areas for breeding and nursing activities, and (4) 13 years of comprehensive monitoring data indicating a lack of obvious observable effects such as direct mortalities or strandings occurring in marine mammal populations as a result of Navy training and testing activities. Consistent with the presentation in the 2015 MITT Final EIS/OEIS, the evidence to date and since 2015 continues to suggest the viability of marine mammal populations where the Navy trains and tests and does not show any direct evidence suggesting Navy training and testing has had or may have any long-term consequences to marine mammal populations. Barring any evidence to the contrary, therefore, what limited evidence there is from monitoring reports and additional other focused scientific investigations should be considered in the analysis of impacts on marine mammals. For the Study Area in particular and since the analysis in 2015, examples include

- the most current information suggesting that the ESA-listed blue whale population in the Pacific, which includes the Study Area as part of their habitat, may have recovered and been at a stable level based on recent surveys and scientific findings (Barlow, 2016; Campbell et al., 2015; Carretta et al., 2017d; Monnahan et al., 2015; Rockwood et al., 2017; Širović et al., 2015); and
- humpback whales continue to use Northern Mariana Islands as a winter calving area (Fulling et al., 2011; Hill et al., 2016a; Hill et al., 2017b; Hill et al., 2018a).

To summarize and bring up to date the findings from the 2015 MITT Final EIS/OEIS, the evidence from reporting, monitoring, and research across the Pacific over more than a decade indicates that while the Proposed Action would result in harassment of marine mammals and may include injury to some individuals, these impacts are expected to be inconsequential at the level of their marine mammal populations. Monitoring of Navy training and testing will continue to confirm this expectation, as it has in the past in locations where Navy training and testing occurs. Across this past monitoring as well as the broader scientific literature, no direct evidence exists that routine Navy training and testing spanning decades has negatively impacted marine mammal populations at any Navy Range Complex or the Study Area. In particular, there is no evidence that would directly contradict the analysis in the 2015 MITT Final EIS/OEIS or this SEIS/OEIS, such as the regular observation of strandings, injuries, or mortalities associated temporarily and spatially with Navy training and testing events.

For some of the most intensively used Navy training and testing areas, evidence such as the continued multi-year presence of long-term resident individual animals and small populations (Baird et al., 2015; Baird et al., 2016; Baird et al., 2017; Schorr et al., 2014; Schorr et al., 2018; U.S. Department of the Navy, 2017b), resident females documented with and without calves from year to year, and high abundances on the Navy ranges for some species in comparison to other off-range locations (Moore & Barlow, 2017; Schorr et al., 2018; U.S. Department of the Navy, 2017b), indicates generally healthy marine mammal populations. Therefore, based on the best available science, including data developed in exercise and monitoring reports submitted to NMFS for over a decade, long-term consequences for marine mammal populations are unlikely to result from Navy training and testing activities in the Study Area.

#### **3.4.4 Marine Mammal Protection Act Determinations**

As required by Section 101(a)(5)(A) of the MMPA, the Navy is seeking a Letter of Authorization from NMFS for the use of sonar and other transducers and explosives during Navy activities under Alternative 1 or Alternative 2 of the Proposed Action. The use of sonar and other transducers may result in Level A and Level B harassment of certain marine mammals. The use of explosives may result in Level A

harassment, Level B harassment. Refer to Section 3.4.2.1.2 (Impacts from Sonar and Other Transducer Stressors) for details on the estimated impacts from sonar and other transducers and Section 3.4.2.2.2 (Impacts from Explosive Stressors) for impacts from explosives. Based on best available science, the Navy concludes that impacts from sonar and other transducers and from explosives to marine mammal species and stocks would result in only short-term effects on most individuals exposed and would not affect annual rates of recruitment or survival for the following reasons:

- Most acoustic exposures are within the non-injurious temporary threshold shift or behavioral effects zones (Level B harassment).
- Although the numbers presented in Appendix E (Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities) represent estimated harassment under the MMPA and they are conservative estimates (i.e., overpredictions) of harassment, primarily by behavioral disturbance.
- The Navy Acoustic Effects Model calculates harassment without taking into consideration mitigation measures, and is not indicative of a likelihood of either injury or harm. Additionally, the mitigation measures described in Chapter 5 (Mitigation) are designed to avoid or reduce sound exposure and explosive effects on marine mammals to levels below those that may cause injury and to achieve the least practicable adverse effect on marine mammal species or stocks.

Weapons noise, vessel noise, aircraft noise, the use of in-water electromagnetic devices, in-air electromagnetic devices, high-energy lasers, vessels, in-water devices, seafloor devices, wires and cables, decelerators/parachutes, and military expended materials are not expected to result in Level A or Level B harassment of any marine mammals.

### 3.4.5 Endangered Species Act Determinations

Pursuant to the ESA, Navy training and testing activities presented in this SEIS/OEIS may affect ESA-listed marine mammals. There is no designated critical habitat for any marine mammal species in the MITT Study Area. The Navy will consult with NMFS as required by section 7(a)(2) of the ESA. The outcome of those consultations pursuant to ESA will be described in the MITT Final SEIS/OEIS.

### 3.4.6 Public Scoping Comments

The public raised a number of issues during the scoping period for this supplement in regard to marine mammals. The issues are summarized in the list below.

- Concern that there be analysis of impacts on marine mammals from various stressors associated with the Proposed Action (e.g., sonar, explosives, “chemical pollution,” destruction of habitat) – Section 3.4.2 (Environmental Consequences) analyzed potential impacts from sonar and other active acoustic stressors, explosives, electromagnetic energy, physical disturbances and strikes, entanglement, ingestion, and secondary stressors. Impacts from sonar and explosives were reanalyzed for several reasons described in Section 3.4.2.1 (Acoustic Stressors) and Section 3.4.2.2 (Explosive Stressors). The analysis of other stressors is summarized in the Section 3.4.2.3 (Energy Stressors) through Section 3.4.2.7 (Secondary Stressors) and described in detail in the 2015 MITT Final EIS/OEIS.
- **Concerns over the amount of take of marine mammals authorized** – The number and species of marine mammals exposed to sonar and explosives (the only stressors predicted to result in a “take” of a marine mammal) are presented in Section 3.4.2.1.2 (Impacts from Sonar and Other

Transducer Stressors) and Section 3.4.2.2.2 (Impacts from Explosive Stressors). The vast majority of predicted impacts are behavioral reactions to sonar or explosives.

- **The analysis must address direct and cumulative impacts on marine mammals** – Direct impacts on marine mammals are addressed in the stressor sections above. A detailed analysis of impacts from acoustic and explosive stressors is provided in Section 3.4.2.1 (Acoustic Stressors) and Section 3.4.2.2 (Explosive Stressors). Impacts from other stressors are summarized in Section 3.4.2.3 (Energy Stressors) through Section 3.4.2.7 (Secondary Stressors) and described in detail in the 2015 MITT Final EIS/OEIS. Chapter 4 (Cumulative Impacts) addresses potential impacts from the Proposed Action in combination with other past, present, and future activities occurring in the Study Area. Cumulative impacts on marine mammals are likely when considering the variety of stressors (e.g., bycatch) that pose a threat to marine mammal populations. Refer to Section 3.4.1.7 (General Threats).
- **Recommendation that this SEIS/OEIS must evaluate alternatives that include temporal and habitat avoidance or time/area closures including restrictions on activities in areas biologically sensitive or important areas** – Regarding the development of alternatives considered, see Section 2.4 (Action Alternatives Development). With regard to the topic specifically, see Section 2.4.1.3 (Alternatives Including Geographic Mitigation Measures within the Study Area) for a discussion on why alternatives including these types of mitigation measures are not generally feasible. A new Appendix I (Geographic Mitigation Assessment) has been included in this SEIS/OEIS to further evaluate areas specifically identified in public scoping comments as areas to consider for geographic/temporal mitigation. Based on the analysis in Appendix I (Geographic Mitigation Assessment), in Chapter 5 (Mitigation), in association with the MMPA and the ESA permitting processes, and other required regulatory consultations, consideration of geographic mitigation has been considered for implementation under both action alternatives.
- **The supplement must include analysis and description of mitigation measures implemented to reduce impacts on marine mammals** – Chapter 5 (Mitigation) provides a detailed description of mitigation measures associated with training and testing activities that would avoid or reduce potential impacts on marine mammals.
- **In-water surveys within the 3 miles around FDM should be conducted for marine mammals** – In-water surveys of marine resources within the 3 NM danger zone surrounding FDM have been conducted for more than a decade by Navy divers (see Smith and Marx (2009); Smith et al. (2013b); Smith and Marx (2016)). Research funding is allocated via the Integrated Comprehensive Monitoring Program (U.S. Department of the Navy, 2010, 2013a), which provides the overarching framework for coordination of the Navy’s marine species research and monitoring efforts and serves as a planning tool to focus Navy monitoring priorities pursuant to ESA and MMPA requirements. The purpose of the Integrated Comprehensive Monitoring Program is to coordinate monitoring efforts across all regions and to allocate the most appropriate level and type of monitoring effort for each range complex based on a set of standardized objectives, regional expertise, and resource availability. Although the Integrated Comprehensive Monitoring Program does not identify specific field work or individual projects, it is designed to provide a flexible, scalable, and adaptable framework using adaptive management and strategic planning processes that periodically assess progress and reevaluate objectives. The adaptive management review process is anticipated to continue between the Navy, NMFS, and the Marine Mammal Commission through technical review meetings and ongoing discussions.

- **The supplement must include the most recently published science cetacean reports for the Mariana Archipelago** – Relevant literature published since the 2015 MITT Final EIS/OEIS has been used throughout Section 3.4 (Marine Mammals) of this SEIS/OEIS. This includes approximately 160 new references cited in the section, which were published between January 2016 and June 2018 in addition to additional emergent works of science that considered in the analysis although not necessarily cited in this section.
- **The EIS must analyze humpback whale calving areas discussed by Hill et al. (2017b)** – Section 3.4.1.11 (Humpback Whale (*Megaptera novaeangliae*)) presents the current information and references with regard to the presence of humpback whales in the Mariana Islands. Note that Hill et al. (2017b) did not identify the location specific calving areas in the Mariana Islands, but indicated, “that the Marianas are a wintering area.” Subsequent to Hill et al. (2017b) and recently, scientists have confirmed the Mariana Islands as a new breeding location for humpback whales in the western North Pacific (National Oceanic and Atmospheric Administration, 2018c). Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment) contain detailed discussions of potential mitigation measures that were evaluated, including temporal and geographic mitigation for areas where humpback whales have been routinely sighted as detailed in various reports (Fulling et al., 2011; Hill et al., 2015a; Hill et al., 2015b; Hill et al., 2016a; Hill et al., 2016b; Hill et al., 2017a; Hill et al., 2017b; Hill et al., 2018a; National Oceanic and Atmospheric Administration, 2018c; Uyeyama, 2014). Based on the analysis in Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment), the MMPA and the ESA permitting processes, and other required regulatory consultations, practical science-based mitigation measures, including temporal or geographic constraints within the Study Area, may be implemented under Alternative 1 or Alternative 2.

## REFERENCES

- Abecassis, M., J. Polovina, R. W. Baird, A. Copeland, J. C. Drazen, R. Domokos, E. Oleson, Y. Jia, G. S. Schorr, D. L. Webster, and R. D. Andrews. (2015). Characterizing a foraging hotspot for short-finned pilot whales and Blainville's beaked whales located off the west side of Hawaii island by using tagging and oceanographic data. *PLoS ONE*, 10(11), e0142628.
- Acevedo-Whitehouse, K., A. Rocha-Gosselin, and D. Gendron. (2010). A novel non-invasive tool for disease surveillance of freeranging whales and its relevance to conservation programs. *Animal Conservation*, 13, 217–225.
- Acevedo, A. (1991). Interactions between boats and bottlenose dolphins, *Tursiops truncatus*, in the entrance to Ensenada De La Paz, Mexico. *Aquatic Mammals*, 17(3), 120–124.
- Afsal, V. V., P. P. Manojkumar, K. S. S. M. Yousuf, B. Anoop, and E. Vivekanandan. (2009). The first sighting of Longman's beaked whale, *Indopacetus pacificus* in the southern Bay of Bengal. *Marine Biodiversity Records*, 2, 3.
- Aguilar de Soto, N., M. Johnson, P. T. Madsen, P. L. Tyack, A. Bocconcelli, and J. F. Borsani. (2006). Does intense ship noise disrupt foraging in deep-diving Cuvier's beaked whales (*Ziphius cavirostris*)? *Marine Mammal Science*, 22(3), 690–789.
- Aguilar de Soto, N., M. P. Johnson, P. T. Madsen, F. Diaz, I. Dominguez, A. Brito, and P. Tyack. (2008). Cheetahs of the deep sea: Deep foraging sprints in short-finned pilot whales off Tenerife (Canary Islands). *The Journal of Animal Ecology*, 77(5), 936–947.
- Akkaya Bas, A., F. Christiansen, A. Amaha Ozturk, B. Ozturk, and C. McIntosh. (2017). The effects of marine traffic on the behaviour of Black Sea harbour porpoises (*Phocoena phocoena relicta*) within the Istanbul Strait, Turkey. *PLoS ONE*, 12(3), e0172970.
- Allen, A. N., J. J. Schanze, A. R. Solow, and P. L. Tyack. (2014). Analysis of a Blainville's beaked whale's movement response to playback of killer whale vocalizations. *Marine Mammal Science*, 30(1), 154–168.
- Allen, S. D., and J. R. Amesbury. (2012). *Commonwealth of the Northern Mariana Islands As a Fishing Community* (NOAA Technical Memorandum NMFS-PIFSC-36). Honolulu, HI: Pacific Island Fisheries Science Center.
- Alter, S. E., M. P. Simmonds, and J. R. Brandon. (2010). Forecasting the consequences of climate-driven shifts in human behavior on cetaceans. *Marine Policy*, 34(5), 943–954.
- Alves, A., R. Antunes, A. Bird, P. L. Tyack, P. J. O. Miller, F. P. A. Lam, and P. H. Kvasdheim. (2014). Vocal matching of naval sonar signals by long-finned pilot whales (*Globicephala melas*). *Marine Mammal Science*, 30(3), 1248–1257.
- Amano, M., N. Miyazaki, and F. Yanagisawa. (1996). Life history of Fraser's dolphin, *Lagenodelphis hosei*, based on a school captured off the Pacific Coast of Japan. *Marine Mammal Science*, 12(2), 199–214.
- Amesbury, S., R. Bonito, R. K. C. Chang, L. Kirkendale, C. Meyer, G. Paulay, R. Ritson-Williams, and T. Rongo. (2001). *Marine Biodiversity Resource Survey and Baseline Reef Monitoring Survey of the Haputo Ecological Reserve Area, COMNAVMARINANAS*. Mangilao, GU: University of Guam.

- Anderson, R. C., R. Clark, P. T. Madsen, C. Johnson, J. Kiszka, and O. Breyse. (2006). Observations of Longman's Beaked Whale (*Indopacetus pacificus*) in the Western Indian Ocean. *Aquatic Mammals*, 32(2), 223–231.
- Anderwald, P., A. Brandecker, M. Coleman, C. Collins, H. Denniston, M. D. Haberlin, M. O'Donovan, R. Pinfield, F. Visser, and L. Walshe. (2013). Displacement responses of a mysticete, an odontocete, and a phocid seal to construction-related vessel traffic. *Endangered Species Research*, 21(3), 231–240.
- Ando-Mizobata, N., K. Ichikawa, N. Arai, and H. Kato. (2014). Does boat noise affect dugong (*Dugong dugon*) vocalization? *Mammal Study*, 39(2), 121–127.
- Andrady, A. (2015). Persistence of plastic litter in the oceans. In M. Bergmann, L. Gutow, & M. Klages (Eds.), *Marine Anthropogenic Litter*. New York, NY: Springer International Publishing.
- Antunes, R., P. H. Kvasdheim, F. P. Lam, P. L. Tyack, L. Thomas, P. J. Wensveen, and P. J. Miller. (2014). High thresholds for avoidance of sonar by free-ranging long-finned pilot whales (*Globicephala melas*). *Marine Pollution Bulletin*, 83(1), 165–180.
- Aragones, L., M. Roque, M. Flores, R. Encomienda, G. Laule, B. Espinos, F. Maniago, G. Diaz, E. Alesna, and R. Braun. (2010). The Philippine marine mammal strandings from 1998 to 2009: Animals in the Philippines in peril? *Aquatic Mammals*, 36(3), 219–233.
- Arcangeli, A., and R. Crosti. (2009). The short-term impact of dolphin-watching on the behaviour of bottlenose dolphins (*Tursiops truncatus*) in western Australia. *Journal of Marine Animals and Their Ecology*, 2(1), 3–9.
- Archer, F. I., S. L. Mesnick, and A. C. Allen. (2010). *Variation and Predictors of Vessel-Response Behavior in a Tropical Dolphin Community* (NOAA Technical Memorandum NMFS-SWFSC-457). La Jolla, CA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.
- Archer, F. I., II, and W. F. Perrin. (1999). *Stenella coeruleoalba*. *American Society of Mammalogists*, 603, 1–9.
- Arnbom, T., V. Papastavrou, S. Weilgart, and H. Whitehead. (1987). Sperm Whales React to an Attack by Killer Whales. *Journal of Mammalogy*, 68(2), 450.
- Atkinson, S., D. Crocker, D. Houser, and K. Mashburn. (2015). Stress physiology in marine mammals: How well do they fit the terrestrial model? *Journal of Comparative Physiology B*, 185, 463–486.
- Au, D. W. K., and W. L. Perryman. (1985). Dolphin habitats in the eastern tropical Pacific. *Fishery Bulletin*, 83, 623–643.
- Au, W. W. L., R. W. Floyd, R. H. Penner, and A. E. Murchison. (1974). Measurement of echolocation signals of the Atlantic bottlenose dolphin, *Tursiops truncatus* Montagu, in open waters. *The Journal of the Acoustical Society of America*, 56(4), 1280–1290.
- Au, W. W. L., and P. W. B. Moore. (1990). Critical ratio and critical bandwidth for the Atlantic bottlenose dolphin. *The Journal of the Acoustical Society of America*, 88(3), 1635–1638.
- Au, W. W. L. (1993). *The Sonar of Dolphins*. New York, NY: Springer-Verlag.
- Au, W. W. L., and M. Green. (2000). Acoustic interaction of humpback whales and whale-watching boats. *Marine Environmental Research*, 49(5), 469–481.

- Avens, L. (2003). Use of multiple orientation cues by juvenile loggerhead sea turtles *Caretta caretta*. *The Journal of Experimental Biology*, 206(23), 4317–4325.
- Ayres, K. L., R. K. Booth, J. A. Hempelmann, K. L. Koski, C. K. Emmons, R. W. Baird, K. Balcomb-Bartok, M. B. Hanson, M. J. Ford, and S. K. Wasser. (2012). Distinguishing the impacts of inadequate prey and vessel traffic on an endangered killer whale (*Orcinus orca*) population. *PLoS ONE*, 7(6), e36842.
- Azzara, A. J., W. M. von Zahren, and J. J. Newcomb. (2013). Mixed-methods analytic approach for determining potential impacts of vessel noise on sperm whale click behavior. *The Journal of the Acoustical Society of America*, 134(6), 4566–4574.
- Bachara, W., and D. Blatchley. (2018). *Blainville's beaked whale strandings in the Philippines* (Report WB2018/2). Unpublished report to Cetal Fauna.
- Bachman, M. J., J. M. Keller, K. L. West, and B. A. Jensen. (2014). Persistent organic pollutant concentrations in blubber of 16 species of cetaceans stranded in the Pacific Islands from 1997 through 2011. *Science of the Total Environment*, 488–489, 115–123.
- Bachman, M. J., K. M. Foltz, J. M. Lynch, K. L. West, and B. A. Jensen. (2015). Using cytochrome P4501A1 expression in liver and blubber to understand effects of persistent organic pollutant exposure in stranded Pacific Island cetaceans. *Environmental Toxicology and Chemistry*, 34(9), 1989–1995.
- Bain, D. E. (2002). *A Model Linking Energetic Effects of Whale Watching to Killer Whale (Orcinus orca) Population Dynamics*. Friday Harbor, WA: Friday Harbor Laboratories, University of Washington.
- Baird, R., D. McSweeney, C. Bane, J. Barlow, D. Salden, L. Antoine, R. LeDuc, and D. Webster. (2006a). Killer Whales in Hawaiian Waters: Information on Population Identity and Feeding Habits. *Pacific Science*, 60(4), 523–530.
- Baird, R. (2013). Odontocete Cetaceans Around the Main Hawaiian Islands: Habitat Use and Relative Abundance from Small-Boat Sighting Surveys. *Aquatic Mammals*, 39(3), 253–269.
- Baird, R. W., A. D. Ligon, S. K. Hooker, and A. M. Gorgone. (2001). Subsurface and nighttime behaviour of pantropical spotted dolphins in Hawai'i. *Canadian Journal of Zoology*, 79(6), 988–996.
- Baird, R. W., D. J. McSweeney, D. L. Webster, A. M. Gorgone, and A. D. Ligon. (2003). *Studies of Odontocete Population Structure in Hawaiian Waters: Results of a Survey Through the Main Hawaiian Islands in May and June 2003*. Seattle, WA: National Oceanic and Atmospheric Administration.
- Baird, R. W., and A. M. Gorgone. (2005). False killer whale dorsal fin disfigurements as a possible indicator of long-line fishery interactions in Hawaiian waters. *Pacific Science*, 59(4), 593–601.
- Baird, R. W., D. L. Webster, D. J. McSweeney, A. D. Ligon, and G. S. Schorr. (2005). *Diving Behavior and Ecology of Cuvier's (Ziphius cavirostris) and Blainville's Beaked Whales (Mesoplodon densirostris) in Hawaii*. La Jolla, CA: U.S. Department of Commerce.
- Baird, R. W., D. McSweeney, C. Bane, J. Barlow, D. Salden, L. Antoine, R. LeDuc, and D. Webster. (2006b). Killer whales in Hawaiian waters: Information on population identity and feeding habits. *Pacific Science*, 60(4), 523–530.
- Baird, R. W., G. S. Schorr, D. L. Webster, D. J. McSweeney, and S. D. Mahaffy. (2006c). *Studies of Beaked Whale Diving Behavior and Odontocete Stock Structure in Hawaii in March/April 2006*. Olympia, WA: Cascadia Research Collective.

- Baird, R. W., A. M. Gorgone, D. J. McSweeney, D. B. Webster, D. R. Salden, M. H. Deakos, A. D. Ligon, G. Schorr, J. Barlow, and S. D. Mahaffy. (2008). False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: Long-term site fidelity, inter-island movements, and association patterns. *Marine Mammal Science*, 24(3), 591–612.
- Baird, R. W. (2009a). False killer whale *Pseudorca crassidens*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 405–406). Cambridge, MA: Academic Press.
- Baird, R. W. (2009b). *A Review of False Killer Whales in Hawaiian Waters: Biology, Status, and Risk Factors*. Olympia, WA: Marine Mammal Commission.
- Baird, R. W., A. M. Gorgone, D. J. McSweeney, A. D. Ligon, M. H. Deakos, D. L. Webster, G. S. Schorr, K. K. Martien, D. R. Salden, and S. D. Mahaffy. (2009a). Population structure of island-associated dolphins: Evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the Main Hawaiian Islands. *Marine Mammal Science*, 25(2), 251–274.
- Baird, R. W., D. J. McSweeney, G. S. Schorr, S. D. Mahaffy, D. L. Webster, J. Barlow, M. B. Hanson, J. P. Turner, and R. D. Andrews. (2009b). Studies of beaked whales in Hawaii: Population size, movements, trophic ecology, social organization, and behaviour. In S. J. Dolman, C. D. MacLeod, & P. G. H. Evans (Eds.), *Beaked Whale Research* (pp. 23–25). San Sebastián, Spain: European Cetacean Society.
- Baird, R. W., J. M. Aschettino, D. J. McSweeney, D. L. Webster, G. S. Schorr, S. Baumann-Pickering, and S. D. Mahaffy. (2010a). *Melon-headed Whales in the Hawaiian Archipelago: An Assessment of Population Structure and Long-term Site Fidelity based on Photo-Identification*. La Jolla, CA: Southwest Fisheries Science Center, National Marine Fisheries Service.
- Baird, R. W., G. S. Schorr, D. L. Webster, D. J. McSweeney, M. B. Hanson, and R. D. Andrews. (2010b). Movements and habitat use of satellite-tagged false killer whales around the main Hawaiian Islands. *Endangered Species Research*, 10, 107–121.
- Baird, R. W. (2011). Short Note: Open-Ocean Movements of a Satellite-Tagged Blainville’s Beaked Whale (*Mesoplodon densirostris*): Evidence for an Offshore Population in Hawai’i? *Aquatic Mammals*, 37(4), 506–511.
- Baird, R. W. (2012). *Preliminary Results from Photo-Identification and Satellite-Tagging of False Killer Whales Off the Island of Kaua’i in June 2012*. Olympia, WA: Cascadia Research Collective.
- Baird, R. W., E. M. Oleson, J. Barlow, A. D. Ligon, A. M. Gorgone, and S. D. Mahaffy. (2013a). Evidence of an Island-Associated Population of False Killer Whales (*Pseudorca crassidens*) in the Northwestern Hawaiian Islands. *Pacific Science*, 67(4), 513–521.
- Baird, R. W., J. A. Shaffer, D. L. Webster, S. D. Fisher, J. M. Aschettino, A. M. Gorgone, B. K. Rone, S. D. Mahaffy, and D. J. Moretti. (2013b). *Odontocete Studies Off the Pacific Missile Range Facility in February 2013: Satellite-Tagging, Photo Identification, and Passive Acoustic Monitoring for Species Verification*. Olympia, WA and Newport, RI: U.S. Navy Pacific Fleet.
- Baird, R. W., D. L. Webster, S. D. Mahaffy, G. S. Schorr, J. M. Aschettino, and A. M. Gorgone. (2013c). *Movements and Spatial Use of Odontocetes in the Western Main Hawaiian Islands: Results of a Three-year Study off Oahu and Kauai*. Olympia, WA: Cascadia Research Collective.
- Baird, R. W., S. M. Jarvis, D. L. Webster, B. K. Rone, J. A. Shaffer, S. D. Mahaffy, A. M. Gorgone, and D. J. Moretti. (2014). *Odontocete Studies on the Pacific Missile Range Facility in July/August 2013*:

- Satellite-Tagging, Photo Identification, and Passive Acoustic Monitoring*. Olympia, WA and Newport, RI: U.S. Navy Pacific Fleet.
- Baird, R. W., D. Cholewiak, D. L. Webster, G. S. Schorr, S. D. Mahaffy, C. Curtice, J. Harrison, and S. M. Van Parijs. (2015). Biologically Important Areas for Cetaceans within U.S. Waters—Hawaii region. In S. M. Van Parijs, C. Curtice, & M. C. Ferguson (Eds.), *Biologically Important Areas for Cetaceans Within U.S. Waters* (Vol. 41, pp. 54–64). Olympia, WA: Cascadia Research Collective.
- Baird, R. W., D. L. Webster, S. Watwood, R. Morrissey, B. K. Rone, S. D. Mahaffy, A. M. Gorgone, D. B. Anderson, and D. J. Moretti. (2016). *Odontocete Studies on the Pacific Missile Range Facility in February 2015: Satellite-Tagging, Photo-Identification, and Passive Acoustic Monitoring. Final Report*. Olympia, WA: HDR Environmental Inc.
- Baird, R. W., S. W. Martin, R. Manzano-Roth, D. L. Webster, and B. L. Southall. (2017). *Assessing Exposure and Response of Three Species of Odontocetes to Mid-frequency Active Sonar During Submarine Commanders Courses at the Pacific Missile Range Facility: August 2013 through February 2015. Draft Report*. Honolulu, HI: HDR, Inc.
- Baker, A. N., and B. Madon. (2007). Bryde's whales (*Balaenoptera cf. brydei*) in the Hauraki Gulf and northeastern New Zealand waters. *Science for Conservation*, 272, 4–14.
- Baker, C. S., L. M. Herman, B. G. Bays, and G. B. Bauer. (1983). *The Impact of Vessel Traffic on the Behavior of Humpback Whales in Southeast Alaska: 1982 Season*. Honolulu, HI: Kewalo Basin Marine Mammal Laboratory, University of Hawaii.
- Baker, C. S., V. Lukoschek, S. Lavery, M. L. Dalebout, M. Yong-un, T. Endo, and N. Funahashi. (2006a). Incomplete reporting of whale, dolphin and porpoise 'bycatch' revealed by molecular monitoring of Korean markets. *Animal Conservation*, 9(4), 474–482.
- Baker, J. D., A. L. Harting, and T. C. Johanos. (2006b). Use of discovery curves to assess abundance of Hawaiian monk seals. *Marine Mammal Science*, 22(4), 847–861.
- Bakhchina, A. V., L. M. Mukhametov, V. V. Rozhnov, and O. I. Lyamin. (2017). Spectral analysis of heart rate variability in the beluga (*Delphinapterus leucas*) during exposure to acoustic noise. *Journal of Evolutionary Biochemistry and Physiology*, 53(1), 60–65.
- Barlow, J. (2006). Cetacean abundance in Hawaiian waters estimated from a Summer–Fall survey in 2002. *Marine Mammal Science*, 22(2), 446–464.
- Barlow, J., J. Calambokidis, E. A. Falcone, C. S. Baker, A. M. Burdin, P. J. Clapham, J. K. B. Ford, C. M. Gabriele, R. LeDuc, D. K. Mattila, T. J. Quinn, II, L. Rojas-Bracho, J. M. Straley, B. L. Taylor, J. Urbán R, P. Wade, D. Weller, B. H. Witteveen, and M. Yamaguchi. (2011). Humpback whale abundance in the North Pacific estimated by photographic capture-recapture with bias correction from simulation studies. *Marine Mammal Science*, 27(4), 793–818.
- Barlow, J. (2016). *Cetacean Abundance in the California Current Estimated from Ship-based Line-transect Surveys in 1991–2014*. (NOAA Administrative Report NMFS-SWFSC-LJ-1601). La Jolla, CA: Southwest Fisheries Science Center.
- Bassett, C., J. Thomson, and B. Polagye. (2010). *Characteristics of Underwater Ambient Noise at a Proposed Tidal Energy Site in Puget Sound*. Seattle, WA: Northwest National Marine Renewable Energy Center.
- Baulch, S., and C. Perry. (2014). Evaluating the impacts of marine debris on cetaceans. *Marine Pollution Bulletin*, 80(1–2), 210–221.

- Baumann-Pickering, S., A. E. Simonis, M. A. Roch, M. A. McDonald, A. Solsona-Berga, E. M. Oleson, S. M. Wiggins, R. L. Brownell, Jr., and J. A. Hildebrand. (2012). *Spatio-temporal patterns of beaked whale echolocation signals in the North Pacific* (2012 Marine Mammal & Biology Program Review). Arlington, VA: Office of Naval Research.
- Baumann-Pickering, S., A. J. Debich, J. T. Trickey, A. Širović, R. Gresalfi, M. A. Roche, S. M. Wiggins, J. A. Hildebrand, and J. A. Carretta. (2013). *Examining Explosions in Southern California and Their Potential Impact on Cetacean Acoustic Behavior*. La Jolla, CA: Southwest Fisheries Science Center.
- Bejder, L., A. Samuels, H. Whitehead, and N. Gales. (2006a). Interpreting short-term behavioural responses to disturbance within a longitudinal perspective. *Animal Behaviour*, 72, 1149–1158.
- Bejder, L., A. Samuels, H. Whitehead, N. Gales, J. Mann, R. Connor, M. Heithaus, J. Waston-Capps, C. Flaherty, and M. Krützen. (2006b). Decline in relative abundance of bottlenose dolphins exposed to long-term disturbance. *Conservation Biology*, 20(6), 1791–1798.
- Benoit-Bird, K. J., W. W. L. Au, R. E. Brainard, and M. O. Lammers. (2001). Diel horizontal migration of the Hawaiian mesopelagic boundary community observed acoustically. *Marine Ecology Progress Series*, 217, 1–14.
- Benoit-Bird, K. J., and W. W. L. Au. (2003). Prey dynamics affect foraging by a pelagic predator (*Stenella longirostris*) over a range of spatial and temporal scales. *Behavioral Ecology and Sociobiology*, 53, 364–373.
- Benoit-Bird, K. J. (2004). Prey caloric value and predator energy needs: Foraging predictions for wild spinner dolphins. *Marine Biology*, 145(3), 435–444.
- Benoit-Bird, K. J., and W. W. L. Au. (2004). Diel migration dynamics of an island-associated sound-scattering layer. *Deep-Sea Research I*, 51, 707–719.
- Benoit-Bird, K. J., and W. W. L. Au. (2009). Cooperative prey herding by the pelagic dolphin, *Stenella longirostris*. *The Journal of the Acoustical Society of America*, 125(1), 125–137.
- Bergström, L., L. Kautsky, T. Malm, R. Rosenberg, M. Wahlberg, N. Åstrand Capetillo, and D. Wilhelmsson. (2014). Effects of offshore wind farms on marine wildlife—A generalized impact assessment. *Environmental Research Letters*, 9(3), 12.
- Berini, C. R., L. M. Kracker, and W. E. McFee. (2015). *Modeling Pygmy Sperm Whale (Kogia breviceps) Strandings Along the Southeast Coast of the United States from 1992 to 2006 in Relation to Environmental Factors* (NOAA Technical Memorandum NOS-NCCOS-203). Charleston, SC: National Oceanic and Atmospheric Administration.
- Berman-Kowalewski, M., F. M. D. Gulland, S. Wilkin, J. Calambokidis, B. Mate, J. Cordaro, D. Rotstein, J. S. Leger, P. Collins, K. Fahy, and S. Dover. (2010). Association Between Blue Whale (*Balaenoptera musculus*) Mortality and Ship Strikes Along the California Coast. *Aquatic Mammals*, 36(1), 59–66.
- Bernaldo de Quiros, Y., O. Gonzalez-Diaz, M. Arbelo, E. Sierra, S. Sacchini, and A. Fernandex. (2012). Decompression vs. decomposition: Distribution, amount, and gas composition of bubbles in stranded marine mammals. *Frontiers in Physiology*, 3 Article 177, 19.
- Bernaldo de Quiros, Y., O. Gonzalez-Diaz, A. Mollerlokken, A. O. Brubakk, A. Hjelde, P. Saavedra, and A. Fernandez. (2013a). Differentiation at autopsy between in vivo gas embolism and putrefaction using gas composition analysis. *International Journal of Legal Medicine*, 127(2), 437–445.

- Bernaldo de Quiros, Y., J. S. Seewald, S. P. Sylva, B. Greer, M. Niemeyer, A. L. Bogomolni, and M. J. Moore. (2013b). Compositional discrimination of decompression and decomposition gas bubbles in bycaught seals and dolphins. *PLoS ONE*, 8(12), e83994.
- Bernard, H. J., and S. B. Reilly. (1999). Pilot whales, *Globicephala* Lesson, 1828. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 6, pp. 245–280). San Diego, CA: Academic Press.
- Bernasconi, M., R. Patel, and L. Nøttestad. (2012). Behavioral observations of baleen whales in proximity of a modern fishing vessel. In A. N. Popper & A. Hawkins (Eds.), *The Effects of Noise on Aquatic Life*. New York, NY: Springer.
- Berrow, S. D., and B. Holmes. (1999). Tour boats and dolphins: A note on quantifying the activities of whalewatching boats in the Shannon Estuary, Ireland. *Journal of Cetacean Research and Management*, 1(2), 199–204.
- Besseling, E., E. M. Foekema, J. A. Van Franeker, M. F. Leopold, S. Kuhn, E. L. B. Rebolledo, E. Hebe, L. Mielke, J. Ijzer, P. Kamminga, and A. A. Koelmans. (2015). Microplastic in a macro filter feeder: Humpback whale *Megaptera novaeangliae*. *Marine Pollution Bulletin*, 95(1), 248–252.
- Bettridge, S., C. S. Baker, J. Barlow, P. J. Clapham, M. Ford, D. Gouveia, D. K. Mattila, R. M. Pace, III, P. E. Rosel, G. K. Silber, and P. R. Wade. (2015). *Status Review of the Humpback Whale (Megaptera novaeangliae) under the Endangered Species Act* (NOAA Technical Memorandum NMFS-SWFSC-540). La Jolla, CA: Southwest Fisheries Science Center.
- Blackwell, S. B., J. W. Lawson, and M. T. Williams. (2004). Tolerance by ringed seals (*Phoca hispida*) to impact pipe-driving and construction sounds at an oil production island. *The Journal of the Acoustical Society of America*, 115(5 [Pt. 1]), 2346–2357.
- Blackwell, S. B., C. S. Nations, T. L. McDonald, C. R. Greene, A. M. Thode, M. Guerra, and A. M. Macrander. (2013). Effects of airgun sounds on bowhead whale calling rates in the Alaskan Beaufort Sea. *Marine Mammal Science*, 29, E342–E365.
- Blackwell, S. B., C. S. Nations, T. L. McDonald, A. M. Thode, D. Mathias, K. H. Kim, C. R. Greene, Jr., and A. M. Macrander. (2015). Effects of airgun sounds on bowhead whale calling rates: Evidence for two behavioral thresholds. *PLoS ONE*, 10(6), e0125720.
- Blackwell, S. B., C. S. Nations, A. M. Thode, M. E. Kauffman, A. S. Conrad, R. G. Norman, and K. H. Kim. (2017). Effects of tones associated with drilling activities on bowhead whale calling rates. *PLoS ONE*, 12(11), e0188459.
- Bonito, L. T., A. Hamdoun, and S. A. Sandin. (2016). Evaluation of the global impacts of mitigation on persistent, bioaccumulative and toxic pollutants in marine fish. *PeerJ*, 4, e1573.
- Bowles, A. E., M. Smultea, B. Wursig, D. P. DeMaster, and D. Palka. (1994). Relative abundance and behavior of marine mammals exposed to transmissions from the Heard Island Feasibility Test. *The Journal of the Acoustical Society of America*, 96, 2469–2484.
- Boyd, I., D. Claridge, C. Clark, and B. Southall. (2008). *BRS 2008 Preliminary Report*. Washington, DC: U.S. Navy NAVSEA PEO IWS 5, ONR, U.S. Navy Environmental Readiness Division, National Oceanic and Atmospheric Administration, Strategic Environmental Research and Development Program.
- Boyd, J. D., and D. J. Brightsmith. (2013). Error properties of Argos satellite telemetry locations using least squares and Kalman filtering. *PLoS ONE*, 8(5), e63051.

- Bradford, A. L., and K. A. Forney. (2014). *Injury Determinations for Cetaceans Observed Interacting with Hawaii and American Samoa Longline Fisheries During 2008–2012* (NOAA Technical Memorandum NMFS-PIFSC-41). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Bradford, A. L., K. A. Forney, E. M. Oleson, and J. Barlow. (2014). Accounting for subgroup structure in line-transect abundance estimates of false killer whales (*Pseudorca crassidens*) in Hawaiian waters. *PLoS ONE*, 9(2), e90464.
- Bradford, A. L., and E. Lyman. (2015). *Injury Determinations for Humpback Whales and Other Cetaceans Reported to NOAA Response Networks in the Hawaiian Islands During 2007–2012* (NOAA Technical Memorandum NMFS-PIFSC-45). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Bradford, A. L., E. A. Oleson, R. W. Baird, C. H. Boggs, K. A. Forney, and N. C. Young. (2015). *Revised Stock Boundaries for False Killer Whales (Pseudorca crassidens) in Hawaiian Waters* (NOAA Technical Memorandum NMFS-PIFSC-47). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Bradford, A. L., and K. A. Forney. (2016). *Injury Determinations for Marine Mammals Observed Interacting with Hawaii and American Samoa Longline Fisheries During 2009–2013* (NOAA Technical Memorandum NMFS-PIFSC-50). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Bradford, A. L., and K. A. Forney. (2017). *Injury Determinations for Marine Mammals Observed Interacting with Hawaii and American Samoa Longline Fisheries During 2010–2014* (NOAA Technical Memorandum NMFS-PIFSC-62). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Bradford, A. L., K. A. Forney, E. M. Oleson, and J. Barlow. (2017). Abundance estimates of cetaceans from a line-transect survey within the U.S. Hawaiian Islands Exclusive Economic Zone. *Fishery Bulletin*, 115(2), 129–142.
- Bradshaw, C. J. A., K. Evans, and M. A. Hindell. (2006). Mass cetacean strandings—A plea for empiricism. *Conservation Biology*, 20(2), 584–586.
- Branch, T. A. (2007). Abundance of Antarctic blue whales south of 60°S from three complete circumpolar sets of surveys. *Journal of Cetacean Research and Management*, 9(3), 253–262.
- Brandt, M. J., A. Diederichs, K. Betke, and G. Nehls. (2011). Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series*, 421, 205–216.
- Branstetter, B. K., and J. J. Finneran. (2008). Comodulation masking release in bottlenose dolphins (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 1, 625–633.
- Branstetter, B. K., J. S. Trickey, K. Bakhtiari, A. Black, H. Aihara, and J. J. Finneran. (2013). Auditory masking patterns in bottlenose dolphins (*Tursiops truncatus*) with natural, anthropogenic, and synthesized noise. *The Journal of the Acoustical Society of America*, 133(3), 1811–1818.
- Branstetter, B. K., K. Bakhtiari, A. Black, J. S. Trickey, J. J. Finneran, and H. Aihara. (2016). Energetic and informational masking of complex sounds by a bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, 140(3), 1904–1917.
- Branstetter, B. K., J. St. Leger, D. Acton, J. Stewart, D. Houser, J. J. Finneran, and K. Jenkins. (2017a). Killer whale (*Orcinus orca*) behavioral audiograms. *The Journal of the Acoustical Society of America*, 141, 2387–2398.
- Branstetter, B. K., K. R. Van Alstyne, T. A. Wu, R. A. Simmons, L. D. Curtis, and M. J. Xitco, Jr. (2017b). Critical ratio functions for odontocete cetaceans. *The Journal of the Acoustical Society of America*, 142, 1897–1900.

- Briggs, C., S. M. Shjegstad, J. A. K. Silva, and M. H. Edwards. (2016). Distribution of chemical warfare agent, energetics, and metals in sediments at a deep-water discarded military munitions site. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 63–69.
- Browne, M. A., P. Crump, S. J. Niven, E. Teuten, A. Tonkin, T. Galloway, and R. Thompson. (2011). Accumulation of microplastic on shorelines worldwide: Sources and sinks. *Environmental Science & Technology*, 45(21), 9175–9179.
- Brownell, R. L., Jr., K. Ralls, S. Baumann-Pickering, and M. M. Poole. (2009a). Behavior of melon-headed whales, *Peponocephala electra*, near oceanic islands. *Marine Mammal Science*, 25(3), 639–658.
- Brownell, R. L., Jr., C.-J. Yao, C.-S. Lee, and M.-C. Wang. (2009b). *Worldwide Review Of Pygmy Killer Whales, Feresa attenuate, Mass Strandings Reveals Taiwan Hot Spot* (Paper 141). Washington, DC: U.S. Department of Commerce.
- Brumm, H., and H. Slabbekoorn. (2005). Acoustic communication in noise. *Advances in the Study of Behavior*, 35, 151–209.
- Bryant, P. J., C. M. Lafferty, and S. K. Lafferty. (1984). Reoccupation of Laguna Guerrero Negro, Baja California, Mexico, by Gray Whales. In M. L. Jones, S. L. Swartz, & S. Leatherwood (Eds.), *The Gray Whale: Eschrichtius robustus* (pp. 375–387). Orlando, FL: Academic Press.
- Bull, J. C., P. D. Jepson, R. K. Ssuna, R. Deaville, C. R. Allchin, R. J. Law, and A. Fenton. (2006). The relationship between polychlorinated biphenyls in blubber and levels of nematode infestations in harbour porpoises, *Phocoena phocoena*. *Parasitology*, 132(Pt 4), 565–573.
- Burkholder, J., D. Eggleston, H. Glasgow, C. Brownie, R. Reed, G. Janowitz, M. Posey, G. Mella, C. Kinder, R. Corbett, D. Toms, T. Alphin, N. Deamer, and J. Springer. (2004). Comparative impacts of two major hurricane seasons on the Neuse River and western Pamlico Sound ecosystems. *Proceedings of the National Academy of Sciences*, 101(25), 9291–9296.
- Calambokidis, J., E. A. Falcone, T. J. Quinn, A. M. Burdin, P. J. Clapham, J. K. B. Ford, C. M. Gabriele, R. LeDuc, D. Mattila, L. Rojas-Bracho, J. M. Straley, B. L. Taylor, J. Urbán R., D. Weller, B. H. Witteveen, M. Yamaguchi, A. Bendlin, D. Camacho, K. Flynn, A. Havron, J. Huggins, and N. Maloney. (2008). *SPLASH: Structure of Populations, Levels of Abundance and Status of Humpback Whales in the North Pacific*. Olympia, WA: Cascadia Research.
- Calambokidis, J., J. Barlow, J. K. B. Ford, T. E. Chandler, and A. B. Douglas. (2009a). Insights into the population structure of blue whales in the Eastern North Pacific from recent sightings and photographic identification. *Marine Mammal Science*, 25(4), 816–832.
- Calambokidis, J., E. M. Oleson, M. F. McKenna, and J. A. Hildebrand. (2009b). *Blue whale behavior in shipping lanes and response to ships*. Paper presented at the 2009 Office of Naval Research Marine Mammal Program Review. Alexandria, VA.
- Calambokidis, J., J. L. Laake, and A. Klimek. (2010). *Abundance and Population Structure of Seasonal Gray Whales in the Pacific Northwest, 1998–2008*. Washington, DC: International Whaling Commission Scientific Committee.
- Calambokidis, J. (2012). *Summary of Ship-Strike Related Research on Blue Whales in 2011*. Olympia, WA: Cascadia Research.
- Caldwell, D. K., and M. C. Caldwell. (1989). Pygmy sperm whale, *Kogia breviceps* (de Blainville, 1838): Dwarf sperm whale *Kogia simus* Owen, 1866. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 4, pp. 234–260). San Diego, CA: Academic Press.

- Campbell, G. S., D. W. Weller, and J. A. Hildebrand. (2010). *SIO Small Boat Based Marine Mammal Surveys in Southern California: Report of Results for August 2009–July 2010: Annual Range Complex Monitoring Report for Hawaii and Southern California. Draft submission to the National Marine Fisheries Service September 15, 2010.* San Diego, CA: U.S. Department of the Navy.
- Campbell, G. S., L. Thomas, K. Whitaker, A. B. Douglas, J. Calambokidis, and J. A. Hildebrand. (2015). Inter-annual and seasonal trends in cetacean distribution, density and abundance off southern California. *Deep Sea Research Part II: Topical Studies in Oceanography*, 112, 143–157.
- Cañadas, A., R. Sagarminaga, and S. García-Tiscar. (2002). Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. *Deep-Sea Research I*, 49, 2053–2073.
- Carrera, M. L., E. G. P. Favaro, and A. Souto. (2008). The response of marine tucuxis (*Sotalia fluviatilis*) towards tourist boats involves avoidance behaviour and a reduction in foraging. *Animal Welfare*, 17, 117–123.
- Carretta, J. V., J. Barlow, and L. Enriquez. (2008). Acoustic pingers eliminate beaked whale bycatch in a gill net fishery. *Marine Mammal Science*, 24(4), 2053–2073.
- Carretta, J. V. (2013). *A Review of Sperm Whale (Physeter macrocephalus) Bycatch in the California Swordfish and Thresher Shark Drift Gillnet Fishery.* La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., S. M. Wilkin, M. M. Muto, and K. Wilkinson. (2013). *Sources of Human-Related Injury and Mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2007–2011* (NOAA Technical Memorandum NMFS-SWFSC-514). La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., S. M. Wilkin, M. M. Muto, K. Wilkinson, and J. Rustin. (2014). *Sources of Human-Related Injury and Mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2008–2012.* (NOAA-TM-NMFS-SWFSC-533). La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., K. Danil, S. J. Chivers, D. W. Weller, D. S. Janiger, M. Berman-Kowalewski, K. M. Hernandez, J. T. Harvey, R. C. Dunkin, D. R. Casper, S. Stoudt, M. Flannery, K. Wilkinson, J. Huggins, and D. M. Lambourn. (2016a). Recovery rates of bottlenose dolphin (*Tursiops truncatus*) carcasses estimated from stranding and survival rate data. *Marine Mammal Science*, 32(1), 349–362.
- Carretta, J. V., M. M. Muto, S. Wilkin, J. Greenman, K. Wilkinson, M. DeAngelis, J. Viezbicke, D. Lawson, and J. Jannot. (2016b). *Sources of Human-Related Injury and Mortality for U.S. Pacific West Coast Marine Mammal Stock Assessments, 2010–2014.* La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., E. M. Oleson, J. Baker, D. W. Weller, A. R. Lang, K. A. Forney, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, J. E. Moore, D. Lynch, L. Carswell, and R. L. Brownell Jr. (2016c). *U.S. Pacific Marine Mammal Stock Assessments: 2015.* La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., J. E. Moore, and K. A. Forney. (2017a). *Regression Tree and Ratio Estimates of Marine Mammal, Sea Turtle, and Seabird Bycatch in the California Drift Gillnet Fishery: 1990–2015.* La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., M. M. Muto, J. Greenman, K. Wilkinson, D. Lawson, J. Viezbicke, and J. Jannot. (2017b). *Sources of Human-Related Injury and Mortality for U.S. Pacific West Coast Marine Mammal*

- Stock Assessments, 2011–2015* (NOAA Technical Memorandum NMFS-SWFSC-579). La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., E. M. Oleson, J. Baker, D. W. Weller, A. R. Lang, K. A. Forney, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, J. E. Moore, D. Lynch, L. Carswell, and R. L. Brownell, Jr. (2017c). *U.S. Pacific Marine Mammal Stock Assessments: 2016* (NOAA Technical Memorandum NMFS-SWFSC-561). La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., E. M. Oleson, K. A. Forney, J. Baker, J. E. Moore, D. W. Weller, A. R. Lang, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, D. Lynch, L. Carswell, and R. L. Brownell, Jr. (2017d). *U.S. Pacific Draft Marine Mammal Stock Assessments: 2017* (NOAA Technical Memorandum NMFS-SWFSC-602). La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., K. A. Forney, E. M. Oleson, D. W. Weller, A. R. Lang, J. Baker, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, J. E. Moore, D. Lynch, L. Carswell, and R. L. Brownell, Jr. (2018a). *U.S. Pacific Marine Mammal Stock Assessments: 2017*. La Jolla, CA: Southwest Fisheries Science Center.
- Carretta, J. V., K. A. Forney, E. M. Oleson, D. W. Weller, A. R. Lang, J. Baker, M. M. Muto, B. Hanson, A. J. Orr, H. Huber, M. S. Lowry, J. Barlow, J. E. Moore, D. Lynch, L. Carswell, and R. L. Brownell, Jr. (2018b). *U.S. Pacific Draft Marine Mammal Stock Assessments: 2018* (NOAA Technical Memorandum NMFS-SWFSC-XXX). La Jolla, CA: National Marine Fisheries Service, Southwest Fisheries Science Center.
- Castellote, M., C. W. Clark, and M. O. Lammers. (2012). Acoustic and behavioral changes by fin whales (*Balaenoptera physalus*) in responses to shipping and airgun noise. *Biological Conservation*, 147, 115–122.
- Castellote, M., T. A. Mooney, L. Quakenbush, R. Hobbs, C. Goertz, and E. Gaglione. (2014). Baseline hearing abilities and variability in wild beluga whales (*Delphinapterus leucas*). *The Journal of Experimental Biology*, 217(Pt 10), 1682–1691.
- Cecchetti, A., K. A. Stockin, J. Gordon, and J. M. N. Azevedo. (2017). Short-term effects of tourism on the behaviour of common dolphins (*Delphinus delphis*) in the Azores. *Journal of the Marine Biological Association of the United Kingdom*, 98(5), 1187–1196.
- Cerchio, S., S. Strindberg, T. Collins, C. Bennett, and H. Rosenbaum. (2014). Seismic surveys negatively affect humpback whale singing activity off northern Angola. *PLoS ONE*, 9(3), e86464.
- Cerchio, S., B. Andrianantenaina, A. Lindsay, M. Rekdahl, N. Andrianarivelo, and T. Rasoloarijao. (2015). Omura's whales (*Balaenoptera omurai*) off northwest Madagascar: Ecology, behaviour and conservation needs. *Royal Society Open Science*, 2(10), 150301.
- Charif, R. A., C. S. Oedekoven, A. Rahaman, B. J. Estabrook, L. Thomas, and A. N. Rice. (2015). *Development of Statistical Methods for Assessing Changes in Whale Vocal Behavior in Response to Mid-Frequency Active Sonar. Final Report*. Virginia Beach, VA: U.S. Fleet Forces Command.
- Chivers, S. J., R. W. Baird, D. J. McSweeney, D. L. Webster, N. M. Hedrick, and J. C. Salinas. (2007). Genetic variation and evidence for population structure in eastern North Pacific false killer whales (*Pseudorca crassidens*). *Canadian Journal of Zoology*, 85(7), 783–794.
- Cholewiak, D., A. I. DeAngelis, D. Palka, P. J. Corkeron, and S. M. Van Parijs. (2017). Beaked whales demonstrate a marked acoustic response to the use of shipboard echosounders. *Royal Society Open Science*, 4(12), 170940.

- Cholewiak, D., C. W. Clark, D. Ponirakis, A. Frankel, L. T. Hatch, D. Risch, J. E. Stanistreet, M. Thompson, E. Vu, and S. M. Van Parijs. (2018). Communicating amidst the noise: Modeling the aggregate influence of ambient and vessel noise on baleen whale communication space in a national marine sanctuary. *Endangered Species Research*, 36, 59–75.
- Choy, C. A., and J. C. Drazen. (2013). Plastic for dinner? Observations of frequent debris ingestion by pelagic predatory fishes from the central North Pacific. *Marine Ecology Progress Series*, 485, 155–163.
- Christian, E. A., and J. B. Gaspin. (1974). *Swimmer Safe Standards from Underwater Explosions*. Navy Science Assistance Program Project No. PHP-11-73. White Oak, MD: Naval Ordnance Laboratory.
- Christiansen, F., D. Lusseau, E. Stensland, and P. Berggren. (2010). Effects of tourist boats on the behaviour of Indo-Pacific bottlenose dolphins off the south coast of Zanzibar. *Endangered Species Research*, 11, 91–99.
- Christiansen, F., M. Rasmussen, and D. Lusseau. (2013). Whale watching disrupts feeding activities of minke whales on a feeding ground. *Marine Ecology Progress Series*, 478, 239–251.
- Christiansen, F., M. Rasmussen, and D. Lusseau. (2014). Inferring energy expenditure from respiration rates in minke whales to measure the effects of whale watching boat interactions. *Journal of Experimental Marine Biology and Ecology*, 459, 96–104.
- Christiansen, F., and D. Lusseau. (2015). Linking behavior to vital rates to measure the effects of non-lethal disturbance on wildlife. *Conservation Letters*, 8(6), 424–431.
- Christiansen, F., A. M. Dujon, K. R. Sprogis, J. P. Y. Arnould, and L. Bejder. (2016a). Noninvasive unmanned aerial vehicle provides estimates of the energetic cost of reproduction in humpback whales. *Ecosphere*, 7(10), e01468.
- Christiansen, F., L. Rojano-Doñate, P. T. Madsen, and L. Bejder. (2016b). Noise levels of multi-rotor unmanned aerial vehicles with implications for potential underwater impacts on marine mammals. *Frontiers in Marine Science*, 3(277), 1–9.
- Clapham, P. J., and J. G. Mead. (1999). *Megaptera novaeangliae*. *American Society of Mammalogists*, 604, 1–9.
- Claridge, D., D. Charlotte, and J. Durban. (2009). *Abundance and movement patterns of Blainville's beaked whales at the Atlantic Undersea Test and Evaluation Center*. Paper presented at the 2009 Office of Naval Research Marine Mammal Program Review. Alexandria, VA.
- Clark, C. (2015). *Potential Acoustic Impacts of Vessel Traffic from the Trans Mountain Expansion Project on Southern Resident Killer Whales*. Sidney, Canada: Prepared for Raincoast Conservation Foundation.
- Clark, C. W., and K. M. Fristrup. (2001). Baleen whale responses to low-frequency human-made underwater sounds. *The Journal of the Acoustical Society of America*, 110(5), 2751.
- Clark, C. W., W. T. Ellison, B. L. Southall, L. Hatch, S. M. Van Parijs, A. Frankel, and D. Ponirakis. (2009). Acoustic masking in marine ecosystems: Intuitions, analysis, and implication. *Marine Ecology Progress Series*, 395, 201–222.
- Clark, S. L., and J. W. Ward. (1943). The effects of rapid compression waves on animals submerged in water. *Surgery, Gynecology & Obstetrics*, 77, 403–412.

- Clarke, M. R. (1996). Cephalopods as prey III. Cetaceans. *Philosophical Transactions of the Royal Society of London*, 351, 1053–1065.
- Constantine, R., T. Iwata, S. L. Nieukirk, and G. S. Penry. (2018). Future Directions in Research on Byrde's Whales. *Frontiers in Marine Science*, 5(333), 1–7.
- Costa, D. P. (1993). The relationship between reproductive and foraging energetics and the evolution of the Pinnipedia. *Symposium of the Zoological Society of London*, 66, 293–314.
- Costa, D. P., and B. A. Block. (2009). *Use of Electronic Tag Data and Associated Analytical Tools to Identify and Predict Habitat Utilization of Marine Predators* (Marine Mammals & Biological Oceanography Annual Reports: FY09). Santa Cruz, CA and Stanford, CA: Office of Naval Research.
- Costa, D. P., L. A. Hückstädt, L. K. Schwarz, A. S. Friedlaender, B. R. Mate, A. N. Zerbini, A. Kennedy, and N. J. Gales. (2016a). *Assessing the exposure of animals to acoustic disturbance: Towards an understanding of the population consequences of disturbance*. Paper presented at the Fourth International Conference on the Effects of Noise on Aquatic Life. Dublin, Ireland.
- Costa, D. P., L. Schwarz, P. Robinson, R. S. Schick, P. A. Morris, R. Condit, D. E. Crocker, and A. M. Kilpatrick. (2016b). A bioenergetics approach to understanding the population consequences of disturbance: Elephant seals as a model system. In *The Effects of Noise on Aquatic Life II* (pp. 116–169). New York, NY: Springer.
- Costidis, A. M., and S. A. Rommel. (2016). The extracranial venous system in the heads of beaked whales, with implications on diving physiology and pathogenesis. *Journal of Morphology*, 277(1), 34–64.
- Courbis, S., and G. Timmel. (2008). Effects of vessels and swimmers on behavior of Hawaiian spinner dolphins (*Stenella longirostris*) in Kealake'akua, Honaunau, and Kauhako bays, Hawai'i. *Marine Mammal Science*, 25(2), 430–440.
- Cox, T. M., T. J. Ragen, A. J. Read, E. Vox, R. W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D'Spain, A. Fernandez, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J. Hildebrand, D. Houser, T. Hullar, P. D. Jepson, D. Ketten, C. D. MacLeod, P. Miller, S. Moore, D. C. Mountain, D. Palka, P. Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Mead, and L. Benner. (2006). Understanding the impacts of anthropogenic sound on beaked whales. *Journal of Cetacean Research and Management*, 7(3), 177–187.
- Cozar, A., F. Echevarria, J. I. Gonzalez-Gordillo, X. Irigoien, B. Ubeda, S. Hernandez-Leon, A. T. Palma, S. Navarro, J. Garcia-de-Lomas, A. Ruiz, M. L. Fernandez-de-Puelles, and C. M. Duarte. (2014). Plastic debris in the open ocean. *Proceedings of the National Academy of Science of the United States of America*, 111(28), 10239–10244.
- Croll, D. A., C. W. Clark, J. Calambokidis, W. T. Ellison, and B. R. Tershy. (2001). Effect of anthropogenic low-frequency noise on the foraging ecology of *Balaenoptera* whales. *Animal Conservation*, 4, 13–27.
- Crowell, S. E., A. M. Wells-Berlin, R. E. Therrien, S. E. Yannuzzi, and C. E. Carr. (2016). In-air hearing of a diving duck: A comparison of psychoacoustic and auditory brainstem response thresholds. *The Journal of the Acoustical Society of America*, 139(5), 3001.
- Crum, L., and Y. Mao. (1996). Acoustically enhanced bubble growth at low frequencies and its implications for human diver and marine mammal safety. *The Journal of the Acoustical Society of America*, 99(5), 2898–2907.

- Crum, L., M. Bailey, J. Guan, P. Hilmo, S. Kargl, and T. Matula. (2005). Monitoring bubble growth in supersaturated blood and tissue *ex vivo* and the relevance to marine mammal bioeffects. *Acoustics Research Letters Online*, 6(3), 214–220.
- Cruz-Uribe, O., D. P. Cheney, and G. L. Rorrer. (2007). Comparison of TNT removal from seawater by three marine macroalgae. *Chemosphere*, 67, 1469–1476.
- Culik, B. M., S. Koschinski, N. Tregenza, and G. M. Ellis. (2001). Reactions of harbor porpoises *Phocoena phocoena* and herring *Clupea harengus* to acoustic alarms. *Marine Ecological Progress Series*, 211, 255–260.
- Culik, B. M. (2004). *Review of Small Cetaceans Distribution, Behaviour, Migration and Threats*. Bonn, Germany: United National Environment Programme and the Secretariat of the Convention on the Conservation of Migratory Species of Wild Animals.
- Cummings, W. C., and P. O. Thompson. (1971). Gray whales, *Eschrichtius robustus*, avoid the underwater sounds of killer whales, *Orcinus orca*. *Fishery Bulletin*, 69(3), 525–530.
- Cunningham, K. A., B. L. Southall, and C. Reichmuth. (2014). Auditory sensitivity of seals and sea lions in complex listening scenarios. *The Journal of the Acoustical Society of America*, 136(6), 3410–3421.
- Curé, C., R. Antunes, F. Samarra, A. C. Alves, F. Visser, P. H. Kvadsheim, and P. J. Miller. (2012). Pilot whales attracted to killer whale sounds: Acoustically-mediated interspecific interactions in cetaceans. *PLoS ONE*, 7(12), e52201.
- Curé, C., L. D. Sivle, F. Visser, P. J. Wensveen, S. Isojunno, C. M. Harris, P. H. Kvadsheim, F. P. A. Lam, and P. J. O. Miller. (2015). Predator sound playbacks reveal strong avoidance responses in a fight strategist baleen whale. *Marine Ecology Progress Series*, 526, 267–282.
- Curé, C., S. Isojunno, F. Visser, P. J. Wensveen, L. D. Sivle, P. H. Kvadsheim, F. P. A. Lam, and P. J. O. Miller. (2016). Biological significance of sperm whale responses to sonar: Comparison with anti-predator responses. *Endangered Species Research*, 31, 89–102.
- Currie, J. J., S. H. Stack, J. A. McCordic, and G. D. Kaufman. (2017). Quantifying the risk that marine debris poses to cetaceans in coastal waters of the 4-island region of Maui. *Marine Pollution Bulletin*, 121(1–2), 69–77.
- D'Vincent, C. G., R. M. Nilson, and R. E. Hanna. (1985). Vocalization and coordinated feeding behavior of the humpback whale in southeastern Alaska. *Scientific Reports of the Whales Research Institute*, 36, 41–47.
- Dähne, M., V. Peschko, A. Gilles, K. Lucke, S. Adler, K. Ronnenberg, and U. Siebert. (2014). Marine mammals and windfarms: Effects of alpha ventus on harbour porpoises. In *Ecological Research at the Offshore Windfarm alpha ventus* (pp. 133–149). New York, NY: Springer Publishing.
- Dähne, M., J. Tougaard, J. Carstensen, A. Rose, and J. Nabe-Nielsen. (2017). Bubble curtains attenuate noise from offshore wind farm construction and reduce temporary habitat loss for harbour porpoises. *Marine Ecology Progress Series*, 580, 221–237.
- Dalebout, M. L., G. M. Lento, F. Cipriano, N. Funahashi, and C. S. Baker. (2002a). How many protected minke whales are sold in Japan and Korea? A census by microsatellite. *Animal Conservation*, 5, 143–152.

- Dalebout, M. L., J. G. Mead, C. S. Baker, A. N. Baker, and A. L. van Helden. (2002b). A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine Mammal Science*, 18(3), 577–608.
- Dalebout, M. L., G. J. B. Ross, C. S. Baker, R. C. Anderson, P. B. Best, V. G. Cockcroft, H. L. Hinsz, V. Peddemors, and R. L. Pitman. (2003). Appearance, distribution and genetic distinctiveness of Longman's beaked whale, *Indopacetus pacificus*. *Marine Mammal Science*, 19(3), 421–461.
- Danil, K., and J. A. St Leger. (2011). Seabird and dolphin mortality associated with underwater detonation exercises. *Marine Technology Society Journal*, 45(6), 89–95.
- Darling, J. D., J. Calambokidis, K. C. Balcomb, P. Bloedel, K. Flynn, A. Mochizuki, K. Mori, F. Sato, H. Sukanuma, and M. Yamaguchi. (1996). Movement of a humpback whale (*Megaptera Novaeangliae*) from Japan to British Columbia and return. *Marine Mammal Science*, 12(2), 281–287.
- Davis, R. W., N. Jaquet, D. Gendron, U. Markaida, G. Bazzino, and W. Gilly. (2007). Diving behavior of sperm whales in relation to behavior of a major prey species, the jumbo squid, in the Gulf of California, Mexico. *Marine Ecology Progress Series*, 333, 291–302.
- De Pierrepont, J. F., B. Dubois, S. Desormonts, M. B. Santos, and J. P. Robin. (2005). Stomach contents of English Channel cetaceans stranded on the coast of Normandy. *Journal of the Marine Biological Association of the United Kingdom*, 85, 1539–1546.
- De Silva, R., K. Grellier, G. Lye, N. McLean, and P. Thompson. (2014). *Use of population viability analysis to assess the potential for long term impacts from piling noise on marine mammal populations - a case study from the Scottish east coast*. Paper presented at the Proceedings of the 2nd International Conference on Environmental Interactions of Marine Renewable Energy Technologies (EIMR2014). Stornoway. Isle of Lewis, Outer Hebrides, Scotland.
- de Vos, A. (2017). First record of Omura's whale, *Balaenoptera omurai*, in Sri Lankan waters. *Marine Biodiversity Records*, 10(18), 1–4.
- Deakos, M. H., and M. F. Richlen. (2015). *Vessel-Based Marine Mammal Survey on the Navy Range off Kauai in Support of Passive Acoustic Monitoring and Satellite-Tagging Efforts: 1–9 February 2014*. Honolulu, HI: HDR Inc.
- Deakos, M. H., R. K. Uyeyama, M. W. Richie, M. F. Richlen, and J. M. Aschettino. (2016). *Guam Marine Species Monitoring Survey, Shore Station Study: May 2013 and March 2015. Final Field Report*. Honolulu, HI: HDR Inc.
- Deecke, V. B., P. J. B. Slater, and J. K. B. Ford. (2002). Selective habituation shapes acoustic predator recognition in harbour seals. *Nature*, 420(November 14), 171–173.
- Defence Science and Technology Laboratory. (2007). *Observations of Marine Mammal Behaviour in Response of Active Sonar*. Salisbury, United Kingdom: Ministry of Defence.
- DeForges, J. P. W., M. Galbraith, N. Dangerfield, and P. S. Ross. (2014). Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean. *Marine Pollution Bulletin*, 79, 94–99.
- Deng, Z. D., B. L. Southall, T. J. Carlson, J. Xu, J. J. Martinez, M. A. Weiland, and J. M. Ingraham. (2014). 200 kHz commercial sonar systems generate lower frequency side lobes audible to some marine mammals. *PLoS ONE*, 9(4), e95315.

- Dennison, S., M. J. Moore, A. Fahlman, K. Moore, S. Sharp, C. T. Harry, J. Hoppe, M. Niemeyer, B. Lentell, and R. S. Wells. (2012). Bubbles in live-stranded dolphins. *Proceedings of the Royal Society B: Biological Sciences*, 279(1732), 1396–1404.
- Derraik, J. G. B. (2002). The pollution of the marine environment by plastic debris: A review. *Marine Pollution Bulletin*, 44, 842–852.
- DeRuiter, S. L., I. L. Boyd, D. E. Claridge, C. W. Clark, C. Gagon, B. L. Southall, and P. L. Tyack. (2013a). Delphinid whistle production and call matching during playback of simulated military sonar. *Marine Mammal Science*, 29(2), E46–59.
- DeRuiter, S. L., B. L. Southall, J. Calambokidis, W. M. Zimmer, D. Sadykova, E. A. Falcone, A. S. Friedlaender, J. E. Joseph, D. Moretti, G. S. Schorr, L. Thomas, and P. L. Tyack. (2013b). First direct measurements of behavioural responses by Cuvier's beaked whales to mid-frequency active sonar. *Biology Letters*, 9(4), 20130223.
- DeRuiter, S. L., R. Langrock, T. Skirbutas, J. A. Goldbogen, J. Calambokidis, A. S. Friedlaender, and B. L. Southall. (2017). A multivariate mixed hidden Markov model for blue whale behaviour and responses to sound exposure. *The Annals of Applied Statistics*, 11(1), 362–392.
- Desforges, J. P., C. Sonne, M. Levin, U. Siebert, S. De Guise, and R. Dietz. (2016). Immunotoxic effects of environmental pollutants in marine mammals. *Environment International*, 86, 126–139.
- Di Clemente, J., F. Christiansen, E. Pirota, D. Steckler, M. Wahlberg, and H. C. Pearson. (2018). Effects of whale watching on the activity budgets of humpback whales, *Megaptera novaeangliae* (Borowski, 1781), on a feeding ground. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28(4), 810–820.
- Di Lorenzo, E., K. M. Cobb, J. C. Furtado, N. Schneider, B. T. Anderson, A. Bracco, M. A. Alexander, and D. J. Vimont. (2010). Central Pacific El Niño and decadal climate change in the North Pacific Ocean. *Nature Geoscience*, 3(11), 762–765.
- Di Lorio, L., and C. W. Clark. (2010). Exposure to seismic survey alters blue whale acoustic communication. *Biology Letters*, 6, 51–54.
- Dolar, M. L. L. (2009). Fraser's dolphin, *Lagenodelphis hosei*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 485–487). Cambridge, MA: Academic Press.
- Donaldson, T. J. (1983). Further investigations of the whales *Peponocephala electra* and *Globicephala macrorhynchus* reported from Guam. *Micronesica*, 19(1-2), 173–181.
- Doney, S. C., M. Ruckelshaus, D. J. Emmett, J. P. Barry, F. Chan, C. A. English, H. M. Galindo, J. M. Grebmeier, A. B. Hollowed, N. Knowlton, J. Polovina, N. N. Rabalais, W. J. Sydeman, and L. D. Talley. (2012). Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4(1), 11–37.
- Doyle, L. R., B. McCowan, S. F. Hanser, C. Chyba, T. Bucci, and E. J. Blue. (2008). Applicability of information theory to the quantification of responses to anthropogenic noise by southeast Alaskan humpback whales. *Entropy*, 10, 33–46.
- Dunlop, R. A., D. H. Cato, and M. J. Noad. (2010). Your attention please: Increasing ambient noise levels elicits a change in communication behaviour in humpback whales (*Megoptera novaeangliae*). *Proceedings of the Royal Society B: Biological Sciences*, 277, 2521–2529.

- Dunlop, R. A., M. J. Noad, D. H. Cato, E. Kniest, P. J. Miller, J. N. Smith, and M. D. Stokes. (2013). Multivariate analysis of behavioural response experiments in humpback whales (*Megaptera novaeangliae*). *The Journal of Experimental Biology*, 216(5), 759–770.
- Dunlop, R. A., D. H. Cato, and M. J. Noad. (2014). Evidence of a Lombard response in migrating humpback whales (*Megaptera novaeangliae*). *The Journal of the Acoustical Society of America*, 136(1), 430–437.
- Dunlop, R. A., M. J. Noad, R. D. McCauley, E. Kniest, D. Paton, and D. H. Cato. (2015). The behavioural response of humpback whales (*Megaptera novaeangliae*) to a 20 cubic inch air gun. *Aquatic Mammals*, 41(4), 412.
- Dunlop, R. A. (2016). The effect of vessel noise on humpback whale, *Megaptera novaeangliae*, communication behaviour. *Animal Behaviour*, 111, 13–21.
- Dunlop, R. A., M. J. Noad, R. D. McCauley, E. Kniest, R. Slade, D. Paton, and D. H. Cato. (2016). Response of humpback whales (*Megaptera novaeangliae*) to ramp-up of a small experimental air gun array. *Marine Pollution Bulletin*, 103(1–2), 72–83.
- Dunlop, R. A., M. J. Noad, R. D. McCauley, L. Scott-Hayward, E. Kniest, R. Slade, D. Paton, and D. H. Cato. (2017). Determining the behavioural dose-response relationship of marine mammals to air gun noise and source proximity. *The Journal of Experimental Biology*, 220(16), 2878–2886.
- Dunn, C. A., D. E. Claridge, and T. L. Pusser. (2007). Killer whale (*Orcinus orca*) occurrence and predation in the Bahamas. *Journal of the Marine Biological Association of the United Kingdom* 94(06), 1305–1309.
- Dunphy-Daly, M. M., M. R. Heithaus, and D. E. Claridge. (2008). Temporal variation in dwarf sperm whale (*Kogia sima*) habitat use and group size off Great Abaco Island, Bahamas. *Marine Mammal Science*, 24(1), 171–182.
- Durban, J. W., H. Fearnbach, L. G. Barrett-Lennard, W. L. Perryman, and D. J. Leroi. (2015). Photogrammetry of killer whales using a small hexacopter launched at sea. *Journal of Unmanned Vehicle Systems*, 3(3), 131–135.
- Dyndo, M., D. M. Wisniewska, L. Rojano-Donate, and P. T. Madsen. (2015). Harbour porpoises react to low levels of high frequency vessel noise. *Scientific Reports*, 5, 11083.
- Edds-Walton, P. L. (1997). Acoustic communication signals of mysticete whales. *Bioacoustics*, 8, 47–60.
- Edwards, H. H. (2013). Potential impacts of climate change on warmwater megafauna: The Florida manatee example (*Trichechus manatus latirostris*). *Climatic Change*, 121(4), 727–738.
- Edwards, M., and J. Beldowski. (2016). Chemical munitions dumped at sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 1–3.
- Edwards, M. H., D. J. Fornari, M. R. Rognstad, C. D. Kelley, C. L. Mah, L. K. Davis, K. R. M. Flores, E. L. Main, and N. L. Bruso. (2016a). Time-lapse camera studies of sea-disposed chemical munitions in Hawaii. *Deep-Sea Research II: Topical Studies in Oceanography*, 128, 25–33.
- Edwards, M. H., S. M. Shjegstad, R. Wilkens, J. C. King, G. Carton, D. Bala, B. Bingham, M. C. Bissonnette, C. Briggs, N. S. Bruso, R. Camilli, M. Cremer, R. B. Davis, E. H. DeCarlo, C. DuVal, D. J. Fornari, I. Kaneakua-Pia, C. D. Kelley, S. Koide, C. L. Mah, T. Kerby, G. J. Kurras, M. R. Rognstad, L. Sheild, J. Silva, B. Wellington, and M. V. Woerkom. (2016b). The Hawaii undersea military munitions assessment. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 4–13.

- Efroymson, R. A., W. H. Rose, and G. W. Suter, II. (2001). *Ecological Risk Assessment Framework for Low-altitude Overflights by Fixed-Wing and Rotary-Wing Military Aircraft*. Oak Ridge, TN: Oak Ridge National Laboratory.
- Eisenhardt, E. (2014). *Recent Trends of Vessel Activities in Proximity to Cetaceans in the Central Salish Sea*. Paper presented at the Salish Sea Ecosystem Conference. Seattle, WA.
- Eldredge, L. G. (1991). Annotated checklist of the marine mammals of Micronesia. *Micronesica*, 24(2), 217–230.
- Eldredge, L. G. (2003). A retrospective look at Guam's marine biodiversity. *Micronesica*, 35-36, 26–37.
- Ellison, W. T., B. L. Southall, C. W. Clark, and A. S. Frankel. (2011). A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology*, 26(1), 21–28.
- Engelhardt, R. (1983). Petroleum effects on marine mammals. *Aquatic Toxicology*, 4, 199–217.
- Erbe, C. (2002). Underwater noise of whale-watching boats and potential effects on killer whales (*Orcinus orca*), based on an acoustic impact model. *Marine Mammal Science*, 18(2), 394–418.
- Erbe, C., A. MacGillivray, and R. Williams. (2012). Mapping cumulative noise from shipping to inform marine spatial planning. *The Journal of the Acoustical Society of America*, 132(5), EL423–EL428.
- Erbe, C., R. Williams, D. Sandilands, and E. Ashe. (2014). Identifying modeled ship noise hotspots for marine mammals of Canada's Pacific region. *PLoS ONE*, 9(3), e89820.
- Erbe, C., C. Reichmuth, K. Cunningham, K. Lucke, and R. Dooling. (2015). Communication masking in marine mammals: a review and research strategy. *Marine Pollution Bulletin*, 1–24.
- Erbe, C., C. Reichmuth, K. Cunningham, K. Lucke, and R. Dooling. (2016). Communication masking in marine mammals: A review and research strategy. *Marine Pollution Bulletin*, 103(1–2), 15–38.
- Evans, P. G. H., and L. A. Miller. (2003). *Proceedings of the workshop on active sonar and cetaceans* (European Cetacean Society newsletter, No. 42—Special Issue). Las Palmas, Gran Canaria: European Cetacean Society.
- Fahlman, A., A. Olszowka, B. Bostrom, and D. R. Jones. (2006). Deep diving mammals: Dive behavior and circulatory adjustments contribute to bends avoidance. *Respiratory Physiology and Neurobiology*, 153, 66–77.
- Fahlman, A., S. K. Hooker, A. Olszowka, B. L. Bostrom, and D. R. Jones. (2009). Estimating the effect of lung collapse and pulmonary shunt on gas exchange during breath-hold diving: The Scholander and Kooyman legacy. *Respiratory Physiology & Neurobiology*, 165(1), 28–39.
- Fahlman, A., S. H. Loring, S. P. Johnson, M. Haulena, A. W. Trites, V. A. Fravel, and W. G. Van Bonn. (2014a). Inflation and deflation pressure-volume loops in anesthetized pinnipeds confirms compliant chest and lungs. *Frontiers in Physiology*, 5(433), 1–7.
- Fahlman, A., P. L. Tyack, P. J. O. Miller, and P. H. Kvasdheim. (2014b). How man-made interference might cause gas bubble emboli in deep diving whales. *Frontiers in Physiology*, 5(13), 1–6.
- Fair, P. A., J. Adams, G. Mitchum, T. C. Hulse, J. S. Reif, M. Houde, D. Muir, E. Wirth, D. Wetzel, E. Zolman, W. McFee, and G. D. Bossart. (2010). Contaminant blubber burdens in Atlantic bottlenose dolphins (*Tursiops truncatus*) from two southeastern U.S. estuarine areas: Concentrations and patterns of PCBs, pesticides, PBDEs, PFCs, and PAHs. *The Science of the Total Environment*, 408(7), 1577–1597.

- Fair, P. A., A. M. Schaefer, T. A. Romano, G. D. Bossart, S. V. Lamb, and J. S. Reif. (2014). Stress response of wild bottlenose dolphins (*Tursiops truncatus*) during capture-release health assessment studies. *General and Comparative Endocrinology*, 206, 203–212.
- Falcone, E. A., G. S. Schorr, A. B. Douglas, J. Calambokidis, E. Henderson, M. F. McKenna, J. Hildebrand, and D. Moretti. (2009). Sighting characteristics and photo-identification of Cuvier's beaked whales (*Ziphius cavirostris*) near San Clemente Island, California: A key area for beaked whales and the military? *Marine Biology*, 156, 2631–2640.
- Falcone, E. A., and G. S. Schorr. (2012). *Distribution and Demographics of Marine Mammals in SOCAL Through Photo-Identification, Genetics, and Satellite Telemetry: A Summary of Surveys Conducted 1 July 2011 – 15 June 2012*. Monterey, CA: Naval Postgraduate School.
- Falcone, E. A., and G. S. Schorr. (2014). *Distribution and Demographics of Marine Mammals in SOCAL through Photo-Identification, Genetics, and Satellite Telemetry* (Prepared for Chief of Naval Operations Energy and Environmental Readiness Division: NPS-OC-14-005CR). Monterey, CA: Naval Postgraduate School.
- Falcone, E. A., G. S. Schorr, S. L. Watwood, S. L. DeRuiter, A. N. Zerbini, R. D. Andrews, R. P. Morrissey, and D. J. Moretti. (2017). Diving behaviour of Cuvier's beaked whales exposed to two types of military sonar. *Royal Society Open Science*, 4(170629), 1–21.
- Falke, K. J., R. D. Hill, J. Qvist, R. C. Schneider, M. Guppy, G. C. Liggins, P. W. Hochachka, R. E. Elliott, and W. M. Zapol. (1985). Seal lungs collapse during free diving: Evidence from arterial nitrogen tensions. *Science*, 229, 556–558.
- Farak, A. M., M. W. Richie, J. A. Rivers, and R. K. Uyeyama. (2011). *Cruise Report, Marine Species Monitoring and Lookout Effectiveness Study Koa Kai, November 2010, Hawaii Range Complex*. Washington, DC: Commander, U.S. Pacific Fleet.
- Farmer, N. A., D. P. Noren, E. M. Fougères, A. Machernis, and K. Baker. (2018). Resilience of the endangered sperm whale *Physeter macrocephalus* to foraging disturbance in the Gulf of Mexico, USA: A bioenergetic approach. *Marine Ecology Progress Series*, 589, 241–261.
- Fauquier, D. A., M. J. Kinsel, M. D. Dailey, G. E. Sutton, M. K. Stolen, R. S. Wells, and F. M. D. Gulland. (2009). Prevalence and pathology of lungworm infection in bottlenose dolphins, *Tursiops truncatus*, from southwest Florida. *Diseases of Aquatic Organisms*, 88, 85–90.
- Fay, R. R. (1988). *Hearing in Vertebrates: A Psychophysics Databook*. Winnetka, IL: Hill-Fay Associates.
- Fay, R. R., and A. N. Popper. (1994). *Comparative Hearing: Mammals*. New York, NY: Springer-Verlag.
- Ferguson, M. C., J. Barlow, P. Feidler, S. B. Reilly, and T. Gerrodette. (2006a). Spatial models of delphinid (family Delphinidae) encounter rate and group size in the eastern tropical Pacific Ocean. *Ecological Modelling*, 193, 645–662.
- Ferguson, M. C., J. Barlow, S. B. Reilly, and T. Gerrodette. (2006b). Predicting Cuvier's (*Ziphius cavirostris*) and Mesoplodon beaked whale population density from habitat characteristics in the eastern tropical Pacific Ocean. *Journal of Cetacean Research and Management*, 7(3), 287–289.
- Ferguson, S. H., J. W. Higdon, and K. H. Westdal. (2012). Prey items and predation behavior of killer whales (*Orcinus orca*) in Nunavut, Canada based on Inuit hunter interviews. *Aquatic Biosystems*, 8(3).
- Fernandez, A., J. Edwards, F. Rodriguez, A. Espinosa De Los Monteros, P. Herraiez, P. Castro, J. Jaber, V. Martin, and M. Arbelo. (2005). "Gas and fat embolic syndrome" involving a mass stranding of

- beaked whales (Family *Ziphiidae*) exposed to anthropogenic sonar signals. *Veterinary Pathology*, 42(4), 446–457.
- Fernandez, A. (2006). *Beaked Whale (Ziphius cavirostris) Mass Stranding on Almeria's Coasts in Southern Spain*. Las Palmas, Canary Islands: University of Las Palmas de Gran Canaria.
- Fernandez, A., E. Sierra, J. Diaz-Delgado, S. Sacchini, Y. Sanchez-Paz, C. Suarez-Santana, M. Arregui, M. Arbelo, and Y. Bernaldo de Quiros. (2017). Deadly acute decompression sickness in Risso's dolphins. *Scientific Reports*, 7(1), 13621.
- Fertl, D., A. Acevedo-Gutiérrez, and F. L. Darby. (1996). A report of killer whales (*Orcinus orca*) feeding on a carcharhinid shark in Costa Rica. *Marine Mammal Science*, 12(4), 606–611.
- Filadelfo, R., J. Mintz, E. Michlovich, A. D'Amico, and D. R. Ketten. (2009a). Correlating military sonar use with beaked whale mass strandings: What do the historical data show? *Aquatic Mammals*, 35(4), 435–444.
- Filadelfo, R., Y. K. Pinelis, S. Davis, R. Chase, J. Mintz, J. Wolfanger, P. L. Tyack, D. R. Ketten, and A. D'Amico. (2009b). Correlating whale strandings with Navy exercises off Southern California. *Aquatic Mammals*, 35(4), 445–451.
- Finneran, J. J., C. E. Schlundt, D. A. Carder, J. A. Clark, J. A. Young, J. B. Gaspin, and S. H. Ridgway. (2000). Auditory and behavioral responses of bottlenose dolphins (*Tursiops truncatus*) and a beluga whale (*Delphinapterus leucas*) to impulsive sounds resembling distant signatures of underwater explosions. *The Journal of the Acoustical Society of America*, 108(1), 417–431.
- Finneran, J. J., D. A. Carder, and S. H. Ridgway. (2001). Temporary threshold shift (TTS) in bottlenose dolphins (*Tursiops truncatus*) exposed to tonal signals. *The Journal of the Acoustical Society of America*, 110(5), 2749(A).
- Finneran, J. J., C. E. Schlundt, R. Dear, D. A. Carder, and S. H. Ridgway. (2002). Temporary shift in masked hearing thresholds in odontocetes after exposure to single underwater impulses from a seismic watergun. *The Journal of the Acoustical Society of America*, 111(6), 2929–2940.
- Finneran, J. J., D. A. Carder, R. Dear, T. Belting, and S. H. Ridgway. (2003a). Pure-tone audiograms and hearing loss in the white whale (*Delphinapterus leucas*). *The Journal of the Acoustical Society of America*, 114, 2434(A).
- Finneran, J. J., R. Dear, D. A. Carder, and S. H. Ridgway. (2003b). Auditory and behavioral responses of California sea lions (*Zalophus californianus*) to single underwater impulses from an arc-gap transducer. *The Journal of the Acoustical Society of America*, 114(3), 1667–1677.
- Finneran, J. J., and C. E. Schlundt. (2004). *Effects of Intense Pure Tones on the Behavior of Trained Odontocetes*. San Diego, CA: Space and Naval Warfare Systems Center Pacific.
- Finneran, J. J., D. A. Carder, R. Dear, T. Belting, J. McBain, L. Dalton, and S. H. Ridgway. (2005a). Pure tone audiograms and possible aminoglycoside-induced hearing loss in belugas (*Delphinapterus leucas*). *The Journal of the Acoustical Society of America*, 117, 3936–3943.
- Finneran, J. J., D. A. Carder, C. E. Schlundt, and S. H. Ridgway. (2005b). Temporary threshold shift (TTS) in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. *The Journal of the Acoustical Society of America*, 118(4), 2696–2705.
- Finneran, J. J., and D. S. Houser. (2006). Comparison of in-air evoked potential and underwater behavioral hearing thresholds in four bottlenose dolphins (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 119(5), 3181–3192.

- Finneran, J. J., C. E. Schlundt, B. Branstetter, and R. L. Dear. (2007). Assessing temporary threshold shift in a bottlenose dolphin (*Tursiops truncatus*) using multiple simultaneous auditory evoked potentials. *The Journal of the Acoustical Society of America*, 122(2), 1249–1264.
- Finneran, J. J., D. S. Houser, B. Mase-Guthrie, R. Y. Ewing, and R. G. Lingenfelter. (2009). Auditory evoked potentials in a stranded Gervais' beaked whale (*Mesoplodon europaeus*). *The Journal of the Acoustical Society of America*, 126(1), 484–490.
- Finneran, J. J., D. A. Carder, C. E. Schlundt, and R. L. Dear. (2010a). Temporary threshold shift in a bottlenose dolphin (*Tursiops truncatus*) exposed to intermittent tones. *The Journal of the Acoustical Society of America*, 127(5), 3267–3272.
- Finneran, J. J., D. A. Carder, C. E. Schlundt, and R. L. Dear. (2010b). Growth and recovery of temporary threshold shift at 3 kHz in bottlenose dolphins: Experimental data and mathematical models. *The Journal of the Acoustical Society of America*, 127(5), 3256–3266.
- Finneran, J. J., and B. K. Branstetter. (2013). Effects of Noise on Sound Perception in Marine Mammals *Animal Communication and Noise* (Vol. 2, pp. 273–308). Berlin, Germany: Springer Berlin Heidelberg.
- Finneran, J. J., and C. E. Schlundt. (2013). Effects of fatiguing tone frequency on temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 133(3), 1819–1826.
- Finneran, J. J. (2015). Noise-induced hearing loss in marine mammals: A review of temporary threshold shift studies from 1996 to 2015. *The Journal of the Acoustical Society of America*, 138(3), 1702–1726.
- Finneran, J. J., C. E. Schlundt, B. K. Branstetter, J. S. Trickey, V. Bowman, and K. Jenkins. (2015). Effects of multiple impulses from a seismic air gun on bottlenose dolphin hearing and behavior. *The Journal of the Acoustical Society of America*, 137(4), 1634–1646.
- Finneran, J. J., J. Mulsow, D. S. Houser, and R. F. Burkard. (2016). Place specificity of the click-evoked auditory brainstem response in the bottlenose dolphin (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 140(4), 2593–2602.
- Finneran, J. J. (2018). Conditioned attenuation of auditory brainstem responses in dolphins warned of an intense noise exposure: Temporal and spectral patterns. *The Journal of the Acoustical Society of America*, 143(2), 795.
- Fish, J. F., and J. S. Vania. (1971). Killer whale, *Orcinus orca*, sounds repel white whales, *Delphinapterus leucas*. *Fishery Bulletin*, 69(3), 531–535.
- Fitch, R., J. Harrison, and J. Lewandowski. (2011). *Marine Mammal and Sound Workshop July 13th and 14th, 2010: Report to the National Ocean Council Ocean Science and Technology Interagency Policy Committee*. Washington, DC: Bureau of Ocean Energy Management, Department of the Navy, National Oceanic and Atmospheric Administration.
- Fleming, A. H., C. T. Clark, J. Calambokidis, and J. Barlow. (2016). Humpback whale diets respond to variance in ocean climate and ecosystem conditions in the California Current. *Global Change Biology*, 22(3), 1214–1224.
- Foltz, K. M., R. W. Baird, G. M. Ylitalo, and B. A. Jensen. (2014). Cytochrome P4501A1 expression in blubber biopsies of endangered false killer whales (*Pseudorca crassidens*) and nine other odontocete species from Hawaii. *Ecotoxicology*, 23(9), 1607–1618.

- Foote, A. D., R. W. Osborne, and A. R. Hoelzel. (2004). Whale-call response to masking boat noise. *Nature*, 428, 910.
- Ford, J. K. B. (2008). Killer whale, *Orcinus orca*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 650–657). San Diego, CA: Academic Press.
- Ford, J. K. B., E. H. Stredulinsky, J. R. Towers, and G. M. Ellis. (2013). *Information in Support of the Identification of Critical Habitat for Transient Killer Whales (Orcinus orca) off the West Coast of Canada DFO* (Science Advisory Report 2013/025). Nanaimo, Canada: Canadian Science Advisory Secretariat.
- Ford, J. K. B., E. H. Stredulinsky, G. M. Ellis, J. W. Durban, and J. F. Pilkington. (2014). *Offshore Killer Whales in Canadian Pacific Waters: Distribution, Seasonality, Foraging Ecology, Population Status and Potential for Recovery*. (Document 2014/088). Ottawa, Canada: Department of Fisheries and Oceans Canada, Canadian Science Advisory, Secretariat.
- Forney, K. A., and P. R. Wade. (2006). Worldwide Distribution and Abundance of Killer Whales. In J. A. Estes, R. L. Brownell, Jr., D. P. DeMaster, D. F. Doak, & T. M. Williams (Eds.), *Whales, Whaling and Ocean Ecosystems* (pp. 145–162). Berkeley, CA: University of California Press.
- Forney, K. A., B. L. Southall, E. Slooten, S. Dawson, A. J. Read, R. W. Baird, and R. L. Brownell, Jr. (2017). Nowhere to go: Noise impact assessments for marine mammal populations with high site fidelity. *Endangered Species Research*, 32, 391–413.
- Frankel, A. S., and C. W. Clark. (2000). Behavioral responses of humpback whales (*Megaptera novaeangliae*) to full-scale ATOC signals. *The Journal of the Acoustical Society of America*, 108(4), 1930–1937.
- Friedlaender, A. S., E. L. Hazen, J. A. Goldbogen, A. K. Stimpert, J. Calambokidis, and B. L. Southall. (2016). Prey-mediated behavioral responses of feeding blue whales in controlled sound exposure experiments. *Ecological Applications*, 26(4), 1075–1085.
- Frisk, G. V. (2012). Noiseconomics: The relationship between ambient noise levels in the sea and global economic trends. *Scientific Reports*, 2(437), 1–4.
- Fristrup, K. M., L. T. Hatch, and C. W. Clark. (2003). Variation in humpback whale (*Megaptera novaeangliae*) song length in relation to low-frequency sound broadcasts. *The Journal of the Acoustical Society of America*, 113(6), 3411–3424.
- Fromm, D. M., J. R. Mobley, S. W. Martin, and P. E. Nachtigall. (2006). Analysis of melon-headed whale aggregation in Hanalei Bay, July 2004. *Animal Bioacoustics: Marine Mammal Acoustics II*, 120(5), 3266.
- Fromm, D. M. (2009). *Reconstruction of Acoustic Exposure on Orcas in Haro Strait (Acoustics)*. Washington, DC: U.S. Naval Research Laboratory.
- Fulling, G. L., P. H. Thorson, and J. Rivers. (2011). Distribution and Abundance Estimates for Cetaceans in the Waters off Guam and the Commonwealth of the Northern Mariana Islands. *Pacific Science*, 65(3), 321–343.
- Fulling, G. L., T. A. Jefferson, D. Fertl, J. C. S. Vega, C. S. Oedekoven, and S. A. Kuczaj, II. (2017). Sperm Whale (*Physeter macrocephalus*) Collision with a Research Vessel: Accidental Collision or Deliberate Ramming? *Aquatic Mammals*, 43(4), 421–429.

- Gailey, G., B. Wursig, and T. L. McDonald. (2007). Abundance, behavior, and movement patterns of western gray whales in relation to a 3-D seismic survey, Northeast Sakhalin Island, Russia. *Environmental Monitoring and Assessment*, 134, 75–91.
- Gailey, G., O. Sychenko, T. McDonald, R. Racca, A. Rutenko, and K. Bröker. (2016). Behavioural responses of western gray whales to a 4-D seismic survey off northeastern Sakhalin Island, Russia. *Endangered Species Research*, 30, 53–71.
- Gannier, A. (2002). Cetaceans of the Marquesas Islands (French Polynesia): Distribution and relative abundance as obtained from a small boat dedicated survey. *Aquatic Mammals*, 28(2), 198–210.
- Gannier, A., and K. L. West. (2005). Distribution of the rough-toothed dolphin (*Steno bredanensis*) around the Windward Islands (French Polynesia). *Pacific Science*, 59, 17–24.
- Gannon, D. P., N. B. Barros, D. P. Nowacek, A. J. Read, D. M. Waples, and R. S. Wells. (2005). Prey detection by bottlenose dolphins, *Tursiops truncatus*: An experimental test of the passive listening hypothesis. *Animal Behaviour*, 69, 709–720.
- Garcia Parraga, D., M. Moore, and A. Fahlman. (2018). Pulmonary ventilation-perfusion mismatch: A novel hypothesis for how diving vertebrates may avoid the bends. *Proceedings of the Royal Society B: Biological Sciences*, 285(1877).
- Garibaldi, F., and M. Podesta. (2014). Stomach contents of a sperm whale (*Physeter macrocephalus*) stranded in Italy (Ligurian Sea, north-western Mediterranean). *Journal of the Marine Biological Association of the United Kingdom*, 94(6), 1087–1091.
- Garrigue, C., M. Oremus, R. Dodémont, P. Bustamante, O. Kwiatek, G. Libeau, C. Lockyer, J. C. Vivier, and M. L. Dalebout. (2016). A mass stranding of seven Longman's beaked whales (*Indopacetus pacificus*) in New Caledonia, South Pacific. *Marine Mammal Science*, 32(3), 884–910.
- Gaspard, J. C., G. B. Bauer, R. L. Reep, K. Dziuk, A. Cardwell, L. Read, and D. A. Mann. (2012). Audiogram and auditory critical ratios of two Florida manatees (*Trichechus manatus latirostris*). *The Journal of Experimental Biology*, 215(Pt 9), 1442–1447.
- Gedamke, J., M. Ferguson, J. Harrison, L. Hatch, L. Henderson, M. B. Porter, B. L. Southall, and S. Van Parijs. (2016). Predicting Anthropogenic Noise Contributions to U.S. Waters. *Advances in Experimental Medicine and Biology*, 875, 341–347.
- Geijer, C. K. A., and A. J. Read. (2013). Mitigation of marine mammal bycatch in U.S. fisheries since 1994. *Biological Conservation*, 159, 54–60.
- Gende, S. M., A. N. Hendrix, K. R. Harris, B. Eichenlaub, J. Nielsen, and S. Pyare. (2011). A Bayesian approach for understanding the role of ship speed in whale-ship encounters. *Ecological Applications*, 21(6), 2232–2240.
- Geraci, J., J. Harwood, and V. Lounsbury. (1999). Marine Mammal Die-Offs: Causes, Investigations, and Issues. In J. Twiss & R. Reeves (Eds.), *Conservation and Management of Marine Mammals* (pp. 367–395). Washington, DC: Smithsonian Institution Press.
- Geraci, J., and V. Lounsbury. (2005). *Marine Mammals Ashore: A Field Guide for Strandings* (Second ed.). Baltimore, MD: National Aquarium in Baltimore.
- Gervaise, C., Y. Simard, N. Roy, B. Kinda, and N. Menard. (2012). Shipping noise in whale habitat: Characteristics, sources, budget, and impact on belugas in Saguenay–St. Lawrence Marine Park hub. *The Journal of the Acoustical Society of America*, 132(1), 76–89.

- Ghoul, A., and C. Reichmuth. (2014). Hearing in the sea otter (*Enhydra lutris*): Auditory profiles for an amphibious marine carnivore. *Journal of Comparative Physiology A: Neuroethology, Sensory Neural, and Behavioral Physiology*, 200(11), 967–981.
- Gill, A. B., I. Gloyne-Philips, J. Kimber, and P. Sigray. (2014). Marine Renewable Energy, Electromagnetic (EM) Fields and EM-Sensitive Animals. In M. Shields & A. Payne (Eds.), *Marine Renewable Energy Technology and Environmental Interactions. Humanity and the Sea* (pp. 61–79). Dordrecht, Netherlands: Springer.
- Giorli, G., and W. W. L. Au. (2017). Spatio-temporal variation and seasonality of Odontocetes' foraging activity in the leeward side of the island of Hawaii. *Deep-Sea Research I*, 121, 202–209.
- Godard-Coding, C. A. J., R. Clark, M. C. Fossi, L. Marsili, S. Maltese, A. G. West, L. Valenzuela, V. Rowntree, I. Polyak, J. C. Cannon, K. Pinkerton, N. Rubio-Cisneros, S. L. Mesnick, S. B. Cox, I. Kerr, R. Payne, and J. J. Stegeman. (2011). Pacific Ocean–Wide Profile of CYP1A1 Expression, Stable Carbon and Nitrogen Isotope Ratios, and Organic Contaminant Burden in Sperm Whale Skin Biopsies. *Environmental Health Perspectives*, 119(3), 337–343.
- Goertner, J. F. (1982). *Prediction of Underwater Explosion Safe Ranges for Sea Mammals*. Dahlgren, VA: Naval Surface Weapons Center.
- Goldbogen, J. A., J. Calambokidis, R. E. Shadwick, E. M. Oleson, M. A. McDonald, and J. A. Hildebrand. (2006). Kinematics of foraging dives and lunge-feeding in fin whales. *The Journal of Experimental Biology*, 209(Pt 7), 1231–1244.
- Goldbogen, J. A., A. S. Friedlaender, J. Calambokidis, M. F. McKenna, M. Simon, and D. Nowacek. (2013a). Integrative Approaches to the Study of Baleen Whale Diving Behavior, Feeding Performance, and Foraging Ecology. *BioScience*, 63(2), 90–100.
- Goldbogen, J. A., B. L. Southall, S. L. DeRuiter, J. Calambokidis, A. S. Friedlaender, E. L. Hazen, E. A. Falcone, G. S. Schorr, A. Douglas, D. J. Moretti, C. Kyburg, M. F. McKenna, and P. L. Tyack. (2013b). Blue whales respond to simulated mid-frequency military sonar. *Proceedings of the Royal Society B: Biological Sciences*, 280(1765), 20130657.
- Gong, Z., A. D. Jain, D. Tran, D. H. Yi, F. Wu, A. Zorn, P. Ratilal, and N. C. Makris. (2014). Ecosystem scale acoustic sensing reveals humpback whale behavior synchronous with herring spawning processes and re-evaluation finds no effect of sonar on humpback song occurrence in the Gulf of Maine in fall 2006. *PLoS ONE*, 9(10), e104733.
- Gordon, J., D. Gillespie, J. Potter, A. Frantzis, M. P. Simmonds, R. Swift, and D. Thompson. (2003). A review of the effects of seismic surveys on marine mammals. *Marine Technology Society Journal*, 37(4), 16–34.
- Gospić, N. R., and M. Picciulin. (2016). Changes in whistle structure of resident bottlenose dolphins in relation to underwater noise and boat traffic. *Marine Pollution Bulletin*, 105, 193–198.
- Graham, I. M., E. Pirotta, N. D. Merchant, A. Farcas, T. R. Barton, B. Cheney, G. D. Hastie, and P. M. Thompson. (2017). Responses of bottlenose dolphins and harbor porpoises to impact and vibration piling noise during harbor construction. *Ecosphere*, 8(5), 1–16.
- Greaves, F. C., R. H. Draeger, O. A. Brines, J. S. Shaver, and E. L. Corey. (1943). An experimental study of concussion. *United States Naval Medical Bulletin*, 41(1), 339–352.
- Green, D. M. (1994). Sound's effects on marine mammals need investigation. *Eos*, 75(27), 305–306.

- Green, D. M., H. DeFerrari, D. McFadden, J. Pearse, A. Popper, W. J. Richardson, S. H. Ridgway, and P. Tyack. (1994). *Low-Frequency Sound and Marine Mammals: Current Knowledge and Research Needs*. Washington, DC: Ocean Studies Board, Commission on Geosciences, Environment, and Resources, National Research Council.
- Green, G. A., J. J. Brueggeman, R. A. Grotefendt, C. E. Bowlby, M. L. Bonnell, and K. C. Balcomb, III. (1992). *Cetacean Distribution and Abundance off Oregon and Washington, 1989–1990*. Los Angeles, CA: U.S. Department of the Interior, Minerals Management Service.
- Gregory, P. R., and A. A. Rowden. (2001). Behaviour patterns of bottlenose dolphins (*Tursiops truncatus*) relative to tidal state, time-of-day, and boat traffic in Cardigan Bay, West Wales. *Aquatic Mammals*, 27.2, 105–114.
- Guan, S., B. L. Southall, J. F. Vignola, J. A. Judge, and D. Turo. (2017). Sonar inter-ping noise field characterization during cetacean behavioral response studies off Southern California. *Acoustical Physics*, 63(2), 204–215.
- Guerra, M., S. M. Dawson, T. E. Brough, and W. J. Rayment. (2014). Effects of boats on the surface and acoustic behaviour of an endangered population of bottlenose dolphins. *Endangered Species Research*, 24(3), 221–236.
- Haelters, J., V. Dulière, L. Vigin, and S. Degraer. (2014). Towards a numerical model to simulate the observed displacement of harbour porpoises *Phocoena phocoena* due to pile driving in Belgian waters. *Hydrobiologia*, 756(1), 105–116.
- Hall, A., K. Hugunin, R. Deaville, R. Law, C. R. Allchin, and P. Jepson. (2006). The Risk of Infection from Polychlorinated Biphenyl Exposure in the Harbor Porpoise (*Phocoena phocoena*): A Case-Control Approach. *Environmental Health Perspectives*, 114, 704–711.
- Hamer, D. J., S. J. Childerhouse, and N. J. Gales. (2010). *Mitigating Operational Interactions Between Odontocetes and the Longline Fishing Industry: A Preliminary Global Review of the Problem and of Potential Solutions*. Tasmania, Australia: International Whaling Commission.
- Hamilton, T. A., J. V. Redfern, J. Barlow, L. T. Ballance, T. Gerrodette, R. S. Holt, K. A. Forney, and B. L. Taylor. (2009). *Atlas of Cetacean Sightings for Southwest Fisheries Science Center Cetacean and Ecosystem Surveys: 1986–2005* (NOAA Technical Memorandum NMFS-SWFSC-440). La Jolla, CA: Southwest Fisheries Science Center.
- Hance, A. J., E. D. Robin, J. B. Halter, N. Lewiston, D. A. Robin, L. Cornell, M. Caligiuri, and J. Theodore. (1982). Hormonal changes and enforced diving in the harbor seal *Phoca vitulina* II. Plasma catecholamines. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, 242(5), R528–R532.
- Hansen, A. M. K., C. E. Bryan, K. West, and B. A. Jensen. (2015). Trace Element Concentrations in Liver of 16 Species of Cetaceans Stranded on Pacific Islands from 1997 through 2013. *Archives of Environmental Contamination and Toxicology*, 70(1), 75–95.
- Harcourt, R., V. Pirodda, G. Heller, V. Peddemors, and D. Slip. (2014). A whale alarm fails to deter migrating humpback whales: An empirical test. *Endangered Species Research*, 25(1), 35–42.
- Hardesty, B. D., and C. Wilcox. (2017). A risk framework for tackling marine debris. *Royal Society of Chemistry*, 9, 1429–1436.

- Harris, C., and L. Thomas. (2015). *Status and Future of Research on the Behavioral Responses of Marine Mammals to U.S. Navy Sonar* (Centre for Research into Ecological & Environmental Modelling Technical Report 2015-3). St. Andrews, United Kingdom: University of St. Andrews.
- Harris, C. M., D. Sadykova, S. L. DeRuiter, P. L. Tyack, P. J. O. Miller, P. H. Kvasdheim, F. P. A. Lam, and L. Thomas. (2015). Dose response severity functions for acoustic disturbance in cetaceans using recurrent event survival analysis. *Ecosphere*, 6(11), art236.
- Harris, C. M., L. Thomas, E. A. Falcone, J. Hildebrand, D. Houser, P. H. Kvasdheim, F.-P. A. Lam, P. J. O. Miller, D. J. Moretti, A. J. Read, H. Slabbekoorn, B. L. Southall, P. L. Tyack, D. Wartzok, V. M. Janik, and J. Blanchard. (2018). Marine mammals and sonar: Dose-response studies, the risk-disturbance hypothesis and the role of exposure context. *Journal of Applied Ecology*, 55(1), 396–404.
- Harwood, J., and S. L. King. (2014). *The Sensitivity of UK Marine Mammal Populations to Marine Renewables Developments*. Submitted to the Natural Environment Research Council (unpublished).
- Hastie, G. D., C. Donovan, T. Gotz, and V. M. Janik. (2014). Behavioral responses by grey seals (*Halichoerus grypus*) to high frequency sonar. *Marine Pollution Bulletin*, 79(1-2), 205–210.
- Hatch, L. T., and A. J. Wright. (2007). A brief review of anthropogenic sound in the oceans. *International Journal of Comparative Psychology*, 20, 121–133.
- Hatch, L. T., C. W. Clark, S. M. Van Parijs, A. S. Frankel, and D. W. Ponirakis. (2012). Quantifying loss of acoustic communication space for right whales in and around a U.S. National Marine Sanctuary. *Conservation Biology*, 26(6), 983–994.
- Haviland-Howell, G., A. S. Frankel, C. M. Powell, A. Bocconcelli, R. L. Herman, and L. S. Sayigh. (2007). Recreational boating traffic: A chronic source of anthropogenic noise in the Wilmington, North Carolina Intracoastal Waterway. *The Journal of the Acoustical Society of America*, 122(1), 151–160.
- Hawaii Pacific University. (2012). *HPU scientists pioneer research on rare whale*. Retrieved from <http://www.hpu.edu/HPUNews/2012/10/whale-research.html>.
- HDR. (2011a). *Guam Marine Species Monitoring Survey: Vessel-Based Monitoring Surveys Winter 2011*. Mariana Islands, Guam: U.S. Navy Marine Species Monitoring Program.
- HDR. (2011b). *Jacksonville Southeast Anti-Submarine Warfare Integration Training Initiative Marine Species Monitoring Aerial Monitoring Surveys Trip Report, 3–5 December 2010*. Jacksonville, FL: U.S. Navy Marine Species Monitoring Program.
- HDR. (2012). *Summary Report: Compilation of Visual Survey Effort and Sightings for Marine Species Monitoring in the Hawaii Range Complex, 2005–2012*. Pearl Harbor, HI: U.S. Pacific Fleet.
- HDR EOC. (2012). *Guam and Saipan Marine Species Monitoring Winter-Spring Survey, March 2012*. Pearl Harbor, HI: Naval Facilities Engineering Command.
- Heenehan, H. L., D. W. Johnston, S. M. Van Parijs, L. Bejder, and J. A. Tyne. (2016). *Acoustic response of Hawaiian spinner dolphins to human disturbances*. Paper presented at the Meetings on Acoustics. Dublin, Ireland.
- Heenehan, H. L., S. M. Van Parijs, L. Bejder, J. A. Tyne, and D. W. Johnston. (2017a). Using acoustics to prioritize management decisions to protect coastal dolphins: A case study using Hawaiian spinner dolphins. *Marine Policy*, 75, 84–90.

- Heenehan, H. L., S. M. Van Parijs, L. Bejder, J. A. Tyne, B. L. Southall, H. Southall, and D. W. Johnston. (2017b). Natural and anthropogenic events influence the soundscapes of four bays on Hawaii Island. *Marine Pollution Bulletin*, 124(1), 9–20.
- Heffner, R. S., and H. E. Heffner. (1982). Hearing in the elephant (*Elephas maximus*): Absolute sensitivity, frequency discrimination, and sound localization. *Journal of Comparative and Physiological Psychology*, 96(6), 926–944.
- Heinis, F., C. A. F. De Jong, and Rijkswaterstaat Underwater Sound Working Group. (2015). *Framework for Assessing Ecological and Cumulative Effects of Offshore Wind Farms: Cumulative Effects of Impulsive Underwater Sound on Marine Mammals* (TNO Report R10335-A). The Hague, Netherlands: Rijkswaterstaat Zee en Delta.
- Heithaus, M. (2001a). Predator-prey and competitive interactions between sharks (order Selachii) and dolphins (suborder Odontoceti): A review. *Journal of the Zoological Society of London*, 253, 53–68.
- Heithaus, M. R. (2001b). Shark attacks on bottlenose dolphins (*Tursiops aduncus*) in Shark Bay, Western Australia: Attack rate, bite scar frequencies and attack seasonality. *Marine Mammal Science*, 17(3), 526–539.
- Helber, R. W., and R. H. Weisberg. (2001). Equatorial upwelling in the western Pacific warm pool. *Journal of Geophysical Research*, 106(C5), 8989–9003.
- Helker, V. T., B. M. Allen, and L. A. Jemison. (2015). *Human-Caused Injury and Mortality of National Marine Fisheries Service-Managed Alaska Marine Mammal Stocks, 2009–2013*. Seattle, WA, and Dillingham, AK: Alaska Fisheries Science Center.
- Helker, V. T., M. M. Muto, K. Savage, S. Teerlink, L. A. Jemison, K. Wilkinson, and J. Jannot. (2017). *Human-Caused Mortality and Injury of NMFS-Managed Alaska Marine Mammal Stocks, 2011–2015* (NOAA Technical Memorandum NMFS-AFSC-354). Seattle, WA: Alaska Fisheries Science Center.
- Henderson, E. E., M. H. Smith, M. Gassmann, S. M. Wiggins, A. B. Douglas, and J. A. Hildebrand. (2014). Delphinid behavioral responses to incidental mid-frequency active sonar. *The Journal of the Acoustical Society of America*, 136(4), 2003–2014.
- Henderson, E. E., R. Manzano-Roth, S. W. Martin, and B. Matsuyama. (2015a). *Behavioral Responses of Beaked Whales to Mid-Frequency Active Sonar on the Pacific Missile Range Facility, Hawaii*. Paper presented at the Society for Marine Mammalogy 20th Biennial Conference. Dunedin, New Zealand.
- Henderson, E. E., R. A. Manzano-Roth, S. W. Martin, and B. Matsuyama. (2015b). *Impacts of U.S. Navy Training Events on Beaked Whale Foraging Dives in Hawaiian Waters: Update*. San Diego, CA: Space and Naval Warfare Systems Command Systems Center Pacific.
- Henderson, E. E., S. W. Martin, R. Manzano-Roth, and B. M. Matsuyama. (2016). Occurrence and habitat use of foraging Blainville's beaked whales (*Mesoplodon densirostris*) on a U.S. Navy range in Hawai'i. *Aquatic Mammals*, 42(4), 549–562.
- Hermanssen, L., K. Beedholm, J. Tougaard, and P. T. Madsen. (2014). High frequency components of ship noise in shallow water with a discussion of implications for harbor porpoises (*Phocoena phocoena*). *The Journal of the Acoustical Society of America*, 136(4), 1640–1653.

- Hewitt, R. P. (1985). Reaction of dolphins to a survey vessel: Effects on census data. *Fishery Bulletin*, 83(2), 187–193.
- Heyning, J. E., and J. G. Mead. (1996). Suction feeding in beaked whales: Morphological and observational evidence. *Contributions in Science*, 464, 1–12.
- Heyning, J. E., and J. G. Mead. (2009). Cuvier's beaked whale *Ziphius cavirostris*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 294–295). Cambridge, MA: Academic Press.
- Hickmott, L. S. (2005). *Diving behaviour and foraging ecology of Blainville's and Cuvier's beaked whales in the Northern Bahamas*. (Unpublished master's thesis). University of St. Andrews, Scotland, United Kingdom.
- Hidalgo-Ruz, V., L. Gutow, R. C. Thompson, and M. Thiel. (2012). Microplastics in the marine environment: A review of methods used for identification and quantification. *Environmental Science and Technology*, 46, 3060–3075.
- Hildebrand, J. (2009). Anthropogenic and natural sources of ambient noise in the ocean. *Marine Ecology Progress Series*, 395, 5–20.
- Hildebrand, J. A. (2005). Impacts of anthropogenic sound. In J. E. Reynolds, III, W. F. Perrin, R. R. Reeves, T. J. Ragen, & S. Montgomery (Eds.), *Marine Mammal Research: Conservation Beyond Crisis* (pp. 101–123). Baltimore, MD: The John Hopkins University Press.
- Hill, M., A. D. Ligon, M. H. Deakos, U. Adam, E. Norris, and E. M. Oleson. (2011). *Cetacean Surveys of Guam and CNMI Waters: August–September, 2011* (MIRC Survey Report FY2011). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Hill, M., A. Ligon, M. Deakos, A. Ü, A. Milette-Winfree, and E. Oleson. (2013a). *Cetacean Surveys of Guam and CNMI Waters: May–July, 2012: Including Individual Photo-Identification of Pilot Whales, Spinner Dolphins and Bottlenose Dolphins (2010–2012)* (PIFSC Data Report). Pearl Harbor, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hill, M., A. Ligon, A. Ü, and A. Bradford. (2015a). *Humpback Whales in the Marianas*. Honolulu, HI: National Oceanic and Atmospheric Administration, Pacific Islands Fisheries Science Center.
- Hill, M. C., A. D. Ligon, M. H. Deakos, A. C. U, and E. M. Oleson. (2013b). *Cetacean Surveys of Guam and SNMI Waters: June–July 2013*. Pearl Harbor, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hill, M. C., A. D. Ligon, M. H. Deakos, A. C. Ü, A. Milette-Winfree, A. R. Bendlin, and E. M. Oleson. (2014). *Cetacean Surveys in the Waters of the Southern Mariana Archipelago (February 2010–April 2014)*. Honolulu, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hill, M. C., E. M. Oleson, A. D. Ligon, K. K. Martien, F. I. Archer, S. Baumann-Pickering, A. R. Bendlin, L. Dolar, K. P. B. Merkens, A. Milette-Winfree, P. A. Morin, A. Rice, K. M. Robertson, J. S. Trickey, A. C. Ü, A. Van Cise, and S. M. Woodman. (2015b). *Cetacean Monitoring in the Mariana Islands Range Complex, 2014*. Honolulu, HI: U.S. Pacific Fleet.
- Hill, M. C., A. L. Bradford, A. D. Ligon, A. C. U, J. Rivers, R. K. Uyeyama, R. L. Brownell, Jr., and E. M. Oleson. (2016a). *Are Humpback Whales (Megaptera novaeangliae) Breeding and Calving in the Mariana Islands?* Cambridge, United Kingdom: International Whaling Commission.

- Hill, M. C., E. M. Oleson, S. Baumann-Pickering, A. M. VanCise, A. D. Ligon, A. R. Bendlin, A. C. Ü, J. S. Trickey, and A. L. Bradford. (2016b). *Cetacean Monitoring in the Mariana Islands Range Complex, 2015*. Honolulu, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hill, M. C., A. R. Bendlin, A. C. Ü, K. M. Yano, A. L. Bradford, A. D. Ligon, and E. M. Oleson. (2017a). *Cetacean Monitoring in the Mariana Islands Range Complex, 2016* (PIFSC Data Report DR-17-002). Honolulu, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hill, M. C., A. L. Bradford, A. D. Ligon, A. C. Ü, C. S. Baker, D. Dietrich-Steel, J. Rivers, R. K. Uyeyama, and E. M. Oleson. (2017b). *Discovery of a Western North Pacific Humpback Whale (Megaptera novaeangliae) Wintering Area in the Mariana Archipelago (Poster)*. Paper presented at the Society for Marine Mammalogy Conference. Halifax, Nova Scotia.
- Hill, M. C., A. L. Bradford, A. D. Ligon, A. C. Ü, and E. M. Oleson. (2018a). *Cetacean Monitoring in the Mariana Islands Range Complex, 2017* (PIFSC Data Report DR-18-002). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Hill, M. C., E. M. Oleson, A. L. Bradford, K. K. Martien, D. Steel, and C. S. Baker. (2018b). *Draft Pacific Islands Fisheries Science Center Mariana Archipelago Cetacean Surveys: A Review of Available Data and Analyses Through March 2018*. (PIFSC Data Report DR-18-xxx). Pearl Harbor, HI: U.S. Pacific Fleet Environmental Readiness Office.
- Hochachka, P. W., G. C. Liggins, G. P. Guyton, R. C. Schneider, K. S. Stanek, W. E. Hurford, R. K. Creasy, D. G. Zapol, and W. M. Zapol. (1995). Hormonal regulatory adjustments during voluntary diving in Weddell seals. *Comparative Biochemistry and Physiology B*, 112, 361–375.
- Holst, M., C. Greene, J. Richardson, T. McDonald, K. Bay, S. Schwartz, and G. Smith. (2011). Responses of pinnipeds to Navy missile launches at San Nicolas Island, California. *Aquatic Animals*, 37(2), 139–150.
- Holt, M. M., D. P. Noren, V. Veirs, C. K. Emmons, and S. Veirs. (2008). Speaking up: Killer whales (*Orcinus orca*) increase their call amplitude in response to vessel noise. *The Journal of the Acoustical Society of America*, 125(1), EL27–EL32.
- Holt, M. M., D. P. Noren, and C. K. Emmons. (2011). Effects of noise levels and call types on the source levels of killer whale calls. *The Journal of the Acoustical Society of America*, 130(5), 3100–3106.
- Holt, M. M., D. P. Noren, R. C. Dunkin, and T. M. Williams. (2015). Vocal performance affects metabolic rate in dolphins: Implications for animals communicating in noisy environments. *The Journal of Experimental Biology*, 218(Pt 11), 1647–1654.
- Holt, M. M., M. B. Hanson, D. A. Giles, C. K. Emmons, and J. T. Hogan. (2017). Noise levels received by endangered killer whales *Orcinus orca* before and after implementation of vessel regulations. *Endangered Species Research*, 34, 15–26.
- Hooker, S. K., R. W. Baird, and A. Fahlman. (2009). Could beaked whales get the bends? Effect of diving behaviour and physiology on modelled gas exchange for three species: *Ziphius cavirostris*, *Mesoplodon densirostris* and *Hyperoodon ampullatus*. *Respiratory Physiology & Neurobiology*, 167(3), 235–246.
- Hooker, S. K., A. Fahlman, M. J. Moore, N. A. de Soto, Y. B. de Quiros, A. O. Brubakk, D. P. Costa, A. M. Costidis, S. Dennison, K. J. Falke, A. Fernandez, M. Ferrigno, J. R. Fitz-Clarke, M. M. Garner, D. S. Houser, P. D. Jepson, D. R. Ketten, P. H. Kvaldsheim, P. T. Madsen, N. W. Pollock, D. S. Rotstein, T. K. Rowles, S. E. Simmons, W. Van Bonn, P. K. Weathersby, M. J. Weise, T. M. Williams, and P. L.

- Tyack. (2012). Deadly diving? Physiological and behavioural management of decompression stress in diving mammals. *Proceedings of the Royal Society B: Biological Sciences*, 279(1731), 1041–1050.
- Horton, T. W., N. Hauser, A. N. Zerbini, M. P. Francis, M. L. Domeier, A. Andriolo, D. P. Costa, P. W. Robinson, C. A. J. Duffy, N. Nasby-Lucas, R. N. Holdaway, and P. J. Clapham. (2017). Route fidelity during marine megafauna migration. *Frontiers in Marine Science*, 4, 1–21.
- Horwood, J. (1987). *The Sei Whale: Population Biology, Ecology, and Management*. New York, NY: Croom Helm.
- Horwood, J. (2009). Sei whale, *Balaenoptera borealis*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 1001–1003). Cambridge, MA: Academic Press.
- Hotchkin, C., and S. Parks. (2013). The Lombard effect and other noise-induced vocal modifications: Insight from mammalian communication systems. *Biological Reviews of the Cambridge Philosophical Society*, 88(4), 809–824.
- Houser, D. S., R. Howard, and S. Ridgway. (2001). Can diving-induced tissue nitrogen supersaturation increase the chance of acoustically driven bubble growth in marine mammals? *Journal of Theoretical Biology*, 213, 183–195.
- Houser, D. S., L. A. Dankiewicz-Talmadge, T. K. Stockard, and P. J. Ponganis. (2009). Investigation of the potential for vascular bubble formation in a repetitively diving dolphin. *The Journal of Experimental Biology*, 213, 52–62.
- Houser, D. S., L. C. Yeates, and D. E. Crocker. (2011). Cold stress induces an adrenocortical response in bottlenose dolphins (*Tursiops truncatus*). *Journal of Zoo and Wildlife Medicine*, 42(4), 565–571.
- Houser, D. S., S. W. Martin, and J. J. Finneran. (2013). Behavioral responses of California sea lions to mid-frequency (3250-3450 Hz) sonar signals. *Marine Environmental Research*, 92, 268–278.
- Houser, D. S., J. J. Finneran, E. Everaarts, J. Meerbeek, R. Dietz, S. Sveegaard, and J. Teilmann. (2016). Assessing auditory evoked potentials of wild harbor porpoises (*Phocoena phocoena*). *The Journal of the Acoustical Society of America*, 140(1), 442–452.
- Houser, D. S., W. Yost, R. F. Burkard, J. J. Finneran, C. Reichmuth, and J. Mulsow. (2017). A review of the history, development and application of auditory weighting functions in humans and marine mammals. *The Journal of the Acoustical Society of America*, 141(3), 1371–1413.
- Huggins, J. L., S. A. Raverty, S. A. Norman, J. Calambokidis, J. K. Gaydos, D. A. Duffield, D. M. Lambourn, J. M. Rice, B. Hanson, K. Wilkinson, S. J. Jeffries, B. Norberg, and L. Barre. (2015). Increased harbor porpoise mortality in the Pacific Northwest, USA: Understanding when higher levels may be normal. *Diseases of Aquatic Organisms*, 115(2), 93–102.
- Hurford, W. E., P. W. Hochachka, R. C. Schneider, G. P. Guyton, K. S. Stanek, D. G. Zapol, G. C. Liggins, and W. M. Zapol. (1996). Splenic contraction, catecholamine release, and blood volume redistribution during diving in the Weddell seal. *Journal of Applied Physiology*, 80(1), 298–306.
- Ilyashenko, V., P. J. Chapham, and R. L. Brownell. (2013). Soviet catches of whales in the North Pacific: Revised totals. *Journal of Cetacean Resource Management*, 13(1), 59–71.
- Ilyashenko, V., R. L. Brownell, and P. J. Chapham. (2014). Distribution of Soviet catches of sperm whales (*Physeter macrocephalus*) in the North Pacific. *Endangered Species Research*, 25, 249–263.

- Ilyashenko, V., and P. J. Chapham. (2014). Too much is never enough: The cautionary tale of Soviet illegal whaling. *Marine Fisheries Review*, 76(1–2), 21.
- Ilyashenko, V., P. J. Chapham, and R. L. Brownell. (2015). *New Data on Soviet Blue Whale Catches in the Eastern North Pacific in 1972*. Cambridge, United Kingdom: International Whaling Committee Scientific Committee.
- International Whaling Commission. (2016). Report of the Scientific Committee. *Journal of Cetacean Research and Management*, 17, 1–92.
- Isojunno, S., and P. J. O. Miller. (2015). Sperm whale response to tag boat presence: Biologically informed hidden state models quantify lost feeding opportunities. *Ecosphere*, 6(1), 1–6.
- Isojunno, S., C. Curé, P. H. Kvalsheim, F. A. Lam, P. L. Tyack, P. Jacobus, P. J. Wensveen, and P. J. O. Miller. (2016). Sperm whales reduce foraging effort during exposure to 1–2 kHz sonar and killer whale sounds. *Ecological Applications*, 26(1), 77–93.
- Jacobsen, J. K., L. Massey, and F. Gulland. (2010). Fatal ingestion of floating net debris by two sperm whales (*Physeter macrocephalus*). *Marine Pollution Bulletin*, 60(5), 765–767.
- Jahoda, M., C. L. Lafortuna, N. Biassoni, C. Almirante, A. Azzellino, S. Panigada, M. Zanardelli, and G. N. Di Sciara. (2003). Mediterranean fin whale's (*Balaenoptera physalus*) response to small vessels and biopsy sampling assessed through passive tracking and timing of respiration. *Marine Mammal Science*, 19(1), 96–110.
- Jambeck, J. R., R. Geyer, C. Wilcox, T. R. Siegler, M. Perryman, A. Andrady, R. Narayan, and K. L. Law. (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768–771.
- Janik, V. M., and P. M. Thompson. (1996). Changes in surfacing patterns of bottlenose dolphins in response to boat traffic. *Marine Mammal Science*, 12(4), 597–602.
- Jefferson, T. A., and S. Leatherwood. (1994). *Lagenodelphis hosei*. *American Society of Mammalogists*, 470, 1–5.
- Jefferson, T. A., and B. E. Curry. (1996). Acoustic methods of reducing or eliminating marine mammal-fishery interactions: Do they work? *Ocean & Coastal Management*, 31(1), 41–70.
- Jefferson, T. A., and N. B. Barros. (1997). *Peponocephala electra*. *Mammalian Species*, 553, 1–6.
- Jefferson, T. A., D. Fertl, M. Michael, and T. D. Fagin. (2006). An unusual encounter with a mixed school of melon-headed whales (*Peponocephala electra*) and rough-toothed dolphins (*Steno bredanensis*) at Rota, Northern Mariana Islands. *Micronesica*, 38(2), 23–244.
- Jefferson, T. A., M. A. Webber, and R. L. Pitman. (2008). *Marine Mammals of the World: A Comprehensive Guide to Their Identification*. London, United Kingdom: Elsevier.
- Jefferson, T. A., M. A. Webber, and R. L. Pitman. (2015). *Marine Mammals of the World: A Comprehensive Guide to Their Identification* (2nd ed.). Cambridge, MA: Academic Press.
- Jepson, P. D., M. Arbelo, R. Deaville, I. A. R. Patterson, P. Castro, J. R. Baker, E. Degollada, H. M. Ross, P. Herráez, A. M. Pocknell, F. Rodriguez, F. E. Howie, A. Espinosa, R. J. Reid, J. R. Jaber, V. Martin, A. A. Cunningham, and A. Fernandez. (2003). Gas-bubble lesions in stranded cetaceans: Was sonar responsible for a spate of whale deaths after an Atlantic military exercise? *Nature*, 425, 575–576.

- Jepson, P. D., P. M. Bennett, R. Deaville, C. R. Allchin, J. R. Baker, and R. J. Law. (2005). Relationships between polychlorinated biphenyls and health status in harbor porpoises (*Phocoena phocoena*) stranded in the United Kingdom. *Environmental Toxicology and Chemistry*, 24(1), 238–248.
- Jepson, P. D., and R. J. Law. (2016). Persistent pollutants, persistent threats; polychlorinated biphenyls remain a major threat to marine apex predators such as orcas. *Science*, 352(6292), 1388–1389.
- Johnson, C. S., M. W. McManus, and D. Skaar. (1989). Masked tonal hearing thresholds in the beluga whale. *The Journal of the Acoustical Society of America*, 85(6), 2651–2654.
- Juhasz, A. L., and R. Naidu. (2007). Explosives: Fate, dynamics, and ecological impact in terrestrial and marine environments. *Reviews of Environmental Contamination and Toxicology*, 191, 163–215.
- Kaluza, P., A. Kölzsch, M. T. Gastner, and B. Blasius. (2010). The complex network of global cargo ship movements. *Proceedings of the Royal Society*, 7(48), 11.
- Kami, H. T. (1982). Recent Beachings of Whales on Guam. *Micronesica*, 18, 133–135.
- Kastak, D., B. L. Southall, R. J. Schusterman, and C. R. Kastak. (2005). Underwater temporary threshold shift in pinnipeds: Effects of noise level and duration. *The Journal of the Acoustical Society of America*, 118(5), 3154–3163.
- Kastak, D., C. Reichmuth, M. M. Holt, J. Mulsow, B. L. Southall, and R. J. Schusterman. (2007). Onset, growth, and recovery of in-air temporary threshold shift in a California sea lion (*Zalophus californianus*). *The Journal of the Acoustical Society of America*, 122(5), 2916–2924.
- Kastelein, R., N. Jennings, W. Verboom, D. de Haan, and N. M. Schooneman. (2006). Differences in the response of a striped dolphin (*Stenella coeruleoalba*) and a harbor porpoise (*Phocoena phocoena*) to an acoustic alarm. *Marine Environmental Research*, 61, 363–378.
- Kastelein, R. A., D. de Haan, N. Vaughan, C. Staal, and N. M. Schooneman. (2001). The influence of three acoustic alarms on the behaviour of harbour porpoises (*Phocoena phocoena*) in a floating pen. *Marine Environmental Research*, 52, 351–371.
- Kastelein, R. A., W. C. Verboom, M. Muijsers, N. V. Jennings, and S. van der Heul. (2005). The influence of acoustic emissions for underwater data transmission on the behaviour of harbour porpoises (*Phocoena phocoena*) in a floating pen. *Marine Environmental Research*, 59, 287–307.
- Kastelein, R. A., and P. J. Wensveen. (2008). Effect of two levels of masking noise on the hearing threshold of a harbor porpoise (*Phocoena phocoena*) for a 4.0 kHz signal. *Aquatic Mammals*, 34(4), 420–425.
- Kastelein, R. A., R. Gransier, L. Hoek, A. Macleod, and J. M. Terhune. (2012a). Hearing threshold shifts and recovery in harbor seals (*Phoca vitulina*) after octave-band noise exposure at 4 kHz. *The Journal of the Acoustical Society of America*, 132(4), 2745–2761.
- Kastelein, R. A., R. Gransier, L. Hoek, and J. Olthuis. (2012b). Temporary threshold shifts and recovery in a harbor porpoise (*Phocoena phocoena*) after octave-band noise at 4 kHz. *The Journal of the Acoustical Society of America*, 132(5), 3525–3537.
- Kastelein, R. A., R. Gransier, L. Hoek, and M. Rambags. (2013a). Hearing frequency thresholds of a harbor porpoise (*Phocoena phocoena*) temporarily affected by a continuous 1.5 kHz tone. *The Journal of the Acoustical Society of America*, 134(3), 2286–2292.

- Kastelein, R. A., D. van Heerden, R. Gransier, and L. Hoek. (2013b). Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to playbacks of broadband pile driving sounds. *Marine Environmental Research*, 92, 206–214.
- Kastelein, R. A., L. Hoek, R. Gransier, C. A. F. de Jong, J. M. Terhune, and N. Jennings. (2014a). Hearing thresholds of a harbor porpoise (*Phocoena phocoena*) for playbacks of seal scarer signals, and effects of the signals on behavior. *Hydrobiologia*, 756(1), 89–103.
- Kastelein, R. A., L. Hoek, R. Gransier, M. Rambags, and N. Claeys. (2014b). Effect of level, duration, and inter-pulse interval of 1–2 kHz sonar signal exposures on harbor porpoise hearing. *The Journal of the Acoustical Society of America*, 136(1), 412–422.
- Kastelein, R. A., J. Schop, R. Gransier, and L. Hoek. (2014c). Frequency of greatest temporary hearing threshold shift in harbor porpoises (*Phocoena phocoena*) depends on the noise level. *The Journal of the Acoustical Society of America*, 136(3), 1410–1418.
- Kastelein, R. A., J. Schop, R. Gransier, N. Steen, and N. Jennings. (2014d). Effect of series of 1 to 2 kHz and 6 to 7 kHz up-sweeps and down-sweeps on the behavior of a harbor porpoise (*Phocoena phocoena*). *Aquatic Mammals*, 40(3), 232–242.
- Kastelein, R. A., R. Gransier, M. A. T. Marijt, and L. Hoek. (2015a). Hearing frequency thresholds of harbor porpoises (*Phocoena phocoena*) temporarily affected by played back offshore pile driving sounds. *The Journal of the Acoustical Society of America*, 137(2), 556–564.
- Kastelein, R. A., R. Gransier, J. Schop, and L. Hoek. (2015b). Effects of exposure to intermittent and continuous 6–7 kHz sonar sweeps on harbor porpoise (*Phocoena phocoena*) hearing. *The Journal of the Acoustical Society of America*, 137(4), 1623–1633.
- Kastelein, R. A., L. Helder-Hoek, G. Janssens, R. Gransier, and T. Johansson. (2015c). Behavioral responses of harbor seals (*Phoca vitulina*) to sonar signals in the 25-kHz range. *Aquatic Mammals*, 41(4), 388–399.
- Kastelein, R. A., I. van den Belt, R. Gransier, and T. Johansson. (2015d). Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to 25.5- to 24.5-kHz sonar down-sweeps with and without side bands. *Aquatic Mammals*, 41(4), 400–411.
- Kastelein, R. A., I. van den Belt, L. Helder-Hoek, R. Gransier, and T. Johansson. (2015e). Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to 25-kHz FM sonar signals. *Aquatic Mammals*, 41(3), 311–326.
- Kastelein, R. A., L. Helder-Hoek, J. Covi, and R. Gransier. (2016). Pile driving playback sounds and temporary threshold shift in harbor porpoises (*Phocoena phocoena*): Effect of exposure duration. *The Journal of the Acoustical Society of America*, 139(5), 2842–2851.
- Kastelein, R. A., J. Huybrechts, J. Covi, and L. Helder-Hoek. (2017). Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to sounds from an acoustic porpoise deterrent. *Aquatic Mammals*, 43(3), 233–244.
- Kastelein, R. A., L. Helder-Hoek, S. Van de Voorde, S. de Winter, S. Janssen, and M. A. Ainslie. (2018). Behavioral responses of harbor porpoises (*Phocoena phocoena*) to sonar playback sequences of sweeps and tones (3.5–4.1 kHz). *Aquatic Mammals*, 44(4), 389–404.
- Kasuya, T. (1971). Consideration of distribution and migration of toothed whales off the Pacific coast of Japan based upon aerial sighting record. *Scientific Reports of the Whales Research Institute*, 23, 37–60.

- Kato, H., and W. F. Perrin. (2009). Bryde's whales, *Balaenoptera edeni/brydei*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 158–163). Cambridge, MA: Academic Press.
- Keck, N., O. Kwiatek, F. Dhermain, F. Dupraz, H. Boulet, C. Danes, C. Laprie, A. Perrin, J. Godenir, L. Micout, and G. Libeau. (2010). Resurgence of *Morbillivirus* infection in Mediterranean dolphins off the French coast. *Veterinary Record*, 166(21), 654–655.
- Kelley, C., G. Carton, M. Tomlinson, and A. Gleason. (2016). Analysis of towed camera images to determine the effects of disposed mustard-filled bombs on the deep water benthic community off south Oahu. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 34–42.
- Kemp, N. J. (1996). Habitat loss and degradation. In M. P. Simmonds & J. D. Hutchinson (Eds.), *The Conservation of Whales and Dolphins* (pp. 263–280). New York, NY: John Wiley & Sons.
- Ketten, D. R., J. Lien, and S. Todd. (1993). Blast injury in humpback whale ears: Evidence and implications. *The Journal of the Acoustical Society of America*, 94(3), 1849–1850.
- Ketten, D. R. (1998). *Marine Mammal Auditory Systems: A Summary of Audiometric and Anatomical Data and Its Implications for Underwater Acoustic Impacts*. La Jolla, CA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.
- Ketten, D. R. (2000). Cetacean Ears. In W. Au, A. N. Popper, & R. R. Fay (Eds.), *Hearing by Whales and Dolphins* (1st ed., pp. 43–108). New York, NY: Springer-Verlag.
- King, S. L., R. S. Schick, C. Donovan, C. G. Booth, M. Burgman, L. Thomas, and J. Harwood. (2015). An interim framework for assessing the population consequences of disturbance. *Methods in Ecology and Evolution*, 6(10), 1150–1158.
- Kishiro, T. (1996). Movements of marked Bryde's whales in the western North Pacific. *Reports of the International Whaling Commission*, 46, 421–428.
- Klinck, H., S. L. Nieukirk, S. Fregosi, K. Klinck, D. K. Mellinger, S. Lastuka, G. B. Shilling, and J. C. Luby. (2016a). *Cetacean Studies on the Kona Coast in March 2010: Passive Acoustic Monitoring of Marine Mammals Using Gliders—Results from an Engineering Test. Final Report*. Honolulu, HI: HDR Inc.
- Klinck, H., S. L. Nieukirk, S. Fregosi, K. Klinck, D. K. Mellinger, S. Lastuka, G. B. Shilling, and J. C. Luby. (2016b). *Final Report Cetacean Studies on the Mariana Islands Range Complex in March–April 2015: Passive Acoustic Monitoring of Marine Mammals Using Gliders* (Submitted to Naval Facilities Engineering Command (NAVFAC) Pacific, Pearl Harbor, Hawaii). Honolulu, HI: HDR Inc.
- Kobayashi, N., H. Okabe, I. Kawazu, N. Higashi, H. Miyahara, H. Kato, and S. Uchida. (2016). Spatial distribution and habitat use patterns of humpback whales in Okinawa, Japan. *Mammal Study*, 41, 207–214.
- Koen-Alonso, M., S. N. Pedraza, A. C. M. Schiavini, R. N. P. Goodall, and E. A. Crespo. (1999). Stomach contents of false killer whales (*Pseudorca crassidens*) stranded on the coasts of the Strait of Magellan, Tierra Del Fuego. *Marine Mammal Science*, 15(3), 712–724.
- Koide, S., J. A. K. Silva, V. Dupra, and M. Edwards. (2015). Bioaccumulation of chemical warfare agents, energetic materials, and metals in deep-sea shrimp from discarded military munitions sites off Pearl Harbor. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 53–62.

- Kooyman, G. L., J. P. Schroeder, D. M. Denison, D. D. Hammond, J. J. Wright, and W. P. Bergman. (1972). Blood nitrogen tensions of seals during simulated deep dives. *American Journal of Physiology*, 223(5), 1016–1020.
- Kooyman, G. L., D. H. Kerem, W. B. Campbell, and J. J. Wright. (1973). Pulmonary gas exchange in freely diving Weddell seals, *Leptonychotes weddelli*. *Respiration Physiology*, 17, 283–290.
- Kooyman, G. L., and E. E. Sinnett. (1982). Pulmonary shunts in harbor seals and sea lions during simulated dives to depth. *Physiological Zoology* 55(1), 105–111.
- Koski, W. R., J. W. Lawson, D. H. Thomson, and W. J. Richardson. (1998). *Point Mugu Sea Range Marine Mammal Technical Report*. San Diego, CA: Naval Air Warfare Center, Weapons Division and Southwest Division, Naval Facilities Engineering Command.
- Koski, W. R., G. Gamage, A. R. Davis, T. Mathews, B. LeBlanc, and S. H. Ferguson. (2015). Evaluation of UAS for photographic re-identification of bowhead whales, *Balaena mysticetus*. *Journal of Unmanned Vehicle Systems*, 3(1), 22–29.
- Krahn, M. M., M. B. Hanson, R. W. Baird, R. H. Boyer, D. G. Burrows, C. K. Emmons, J. K. Ford, L. L. Jones, D. P. Noren, P. S. Ross, G. S. Schorr, and T. K. Collier. (2007). Persistent organic pollutants and stable isotopes in biopsy samples (2004/2006) from Southern Resident killer whales. *Marine Pollution Bulletin*, 54(12), 1903–1911.
- Krahn, M. M., M. B. Hanson, G. S. Schorr, C. K. Emmons, D. G. Burrows, J. L. Bolton, R. W. Baird, and G. M. Ylitalo. (2009). Effects of age, sex and reproductive status on persistent organic pollutant concentrations in "Southern Resident" killer whales. *Marine Pollution Bulletin*, 58(10), 1522–1529.
- Kremers, D., J. Lopez Marulanda, M. Hausberger, and A. Lemasson. (2014). Behavioural evidence of magnetoreception in dolphins: Detection of experimental magnetic fields. *Die Naturwissenschaften*, 101(11), 907–911.
- Kremers, D., A. Celerier, B. Schaal, S. Campagna, M. Trabalon, M. Boye, M. Hausberger, and A. Lemasson. (2016). Sensory Perception in Cetaceans: Part II—Promising Experimental Approaches to Study Chemoreception in Dolphins. *Frontiers in Ecology and Evolution*, 4(50), 1–9.
- Kruse, S. (1991). The interactions between killer whales and boats in Johnstone Strait, B.C. In K. Pryor & K. S. Norris (Eds.), *Dolphin Societies: Discoveries and Puzzles* (pp. 149–159). Berkeley and Los Angeles, CA: University of California Press.
- Kryter, K. D., W. D. Ward, J. D. Miller, and D. H. Eldredge. (1965). Hazardous exposure to intermittent and steady-state noise. *The Journal of the Acoustical Society of America*, 39(3), 451–464.
- Kujawa, S. G., and M. C. Liberman. (2009). Adding insult to injury: Cochlear nerve degeneration after "temporary" noise-induced hearing loss. *The Journal of Neuroscience*, 29(45), 14077–14085.
- Kuningas, S., P. H. Kvalsheim, F. P. A. Lam, and P. J. O. Miller. (2013). Killer whale presence in relation to naval sonar activity and prey abundance in northern Norway. *ICES Journal of Marine Science*, 70(7), 1287–1293.
- Kvalsheim, P. H., E. M. Sevaldsen, L. P. Folkow, and A. S. Blix. (2010). Behavioural and physiological responses of hooded seals (*Cytophoca cristata*) to 1 to 7 kHz sonar signals. *Aquatic Mammals*, 36(3), 239–247.
- Kvalsheim, P. H., P. J. Miller, P. L. Tyack, L. D. Sivle, F. P. Lam, and A. Fahlman. (2012). Estimated Tissue and Blood N<sub>2</sub> Levels and Risk of Decompression Sickness in Deep-, Intermediate-, and Shallow-

- Diving Toothed Whales during Exposure to Naval Sonar. *Frontiers in Physiology*, 3(Article 125), 125.
- Kvadsheim, P. H., S. DeRuiter, L. D. Sivle, J. Goldbogen, R. Roland-Hansen, P. J. O. Miller, F. A. Lam, J. Calambokidis, A. Friedlaender, F. Visser, P. L. Tyack, L. Kleivane, and B. Southall. (2017). Avoidance responses of minke whales to 1-4 kHz naval sonar. *Marine Pollution Bulletin*, 121(1–2), 60–68.
- Kyhn, L. A., P. B. Jørgensen, J. Carstensen, N. I. Bech, J. Tougaard, T. Dabelsteen, and J. Teilmann. (2015). Pingers cause temporary habitat displacement in the harbour porpoise *Phocoena phocoena*. *Marine Ecology Progress Series*, 526, 253–265.
- Laggner, D. (2009). *Blue whale (Baleanoptera musculus) ship strike threat assessment in the Santa Barbara Channel, California*. (Unpublished master's thesis). The Evergreen State College, Olympia, WA. Retrieved from <http://archives.evergreen.edu>.
- Laist, D. W. (1997). Impacts of marine debris: Entanglement of marine life in marine debris including a comprehensive list of species with entanglement and ingestion records. In J. M. Coe & D. B. Rogers (Eds.), *Marine Debris: Sources, Impacts, and Solutions* (pp. 99–140). New York, NY: Springer-Verlag.
- Laist, D. W., A. R. Knowlton, J. G. Mead, A. S. Collet, and M. Podesta. (2001). Collisions between ships and whales. *Marine Mammal Science*, 17(1), 35–75.
- Lammers, M. O. (2004). Occurrence and behavior of Hawaiian spinner dolphins (*Stenella longirostris*) along Oahu's leeward and south shores. *Aquatic Mammals*, 30(2), 237–250.
- Lammers, M. O., M. Howe, E. Zang, M. McElligott, A. Engelhaupt, and L. Munger. (2017). Acoustic monitoring of coastal dolphins and their response to naval mine neutralization exercises. *Royal Society Open Science*, 4(12), e170558.
- Law, R. J. (2014). An overview of time trends in organic contaminant concentrations in marine mammals: Going up or down? *Marine Pollution Bulletin*, 82(1–2), 7–10.
- Leatherwood, S., W. F. Perrin, V. L. Kirby, C. L. Hubbs, and M. Dahlheim. (1980). Distribution and movements of Risso's dolphin, *Grampus griseus*, in the eastern North Pacific. *Fishery Bulletin*, 77(4), 951–963.
- Leatherwood, S., F. T. Awbrey, and J. A. Thomas. (1982). Minke whale response to a transiting survey vessel. *Reports of the International Whaling Commission*, 32, 795–802.
- Lefebvre, K. A., L. Quakenbush, E. Frame, K. B. Huntington, G. Sheffield, R. Stimmelmayer, A. Bryan, P. Kendrick, H. Ziel, T. Goldstein, J. A. Snyder, T. Gelatt, F. Gulland, b. Dickerson, and V. Gill. (2016). Prevalence of algal toxins in Alaskan marine mammals foraging in a changing arctic and subarctic environment. *Harmful Algae*, 55(2016), 13–24.
- Lemonds, D. W., L. N. Kloepper, P. E. Nachtigall, W. W. Au, S. A. Vlachos, and B. K. Branstetter. (2011). A re-evaluation of auditory filter shape in delphinid odontocetes: Evidence of constant-bandwidth filters. *The Journal of the Acoustical Society of America*, 130(5), 3107–3114.
- Lent, R., and D. Squires. (2017). Reducing marine mammal bycatch in global fisheries: An economics approach. *Deep-Sea Research II: Topical Studies in Oceanography*, 140, 268–277.
- Lesage, V., C. Barrette, M. C. S. Kingsley, and B. Sjare. (1999). The effect of vessel noise on the vocal behavior of belugas in the St. Lawrence River estuary, Canada. *Marine Mammal Science*, 15(1), 65–84.

- Lesage, V., A. Omrane, T. Daniol-Valcroze, and A. Mosnier. (2017). Increased proximity of vessels reduces feeding opportunities of blue whales in the St. Lawrence Estuary, Canada. *Endangered Species Research*, 32, 351–361.
- Leslie, M. S., and P. A. Morin. (2018). Structure and phylogeography of two tropical predators, spinner (*Stenella longirostris*) and pantropical spotted (*S. attenuata*) dolphins, from SNP data. *Royal Society Open Science*, 5(4), 171615.
- Li, S., T. Akamatsu, D. Wang, K. Wang, S. Dong, X. Zhao, Z. Wei, X. Zhang, B. Taylor, L. A. Barrett, S. T. Turvey, R. R. Reeves, B. S. Stewart, M. Richlen, and J. R. Brandon. (2008). Indirect evidence of boat avoidance behavior of Yangtze finless porpoises. *Bioacoustics*, 17, 174–176.
- Li, S., H. Wu, Y. Xu, C. Peng, L. Fang, M. Lin, L. Xing, and P. Zhang. (2015). Mid- to high-frequency noise from high-speed boats and its potential impacts on humpback dolphins. *The Journal of the Acoustical Society of America*, 138(2), 942–952.
- Ligon, A. D., M. H. Deakos, and C. U. Adam. (2011). *Small-boat cetacean surveys off Guam and Saipan, Mariana Islands, February - March 2010*. Honolulu, HI: Pacific Island Fisheries Science Center.
- Lin, H. W., A. C. Furman, S. G. Kujawa, and M. C. Liberman. (2011). Primary neural degeneration in the guinea pig cochlea after reversible noise-induced threshold shift. *Journal of the Association for Research in Otolaryngology*, 12(5), 605–616.
- Liu, M., L. Dong, M. Lin, and S. Li. (2017). Broadband ship noise and its potential impacts on Indo-Pacific humpback dolphins: Implications for conservation and management. *The Journal of the Acoustical Society of America*, 142(5), 2766.
- Long, D. J. (1991). Apparent Predation by a White Shark *Carcharodon carcharias* on a Pygmy Sperm Whale *Kogia breviceps*. *Fishery Bulletin*, 89, 538–540.
- Lucke, K., U. Siebert, P. A. Lepper, and M. Blanchet. (2009). Temporary shift in masked hearing thresholds in a harbor porpoise (*Phocoena phocoena*) after exposure to seismic airgun stimuli. *The Journal of the Acoustical Society of America*, 125(6), 4060–4070.
- Luís, A. R., M. N. Couchinho, and M. E. dos Santos. (2014). Changes in the acoustic behavior of resident bottlenose dolphins near operating vessels. *Marine Mammal Science*, 30(4), 1417–1426.
- Lukoschek, V., N. Funahashi, S. Lavery, M. L. Dalebout, F. Cipriano, and C. S. Baker. (2009). The rise of commercial 'by-catch whaling' in Japan and Korea. *Animal Conservation*, 12, 398–399.
- Luksenburg, J. A., and E. C. M. Parsons. (2009). *The effects of aircraft on cetaceans: Implications for aerial whalewatching*. Paper presented at the 61st Meeting of the International Whaling Commission. Madeira, Portugal.
- Lusseau, D. (2004). The hidden cost of tourism: Detecting long-term effects of tourism using behavioral information. *Ecology and Society*, 9(1), 2.
- Lusseau, D. (2006). The short-term behavioral reactions of bottlenose dolphins to interactions with boats in Doubtful Sound, New Zealand. *Marine Mammal Science*, 22(4), 802–818.
- Lusseau, D., and L. Bejder. (2007). The long-term consequences of short-term responses to disturbance experiences from whalewatching impact assessment. *International Journal of Comparative Psychology*, 20, 228–236.
- Lusseau, D., D. E. Bain, R. Williams, and J. C. Smith. (2009). Vessel traffic disrupts the foraging behavior of southern resident killer whales, *Orcinus orca*. *Endangered Species Research*, 6, 211–221.

- Lyamin, O. I., S. M. Korneva, V. V. Rozhnov, and L. M. Mukhametov. (2011). Cardiorespiratory changes in beluga in response to acoustic noise. *Doklady Biological Sciences*, 440(5), 704–707.
- MacLeod, C., W. F. Perrin, R. Pitman, J. Barlow, L. Ballance, A. D'Amico, T. Gerrodette, G. Joyce, K. D. Mullin, D. L. Palka, and G. T. Waring. (2006). Known and inferred distributions of beaked whale species (family Ziphiidae; Order Cetacea). *Journal of Cetacean Research and Management*, 7(3), 271–286.
- MacLeod, C. D. (2000). Review of the distribution of *Mesoplodon* species (order Cetacea, family Ziphiidae) in the North Atlantic. *Mammal Review*, 30(1), 1–8.
- MacLeod, C. D., N. Hauser, and H. Peckham. (2004). Diversity, relative density and structure of the cetacean community in summer months east of Great Abaco, Bahamas. *Journal of the Marine Biological Association of the United Kingdom*, 84, 469–474.
- MacLeod, C. D., and A. D'Amico. (2006). A review of beaked whale behaviour and ecology in relation to assessing and mitigating impacts of anthropogenic noise. *Journal of Cetacean Research and Management*, 7(3), 211–222.
- MacLeod, C. D., and G. Mitchell. (2006). Key areas for beaked whales worldwide. *Journal of Cetacean Research and Management*, 7(3), 309–322.
- Madsen, P., M. Johnson, P. Miller, N. Soto, J. Lynch, and P. Tyack. (2006). Quantitative measures of air-gun pulses recorded on sperm whales (*Physeter macrocephalus*) using acoustic tags during controlled exposure experiments. *The Journal of the Acoustical Society of America*, 120(4), 2366–2379.
- Madsen, P. T., D. A. Carder, K. Bedholm, and S. H. Ridgway. (2005). Porpoise clicks from a sperm whale nose—Convergent evolution of 130 kHz pulses in toothed whale sonars? *Bioacoustics*, 15, 195–206.
- Magalhães, S., R. Prieto, M. A. Silva, J. Gonçalves, M. Afonso-Dias, and R. S. Santos. (2002). Short-term reactions of sperm whales (*Physeter macrocephalus*) to whale-watching vessels in the Azores. *Aquatic Mammals*, 28(3), 267–274.
- Malme, C. I., B. Würsig, J. E. Bird, and P. Tyack. (1986). *Behavioral responses of gray whales to industrial noise: Feeding observations and predictive modelling* (Outer Continental Shelf Environmental Assessment Program, Final Report of Principal Investigators MMS 88-0048). Anchorage, AK: Bolt Beranek, & Newman, Inc.
- Malme, C. I., B. Würsig, J. E. Bird, and P. Tyack. (1988). Observations of feeding gray whale responses to controlled industrial noise exposure. In W. M. Sackinger, M. O. Jeffries, J. L. Imm, & S. D. Tracey (Eds.), *Port and Ocean Engineering Under Arctic Conditions* (Vol. 2, pp. 55–73). Fairbanks, AK: Geophysical Institute, University of Alaska.
- Manci, K. M., D. N. Gladwin, R. Villella, and M. G. Cavendish. (1988). *Effects of Aircraft Noise and Sonic Booms on Domestic Animals and Wildlife: A Literature Synthesis* (NERC-88/29). Fort Collins, CO: U.S. Fish and Wildlife Service, National Ecology Research Center.
- Manzano-Roth, R., E. E. Henderson, S. W. Martin, C. Martin, and B. M. Matsuyama. (2016). Impacts of U.S. Navy training events on Blainville's beaked whale (*Mesoplodon densirostris*) foraging dives in Hawaiian waters. *Aquatic Mammals*, 42(4), 507–518.

- Marcoux, M., H. Whitehead, and L. Rendell. (2007). Sperm whale feeding variations by location, year, social group and clan: Evidence from stable isotopes. *Marine Ecology Progress Series*, 333, 309–314.
- Marine Mammal Commission. (2010). *The Marine Mammal Commission Annual Report to Congress 2009*. Bethesda, MD: Marine Mammal Commission.
- Marsh, H. E. (1989). Mass stranding of dugongs by a tropical cyclone in northern Australia. *Marine Mammal Science*, 5(1), 78–84.
- Marten, K., and S. Psarakos. (1999). Long-term site fidelity and possible long-term associations of wild spinner dolphins (*Stenella longirostris*) seen off Oahu, Hawaii. *Marine Mammal Science*, 15(4), 1329–1336.
- Martien, K. K., R. W. Baird, N. M. Hedrick, A. M. Gorgone, J. L. Thieleking, D. J. McSweeney, K. M. Robertson, and D. L. Webster. (2012). Population structure of island-associated dolphins: Evidence from mitochondrial and microsatellite markers for common bottlenose dolphins (*Tursiops truncatus*) around the main Hawaiian Islands. *Marine Mammal Science*, 28(3), E208–E232.
- Martien, K. K., S. J. Chivers, R. W. Baird, F. I. Archer, A. M. Gorgone, B. L. Hancock-Hanser, D. Mattila, D. J. McSweeney, E. M. Oleson, C. Palmer, V. L. Pease, K. M. Robertson, G. S. Schorr, M. B. Schultz, D. L. Webster, and B. L. Taylor. (2014a). Nuclear and mitochondrial patterns of population structure in North Pacific false killer whales (*Pseudorca crassidens*). *The Journal of Heredity*, 105(5), 611–626.
- Martien, K. K., M. C. Hill, A. M. Van Cise, K. M. Robertson, S. M. Woodman, L. Dollar, V. L. Pease, and E. M. Oleson. (2014b). *Genetic Diversity and Population Structure in Four Species of Cetaceans Around the Mariana Islands* (NOAA Technical Memorandum NMFS-SWFSC-536). La Jolla, CA: National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.
- Martin, C. R., S. W. Martin, E. E. Henderson, T. A. Helble, R. A. Manzano-Roth, B. M. Matsuyama, and G. C. Alongi. (2017). *SSC Pacific FY16 annual report on PMRF Marine Mammal Monitoring. Final Report*. San Diego, CA: National Marine Mammal Foundation and Space and Naval Warfare Systems Center Pacific.
- Martin, S. L., K. S. Van Houtan, T. T. Jones, C. F. Aguon, J. T. Gutierrez, R. B. Tibbatts, S. B. Wusstig, and J. D. Bass. (2016). Five decades of marine megafauna surveys from Micronesia. *Frontiers in Marine Science*, 2(116), 1–13.
- Martin, S. W., C. R. Martin, B. M. Matsuyama, and E. E. Henderson. (2015). Minke whales (*Balaenoptera acutorostrata*) respond to navy training. *The Journal of the Acoustical Society of America*, 137(5), 2533–2541.
- Masaki, Y. (1972). Tagging Investigations of Whales in the Ogasawara and Mariana Islands. *Japan Whaling Association Institute of Cetacean Research*, 249, 35–42.
- Matkin, C. O., E. L. Saulitis, G. M. Ellis, P. Olesiuk, and S. D. Rice. (2008). Ongoing population-level impacts on killer whales, *Orcinus orca*, following the ‘Exxon Valdez’ oil spill in Prince William Sound, Alaska. *Marine Ecology Progress Series*, 356, 269–281.

- Mattson, M. C., J. A. Thomas, and D. St. Aubin. (2005). Effects of boat activity on the behavior of bottlenose dolphins (*Tursiops truncatus*) in waters surrounding Hilton Head Island, South Carolina. *Aquatic Mammals*, 31(1), 133–140.
- May-Collado, L. J., and D. Wartzok. (2008). A comparison of bottlenose dolphin whistles in the Atlantic Ocean: Factors promoting whistle variation. *Journal of Mammalogy*, 89(5), 1229–1240.
- McAlpine, D. F. (2009). Pygmy and dwarf sperm whales *Kogia breviceps* and *K. sima*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 936–938). Cambridge, MA: Academic Press.
- McCarthy, E., D. Moretti, L. Thomas, N. DiMarzio, R. Morrissey, S. Jarvis, J. Ward, A. Izzi, and A. Dilley. (2011). Changes in spatial and temporal distribution and vocal behavior of Blainville's beaked whales (*Mesoplodon densirostris*) during multiship exercises with mid-frequency sonar. *Marine Mammal Science*, 27(3), E206–E226.
- McCauley, R. D., M. N. Jenner, C. Jenner, K. A. McCabe, and J. Murdoch. (1998). The response of humpback whales (*Megaptera novaeangliae*) to offshore seismic survey noise: Preliminary results of observations about a working seismic vessel and experimental exposures. *Australian Petroleum Production and Exploration Association Journal*, 38, 692–706.
- McCauley, R. D., J. Fewtrell, A. J. Duncan, C. Jenner, M.-N. Jenner, J. D. Penrose, R. I. T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. (2000). Marine seismic surveys: A study of environmental implications. *Australian Petroleum Production Exploration Association Journal*, 2000, 692–708.
- McDonald, B. I., and P. J. Ponganis. (2012). Lung collapse in the diving sea lion: Hold the nitrogen and save the oxygen. *Biology Letters*, 8, 1047–1049.
- McDonald, M., J. Hildebrand, S. Wiggins, and D. Ross. (2008). A 50 year comparison of ambient ocean noise near San Clemente Island: A bathymetrically complex coastal region off Southern California. *The Journal of the Acoustical Society of America*, 124(4), 1985–1992.
- McDonald, M. A., J. A. Hildebrand, and S. C. Webb. (1995). Blue and fin whales observed on a seafloor array in the Northeast Pacific. *The Journal of the Acoustical Society of America*, 98(2), 712–721.
- McDonald, M. A., J. A. Hildebrand, S. M. Wiggins, D. W. Johnston, and J. J. Polovina. (2009). An acoustic survey of beaked whales at Cross Seamount near Hawaii. *The Journal of the Acoustical Society of America*, 125(2), 624–627.
- McHuron, E. A., L. K. Schwarz, D. P. Costa, and M. Mangel. (2018). A state-dependent model for assessing the population consequences of disturbance on income-breeding mammals. *Ecological Modelling*, 385, 133–144.
- McKenna, M. F., D. Ross, S. M. Wiggins, and J. A. Hildebrand. (2012). Underwater radiated noise from modern commercial ships. *The Journal of the Acoustical Society of America*, 131(1), 92–103.
- McSweeney, D. J., R. W. Baird, and S. D. Mahaffy. (2007). Site fidelity, associations, and movements of Cuvier's (*Ziphius Cavirostris*) and Blainville's (*Mesoplodon Densirostris*) beaked whales off the island of Hawaii. *Marine Mammal Science*, 23(3), 666–687.
- Mead, J. G., and C. W. Potter. (1995). Recognizing two populations of the bottlenose dolphin (*Tursiops truncatus*) off the Atlantic Coast of North America: Morphologic and ecologic considerations. *IBI Reports*, 5, 31–44.
- Meissner, A. M., F. Christiansen, E. Martinez, M. D. Pawley, M. B. Orams, and K. A. Stockin. (2015). Behavioural effects of tourism on oceanic common dolphins, *Delphinus* sp., in New Zealand: The

- effects of Markov analysis variations and current tour operator compliance with regulations. *PLoS ONE*, 10(1), e0116962.
- Melcón, M. L., A. J. Cummins, S. M. Kerosky, L. K. Roche, S. M. Wiggins, and J. A. Hildebrand. (2012). Blue whales respond to anthropogenic noise. *PLoS ONE*, 7(2).
- Mesnick, S. L., B. L. Taylor, F. I. Archer, K. K. Martien, S. E. Trevino, B. L. Hancock-Hanser, S. C. M. Medina, V. L. Pease, K. M. Robertson, J. M. Straley, R. W. Baird, J. Calambokidis, G. S. Schorr, P. Wade, V. Burkanov, C. R. Lunsford, L. Rendell, and P. A. Morin. (2011). Sperm whale population structure in the eastern and central North Pacific inferred by the use of single-nucleotide polymorphisms, microsatellites and mitochondrial DNA. *Molecular Ecology Resources*, 11 (Supplement 1), 278–298.
- Mignucci-Giannoni, A. A., B. Pinto-Rodríguez, M. Velasco-Escudero, R. A. Montoya-Ospina, N. M. Jiménez-Marrero, M. A. Rodríguez-López, E. H. Williams, Jr., and D. K. Odell. (1999). Cetacean strandings in Puerto Rico and the Virgin Islands. *Journal of Cetacean Research and Management*, 1(2), 191–198.
- Mikkelsen, L., L. Hermannsen, K. Beedholm, P. T. Madsen, and J. Tougaard. (2017). Simulated seal scarer sounds scare porpoises, but not seals: Species-specific responses to 12 kHz deterrence sounds. *Royal Society Open Science*, 4(7), 170286.
- Miksis-Olds, J. L., and S. M. Nichols. (2015). Is low frequency ocean sound increasing globally? *The Journal of the Acoustical Society of America*, 139(1), 501–511.
- Miksis, J. L., R. C. Connor, M. D. Grund, D. P. Nowacek, A. R. Solow, and P. L. Tyack. (2001). Cardiac responses to acoustic playback experiments in the captive bottlenose dolphin (*Tursiops truncatus*). *Journal of Comparative Psychology*, 115(3), 227–232.
- Miller, J. D., C. S. Watson, and W. P. Covell. (1963). Deafening effects of noise on the cat. *Acta Oto-Laryngologica, Supplement 176*, 1–88.
- Miller, L. J., A. D. Mackey, T. Hoffland, and S. A. Kuczaj, II. (2010). Potential effects of a major hurricane on Atlantic bottlenose dolphin (*Tursiops truncatus*) reproduction in the Mississippi Sound. *Marine Mammal Science*, 26(3), 707–715.
- Miller, P., M. Johnson, P. Madsen, N. Biassoni, M. Quero, and P. Tyack. (2009). Using at-sea experiments to study the effects of airguns on the foraging behavior of sperm whales in the Gulf of Mexico. *Deep Sea Research I*, 56(7), 1168–1181.
- Miller, P., R. Antunes, A. C. Alves, P. Wensveen, P. Kvadsheim, L. Kleivane, N. Nordlund, F.-P. Lam, S. van IJsselmuide, F. Visser, and P. Tyack. (2011). *The 3S experiments: Studying the behavioural effects of naval sonar on killer whales (Orcinus orca), sperm whales (Physeter macrocephalus), and long-finned pilot whales (Globicephala melas) in Norwegian waters* (Technical Report SOI-2011-001). St. Andrews, United Kingdom: Scottish Oceans Institute.
- Miller, P. (2012). The severity of behavioral changes observed during experimental exposures of killer (*Orcinus orca*), long-finned pilot (*Globicephala melas*), and sperm (*Physeter macrocephalus*) whales to naval sonar. *Aquatic Mammals*, 38(4), 362–401.
- Miller, P. J., R. N. Antunes, P. J. Wensveen, F. I. Samarra, A. C. Alves, P. L. Tyack, P. H. Kvadsheim, L. Kleivane, F. P. Lam, M. A. Ainslie, and L. Thomas. (2014). Dose-response relationships for the onset of avoidance of sonar by free-ranging killer whales. *The Journal of the Acoustical Society of America*, 135(2), 975–993.

- Miller, P. J., P. H. Kvadsheim, F. P. Lam, P. L. Tyack, C. Cure, S. L. DeRuiter, L. Kleivane, L. D. Sivle, I. S. P. van, F. Visser, P. J. Wensveen, A. M. von Benda-Beckmann, L. M. Martin Lopez, T. Narazaki, and S. K. Hooker. (2015). First indications that northern bottlenose whales are sensitive to behavioural disturbance from anthropogenic noise. *Royal Society Open Science*, 2(6), 140484.
- Miller, P. J. O., N. Biassoni, A. Samuels, and P. L. Tyack. (2000). Whale songs lengthen in response to sonar. *Nature*, 405(6789), 903.
- Miyashita, T. (1993a). Abundance of dolphin stocks in the western North Pacific taken by the Japanese drive fishery. *Reports of the International Whaling Commission*, 43, 417–437.
- Miyashita, T. (1993b). Distribution and abundance of some dolphins taken in the North Pacific driftnet fisheries. *International North Pacific Fisheries Commission Bulletin*, 53(3), 435–450.
- Miyashita, T., H. Kato, and T. Kasuya. (1995). *Worldwide Map of Cetacean Distribution Based on Japanese Sighting Data*. Shimizu, Japan: National Research Institute of Far Seas Fisheries.
- Miyashita, T., T. Kishiro, N. Higashi, F. Sato, K. Mori, and H. Kato. (1996). Winter distribution of cetaceans in the western North Pacific inferred from sighting cruises 1993–1995. *Reports of the International Whaling Commission*, 46, 437–442.
- Miyazaki, N., and W. F. Perrin. (1994). Rough-toothed dolphin, *Steno bredanensis* (Lesson, 1828). In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 1–21). San Diego, CA: Academic Press.
- Mizroch, S. A., D. W. Rice, D. Zwiefelhofer, J. M. Waite, and W. L. Perryman. (2009). Distribution and movements of fin whales in the North Pacific Ocean. *Mammal Review*, 39(3), 193–227.
- Mizroch, S. A., and D. W. Rice. (2013). Ocean nomads: Distribution and movements of sperm whales in the North Pacific shown by whaling data and discovery marks. *Marine Mammal Science*, 29(2), E136–E165.
- Moberg, G. P., and J. A. Mench. (2000). *The Biology of Animal Stress; Basic Principles and Implications for Animal Welfare*. London, United Kingdom: CAB International.
- Mobley, J. R. (2007). *Marine Mammal Monitoring Surveys in Support of "Valiant Shield" Training Exercises (Aug. 13-17, 2007)—Final Report*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Mobley, J. R., S. W. Martin, D. Fromm, and P. E. Nachtigall. (2007). *Lunar influences as possible cause for simultaneous aggregations of melon-headed whales in Hanalei Bay, Kaua'i, and Sasanhaya Bay, Rota*. Paper presented at the Abstract, 17th Biennial Meeting on the Biology of Marine Mammals, Society for Marine Mammalogy. Cape Town, South Africa.
- Mobley, J. R., and A. Milette. (2010). *Aerial Survey Monitoring for Marine Mammals and Sea Turtles in the Hawaiian Range Complex in Conjunction with a Navy Training Event, SCC February 16–21, 2010, Final Field Report*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Mobley, J. R. (2011). *Aerial Survey Monitoring for Marine Mammals and Sea Turtles in the Hawaii Range Complex in Conjunction with Two Navy Training Events. SCC and USWEX February 16–March 5, 2011. Final Field Report*. San Diego, CA: HDR Inc.
- Mobley, J. R., and A. F. Pacini. (2012). *Aerial Survey Monitoring for Marine Mammals and Sea Turtles in the Hawaii Range Complex in Conjunction with a Navy Training Event, SCC February 15–25, 2012, Final Field Report*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.

- Mobley, J. R., M. A. Smultea, C. E. Bacon, and A. S. Frankel. (2012). *Preliminary Report: Aerial Survey Monitoring for Marine Mammals and Sea Turtles in the Hawaii Range Complex—Summary of Focal Follow Analysis for 2008–2012 SCC Events: Preliminary Report*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Mobley, J. R., and M. H. Deakos. (2015). *Aerial Shoreline Surveys for Marine Mammals and Sea Turtles in the Hawaii Range Complex, Conducted after Navy Training Events. Koa Kai Surveys: 31 January and 5 February 2014. RIMPAC Surveys: 1 and 4–6 July 2014*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Møller, A. R. (2013). *Hearing: Anatomy, Physiology, and Disorders of the Auditory System*. San Diego, CA: Plural Publishing.
- Monnahan, C. C. (2013). *Population Trends of the Eastern North Pacific Blue Whale*. (Unpublished master's thesis). University of Washington, Seattle, WA. Retrieved from <http://digital.lib.washington.edu>.
- Monnahan, C. C., T. A. Branch, K. M. Stafford, Y. V. Ivashchenko, and E. M. Oleson. (2014). Estimating historical eastern North Pacific blue whale catches using spatial calling patterns. *PLoS ONE*, 9(6), e98974.
- Monnahan, C. C., T. A. Branch, and A. E. Punt. (2015). Do ship strikes threaten the recovery of endangered eastern North Pacific blue whales? *Marine Mammal Science*, 31(1), 279–297.
- Montie, E. W., C. A. Manire, and D. A. Mann. (2011). Live CT imaging of sound reception anatomy and hearing measurements in the pygmy killer whale, *Feresa attenuata*. *The Journal of Experimental Biology*, 214, 945–955.
- Moon, H. B., K. Kannan, M. Choi, J. Yu, H. G. Choi, Y. R. An, S. G. Choi, J. Y. Park, and Z. G. Kim. (2010). Chlorinated and brominated contaminants including PCBs and PBDEs in minke whales and common dolphins from Korean coastal waters. *Journal of Hazardous Materials*, 179(1–3), 735–741.
- Mooney, T. A., P. E. Nachtigall, M. Breese, S. Vlachos, W. Whitlow, and L. Au. (2009a). Predicting temporary threshold shifts in a bottlenose dolphin (*Tursiops truncatus*): The effects of noise level and duration. *The Journal of the Acoustical Society of America*, 125(3), 1816–1826.
- Mooney, T. A., P. E. Nachtigall, and S. Vlachos. (2009b). Sonar-induced temporary hearing loss in dolphins. *Biology Letters*, 5(4), 565–567.
- Mooney, T. A., M. Yamato, and B. K. Branstetter. (2012). *Hearing in Cetaceans: From Natural History to Experimental Biology*. Woods Hole, MA: Woods Hole Oceanographic Institution and the National Marine Mammal Foundation.
- Moore, C. J. (2008). Synthetic polymers in the marine environment: A rapidly increasing, long-term threat. *Environmental Research*, 108(2), 131–139.
- Moore, J., and J. Barlow. (2017). *Population Abundance and Trend Estimates for Beaked Whales and Sperm Whales in the California Current from Ship-Based Visual Line-Transect Survey Data, 1991–2014* (National Oceanic and Atmospheric Administration Technical Memorandum NMFS-SWFSC-585). La Jolla, CA: Southwest Fisheries Science Center.
- Moore, J. C. (1972). More skull characters of the beaked whale, *Indopacetus pacificus*, and comparative measurements of austral relatives. *Fieldiana Zoology*, 62(1), 1–19.

- Moore, J. E., and J. P. Barlow. (2013). Declining abundance of beaked whales (Family Ziphiidae) in the California Current Large Marine Ecosystem. *PLoS ONE*, 8(1), e52770.
- Moore, M. J., and G. A. Early. (2004). Cumulative sperm whale bone damage and the bends. *Science*, 306, 2215.
- Moore, M. J., A. L. Bogomolni, S. E. Dennison, G. Early, M. M. Garner, B. A. Hayward, B. J. Lentell, and D. S. Rotstein. (2009). Gas bubbles in seals, dolphins, and porpoises entangled and drowned at depth in gillnets. *Veterinary Pathology* 46, 536–547.
- Moore, M. J., J. van der Hoop, S. G. Barco, A. M. Costidis, F. M. Gulland, P. D. Jepson, K. T. Moore, S. Raverty, and W. A. McLellan. (2013). Criteria and case definitions for serious injury and death of pinnipeds and cetaceans caused by anthropogenic trauma. *Diseases of Aquatic Organisms*, 103(3), 229–264.
- Moretti, D., N. DiMarzio, R. Morrissey, E. McCarthy, and S. Jarvis. (2009). *An opportunistic study of the effect of sonar on marine mammals, marine mammal monitoring on Navy ranges (M3R)*. Paper presented at the 2009 ONR Marine Mammal Program Review. Alexandria, VA.
- Moretti, D., L. Thomas, T. Marques, J. Harwood, A. Dilley, B. Neales, J. Shaffer, E. McCarthy, L. New, S. Jarvis, and R. Morrissey. (2014). A risk function for behavioral disruption of Blainville's beaked whales (*Mesoplodon densirostris*) from mid-frequency active sonar. *PLoS ONE*, 9(1), e85064.
- Moretti, D. (2017). *Marine Mammal Monitoring on Navy Ranges (M3R) Passive Acoustic Monitoring of Abundance on the Pacific Missile Range Facility (PMRF) and Southern California Offshore Range (SCORE)*. Newport, RI: Naval Undersea Warfare Center.
- Moretti, D. J., and R. W. Baird. (2015). *Marine Mammal Monitoring on Navy Ranges (M3R), An Opportunistic Study of the Effect of Sonar on Marine Mammals on the Pacific Missile Range Facility (PMRF): FY14 Summary*. Newport, RI: Cascadia Research Collective, Naval Undersea Warfare Center.
- Mulsow, J., and C. Reichmuth. (2010). Psychophysical and electrophysiological aerial audiograms of a Steller sea lion (*Eumetopias jubatus*). *The Journal of the Acoustical Society of America*, 127(4), 2692–2701.
- Mulsow, J., C. E. Schlundt, L. Brandt, and J. J. Finneran. (2015). Equal latency contours for bottlenose dolphins (*Tursiops truncatus*) and California sea lions (*Zalophus californianus*). *The Journal of the Acoustical Society of America*, 138(5), 2678.
- Mulsow, J. L., J. J. Finneran, and D. S. Houser. (2011). California sea lion (*Zalophus californianus*) aerial hearing sensitivity measured using auditory steady-state response and psychophysical methods. *The Journal of the Acoustical Society of America*, 129(4), 2298–2306.
- Munger, L. M., M. O. Lammers, and W. W. L. Au. (2014). *Passive Acoustic Monitoring for Cetaceans within the Marianas Islands Range Complex. Preliminary Report*. Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- Munger, L. M., M. O. Lammers, J. N. Oswald, T. M. Yack, and W. W. L. Au. (2015). *Passive Acoustic Monitoring of Cetaceans within the Mariana Islands Range Complex Using Ecological Acoustic Recorders. Final Report*. Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- Murase, H., T. Tamura, S. Otani, and S. Nishiwaki. (2015). Satellite tracking of Bryde's whales *Balaenoptera edeni* in the offshore western North Pacific in summer 2006 and 2008. *Fisheries Science*, 82(1), 35–45.

- Muto, M. M., V. T. Helker, R. P. Angliss, B. A. Allen, P. L. Boveng, J. M. Breiwick, M. F. Cameron, P. J. Clapham, S. P. Dahle, M. E. Dahlheim, B. S. Fadely, M. C. Ferguson, L. W. Fritz, R. C. Hobbs, Y. V. Ivashchenko, A. S. Kennedy, J. M. London, S. A. Mizroch, R. R. Ream, E. L. Richmond, K. E. W. Shelden, R. G. Towell, P. R. Wade, J. M. Waite, and A. R. Zerbini. (2017a). *Alaska Marine Mammal Stock Assessments, 2016* (NOAA Technical Memorandum NMFS-AFSC-323). Seattle, WA: National Marine Mammal Laboratory.
- Muto, M. M., V. T. Helker, R. P. Angliss, B. A. Allen, P. L. Boveng, J. M. Breiwick, M. F. Cameron, P. J. Clapham, S. P. Dahle, M. E. Dahlheim, B. S. Fadely, J. M. London, S. A. Mizroch, R. R. Ream, E. L. Richmond, K. E. W. Shelden, R. G. Towell, P. R. Wade, J. M. Waite, and A. N. Zerbini. (2017b). *Alaska Marine Mammal Stock Assessments, 2017*. Seattle, WA: Alaska Fisheries Science Center.
- Muto, M. M., V. T. Helker, R. P. Angliss, P. L. Boveng, J. M. Breiwick, M. F. Cameron, P. J. Clapham, S. P. Dahle, M. E. Dahlheim, B. S. Fadely, M. C. Ferguson, L. W. Fritz, R. C. Hobbs, Y. V. Ivashchenko, A. S. Kennedy, J. M. London, S. A. Mizroch, R. R. Ream, E. L. Richmond, K. E. W. Shelden, K. L. Sweeney, R. G. Towell, P. R. Wade, J. M. Waite, and A. N. Zerbini. (2018). *Alaska Marine Mammal Stock Assessments, 2018. Draft*. Seattle, WA: National Marine Fisheries Service, Alaska Fisheries Science Center.
- Nabe-Nielsen, J., R. M. Sibly, J. Tougaard, J. Teilmann, and S. Sveegaard. (2014). Effects of noise and by-catch on a Danish harbour porpoise population. *Ecological Modelling*, 272, 242–251.
- Nachtigall, P. E., D. W. Lemonds, and H. L. Roitblat. (2000). Psychoacoustic Studies of Dolphin and Whale Hearing. In W. W. L. Au, R. R. Fay, & A. N. Popper (Eds.), *Hearing by Whales and Dolphins* (pp. 330–363). New York, NY: Springer.
- Nachtigall, P. E., J. L. Pawloski, and W. W. L. Au. (2003). Temporary threshold shifts and recovery following noise exposure in the Atlantic bottlenosed dolphin (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 113(6), 3425–3429.
- Nachtigall, P. E., A. Y. Supin, J. Pawloski, and W. W. L. Au. (2004). Temporary threshold shifts after noise exposure in the bottlenose dolphin (*Tursiops truncatus*) measured using evoked auditory potentials. *Marine Mammal Science*, 20(4), 673–687.
- Nachtigall, P. E., A. Y. Supin, M. Amundin, B. Roken, T. Møller, T. A. Mooney, K. A. Taylor, and M. Yuen. (2007). Polar bear, *Ursus maritimus*, hearing measured with auditory evoked potentials. *The Journal of Experimental Biology*, 210(7), 1116–1122.
- Nachtigall, P. E., T. A. Mooney, K. A. Taylor, L. A. Miller, M. H. Rasmussen, T. Akamatsu, J. Teilmann, M. Linnenschmidt, and G. A. Vikingsson. (2008). Shipboard Measurements of the Hearing of the White-Beaked Dolphin, *Lagenorhynchus albirostris*. *The Journal of Experimental Biology*, 211, 642–647.
- Nachtigall, P. E., A. Y. Supin, A. F. Pacini, and R. A. Kastelein. (2018). Four odontocete species change hearing levels when warned of impending loud sound. *Integrative Zoology*, 13, 2–20.
- National Academies of Sciences Engineering and Medicine. (2017). *Approaches to Understanding the Cumulative Effects of Stressors on Marine Mammals*. Washington, DC: The National Academies Press Series.
- National Marine Fisheries Service. (2005). *Assessment of Acoustic Exposures on Marine Mammals in Conjunction with USS Shoup Active Sonar Transmissions in the Eastern Strait of Juan de Fuca and Haro Strait, Washington (5 May 2003)*. Seattle, WA: National Marine Fisheries Service.

- National Marine Fisheries Service. (2007a). *Biological Opinion on the U.S. Navy's Proposed Undersea Warfare Training Exercises in the Hawaii Range Complex from January 2007 Through January 2009*. Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.
- National Marine Fisheries Service. (2007b). *Report to Congress on the Impact of Hurricanes Katrina, Rita, and Wilman on Commercial and Recreational Fisheries Habitat of Alabama, Florida, Louisiana, Mississippi, and Texas*. Silver Spring, MD: U.S. Department of Commerce.
- National Marine Fisheries Service. (2011a). *Hawaiian Monk Seal Recovery; 2009–2010 Program Update and Accomplishments Report*. Honolulu, HI: National Oceanic and Atmospheric Administration Pacific Service; Pacific Islands Region.
- National Marine Fisheries Service. (2011b). *Southwest Region Stranding Database Excel file containing stranding from Southwest Region. Provided to Navy, Manuscript on file*. Provided to Navy, Manuscript on file.
- National Marine Fisheries Service. (2011c). Pacific Science Center Stranding Data. Excel file containing stranding from the Hawaiian Islands, manuscript on file.
- National Marine Fisheries Service. (2011d). *A Decade of Support to Save and Conserve Stranded Marine Mammals*. Retrieved from <http://www.nmfs.noaa.gov/pr/health/>.
- National Marine Fisheries Service. (2011e). *Marine Mammal Health and Stranding Response Program (MMHSRP)*. Retrieved from <https://www.fisheries.noaa.gov/national/marine-life-distress/marine-mammal-health-and-stranding-response-program>.
- National Marine Fisheries Service. (2015a). *Sperm Whale (Physeter macrocephalus) 5-Year Review: Summary and Evaluation, June 2015*. Silver Spring, MD: Office of Protected Resources.
- National Marine Fisheries Service. (2015b). *National Marine Fisheries Service Endangered Species Act Section 7 Consultation Biological Opinion and Conference Report for U.S. Military Mariana Islands Training and Testing Activities and the National Marine Fisheries Services' promulgation of regulations and issuance of a letter of authorization pursuant to the Marine Mammal Protection Act for the U.S. Navy to "take" marine mammals incidental to Mariana Islands Training and Testing activities from August 2015 through August 2020*. Silver Spring, MD: National Marine Fisheries Service.
- National Marine Fisheries Service. (2015c). *Marine Mammal Non-Lethal Deterrents: Summary of the Technical Expert Workshop on Marine Mammal Non-Lethal Deterrents, 10–12 February 2015*. Seattle, WA: National Oceanic and Atmospheric Administration.
- National Marine Fisheries Service. (2015d). *Programmatic Environmental Assessment for the Implementation of Decisions of the Western and Central Pacific Fisheries Commission on Management of Tropical Tunas in the Western and Central Ocean from 2015–2020*. Honolulu, HI: Pacific Islands Regional Office.
- National Marine Fisheries Service. (2015e). *Environmental Changes Stress West Coast Sea Lions*. Retrieved from [https://www.nwfsc.noaa.gov/news/features/west\\_coast\\_sea\\_lions/index.cfm](https://www.nwfsc.noaa.gov/news/features/west_coast_sea_lions/index.cfm).
- National Marine Fisheries Service. (2016a). *National Marine Fisheries Service, Alaska Region Occurrence of Endangered Species Act (ESA) Listed Humpback Whales off Alaska*. Silver Spring, MD: National Marine Fisheries Service.

- National Marine Fisheries Service. (2016b). *Guidelines for Preparing Stock Assessment Reports Pursuant to Section 117 of the Marine Mammal Protection Act*. Silver Spring, MD: National Oceanic and Atmospheric Administration.
- National Marine Fisheries Service. (2016c). *Fisheries Interactions and Protected Species Bycatch*. Silver Spring, MD: National Oceanic and Atmospheric Administration.
- National Marine Fisheries Service. (2016d). *FAQs: Whale, Dolphin, Seal, and Sea Lion (Marine Mammal) Strandings*. Retrieved from <http://www.nmfs.noaa.gov/pr/health/faq.htm> (accessed in June 2016).
- National Marine Fisheries Service. (2016e). Endangered and Threatened Species; Identification of 14 Distinct Population Segments of the Humpback Whale (*Megaptera novaeangliae*) and Revision of Species-Wide Listing. *Federal Register*, 81(174), 62260–62320.
- National Marine Fisheries Service. (2018a). *Proposed Rule for Western and Central Pacific Fisheries for Highly Migratory Species; Fishing Limits in Purse Seine and Longline Fisheries, Restrictions on the Use of Fish Aggregating Devices in Purse Seine Fisheries, and Transshipment Prohibitions (TIN 0648-BH77). Supplemental Information Report on National Environmental Policy Act Analyses*. Honolulu, HI: Pacific Islands Regional Office.
- National Marine Fisheries Service. (2018b). *Draft Recovery Plan for the Blue Whale (Balaenoptera musculus): Revision*. Silver Spring, MD: National Oceanic and Atmospheric Administration, Office of Protected Resources and West Coast Region.
- National Marine Fisheries Service. (2018c). *#Mlhumpbacks: Humpback Whales of the Mariana Islands*. Honolulu, HI: Pacific Islands Fisheries Science Center.
- National Oceanic and Atmospheric Administration. (2002). *Report of the Workshop on acoustic resonance as a source of tissue trauma in cetaceans*. Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.
- National Oceanic and Atmospheric Administration. (2015a). Endangered and Threatened Species; Identification of 14 Distinct Population Segments of the Humpback Whale (*Megaptera novaeangliae*) and Proposed Revision of Species-Wide Listing; Proposed Rule. *Federal Register*, 80(76), 22304–22356.
- National Oceanic and Atmospheric Administration. (2015b). Takes of marine mammals incidental to specified activities; U.S. Navy training and testing activities in the Mariana Islands Training and Testing Study Area. *Federal Register*, 80(148), 46112–46171.
- National Oceanic and Atmospheric Administration. (2016). *Testing Detects Algal Toxins in Alaska Marine Mammals*. Retrieved from [http://www.nwfsc.noaa.gov/news/features/algal\\_blooms\\_in\\_arctic\\_waters/index.cfm](http://www.nwfsc.noaa.gov/news/features/algal_blooms_in_arctic_waters/index.cfm).
- National Oceanic and Atmospheric Administration. (2017). *2016 West Coast Entanglement Summary*. Seattle, WA: National Marine Fisheries Service.
- National Oceanic and Atmospheric Administration. (2018a). *2015–2018 Guadalupe Fur Seal Unusual Mortality Event in California*. Retrieved from <https://www.fisheries.noaa.gov/national/marine-life-distress/2015-2018-guadalupe-fur-seal-unusual-mortality-event-california>.
- National Oceanic and Atmospheric Administration. (2018b). International Fisheries; Western and Central Pacific Fisheries for Highly Migratory Species; Fishing Limits in Purse Seine and Longline

- Fisheries, Restrictions on the Use of Fish Aggregating Devices in Purse Seine Fisheries, and Transshipment Prohibitions. *Federal Register*, 83(91), 21748–21761.
- National Oceanic and Atmospheric Administration. (2018c). #MIhumpbacks: Humpback Whales of the Mariana Islands. Retrieved from <https://www.fisheries.noaa.gov/feature-story/mihumpbacks-humpback-whales-mariana-islands>.
- National Oceanic and Atmospheric Administration Fisheries. (2014). *Southern Resident Killer Whales: 10 Years of Research & Conservation*. Seattle, WA: Northwest Fisheries Science Center West Coast Region.
- National Oceanic and Atmospheric Administration Fisheries. (2018). *Removal and Research: The Marine Debris Team Strikes Again*. Retrieved from <https://www.fisheries.noaa.gov/feature-story/removal-and-research-marine-debris-team-strikes-again>.
- National Oceanic and Atmospheric Administration Marine Debris Program. (2014a). *Report on the Entanglement of Marine Species in Marine Debris with an Emphasis on Species in the United States*. Silver Spring, MD: National Oceanic and Atmospheric Administration.
- National Oceanic and Atmospheric Administration Marine Debris Program. (2014b). *Report on the Occurrence of Health Effects of Anthropogenic Debris Ingested by Marine Organisms*. Silver Spring, MD: National Ocean Service.
- National Research Council. (2003). *Ocean Noise and Marine Mammals*. Washington, DC: The National Academies Press.
- National Research Council. (2005). *Marine Mammal Populations and Ocean Noise*. Washington, DC: The National Academies Press.
- National Research Council. (2006). *Dynamic Changes in Marine Ecosystems: Fishing, Food Webs, and Future Options, Committee on Ecosystem Effects of Fishing: Phase II—Assessments of the Extent of Change and the Implications for Policy*. Washington, DC: National Research Council.
- Nedwell, J. R., B. Edwards, A. W. H. Turnpenny, and J. Gordon. (2004). *Fish and Marine Mammal Audiograms: A Summary of Available Information*. Hampshire, United Kingdom: Subacoustech Ltd.
- Nemoto, T., and A. Kawamura. (1977). Characteristics of food habits and distribution of baleen whales with special reference to the abundance of North Pacific sei and Bryde's whales. *Reports of the International Whaling Commission, Special Issue 1*, 80–87.
- New, L. F., J. Harwood, L. Thomas, C. Donovan, J. S. Clark, G. Hastie, P. M. Thompson, B. Cheney, L. Scott-Hayward, D. Lusseau, and D. Costa. (2013a). Modelling the biological significance of behavioural change in coastal bottlenose dolphins in response to disturbance. *Functional Ecology*, 27(2), 314–322.
- New, L. F., D. J. Moretti, S. K. Hooker, D. P. Costa, and S. E. Simmons. (2013b). Using energetic models to investigate the survival and reproduction of beaked whales (family Ziphiidae). *PLoS ONE*, 8(7), e68725.
- New, L. F., D. J. Morretti, S. K. Hooker, D. P. Costa, and S. E. Simmons. (2013c). Using Energetic Models to Investigate the Survival and Reproduction of Beaked Whales (family Ziphiidae). *PLoSOne*, 8(7), e68725.
- New, L. F., J. S. Clark, D. P. Costa, E. Fleishman, M. A. Hindell, T. Klanjšček, D. Lusseau, S. Kraus, C. R. McMahon, P. W. Robinson, R. S. Schick, L. K. Schwarz, S. E. Simmons, L. Thomas, P. Tyack, and J.

- Harwood. (2014). Using short-term measures of behaviour to estimate long-term fitness of southern elephant seals. *Marine Ecology Progress Series*, 496, 99–108.
- Ng, S. L., and S. Leung. (2003). Behavioral response of Indo-Pacific humpback dolphin (*Sousa chinensis*) to vessel traffic. *Marine Environmental Research*, 56(5), 555–567.
- Nieukirk, S. L., D. K. Mellinger, S. E. Moore, K. Klinck, R. P. Dziak, and J. Goslin. (2012). Sounds from airguns and fin whales recorded in the mid-Atlantic Ocean, 1999–2009. *The Journal of the Acoustical Society of America*, 131(2), 1102–1112.
- Nieukirk, S. L., S. Fregosi, D. K. Mellinger, and H. Klinck. (2016). A complex baleen whale call recorded in the Mariana Trench Marine National Monument. *The Journal of the Acoustical Society of America*, 140(3), EL274.
- Noren, D. P., A. H. Johnson, D. Rehder, and A. Larson. (2009). Close approaches by vessels elicit surface active behaviors by southern resident killer whales. *Endangered Species Research*, 8(3), 179–192.
- Norris, K. S., and J. H. Prescott. (1961). Observations on Pacific cetaceans of Californian and Mexican waters. *University of California Publications in Zoology*, 63(4), 291–402.
- Norris, K. S., B. Wursig, R. S. Wells, and M. Würsig. (1994). *The Hawaiian Spinner Dolphin*. Berkeley, CA: University of California Press.
- Norris, T. F., J. Oswald, T. Yack, E. Ferguson, C. Hom-Weaver, K. Dunleavy, S. Coates, and T. Dominello. (2012a). *An Analysis of Acoustic Data from the Mariana Islands Sea Turtle and Cetacean Survey (MISTCS)*. Encinitas, CA: Bio-Waves, Inc.
- Norris, T. F., J. O. Oswald, T. M. Yack, and E. L. Ferguson. (2012b). *An Analysis of Marine Acoustic Recording Unit (MARU) Data Collected off Jacksonville, Florida in Fall 2009 and Winter 2009–2010*. Norfolk, VA: Naval Facilities Engineering Command Atlantic.
- Norris, T. F., J. Oswald, T. Yack, E. Ferguson, C. Hom-Weaver, K. Dunleavy, S. Coates, and T. Dominello. (2014). *An Analysis of Acoustic Data from the Mariana Islands Sea Turtle and Cetacean Survey (MISTCS) March 2014 Revision*. Encinitas, CA: Bio-Waves, Inc.
- Norris, T. F., K. J. Dunleavy, T. M. Yack, and E. L. Ferguson. (2017). Estimation of minke whale abundance from an acoustic line transect survey of the Mariana Islands. *Marine Mammal Science*, 33(2), 574–592.
- Northridge, S. (2009). Fishing industry, effects of. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 443–447). Cambridge, MA: Academic Press.
- Nowacek, D., M. Johnson, and P. Tyack. (2004). North Atlantic right whales (*Eubalaena glacialis*) ignore ships but respond to alerting stimuli. *Proceedings of the Royal Society of London*, 271(B), 227–231.
- Nowacek, D., L. H. Thorne, D. Johnston, and P. Tyack. (2007a). Responses of cetaceans to anthropogenic noise. *Mammal Review*, 37(2), 81–115.
- Nowacek, D. P., L. H. Thorne, D. W. Johnston, and P. L. Tyack. (2007b). Responses of cetaceans to anthropogenic noise. *Mammal Review*, 37(2), 81–115.
- Nowacek, D. P., C. W. Clark, D. Mann, P. J. O. Miller, H. C. Rosenbaum, J. S. Golden, M. Jasny, J. Kraska, and B. L. Southall. (2015). Marine seismic surveys and ocean noise: Time for coordinated and prudent planning. *Frontiers in Ecology and the Environment*, 13(7), 378–386.

- Nowacek, D. P., F. Christiansen, L. Bejder, J. A. Goldbogen, and A. S. Friedlaender. (2016). Studying cetacean behaviour: New technological approaches and conservation applications. *Animal Behaviour*, 120, 235–244.
- Oakley, J. A., A. T. Williams, and T. Thomas. (2017). Reactions of harbour porpoise (*Phocoena phocoena*) to vessel traffic in the coastal waters of South West Wales, UK. *Ocean & Coastal Management*, 138, 158–169.
- Obusan, M. C. M., W. L. Rivera, M. A. T. Siringan, and L. V. Aragonés. (2016). Stranding events in the Philippines provide evidence for impacts of human interactions on cetaceans. *Ocean & Coastal Management*, 134, 41–51.
- Ocean Alliance. (2010). *The Voyage of the Odyssey: Executive Summary*. Lincoln, MA: Public Broadcasting System.
- Odell, D. K., and K. M. McClune. (1999). False killer whale—*Pseudorca crassidens* (Owen, 1846). In S. H. Ridgway & S. R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 6, pp. 213–244). New York, NY: Academic Press.
- Office of the Surgeon General. (1991). Conventional warfare ballistic, blast, and burn injuries. In R. Zajitchuk, Col. (Ed.), *U.S.A. Textbook of Military Medicine*. Washington, DC: Office of the Surgeon General.
- Oleson, E. (2017). *Mariana Archipelago Cetacean Survey (MACS) 2015 Cruise Report*. Honolulu, HI: National Marine Fisheries Service, Pacific Islands Fisheries Science Center.
- Oleson, E. M., C. H. Boggs, K. A. Forney, M. B. Hanson, D. R. Kobayashi, B. L. Taylor, P. R. Wade, and G. M. Ylitalo. (2010). *Status Review of Hawaiian Insular False Killer Whales (Pseudorca crassidens) under the Endangered Species Act* (NOAA Technical Memorandum NMFS-PIFSC-22). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Oleson, E. M., and M. C. Hill. (2010a). *2010 Report to PACFLT: Report of Cetacean Surveys in Guam, CNMI, and the High-seas & Follow up on 2009 Main Hawaiian Islands Cetacean Survey*. Honolulu, HI: Pacific Islands Fisheries Science Center.
- Oleson, E. M., and M. C. Hill. (2010b). *2010 Report to PACFLT: Report to Cetacean Surveys in Guam, CNMI, and the High-seas*. Honolulu, HI: National Marine Fisheries Service, Pacific Islands Fisheries Science Center.
- Oleson, E. M., R. W. Baird, K. K. Martien, and B. L. Taylor. (2013). *Island-associated stocks of odontocetes in the main Hawaiian Islands: A synthesis of available information to facilitate evaluation of stock structure* (Pacific Islands Fisheries Science Center Working Paper WP-13-003). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Oleson, E. M., S. Baumann-Pickering, A. Širović, K. P. Merckens, L. M. Munger, J. S. Trickey, and P. Fisher-Pool. (2015). *Analysis of long-term acoustic datasets for baleen whales and beaked whales within the Mariana Islands Range Complex (MIRC) for 2010 to 2013* (Pacific Islands Fisheries Science Center Data Report DR-15-002). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Olson, P. A. (2009). Pilot whales, *Globicephala melas* and *G. macrorhynchus*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 898–903). Cambridge, MA: Academic Press.

- Oregon State University. (2017). *Southern and Central California 2016 Whale Approach Summary from Bruce Mate regarding body condition of blue and fin whales off Southern and Central California*. Corvallis, OR: Oregon State University.
- Owen, M. A., and A. E. Bowles. (2011). In-air auditory psychophysics and the management of a threatened carnivore, the polar bear (*Ursus maritimus*). *International Journal of Comparative Psychology*, 24, 244–254.
- Pack, A. A., L. M. Herman, A. S. Craig, S. S. Spitz, J. O. Waterman, E. Y. K. Herman, M. H. Deakos, S. Hakala, and C. Lowe. (2017). Habitat preferences by individual humpback whale mothers in the Hawaiian breeding grounds vary with the age and size of their calves. *Animal Behaviour*, 133, 131–144.
- Palacios, D. M., and B. R. Mate. (1996). Attack by false killer whales (*Pseudorca crassidens*) on sperm whales (*Physeter macrocephalus*) in the Galápagos Islands. *Marine Mammal Science*, 12(4), 582–587.
- Paniz-Mondolfi, A. E., and L. Sander-Hoffmann. (2009). Lobomycosis in inshore and estuarine dolphins. *Emerging Infectious Diseases*, 15(4), 672–673.
- Papale, E., M. Gamba, M. Perez-Gil, V. M. Martin, and C. Giacomini. (2015). Dolphins adjust species-specific frequency parameters to compensate for increasing background noise. *PLoS ONE*, 10(4), e0121711.
- Parks, S. E., C. W. Clark, and P. L. Tyack. (2007). Short- and long-term changes in right whale calling behavior: The potential effects of noise on acoustic communication. *The Journal of the Acoustical Society of America*, 122(6), 3725–3731.
- Parks, S. E. (2009). *Assessment of acoustic adaptations for noise compensation in marine mammals*. Paper presented at the 2009 Office of Naval Research Marine Mammal Program Review. Alexandria, VA.
- Parks, S. E., M. Johnson, D. Nowacek, and P. L. Tyack. (2011). Individual right whales call louder in increased environmental noise. *Biology Letters*, 7, 33–35.
- Patenaude, N. J., W. J. Richardson, M. A. Smultea, W. R. Koski, G. W. Miller, B. Würsig, and C. R. Greene, Jr. (2002). Aircraft sound and disturbance to bowhead and beluga whales during spring migration in the Alaskan Beaufort Sea. *Marine Mammal Science*, 18(2), 309–335.
- Pavlostathis, S. G., and G. H. Jackson. (2002). Biotransformation of 2, 4, 6-trinitrotoluene in a continuous-flow *Anabaena* sp. system. *Water Research*, 36, 1699–1706.
- Pepper, C. B., M. A. Nascarella, and R. J. Kendall. (2003). A review of the effects of aircraft noise on wildlife and humans, current control mechanisms, and the need for further study. *Environmental Management*, 32(4), 418–432.
- Perkins, J. S., and G. W. Miller. (1983). Mass stranding of *Steno bredanensis* in Belize. *Biotropica*, 15(3), 235–236.
- Perrin, W. F., and J. W. Gilpatrick, Jr. (1994). Spinner dolphin, *Stenella longirostris* (Gray, 1828). In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 99–128). San Diego, CA: Academic Press.
- Perrin, W. F., and A. A. Hohn. (1994). Pantropical spotted dolphin, *Stenella attenuata*. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 71–98). San Diego, CA: Academic Press.

- Perrin, W. F., S. Leatherwood, and A. Collet. (1994a). Fraser's dolphin, *Lagenodelphis hosei* Fraser, 1956. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 225–240). San Diego, CA: Academic Press.
- Perrin, W. F., C. E. Wilson, and F. I. Archer, II. (1994b). Striped dolphin—*Stenella coeruleoalba* (Meyen, 1833). In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 129–159). San Diego, CA: Academic Press.
- Perrin, W. F., and J. R. Geraci. (2002). Stranding. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (pp. 1192–1197). San Diego, CA: Academic Press.
- Perrin, W. F. (2009a). Spinner dolphin, *Stenella longirostris*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 1100–1103). Cambridge, MA: Academic Press.
- Perrin, W. F. (2009b). Pantropical spotted dolphin, *Stenella attenuata*. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 819–821). Cambridge, MA: Academic Press.
- Perrin, W. F., B. Würsig, and J. G. M. Thewissen. (2009). *Encyclopedia of Marine Mammals* (2nd ed.). Cambridge, MA: Academic Press.
- Perryman, W. L., and T. C. Foster. (1980). *Preliminary Report on Predation by Small Whales, Mainly the False Killer Whale, Pseudorca crassidens, on Dolphins (Stenella spp. and Delphinus delphis) in the Eastern Tropical Pacific*. (LJ-80-05). La Jolla, CA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.
- Peterson, S. H., J. L. Hassrick, A. Lafontaine, J. P. Thome, D. E. Crocker, C. Debier, and D. P. Costa. (2014). Effects of age, adipose percent, and reproduction on PCB concentrations and profiles in an extreme fasting North Pacific marine mammal. *PLoS ONE*, 9(4), e96191.
- Peterson, S. H., J. T. Ackerman, and D. P. Costa. (2015). Marine foraging ecology influences mercury bioaccumulation in deep-diving northern elephant seals. *Proceedings of the Royal Society B: Biological Sciences*, 282(20150710), 10.
- Piantadosi, C. A., and E. D. Thalmann. (2004). Whales, sonar and decompression sickness. *Nature*, 425, 575–576.
- Pine, M. K., A. G. Jeffs, D. Wang, and C. A. Radford. (2016). The potential for vessel noise to mask biologically important sounds within ecologically significant embayments. *Ocean & Coastal Management*, 127, 63–73.
- Pirotta, E., R. Milor, N. Quick, D. Moretti, N. Di Marzio, P. Tyack, I. Boyd, and G. Hastie. (2012). Vessel noise affects beaked whale behavior: Results of a dedicated acoustic response study. *PLoS ONE*, 7(8), e42535.
- Pirotta, E., K. L. Brookes, I. M. Graham, and P. M. Thompson. (2014). Variation in harbour porpoise activity in response to seismic survey noise. *Biology Letters*, 10(5), 20131090.
- Pirotta, E., J. Harwood, P. M. Thompson, L. New, B. Cheney, M. Arso, P. S. Hammond, C. Donovan, and D. Lusseau. (2015a). Predicting the effects of human developments on individual dolphins to understand potential long-term population consequences. *Proceedings of the Royal Society B: Biological Sciences*, 282(1818), 20152109.

- Pirotta, E., N. D. Merchant, P. M. Thompson, T. R. Barton, and D. Lusseau. (2015b). Quantifying the effect of boat disturbance on bottlenose dolphin foraging activity. *Biological Conservation*, 181, 82–89.
- Pirotta, E., M. Mangel, D. P. Costa, B. Mate, J. A. Goldbogen, D. M. Palacios, L. A. Hückstädt, E. A. McHuron, L. Schwarz, and L. New. (2018). A Dynamic State Model of Migratory Behavior and Physiology to Assess the Consequences of Environmental Variation and Anthropogenic Disturbance on Marine Vertebrates. *The American Naturalist*, 191(2), 17.
- Pirotta, V., D. Slip, I. D. Jonsen, V. M. Peddemors, D. H. Cato, G. Ross, and R. Harcourt. (2016). Migrating humpback whales show no detectable response to whale alarms off Sydney, Australia. *Endangered Species Research*, 29(3), 201–209.
- Piscitelli, M. A., W. A. McLellan, A. S. Rommel, J. E. Blum, S. G. Barco, and D. A. Pabst. (2010). Lung size and thoracic morphology in shallow and deep-diving cetaceans. *Journal of Morphology*, 271, 654–673.
- Pitman, R. (2009). Mesoplodont whales (*Mesoplodon spp.*). In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 721–726). Cambridge, MA: Academic Press.
- Pitman, R. L., D. W. K. Au, M. D. Scott, and J. M. Cotton. (1988). *Observations of Beaked Whales (Ziphiidae) from the Eastern Tropical Pacific Ocean*. Cambridge, United Kingdom: International Whaling Commission.
- Pitman, R. L., L. T. Ballance, S. L. Mesnick, and S. J. Chivers. (2001). Killer whale predation on sperm whales: Observations and implications. *Marine Mammal Science*, 17(3), 494–507.
- Pitman, R. L., and C. Stinchcomb. (2002). Rough-toothed dolphins (*Steno bredanensis*) as predators of mahimahi (*Coryphaena hippurus*). *Pacific Science*, 56(4), 447–450.
- Pitman, R. L., H. Fearnbach, R. LeDuc, J. W. Gilpatrick, Jr., J. K. B. Ford, and L. T. Ballance. (2007). Killer whales preying on a blue whale calf on the Costa Rica Dome: Genetics, morphometrics, vocalisations and composition of the group. *Journal of Cetacean Research and Management*, 9(2), 151–157.
- Polacheck, T., and L. Thorpe. (1990). The swimming direction of harbor porpoise in relationship to a survey vessel. *Reports of the International Whaling Commission*, 40, 463–470.
- Poloczanska, E. S., M. T. Burrows, C. J. Brown, J. G. Molinos, B. S. Halpern, O. Hoegh-Guldberg, C. V. Kappel, P. J. Moore, A. J. Richardson, D. S. Schoeman, and W. J. Sydeman. (2016). Responses of marine organisms to climate change across oceans. *Frontiers in Marine Science*, 3(62), 1–21.
- Popov, V. V., A. Y. Supin, D. Wang, K. Wang, L. Dong, and S. Wang. (2011). Noise-induced temporary threshold shift and recovery in Yangtze finless porpoises, *Neophocaena phocaenoides asiaeorientalis*. *The Journal of the Acoustical Society of America*, 130(1), 574–584.
- Popov, V. V., A. Y. Supin, V. V. Rozhnov, D. I. Nechaev, E. V. Sysuyeva, V. O. Klishin, M. G. Pletenko, and M. B. Tarakanov. (2013). Hearing threshold shifts and recovery after noise exposure in beluga whales, *Delphinapterus leucas*. *The Journal of Experimental Biology*, 216(9), 1587–1596.
- Popov, V. V., A. Y. Supin, V. V. Rozhnov, D. I. Nechaev, and E. V. Sysueva. (2014). The limits of applicability of the sound exposure level (SEL) metric to temporal threshold shifts (TTS) in beluga whales, *Delphinapterus leucas*. *The Journal of Experimental Biology*, 217(Pt 10), 1804–1810.

- Potter, J. R., M. Thillet, C. Douglas, M. A. Chitre, Z. Doborzynski, and P. J. Seekings. (2007). Visual and passive acoustic marine mammal observations and high-frequency seismic source characteristics recorded during a seismic survey. *IEEE Journal of Oceanic Engineering*, 32(2), 469–483.
- Puig-Lozano, R., Y. Bernaldo de Quirós, J. Díaz-Delgado, N. García-Álvarez, E. Sierra, J. De la Fuente, S. Sacchini, C. M. Suárez-Santana, D. Zucca, N. Câmara, P. Saavedra, J. Almunia, M. A. Rivero, A. Fernáandez, and M. Arbelo. (2018). Retrospective study of foreign body-associated pathology in stranded cetaceans, Canary Islands (2000–2015). *Environmental Pollution*, 243, 519–527.
- Quick, N., L. Scott-Hayward, D. Sadykova, D. Nowacek, and A. Read. (2017). Effects of a scientific echo sounder on the behavior of short-finned pilot whales (*Globicephala macrorhynchus*). *Canadian Journal of Fisheries and Aquatic Sciences*, 74(5), 716–726.
- Ramos, E. A., B. Maloney, M. O. Magnasco, and D. Reiss. (2018). Bottlenose dolphins and Antillean manatees respond to small multi-rotor unmanned aerial systems. *Frontiers in Marine Science*, 5, 316.
- Ramp, C., J. Delarue, P. J. Palsboll, R. Sears, and P. S. Hammond. (2015). Adapting to a warmer ocean—Seasonal shift of baleen whale movements over three decades. *PLoS ONE*, 10(3), e0121374.
- Read, A., P. Drinker, and S. Northridge. (2006). Bycatch of marine mammals in U.S. and global fisheries. *Conservation Biology*, 20(1), 163–169.
- Read, A. J. (2008). The looming crisis: Interactions between marine mammals and fisheries. *Journal of Mammalogy*, 89(3), 541–548.
- Redfern, J. V., L. T. Hatch, C. Caldow, M. L. DeAngelis, J. Gedamke, S. Hastings, L. Henderson, M. F. McKenna, T. J. Moore, and M. B. Porter. (2017). Assessing the risk of chronic shipping noise to baleen whales off Southern California, USA. *Endangered Species Research*, 32, 153–167.
- Reeves, R. R., G. K. Silber, and P. M. Payne. (1998). *Draft recovery plan for the fin whale Balaenoptera physalus and sei whale Balaenoptera borealis*. Silver Spring, MD: Office of Protected Resources, National Marine Fisheries Service, National Oceanic and Atmospheric Administration.
- Reeves, R. R., B. S. Stewart, P. J. Clapham, and J. A. Powell. (2002). *National Audubon Society Guide to Marine Mammals of the World*. New York, NY: Alfred A. Knopf.
- Reeves, R. R., S. Leatherwood, and R. W. Baird. (2009). Evidence of a possible decline since 1989 in false killer whales (*Pseudorca crassidens*) around the main Hawaiian islands. *Pacific Science*, 63(2), 253–261.
- Reichmuth, C., M. M. Holt, J. Mulsow, J. M. Sills, and B. L. Southall. (2013). Comparative assessment of amphibious hearing in pinnipeds. *Journal of Comparative Physiology A: Neuroethology, Sensory Neural, and Behavioral Physiology*, 199(6), 491–507.
- Reilly, S. B. (1990). Seasonal changes in distribution and habitat differences among dolphins in the eastern tropical Pacific. *Marine Ecology Progress Series*, 66, 1–11.
- Rice, A. C., S. Baumann-Pickering, A. Širović, J. A. Hildebrand, A. M. Brewer, A. J. Debich, S. T. Herbert, B. J. Thayre, J. S. Trickey, and S. M. Wiggins. (2015). *Passive Acoustic Monitoring for Marine Mammals in the Gulf of Alaska Temporary Maritime Activities Area 2014-2015*. (W9126G-14-2-0040). La Jolla, CA: Whale Acoustics Laboratory, Marine Physical Laboratory, Scripps Institution of Oceanography.
- Rice, D. W. (1989). Sperm whale *Physeter macrocephalus* Linnaeus, 1758. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 4, pp. 177–234). San Diego, CA: Academic Press.

- Rice, D. W. (1998). *Marine Mammals of the World: Systematics and Distribution* (Society for Marine Mammalogy Special Publication). Lawrence, KS: Society for Marine Mammalogy.
- Richardson, K., D. Haynes, A. Talouli, and M. Donoghue. (2016). Marine pollution originating from purse seine and longline fishing vessel operations in the Western and Central Pacific Ocean, 2003–2015. *Ambio*, 46(2), 190–200.
- Richardson, W. J., M. A. Fraker, B. Würsig, and R. S. Wells. (1985). Behaviour of bowhead whales (*Balaena mysticetus*) summering in the Beaufort Sea: Reactions to industrial activities. *Biological Conservation*, 32, 195–230.
- Richardson, W. J., C. R. Greene, Jr., J. S. Hanna, W. R. Koski, G. W. Miller, N. J. Patenaude, and M. A. Smultea. (1995a). *Acoustic Effects of Oil Production Activities on Bowhead and White Whales Visible during Spring Migration near Pt. Barrow, Alaska – 1991 and 1994 Phases: Sound Propagation and Whale Responses to Playbacks of Icebreaker Noise*. Anchorage, AK: U.S. Minerals Management Service, Procurement Operations.
- Richardson, W. J., C. R. Greene, Jr., C. I. Malme, and D. H. Thomson. (1995b). *Marine Mammals and Noise*. San Diego, CA: Academic Press.
- Richardson, W. J., G. W. Miller, and C. R. Greene. (1999). Displacement of migrating bowhead whales by sounds from seismic surveys in shallow waters of the Beaufort Sea. *The Journal of the Acoustical Society of America*, 106(4), 2281.
- Richmond, D. R., J. T. Yelverton, and E. R. Fletcher. (1973). *Far-Field Underwater-Blast Injuries Produced by Small Charges*. Washington, DC: Lovelace Foundation for Medical Education and Research, Defense Nuclear Agency.
- Richter, C., S. M. Dawson, and E. Slooten. (2003). Sperm whale watching off Kaikoura, New Zealand: Effects of current activities on surfacing and vocalisation patterns. *Science for Conservation*, 219, 78.
- Richter, C., S. Dawson, and E. Slooten. (2006). Impacts of commercial whale watching on male sperm whales at Kaikoura, New Zealand. *Marine Mammal Science*, 22(1), 46–63.
- Ridgway, S. H. (1972). Homeostasis in the Aquatic Environment. In S. H. Ridgway (Ed.), *Mammals of the Sea: Biology and Medicine* (pp. 590–747). Springfield, IL: Charles C. Thomas.
- Ridgway, S. H., and R. Howard. (1979). Dolphin lung collapse and intramuscular circulation during free diving: Evidence from nitrogen washout. *Science*, 206, 1182–1183.
- Ridgway, S. H., D. A. Carder, R. R. Smith, T. Kamolnick, C. E. Schlundt, and W. R. Elsberry. (1997). *Behavioral Responses and Temporary Shift in Masked Hearing Threshold of Bottlenose Dolphins, Tursiops truncatus, to 1-second Tones of 141 to 201 dB re 1  $\mu$ Pa*. (Technical Report 1751, Revision 1). San Diego, CA: U.S. Department of Navy, Naval Command, Control and Ocean Surveillance Center, Research, Development, Test, and Evaluation Division.
- Risch, D., P. J. Corkeron, W. T. Ellison, and S. M. Van Parijs. (2012). Changes in humpback whale song occurrence in response to an acoustic source 200 km away. *PLoS ONE*, 7(1), e29741.
- Risch, D., P. J. Corkeron, W. T. Ellison, and S. M. Van Parijs. (2014). Formal comment to Gong et al.: Ecosystem scale acoustic sensing reveals humpback whale behavior synchronous with herring spawning processes and re-evaluation finds no effect of sonar on humpback song occurrence in the Gulf of Maine in fall 2006. *PLoS ONE*, 9(10), e109225.

- Ritter, F. (2002). Behavioural observations of rough-toothed dolphins (*Steno bredanensis*) off La Gomera, Canary Islands (1995–2000), with special reference to their interactions with humans. *Aquatic Mammals*, 28(1), 46–59.
- Robertson, F. C. (2014). *Effects of Seismic Operations on Bowhead Whale Behavior: Implications for Distribution and Abundance Assessments*. (Unpublished doctoral dissertation). The University of British Columbia, Vancouver, Canada. Retrieved from [http://www.marinemammal.org/wp-content/pdfs/Robertson\\_2014.pdf](http://www.marinemammal.org/wp-content/pdfs/Robertson_2014.pdf).
- Rocha, R. C., P. J. J. Clapham, and Y. V. Ivashchenko. (2014). Emptying the Oceans: A Summary of Industrial Whaling Catches in the 20th Century. *Marine Fisheries Review*, 76(4), 37–48.
- Rock, T. (1993, April 12). Killer whales of the tropics. *Pacific Daily News*, p. 19.
- Rockwood, R. C., J. Calambokidis, and J. Jahncke. (2017). High mortality of blue, humpback and fin whales from modeling of vessel collisions on the U.S. West Coast suggests population impacts and insufficient protection. *PLoS ONE*, 12(8), e0183052.
- Rogers, K. S. (2016). Feral cats are a serious threat to Hawaii's endangered birds, monk seals and dolphins. *Honolulu Magazine*. Retrieved from <http://www.honolulumagazine.com/Honolulu-Magazine/October-2016/Cats-vs-Birds-and-Everyone-Else/>.
- Rolland, R. M., S. E. Parks, K. E. Hunt, M. Castellote, P. J. Corkeron, D. P. Nowacek, S. K. Wasser, and S. D. Kraus. (2012). Evidence that ship noise increases stress in right whales. *Proceedings of the Royal Society B: Biological Sciences*, 279(1737), 2363–2368.
- Roman, J., I. Altman, M. M. Dunphy-Daly, C. Campbell, M. Jasny, and A. J. Read. (2013). The Marine Mammal Protection Act at 40: Status, Recovery, and Future of U.S. Marine Mammals. *Annals of the New York Academy of Sciences*, 1286, 29–49.
- Romano, T. A., M. J. Keogh, C. Kelly, P. Feng, L. Berk, C. E. Schlundt, D. A. Carder, and J. J. Finneran. (2004). Anthropogenic sound and marine mammal health: Measures of the nervous and immune systems before and after intense sound exposures. *Canadian Journal of Fisheries and Aquatic Sciences*, 61, 1124–1134.
- Rosel, P. E., and H. Watts. (2008). Hurricane impacts on bottlenose dolphins in the northern Gulf of Mexico. *Gulf of Mexico Science*, 25(1), 88–94.
- Rosen, G., and G. R. Lotufo. (2010). Fate and effects of composition B in multispecies marine exposures. *Environmental Toxicology and Chemistry*, 29(6), 1330–1337.
- Rosen, Y. (2015). More whales found dead in southern Alaska waters. *Alaska Dispatch News*. Retrieved from <http://www.adn.com/article/20150710/more-whales-found-dead-southern-alaska-waters>.
- Rosowski, J. J. (1994). Outer and Middle Ears. In R. R. Fay & A. N. Popper (Eds.), *Comparative Hearing: Mammals* (pp. 172–247). Berlin, Germany: Springer-Verlag.
- Ross, G. J. B., and A. J. Bass. (1971). Shark attack on an ailing dolphin *Stenella coeruleoalba* (Meyen). *South African Journal of Science*, 67, 413–414.
- Ross, G. J. B., and S. Leatherwood. (1994). Pygmy killer whale *Feresa attenuata* Gray, 1874. In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 5, pp. 387–404). San Diego, CA: Academic Press.
- Rossmann, S., E. B. McCabe, N. B. Barros, H. G. Gandhi, P. H. Ostrom, C. A. Stricker, and R. S. Wells. (2015). Foraging habits in a generalist predator: Sex and age influence habitat selection and

- resource use among bottlenose dolphins (*Tursiops truncatus*). *Marine Mammal Science*, 31(1), 155–168.
- Sabet, S. S., K. Wesdorp, J. Campbell, P. Snelderwaard, and H. Slabbekoorn. (2016). Behavioural responses to sound exposure in captivity by two fish species with different hearing ability. *Animal Behaviour*, 116, 1–11.
- Sadove, S. S., and S. J. Morreale. (1989). *Marine Mammal and Sea Turtle Encounters with Marine Debris in the New York Bight and the Northeast Atlantic*. Paper presented at the Proceedings of the Second International Conference on Marine Debris. Honolulu, HI.
- Saez, L., D. Lawson, M. DeAngelis, S. Wilkin, E. Petras, and C. Fahy. (2012). *Co-occurrence of Large Whales and Fixed Commercial Fishing Gear: California, Oregon, and Washington*. Paper presented at the Southern California Marine Mammal Workshop. Newport Beach, CA.
- Saez, L., D. Lawson, M. DeAngelis, E. Petras, S. Wilkin, and C. Fahy. (2013). *Understanding the Co-Occurrence of Large Whales and Commercial Fixed Gear Fisheries Off the West Coast of the United States* (NOAA Technical Memorandum NMFS-SWR-044). Long Beach, CA: Southwest Regional Office, Protected Resources Division.
- Saipan Tribune. (2011). *Necropsy found abnormalities in kidneys, intestines of beached whale*. Retrieved from <http://www.saipantribune.com/newsstory.aspx?newsID=112203>
- Sairanen, E. E. (2014). *Weather and Ship Induced Sounds and the Effect of Shipping on Harbor Porpoise (Phocoena phocoena) Activity*. (Unpublished master's thesis). University of Helsinki, Helsinki, Finland. Retrieved from [helda.helsinki.fi](http://helda.helsinki.fi).
- Salvadeo, C. J., D. Lluch-Belda, A. Gómez-Gallardo, J. Urbán-Ramírez, and C. D. MacLeod. (2010). Climate change and a poleward shift in the distribution of the Pacific white-sided dolphin in the northeastern Pacific. *Endangered Species Research*, 11, 13–19.
- Salvadeo, C. J., A. Gomez-Gallardo U., M. Najera-Caballero, J. Urban-Ramirez, and D. Lluch-Belda. (2015). The effect of climate variability on gray whales (*Eschrichtius robustus*) within their wintering areas. *PLoS ONE*, 10(8), 1–17.
- Santos, M. B., V. Martin, M. Arbelo, A. Fernandez, and G. J. Pierce. (2007). Insights into the diet of beaked whales from the atypical mass strandings in the Canary Islands in September 2002. *Journal of the Marine Biological Association of the United Kingdom*, 87, 243–251.
- Saunders, K. J., P. R. White, and T. G. Leighton. (2008). Models for Predicting Nitrogen Tensions and Decompression Sickness Risk in Diving Beaked Whales. *Proceedings of the Institute of Acoustics*, 30(5), 1–8.
- Savage, K., D. Fauquier, S. Raverty, K. B. Huntington, J. Moran, M. Migura, P. Cottrell, K. Wynne, B. Witteveen, and F. Van Dolah. (2017). *Abstract: 2015 Gulf of Alaska Large Whale Unusual Mortality Event*. Paper presented at the Kodiak Area Marine Science Symposium April 18–21, 2017. Kodiak, AK. Retrieved from <https://seagrant.uaf.edu/events/2017/kamss/kamssprog-2017.pdf>.
- Scarpaci, C., S. W. Bigger, P. J. Corkeron, and D. Nugegoda. (2000). Bottlenose dolphins (*Tursiops truncatus*) increase whistling in the presence of 'swim-with-dolphin' tour operations. *Journal of Cetacean Research and Management*, 2(3), 183–185.
- Schakner, Z. A., and D. T. Blumstein. (2013). Behavioral biology of marine mammal deterrents: A review and prospectus. *Biological Conservation*, 167, 380–389.

- Scheifele, P. M., S. Andrew, R. A. Cooper, M. Darre, R. E. Musiek, and L. Max. (2005). Indication of a Lombard vocal response in the St. Lawrence River beluga. *The Journal of the Acoustical Society of America*, 117(3), 1486–1492.
- Schlundt, C. E., J. J. Finneran, D. A. Carder, and S. H. Ridgway. (2000). Temporary shift in masked hearing thresholds of bottlenose dolphins, *Tursiops truncatus*, and white whales, *Delphinapterus leucas*, after exposure to intense tones. *The Journal of the Acoustical Society of America*, 107(6), 3496–3508.
- Schlundt, C. E., R. L. Dear, L. Green, D. S. Houser, and J. J. Finneran. (2007). Simultaneously measured behavioral and electrophysiological hearing thresholds in a bottlenose dolphin (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 122(1), 615–622.
- Schorr, G. S., E. A. Falcone, D. J. Moretti, and R. D. Andrews. (2014). First long-term behavioral records from Cuvier's beaked whales (*Ziphius cavirostris*) reveal record-breaking dives. *PLoS ONE*, 9(3), e92633.
- Schorr, G. S., E. A. Falcone, and B. K. Rone. (2017). *Distribution and Demographics of Cuvier's Beaked Whales and Fin Whales in the Southern California Bight* (Annual report for on-water surveys conducted in conjunction with Marine Mammal Monitoring on Navy Ranges). Seabeck, WA: Marine Ecology and Telemetry Research.
- Schorr, G. S., E. A. Falcone, B. K. Rone, and E. L. Keene. (2018). *Distribution and Demographics of Cuvier's Beaked Whales in the Southern California Bight*. Seabeck, WA: Marine Ecology and Telemetry Research.
- Sears, R., and W. F. Perrin. (2009). Blue whale. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 120–124). Cambridge, MA: Academic Press.
- Sekiguchi, K., N. T. W. Klages, and P. B. Best. (1992). Comparative analysis of the diets of smaller odontocete cetaceans along the coast of southern Africa. *South African Journal of Marine Science*, 12, 843–861.
- Serrano, L., C. A. Simeone, K. M. Colegrove, P. J. Duignan, T. Goldstein, and F. M. D. Gulland. (2017). Cetacean Morbillivirus in Odontocetes Stranded Along the Central California Coast, 2000–2015. *Journal of Wildlife Diseases*, 53(2), 1–7.
- Shane, S. H., R. S. Wells, and B. Wursig. (1986). Ecology, behavior and social organization of the bottlenose dolphin: A review. *Marine Mammal Science*, 2(1), 34–63.
- Shane, S. H. (1990). Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. In S. Leatherwood & R. R. Reeves (Eds.), *The Bottlenose Dolphin* (pp. 541–558). San Diego, CA: Academic Press.
- Shirasago-Germán, B., E. L. Pérez-Lezama, E. A. Chávez, and R. García-Morales. (2015). Influence of El Niño-Southern Oscillation on the population structure of a sea lion breeding colony in the Gulf of California. *Estuarine, Coastal and Shelf Science*, 154, 69–76.
- Silber, G. K., M. D. Lettrich, P. O. Thomas, J. D. Baker, M. Baumgartner, E. A. Becker, P. Boveng, D. M. Dick, J. Fiechter, J. Forcada, K. A. Forney, R. B. Griffis, J. A. Hare, A. J. Hobday, D. Howell, K. L. Laidre, N. Mantua, L. Quakenbush, J. A. Santora, K. M. Stafford, P. Spencer, C. Stock, W. Sydeman, K. Van Houtan, and R. S. Waples. (2017). Projecting Marine Mammal Distribution in a Changing Climate. *Frontiers in Marine Science*, 4, 14.

- Sills, J. M., B. L. Southall, and C. Reichmuth. (2017). The influence of temporally varying noise from seismic air guns on the detection of underwater sounds by seals. *The Journal of the Acoustical Society of America*, 141(2), 996–1008.
- Silva, T. L., T. A. Mooney, L. S. Sayigh, P. L. Tyack, R. W. Baird, and J. N. Oswald. (2016). Whistle characteristics and daytime dive behavior in pantropical spotted dolphins (*Stenella attenuata*) in Hawaii measured using digital acoustic recording tags (DTAGs). *The Journal of the Acoustical Society of America*, 140(1), 421–429.
- Simeone, C. A., F. M. Gulland, T. Norris, and T. K. Rowles. (2015). A systematic review of changes in marine mammal health in North America, 1972–2012: The need for a novel integrated approach. *PLoS ONE*, 10(11), e0142105.
- Simmonds, M. P., and W. J. Elliott. (2009). Climate change and cetaceans: Concerns and recent developments. *Journal of the Marine Biological Association of the United Kingdom*, 89(1), 203–210.
- Singh, R., P. Soni, P. Kumar, S. Purohit, and A. Singh. (2009). Biodegradation of high explosive production effluent containing RDX and HMX by denitrifying bacteria. *World Journal of Microbiology and Biotechnology*, 25, 269–275.
- Širović, A., J. A. Hildebrand, S. M. Wiggins, M. A. McDonald, S. E. Moore, and D. Thiele. (2004). Seasonality of blue and fin whale calls and the influence of sea ice in the Western Antarctic Peninsula. *Deep Sea Research II*, 51(17–19), 2327–2344.
- Širović, A., J. A. Hildebrand, and S. M. Wiggins. (2007). Blue and fin whale call source levels and propagation range in the Southern Ocean. *The Journal of the Acoustical Society of America*, 122(2), 1208–1215.
- Širović, A., A. Rice, E. Chou, J. A. Hildebrand, S. M. Wiggins, and M. A. Roch. (2015). Seven years of blue and fin whale call abundance in the Southern California Bight. *Endangered Species Research*, 28, 61–76.
- Sivle, L. D., P. H. Kvasdheim, A. Fahlman, F. P. Lam, P. L. Tyack, and P. J. Miller. (2012). Changes in dive behavior during naval sonar exposure in killer whales, long-finned pilot whales, and sperm whales. *Frontiers in Physiology*, 3, 400.
- Sivle, L. D., P. H. Kvasdheim, C. Curé, S. Isojunno, P. J. Wensveen, F. A. Lam, F. Visser, L. Kleivane, P. L. Tyack, C. M. Harris, and P. J. O. Miller. (2015). Severity of expert-identified behavioural responses of humpback whale, minke whale, and northern bottlenose whale to naval sonar. *Aquatic Mammals*, 41(4), 469–502.
- Sivle, L. D., P. J. Wensveen, P. H. Kvasdheim, F. P. A. Lam, F. Visser, C. Curé, C. M. Harris, P. L. Tyack, and P. J. O. Miller. (2016). Naval sonar disrupts foraging in humpback whales. *Marine Ecology Progress Series*, 562, 211–220.
- Skarke, A., C. Ruppel, M. Kodis, D. Brothers, and E. Lobecker. (2014). Widespread methane leakage from the sea floor on the northern US Atlantic margin. *Nature Geoscience*, 7, 657–661.
- Smith, B. D., G. Braulik, S. Strindberg, R. Mansur, M. A. A. Diyan, and B. Ahmed. (2009). Habitat selection of freshwater-dependent cetaceans and the potential effects of declining freshwater flows and sea-level rise in waterways of the Sundarbans mangrove forest, Bangladesh. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19(2), 209–225.

- Smith, C. E., B. J. Hurley, C. N. Toms, A. D. Mackey, M. Solangi, and S. A. Kuczaj li. (2013a). Hurricane impacts on the foraging patterns of bottlenose dolphins *Tursiops truncatus* in Mississippi Sound. *Marine Ecology Progress Series*, 487, 231–244.
- Smith, C. E., S. T. Sykora–Bodie, B. Bloodworth, S. M. Pack, T. R. Spradlin, and N. R. LeBoeuf. (2016). Assessment of known impacts of unmanned aerial systems (UAS) on marine mammals: Data gaps and recommendations for researchers in the United States. *Journal of Unmanned Vehicle Systems*, 4(1), 31–44.
- Smith, S. D. A. (2012). Marine debris: A proximate threat to marine sustainability in Bootless Bay, Papua New Guinea. *Marine Pollution Bulletin*, 64, 1880–1883.
- Smith, S. H., and D. E. Marx, Jr. (2009). *Assessment of Near Shore Marine Resources at Farallon De Medinilla: 2006, 2007 and 2008*. Capitol Hill, Commonwealth of the Northern Mariana Islands: Pacific Division, Naval Facilities Engineering Command.
- Smith, S. H., J. Marx, D. E., and L. H. Shannon. (2013b). *Calendar Year 2012 Assessment of Near Shore Marine Resources at Farallon de Medinilla, Commonwealth of the Northern Mariana Islands*. Port Hueneme, CA: U.S. Department of the Navy.
- Smith, S. H., and D. E. Marx, Jr. (2016). De-facto marine protection from a Navy bombing range: Farallon de Medinilla, Mariana Archipelago, 1997 to 2012. *Marine Pollution Bulletin*, 102(1), 187–198.
- Smultea, M. A., J. L. Hopkins, and A. M. Zoidis. (2008). *Marine Mammal and Sea Turtle Monitoring Survey in Support of Navy Training Exercises in the Hawai'i Range Complex November 11–17, 2007*. Oakland, CA: Cetos Research Organization.
- Smultea, M. A., and J. R. Mobley, Jr. (2009). *Aerial Survey Monitoring of Marine Mammals and Sea Turtles in Conjunction with SCC OPS Navy Exercises off Kauai, 18–21 August 2008, Final Report, May 2009*. Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.
- Smultea, M. A., J. R. Mobley, Jr., and K. Lomac-MacNair. (2009). *Aerial Survey Monitoring for Marine Mammals and Sea Turtles in the Hawaii Range Complex in Conjunction with a Navy Training Event, SSC OPS February 15–19, 2009, Final Field Report*. Honolulu, HI: Marine Mammal Research Consultants and Issaquah, WA: Smultea Environmental Sciences, LLC.
- Smultea, M. A., C. E. Bacon, and J. S. D. Black. (2011). *Aerial Survey Marine Mammal Monitoring off Southern California in Conjunction with US Navy Major Training Events (MTE), July 27–August 3 and September 23–28, 2010—Final Report, June 2011*. Issaquah, WA: Smultea Environmental Sciences.
- Smultea, M. A., C. E. Bacon, T. F. Norris, and D. Steckler. (2012). *Aerial Surveys Conducted in the SOCAL OPAREA From 1 August 2011–31 July 2012*. San Diego, CA: HDR, Inc.
- Song, K.-J. (2017). Bycatch of cetaceans on Korea fisheries in the East Sea. *Fisheries Research*, 197, 7–9.
- Sousa-Lima, R. S., and C. W. Clark. (2008). Modeling the effect of boat traffic on the fluctuation of humpback whale singing activity in the Abrolhos National Marine Park, Brazil. *Canadian Acoustics*, 36(1), 174–181.
- Southall, B., A. Bowles, W. Ellison, J. Finneran, R. Gentry, C. Greene, D. Kastak, D. Ketten, J. Miller, P. Nachtigall, W. Richardson, J. Thomas, and P. Tyack. (2007). Marine mammal noise exposure criteria: Initial scientific recommendations. *Aquatic Mammals*, 33(4), 122.
- Southall, B., J. Calambokidis, P. Tyack, D. Moretti, J. Hildebrand, C. Kyburg, R. Carlson, A. Friedlaender, E. Falcone, G. Schorr, A. Douglas, S. DeRuiter, J. Goldbogen, and J. Barlow. (2011). *Biological and*

- Behavioral Response Studies of Marine Mammals in Southern California, 2010 ("SOCAL-10") Project Report.* Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- Southall, B., J. Calambokidis, P. Tyack, D. Moretti, A. Friedlaender, S. DeRuiter, J. Goldbogen, E. Falcone, G. Schorr, A. Douglas, A. K. Stimpert, J. Hildebrand, C. Kyburg, R. Carlson, T. Yack, and J. Barlow. (2012). *Biological and Behavioral Response Studies of Marine Mammals in Southern California, 2011 ("SOCAL-11") Final Project Report.* Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- Southall, B., J. Calambokidis, J. Barlow, D. Moretti, A. Friedlaender, A. Stimpert, A. Douglas, K. Southall, S. Arranz, S. DeRuiter, E. Hazen, J. Goldbogen, E. Falcone, and G. Schorr. (2013). *Biological and Behavioral Response Studies of Marine Mammals in Southern California, 2012 ("SOCAL-12") Final Project Report.* Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- Southall, B., J. Calambokidis, J. Barlow, D. Moretti, A. Friedlaender, A. Stimpert, A. Douglas, K. Southall, P. Arranz, S. DeRuiter, J. Goldbogen, E. Falcone, and G. Schorr. (2014). *Biological and Behavioral Response Studies of Marine Mammals in Southern California, 2013 ("SOCAL-13") Final Project Report.* Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- Southall, B., J. Calambokidis, D. Moretti, A. Stimpert, A. Douglas, J. Barlow, R. W. Rankin, K. Southall, A. Friedlaender, E. Hazen, J. Goldbogen, E. Falcone, G. Schorr, G. Gailey, and A. Allen. (2015). *Biological and Behavioral Response Studies of Marine Mammals in Southern California, 2014 ("SOCAL-14") Final Project Report.* Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- Southall, B. L., R. J. Schusterman, and D. Kastak. (2000). Masking in three pinnipeds: Underwater, low-frequency critical ratios. *The Journal of the Acoustical Society of America*, 108(3), 1322–1326.
- Southall, B. L., R. J. Schusterman, and D. Kastak. (2003). Auditory masking in three pinnipeds: Aerial critical ratios and direct critical bandwidth measurements. *The Journal of the Acoustical Society of America*, 114(3), 1660–1666.
- Southall, B. L., R. Braun, F. M. D. Gulland, A. D. Heard, R. W. Baird, S. M. Wilkin, and T. K. Rowles. (2006). *Hawaiian Melon-headed Whales (Peponacephala electra) Mass Stranding Event of July 3–4, 2004* (National Oceanic and Atmospheric Administration Technical Memorandum NMFS-OPR-31). Silver Spring, MD: National Marine Fisheries Service.
- Southall, B. L., P. L. Tyack, D. Moretti, C. Clark, D. Claridge, and I. Boyd. (2009). *Behavioral responses of beaked whales and other cetaceans to controlled exposures of simulated sonar and other sounds.* Paper presented at the 18th Biennial Conference on the Biology of Marine Mammals. Quebec City, Canada.
- Southall, B. L., D. P. Nowacek, P. J. O. Miller, and P. L. Tyack. (2016). Experimental field studies to measure behavioral responses of cetaceans to sonar. *Endangered Species Research*, 31, 293–315.
- Southall, B. L., L. Hatch, A. Scholik-Schlomer, T. Bergmann, M. Jasny, K. Metcalf, L. Weilgart, A. J. Wright, and M. E. Perera. (2018). Reducing Noise from Large Commercial Ships. *Proceedings of the Marine Safety & Security Council*, 75(1), 1–8.
- Spiesberger, J. L., and K. M. Fristrup. (1990). Passive localization of calling animals and sensing of their acoustic environment using acoustic tomography. *The American Naturalist*, 135(1), 107–153.
- Sprogis, K. R., C. King, L. Bejder, and N. R. Loneragan. (2018). Frequency and temporal trends of shark predation attempts on bottlenose dolphins (*Tursiops aduncus*) in temperate Australian waters. *Journal of Experimental Marine Biology and Ecology*, 508(1), 35–43.

- St. Aubin, D., and L. A. Dierauf. (2001a). Stress and Marine Mammals. In L. A. Dierauf & F. M. D. Gulland (Eds.), *Marine Mammal Medicine* (2nd ed., pp. 253–269). Boca Raton, FL: CRC Press.
- St. Aubin, D. J., and J. R. Geraci. (1989). Adaptive changes in hematologic and plasma chemical constituents in captive beluga whales, *Delphinapterus leucas*. *Canadian Journal of Fisheries and Aquatic Sciences*, 46, 796–803.
- St. Aubin, D. J., S. H. Ridgway, R. S. Wells, and H. Rhinehart. (1996). Dolphin thyroid and adrenal hormones: Circulating levels in wild and semidomesticated *Tursiops truncatus*, and influence of sex, age, and season. *Marine Mammal Science*, 12(1), 1–13.
- St. Aubin, D. J., and L. A. Dierauf. (2001b). Stress and Marine Mammals. In L. A. Dierauf & F. M. D. Gulland (Eds.), *Marine Mammal Medicine* (2nd ed., pp. 253–269). Boca Raton, FL: CRC Press.
- Stamation, K. A., D. B. Croft, P. D. Shaughnessy, K. A. Waples, and S. V. Briggs. (2009). Behavioral responses of humpback whales (*Megaptera novaeangliae*) to whale-watching vessels on the southeastern coast of Australia. *Marine Mammal Science*, 26(1), 98–122.
- Stamper, M. A., B. R. Whitaker, and T. D. Schofield. (2006). Case study: Morbidity in a pygmy sperm whale *Kogia breviceps* due to ocean-bourne plastic. *Marine Mammal Science*, 22(3), 719–722.
- State of Hawaii. (2015). *Friends of the Future to help Lapakahi State Park*. Retrieved from <http://www.bigislandvideonews.com/2015/10/21/friends-of-the-future-to-help-lapakahi-state-park/>.
- Steckenreuter, A., R. Harcourt, and L. Moller. (2011). Distance does matter: Close approaches by boats impede feeding and resting behaviour of Indo-Pacific bottlenose dolphins. *Wildlife Research*, 38(6), 455–463.
- Steiger, G. H., J. Calambokidis, J. M. Straley, L. M. Herman, S. Cerchio, D. R. Salden, J. Urban-R, J. K. Jacobsen, O. von Ziegesar, K. C. Balcomb, C. M. Gabriele, M. E. Dahlheim, S. Uchida, J. K. B. Ford, P. Ladron de Guevara-P., M. Yamaguchi, and J. Barlow. (2008). Geographic variation in killer whale attacks on humpback whales in the North Pacific: Implications for predation pressure. *Endangered Species Research*, 4, 247–256.
- Stevens, P. W., D. A. Blewett, and J. P. Casey. (2006). Short-term effects of a low dissolved oxygen event on estuarine fish assemblages following passage of Hurricane Charley. *Estuaries and Coasts*, 29(6A), 997–1003.
- Stimpert, A. K., D. N. Wiley, W. W. Au, M. P. Johnson, and R. Arsenault. (2007). 'Megapclicks': Acoustic click trains and buzzes produced during night-time foraging of humpback whales (*Megaptera novaeangliae*). *Biology Letters*, 3(5), 467–470.
- Stimpert, A. K., S. L. DeRuiter, B. L. Southall, D. J. Moretti, E. A. Falcone, J. A. Goldbogen, A. Friedlaender, G. S. Schorr, and J. Calambokidis. (2014). Acoustic and foraging behavior of a Baird's beaked whale, *Berardius bairdii*, exposed to simulated sonar. *Scientific Reports*, 4, 7031.
- Stockin, K. A., D. Lusseau, V. Binedell, N. Wiseman, and M. B. Orams. (2008). Tourism affects the behavioural budget of the common dolphin *Delphinus* sp. in the Hauraki Gulf, New Zealand. *Marine Ecology Progress Series*, 355, 287–295.
- Sullivan, F. A., and L. G. Torres. (2018). Assessment of vessel disturbance to gray whales to inform sustainable ecotourism. *The Journal of Wildlife Management*, 82(5), 896–905.
- Summers, D. J. (2017). Algal toxins found in Alaska marine mammals for first time. *Alaska Journal of Commerce*(3), 3.

- Supin, A. Y., V. V. Popov, and A. M. Mass. (2001). *The Sensory Physiology of Aquatic Mammals*. Boston, MA: Kluwer Academic Publishers.
- Swisdak, M. M., Jr., and P. E. Montanaro. (1992). *Airblast and Fragmentation Hazards Produced by Underwater Explosions*. Silver Spring, MD: Naval Surface Warfare Center.
- Sydeman, W. J., and S. G. Allen. (1999). Pinniped population dynamics in central California: Correlations with sea surface temperature and upwelling indices. *Marine Mammal Science*, 15(2), 446–461.
- Sylvestre, J.-P. (1988). Note on three Dwarf Sperm Whales *Kogia simus* (Owen 1866) and comments on Kogiids of Japanese Coasts. *Aquatic Mammals*, 14.3, 120–122.
- Tarpley, R. J., and S. Marwitz. (1993). Plastic debris ingestion by cetaceans along the Texas coast: Two case reports. *Aquatic Mammals*, 19(2), 93–98.
- Teilmann, J., J. Tougaard, L. A. Miller, T. Kirketerp, K. Hansen, and S. Brando. (2006). Reactions of captive harbor porpoises (*Phocoena phocoena*) to pinger-like sounds. *Marine Mammal Science*, 22(2), 240–260.
- Tennessen, J. B., and S. E. Parks. (2016). Acoustic propagation modeling indicates vocal compensation in noise improves communication range for North Atlantic right whales. *Endangered Species Research*, 30, 225–237.
- Terhune, J. M., and W. C. Verboom. (1999). Right whales and ship noises. *Marine Mammal Science*, 15(1), 256–258.
- Tetra Tech Inc. (2014). *Marine Mammal Survey Report in Support of the Commonwealth of the Northern Mariana Islands Joint Military Training Environmental Impact Statement/Overseas Environmental Impact Statement. Final (Version 3)*. Oakland, CA: TEC-AECOM Pacific Joint Venture
- Tezanos-Pinto, G., C. S. Baker, K. Russell, K. Martien, R. W. Baird, A. Hutt, G. Stone, A. A. Mignucci-Giannoni, S. Caballero, T. Endo, S. Lavery, M. Oremus, C. Olavarria, and C. Garrigue. (2009). A worldwide perspective on the population structure and genetic diversity of bottlenose dolphins (*Tursiops truncatus*) in New Zealand. *Journal of Heredity*, 100(1), 11–24.
- Thiel, M., G. Luna-Jorquera, R. Álvarez-Varas, C. Gallardo, I. A. Hinojosa, N. Luna, D. Miranda-Urbina, N. Morales, N. Ory, A. S. Pacheco, M. Portflitt-Toro, and C. Zavalaga. (2018). Impacts of Marine Plastic Pollution From Continental Coasts to Subtropical Gyres—Fish, Seabirds, and Other Vertebrates in the SE Pacific. *Frontiers in Marine Science*, 5, 1–16.
- Thomas, J., P. Moore, R. Withrow, and M. Stoermer. (1990a). Underwater audiogram of a Hawaiian monk seal (*Monachus schauinslandi*). *The Journal of the Acoustical Society of America*, 87(1), 417–420.
- Thomas, J. A., R. A. Kastelein, and F. T. Awbrey. (1990b). Behavior and blood catecholamines of captive belugas during playbacks of noise from an oil drilling platform. *Zoo Biology*, 9(5), 393–402.
- Thompson, D., M. Sjoberg, M. E. Bryant, P. Lovell, and A. Bjorge. (1998). *Behavioral and physiological responses of harbour (Phoca vitulina) and grey (Halichoerus grypus) seals to seismic surveys* (Report to European Commission of BROMMAD Project. MAS2 C7940098). Brussels, Belgium: European Commission.
- Thompson, P. M., D. Lusseau, T. Barton, D. Simmons, J. Rusin, and H. Bailey. (2010). Assessing the responses of coastal cetaceans to the construction of offshore wind turbines. *Marine Pollution Bulletin*, 60(8), 1200–1208.

- Thompson, P. M., K. L. Brookes, I. M. Graham, T. R. Barton, K. Needham, G. Bradbury, and N. D. Merchant. (2013). Short-term disturbance by a commercial two-dimensional seismic survey does not lead to long-term displacement of harbour porpoises. *Proceedings of the Royal Society B: Biological Sciences*, 280(1771), 20132001.
- Thompson, R., Y. Olsen, R. Mitchell, A. Davis, S. Rowland, A. John, D. McGonigle, and A. Russell. (2004). Lost at sea: Where is all the plastic? *Science, New Series*, 304(5672), 838.
- Tirard, P., M. J. Manning, I. Jollit, C. Duffy, and P. Borsa. (2010). Records of great white shark (*Carcharodon carcharias*) in New Caledonian waters. *Pacific Science*, 64, 567–576.
- Titova, O. V., O. A. Filatova, I. D. Fedutin, E. N. Ovsyanikova, H. Okabe, N. Kobayashi, J. M. V. Acebes, A. M. Burdin, and E. Hoyt. (2017). Photo-identification matches of humpback whales (*Megaptera novaeangliae*) from feeding areas in Russian Far East seas and breeding grounds in the North Pacific. *Marine Mammal Science*, 34(1), 100–112.
- Tixier, P., N. Gasco, G. Duhamel, and C. Guinet. (2014). Habituation to an acoustic harassment device by killer whales depredating demersal longlines. *ICES Journal of Marine Science*, 72(5), 1673–1681.
- Todd, S., P. Stevick, J. Lien, F. Marques, and D. Ketten. (1996). Behavioural effects of exposure to underwater explosions in humpback whales (*Megaptera novaeangliae*). *Canadian Journal of Zoology*, 74, 1661–1672.
- Tougaard, J., J. Carstensen, J. Teilmann, N. I. Bech, H. Skov, and O. D. Henriksen. (2005). *Effects of the Nysted Offshore Wind Farm on Harbour Porpoises* (Annual Status Report for the T-POD Monitoring Program). Roskilde, Denmark: National Environmental Research Institute.
- Tougaard, J., J. Carstensen, J. Teilmann, H. Skov, and P. Rasmussen. (2009). Pile driving zone of responsiveness extends beyond 20 km for harbor porpoises (*Phocoena phocoena* [L.]). *The Journal of the Acoustical Society of America*, 126(1), 11.
- Townsend, C. H. (1935). The Distribution of Certain Whales As Shown by Logbook Records of American Whaleships. *Zoologica*, XIX(1), 3–40.
- Trianni, M. S., and C. C. Kessler. (2002). Incidence and strandings of the Spinner Dolphin, *Stenella longirostris*, in Saipan Lagoon. *Micronesica*, 34(2), 249–260.
- Trianni, M. S., and M. C. Tenorio. (2012). Summary of recorded cetacean strandings in the Commonwealth of the Northern Mariana Islands. *Micronesica*, 43(1), 1–13.
- Trickey, J. S., B. K. Branstetter, and J. J. Finneran. (2010). Auditory masking of a 10 kHz tone with environmental, comodulated, and Gaussian noise in bottlenose dolphins (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 128(6), 3799–3804.
- Trickey, J. S., S. Baumann-Pickering, A. Širović, J. A. Hildebrand, A. M. Brewer, A. J. Debich, S. Herbert, A. C. Rice, B. Thayre, and S. M. Wiggins. (2015). *Passive Acoustic Monitoring for Marine Mammals in the Northwest Training Range Complex July 2013–April 2014*. La Jolla, CA: Marine Physical Laboratory, Scripps Institution of Oceanography, University of California, San Diego.
- Trites, A. W., and D. E. Bain. (2000). *Short- and long-term effects of whale watching on killer whales (Orcinus orca) in British Columbia*. Adelaide, Australia: International Whaling Commission.
- Twiss, J. R., Jr., and R. R. Reeves. (1999). *Conservation and Management of Marine Mammals*. Washington, DC: Smithsonian Institution Press.

- Tyack, P., W. Zimmer, D. Moretti, B. Southall, D. Claridge, J. Durban, C. Clark, A. D'Amico, N. DiMarzio, S. Jarvis, E. McCarthy, R. Morrissey, J. Ward, and I. Boyd. (2011). Beaked Whales Respond to Simulated and Actual Navy Sonar. *PLoS ONE*, 6(3), 15.
- Tyack, P. L., M. Johnson, N. Aguilar Soto, A. Sturlese, and P. T. Madsen. (2006). Extreme deep diving of beaked whales. *The Journal of Experimental Biology*, 209, 4238–4253.
- Tyack, P. L. (2009). Human-generated sound and marine mammals. *Physics Today*, 39–44.
- Tyne, J. A., K. H. Pollock, D. W. Johnston, and L. Bejder. (2014). Abundance and survival rates of the Hawaii Island associated spinner dolphin (*Stenella longirostris*) stock. *PLoS ONE*, 9(1), e86132.
- Tyne, J. A. (2015). *A scientific foundation for informed management decisions: Quantifying the abundance, important habitat and cumulative exposure of the Hawaii Island spinner dolphin (Stenella longirostris) stock to human activities*. (Unpublished doctoral dissertation in Philosophy). Murdoch University, Murdoch, Australia. Retrieved from [https://www.researchgate.net/publication/311608220\\_A\\_scientific\\_foundation\\_for\\_informed\\_management\\_decisions\\_Quantifying\\_the\\_abundance\\_important\\_habitat\\_and\\_cumulative\\_exposure\\_of\\_the\\_Hawaii\\_Island\\_spinner\\_dolphin\\_Stenella\\_longirostris\\_stock\\_to\\_human\\_](https://www.researchgate.net/publication/311608220_A_scientific_foundation_for_informed_management_decisions_Quantifying_the_abundance_important_habitat_and_cumulative_exposure_of_the_Hawaii_Island_spinner_dolphin_Stenella_longirostris_stock_to_human_).
- Tyne, J. A., D. W. Johnston, R. Rankin, N. R. Loneragan, L. Bejder, and A. Punt. (2015). The importance of spinner dolphin (*Stenella longirostris*) resting habitat: Implications for management. *Journal of Applied Ecology*, 52(3), 621–630.
- Tyne, J. A., D. W. Johnston, F. Christiansen, and L. Bejder. (2017). Temporally and spatially partitioned behaviours of spinner dolphins: Implications for resilience to human disturbance. *Royal Society Open Science*, 4(1), 160626.
- Tyne, J. A., F. Christiansen, H. L. Heenehan, D. W. Johnston, and L. Bejder. (2018). Chronic exposure of Hawaii Island spinner dolphins (*Stenella longirostris*) to human activities. *Royal Society Open Science*, 5, e171506.
- U.S. Department of Commerce, and U.S. Department of the Navy. (2001). *Joint Interim Report Bahamas Marine Mammal Stranding Event of 15–16 March 2000*. Washington, DC: Department of Commerce.
- U.S. Department of the Navy. (2003). *Report on the Results of the Inquiry into Allegations of Marine Mammal Impacts Surrounding the Use of Active Sonar by USS SHOUP (DDG 86) in the Haro Strait on or about 5 May 2003*. Washington, DC: Commander, U.S. Pacific Fleet.
- U.S. Department of the Navy. (2005). *Marine Resources Assessment for the Marianas Operating Area (Final Report)*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- U.S. Department of the Navy. (2007). *Final Supplemental Environmental Impact Statement for Surveillance Towed Array Sensor System Low Frequency Active (SURTASS LFA) Sonar Volume 1 of 2*. Washington, DC: U.S. Department of the Navy.
- U.S. Department of the Navy. (2010). *Navy Integrated Comprehensive Monitoring Plan*. Washington, DC: U.S. Department of the Navy.
- U.S. Department of the Navy. (2011a). *Marine Species Monitoring, Information on Sightings Recorded by U.S. Navy MMOs on Vessels during Sonar Test Events in the Naval Surface Warfare Center Panama City Division (NSWC PCD)*. Norfolk, VA: United States Fleet Forces Command.

- U.S. Department of the Navy. (2011b). *Marine Species Monitoring for the U.S. Navy's Hawaii Range Complex and the Southern California Range Complex, 2011 Annual Report*. Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- U.S. Department of the Navy. (2013a). *U.S. Navy Strategic Planning Process for Marine Species Monitoring*. Washington, DC: Chief of Naval Operations, Energy & Environmental Readiness Division.
- U.S. Department of the Navy. (2013b). *Comprehensive Exercise and Marine Species Monitoring Report for the U.S. Navy's Atlantic Fleet Active Sonar Training (AFAST) and Virginia Capes, Cherry Point, Jacksonville, and Gulf of Mexico Range Complexes 2009–2012*. Norfolk, VA: United States Fleet Forces Command.
- U.S. Department of the Navy. (2013c). *Comprehensive Exercise and Marine Species Monitoring Report for the U.S. Navy's Hawaii Range Complex 2009–2012*. Pearl Harbor, HI: U.S. Navy Pacific Fleet.
- U.S. Department of the Navy. (2014a). *Unclassified Annual Range Complex Exercise Report, 2 August 2012 to 25 November 2013, for the U.S. Navy's Atlantic Fleet Active Sonar Training (AFAST) Study Area*. Silver Spring, MD: National Marine Fisheries Service, Office of Protected Resources.
- U.S. Department of the Navy. (2014b). *Marine Species Monitoring Report for the U.S. Navy's Atlantic Fleet Active Sonar Training (AFAST) and Virginia Capes, Cherry Point, Jacksonville, and Gulf of Mexico Range Complexes - Annual Report 2013*. Norfolk, VA: United States Fleet Forces Command.
- U.S. Department of the Navy. (2015a). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2015b). *Unclassified 2014 Annual Atlantic Fleet Training and Testing (AFTT) Exercise and Testing Report 14 November 2013 to 13 November 2014*. Silver Spring, MD: National Marine Fisheries Service, Office of Protected Resources.
- U.S. Department of the Navy. (2017a). *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)*. San Diego, CA: Space and Naval Warfare System Command, Pacific.
- U.S. Department of the Navy. (2017b). *Hawaii-Southern California Training and Testing Draft Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2017c). *Marine Mammal Strandings Associated with U.S. Navy Sonar Activities*. San Diego, CA: U.S. Navy Marine Mammal Program and SPAWAR Naval Facilities Engineering Command.
- U.S. Department of the Navy. (2018a). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (Technical Report prepared by NUWC Division Newport, Space and Naval Warfare Systems Center Pacific, G2 Software Systems, and the National Marine Mammal Foundation). Newport, RI: Naval Undersea Warfare Center.
- U.S. Department of the Navy. (2018b). *U.S. Navy Marine Species Density Database Phase III for the Mariana Islands Training and Testing Study Area* (Naval Facilities Engineering Command Pacific Technical Report). Pearl Harbor, HI: Naval Facilities Engineering Command Pacific.

- Unger, B., E. L. Rebolledo, R. Deaville, A. Grone, L. L. IJsseldijk, M. F. Leopold, U. Siebert, J. Spitz, P. Wohlsein, and H. Herr. (2016). Large amounts of marine debris found in sperm whales stranded along the North Sea coast in early 2016. *Marine Pollution Bulletin*, 112(1-2), 134–141.
- University of Hawaii. (2010). *Hawaii Undersea Military Munitions Assessment, Final Investigation Report HI-05, South of Pearl Harbor, Oahu, HI*. Honolulu, HI: University of Hawaii at Manoa.
- Uyeyama, R. (2014). *Compilation of Incidental Marine Mammal and Sea Turtle Sightings in the Mariana Islands Range Complex*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Uyeyama, R. K., M. A. Fagan, and L. H. Shannon. (2012). *Cruise Report Marine Mammal and Sea Turtle Observer UNDET Monitoring Hawaii Range Complex: 19 & 26 October and 2 November 2011*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- Vallejo, G. C., K. Grellier, E. J. Nelson, R. M. McGregor, S. J. Canning, F. M. Caryl, and N. McLean. (2017). Responses of two marine top predators to an offshore wind farm. *Ecology and Evolution*, 7(21), 8698–8708.
- van Beest, F. M., L. Kindt-Larsen, F. Bastardie, V. Bartolino, and J. Nabe-Nielsen. (2017). Predicting the population-level impact of mitigating harbor porpoise bycatch with pingers and time-area fishing closures. *Ecosphere*, 8(4), e01785.
- Veirs, S., V. Veirs, and J. Wood. (2015). Ship noise in an urban estuary extends to frequencies used for echolocation by endangered killer whales. *PeerJ*, 4, e1657.
- Vilela, R., U. Pena, R. Esteban, and R. Koemans. (2016). Bayesian spatial modeling of cetacean sightings during a seismic acquisition survey. *Marine Pollution Bulletin*, 109(1), 512–520.
- Villadsgaard, A., M. Wahlberg, and J. Tougaard. (2007). Echolocation signals of wild harbour porpoises, *Phocoena phocoena*. *The Journal of Experimental Biology*, 210, 56–64.
- Villegas-Amtmann, S., L. K. Schwarz, G. Gailey, O. Sychenko, and D. P. Costa. (2017). East or west: The energetic cost of being a gray whale and the consequence of losing energy to disturbance. *Endangered Species Research*, 34, 167–183.
- Vincent, C., B. J. McConnell, V. Ridoux, and M. A. Fedak. (2002). Assessment of Argos location accuracy from satellite tags deployed on captive grey seals. *Marine Mammal Science*, 18(1), 156–166.
- Visser, F., C. Cure, P. H. Kvasdheim, F. P. Lam, P. L. Tyack, and P. J. Miller. (2016). Disturbance-specific social responses in long-finned pilot whales, *Globicephala melas*. *Scientific Reports*, 6, 28641.
- Vogt, S. (2008). *Fiscal Years 2007-2008 Annual Report for 61755NR410 Wildlife Surveys on Military Leased Lands, Farallon de Medinilla, Commonwealth of the Northern Mariana Islands*. Honolulu, HI: Naval Facilities Engineering Command Pacific.
- von Benda-Beckmann, A. M., P. J. Wensveen, P. H. Kvasdheim, F. P. Lam, P. J. Miller, P. L. Tyack, and M. A. Ainslie. (2014). Modeling effectiveness of gradual increases in source level to mitigate effects of sonar on marine mammals. *Conservation Biology*, 28(1), 119–128.
- von Benda-Beckmann, A. M., P. J. Wensveen, P. H. Kvasdheim, F. P. A. Lam, P. J. Miller, P. L. Tyack, and M. A. Ainslie. (2016). Assessing the effectiveness of ramp-up during sonar operations using exposure models. In A. N. Popper & A. Hawkins (Eds.), *The Effects of Noise on Aquatic Life II* (pp. 1197–1203). New York, NY: Springer.
- Wada, S., M. Oishi, and T. K. Yamada. (2003). A newly discovered species of living baleen whale. *Nature*, 426, 278–281.

- Wade, P. R., T. J. Quinn, II, J. Barlow, C. S. Baker, A. M. Burdin, J. Calambokidis, P. J. Clapham, E. A. Falcone, J. K. B. Ford, C. M. Gabriele, D. K. Mattila, L. Rojas-Bracho, J. M. Straley, and B. Taylor. (2016). *Estimates of Abundance and Migratory Destination for North Pacific Humpback Whales in Both Summer Feeding Areas and Winter Mating and Calving Areas* (SC/66b/IA/21). Washington, DC: International Whaling Commission.
- Walker, R. J., E. O. Keith, A. E. Yankovsky, and D. K. Odell. (2005). Environmental correlates of cetacean mass stranding sites in Florida. *Marine Mammal Science*, 21(2), 327–335.
- Walker, S. W., C. L. Osburn, T. J. Boyd, L. J. Hamdan, R. B. Coffin, M. T. Montgomery, J. P. Smith, Q. X. Li, C. Hennessee, F. Monteil, and J. Hawari. (2006). *Mineralization of 2, 4, 6-Trinitrotoluene (TNT) in Coastal Waters and Sediments*. Washington, DC: U.S. Department of the Navy, Naval Research Laboratory.
- Wang, J. Y., S. C. Yang, and H. C. Liao. (2001). Species composition, distribution and relative abundance of cetaceans in the waters of southern Taiwan: Implications for conservation and eco-tourism. *Journal of the National Parks of Taiwan*, 11(2), 136–158.
- Wang, J. Y., and S. C. Yang. (2006). Unusual cetacean stranding events of Taiwan in 2004 and 2005. *Journal of Cetacean Research and Management*, 8(3), 283–292.
- Ward, W. D., A. Glorig, and D. L. Sklar. (1958). Dependence of temporary threshold shift at 4 kc on intensity and time. *The Journal of the Acoustical Society of America*, 30(10), 944–954.
- Ward, W. D., A. Glorig, and D. L. Sklar. (1959). Relation between recovery from temporary threshold shift and duration of exposure. *The Journal of the Acoustical Society of America*, 31(5), 600–602.
- Ward, W. D. (1960). Recovery from high values of temporary threshold shift. *The Journal of the Acoustical Society of America*, 32(4), 497–500.
- Wartzok, D., and D. R. Ketten. (1999). Marine Mammal Sensory Systems. In J. E. Reynolds, III & S. A. Rommel (Eds.), *Biology of Marine Mammals* (pp. 117–175). Washington, DC: Smithsonian Institution Press.
- Wartzok, D., A. N. Popper, J. Gordon, and J. Merrill. (2003). Factors affecting the responses of marine mammals to acoustic disturbance. *Marine Technology Society Journal*, 37(4), 6–15.
- Watkins, W. A., and W. E. Schevill. (1975). Sperm whales (*Physeter catodon*) react to pingers. *Deep-Sea Research*, 22, 123–129.
- Watkins, W. A. (1981). Reaction of three species of whales *Balaenoptera physalus*, *Megaptera novaeangliae*, and *Balaenoptera edeni* to implanted radio tags. *Deep-Sea Research*, 28A(6), 589–599.
- Watkins, W. A., K. E. Moore, and P. Tyack. (1985). Sperm whale acoustic behavior in the southeast Caribbean. *Cetology*, 49, 1–15.
- Watkins, W. A. (1986). Whale reactions to human activities in Cape Cod waters. *Marine Mammal Science*, 2(4), 251–262.
- Watkins, W. A., M. A. Daher, K. M. Fristrup, and G. Notarbartolo di Sciara. (1994). Fishing and acoustic behavior of Fraser's dolphin (*Lagenodelphis hosei*) near Dominica, southeast Caribbean. *Caribbean Journal of Science*, 30, 76–82.

- Watwood, S., M. Fagan, A. D'Amico, and T. Jefferson. (2012). *Cruise Report, Marine Species Monitoring and Lookout Effectiveness Study, Koa Kai, November 2011, Hawaii Range Complex*. Pearl Harbor, HI: Commander, U.S. Pacific Fleet.
- Watwood, S., E. McCarthy, N. DiMarzio, R. Morrissey, S. Jarvis, and D. Moretti. (2017). *Beaked whale foraging behavior before, during, and after sonar exposure on a Navy test range*. Paper presented at the 22nd Biennial Conference on the Biology of Marine Mammals. Halifax, Canada.
- Weaver, A. (2015). Sex difference in bottlenose dolphin sightings during a long-term bridge construction project. *Animal Behavior and Cognition*, 2(1), 1–13.
- Weir, C. R. (2008). Overt responses of humpback whales (*Megaptera novaeangliae*), sperm whales (*Physeter macrocephalus*), and Atlantic spotted dolphins (*Stenella frontalis*) to seismic exploration off Angola. *Aquatic Mammals*, 34(1), 71–83.
- Wellard, R., K. Lightbody, L. Fouda, M. Blewitt, D. Riggs, and C. Erbe. (2016). Killer Whale (*Orcinus orca*) Predation on Beaked Whales (*Mesoplodon* spp.) in the Bremer Sub-Basin, Western Australia. *PLoS ONE*, 11(12).
- Weller, D. W. (2009). Predation on marine mammals. In W. F. Perrin, B. Wursig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 923–931). Cambridge, MA: Academic Press.
- Wells, R. S., and M. D. Scott. (1999). Bottlenose dolphin, *Tursiops truncatus* (Montagu, 1821). In S. H. Ridgway & R. Harrison (Eds.), *Handbook of Marine Mammals* (Vol. 6, pp. 137–182). San Diego, CA: Academic Press.
- Wells, R. S., J. B. Allen, S. Hofmann, K. Bassos-Hull, D. A. Fauquier, N. B. Barros, R. E. DeLynn, G. Sutton, V. Socha, and M. D. Scott. (2008). Consequences of injuries on survival and reproduction of common bottlenose dolphins (*Tursiops truncatus*) along the west coast of Florida. *Marine Mammal Science*, 24(4), 774–794.
- Wells, R. S., and M. D. Scott. (2009). Common bottlenose dolphin, *Tursiops truncatus*. In W. F. Perrin, W. B., & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 249–255). Cambridge, MA: Academic Press.
- Weninger, P. (2010). *FY 2010 Annual Report Wildlife Surveys on Military Leased Lands, Farallon de Medinilla CNMI*. Apra Harbor, GU: U.S. Department of the Navy, Naval Base Guam Public Works Department.
- Wensveen, P. J., A. M. von Benda-Beckmann, M. A. Ainslie, F. P. Lam, P. H. Kvadsheim, P. L. Tyack, and P. J. Miller. (2015). How effectively do horizontal and vertical response strategies of long-finned pilot whales reduce sound exposure from naval sonar? *Marine Environmental Research*, 106, 68–81.
- Wensveen, P. J., P. H. Kvadsheim, F.-P. A. Lam, A. M. Von Benda-Beckmann, L. D. Sivle, F. Visser, C. Curé, P. Tyack, and P. J. O. Miller. (2017). Lack of behavioural responses of humpback whales (*Megaptera novaeangliae*) indicate limited effectiveness of sonar mitigation. *The Journal of Experimental Biology*, 220, 1–12.
- Werth, A. J. (2006a). Mandibular and dental variation and the evolution of suction feeding in Odontoceti. *Journal of Mammalogy*, 87(3), 579–588.
- Werth, A. J. (2006b). Odontocete suction feeding: Experimental analysis of water flow and head shape. *Journal of Morphology*, 267(12), 1415–1428.

- West, K. (2016). *Necropsy Findings, Stomach Contents and Hook Ingestion in IFKW's*. Honolulu, HI: University of Hawaii, Hawaii Pacific University.
- West, K. (2018). [Personal Communication between Kristi West and Julie Rivers regarding spinner dolphin and monk seal strandings].
- West, K. L., W. A. Walker, R. W. Baird, W. White, G. Levine, E. Brown, and D. Schofield. (2009). Diet of pygmy sperm whales (*Kogia breviceps*) in the Hawaiian Archipelago. *Marine Mammal Science*, 25(4), 931–943.
- West, K. L., S. Sanchez, D. Rotstein, K. M. Robertson, S. Dennison, G. Levine, N. Davis, D. Schofield, C. W. Potter, and B. Jensen. (2012). A Longman's beaked whale (*Indopacetus pacificus*) strands in Maui, Hawaii, with first case of morbillivirus in the central Pacific. *Marine Mammal Science*, 29, 767–776.
- West, K. L., G. Levine, J. Jacob, B. Jensen, S. Sanchez, K. Colegrove, and D. Rotstein. (2015). Coinfection and Vertical Transmission of *Brucella* and *Morbillivirus* in a Neonatal Sperm Whale (*Physeter macrocephalus*) in Hawaii, USA. *Journal of Wildlife Diseases*, 51(1), 227–232.
- Williams, R., D. E. Bain, J. K. B. Ford, and A. W. Trites. (2002a). Behavioural responses of male killer whales to a 'leapfrogging' vessel. *Journal of Cetacean Research and Management*, 4(3), 305–310.
- Williams, R., A. W. Trites, and D. E. Bain. (2002b). Behavioural responses of killer whales (*Orcinus orca*) to whale-watching boats: Opportunistic observations and experimental approaches. *Journal of Zoology, London*, 256, 255–270.
- Williams, R., D. Lusseau, and P. S. Hammond. (2006). Estimating relative energetic costs of human disturbance to killer whales (*Orcinus orca*). *Biological Conservation*, 133, 301–311.
- Williams, R., D. E. Bain, J. C. Smith, and D. Lusseau. (2009). Effects of vessels on behaviour patterns of individual southern resident killer whales, *Orcinus orca*. *Endangered Species Research*, 6, 199–209.
- Williams, R., C. W. Clark, D. Ponirakis, and E. Ashe. (2013). Acoustic quality of critical habitats for three threatened whale populations. *Animal Conservation*, 17(2), 174–185.
- Williams, R., E. Ashe, L. Blight, M. Jasny, and L. Nowlan. (2014a). Marine mammals and ocean noise: Future directions and information needs with respect to science, policy and law in Canada. *Marine Pollution Bulletin*, 86(1-2), 29–38.
- Williams, R., C. Erbe, E. Ashe, A. Beerman, and J. Smith. (2014b). Severity of killer whale behavioral responses to ship noise: A dose-response study. *Marine Pollution Bulletin*, 79(1–2), 254–260.
- Williams, T. M., T. L. Kendall, B. P. Richter, C. R. Ribeiro-French, J. S. John, K. L. Odell, B. A. Losch, D. A. Feuerbach, and M. A. Stamper. (2017). Swimming and diving energetics in dolphins: A stroke-by-stroke analysis for predicting the cost of flight responses in wild odontocetes. *The Journal of Experimental Biology*, 220(6), 1135–1145.
- Williamson, M. J., A. S. Kavanagh, M. J. Noad, E. Kniest, and R. A. Dunlop. (2016). The effect of close approaches for tagging activities by small research vessels on the behavior of humpback whales (*Megaptera novaeangliae*). *Marine Mammal Science*, 32(4), 1234–1253.
- Wisniewska, D. M., M. Johnson, J. Teilmann, U. Siebert, A. Galatius, R. Dietz, and P. T. Madsen. (2018). High rates of vessel noise disrupt foraging in wild harbour porpoises (*Phocoena phocoena*). *Proceedings of the Royal Society B: Biological Sciences*, 285(1872), 10.

- Woodworth, P. A., G. S. Schorr, R. W. Baird, D. L. Webster, D. J. McSweeney, M. B. Hanson, R. D. Andrews, and J. J. Polovina. (2012). Eddies as offshore foraging grounds for melon-headed whales (*Peponocephala electra*). *Marine Mammal Science*, 28(3), 638–647.
- Würsig, B., S. K. Lynn, T. A. Jefferson, and K. D. Mullin. (1998). Behaviour of cetaceans in the northern Gulf of Mexico relative to survey ships and aircraft. *Aquatic Mammals*, 24(1), 41–50.
- Würsig, B., and W. J. Richardson. (2009). Noise, effects of. In W. F. Perrin, B. Würsig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 765–773). Cambridge, MA: Academic Press.
- Yack, T. M., T. F. Norris, and N. Novak. (2016). *Acoustic Based Habitat Models for Sperm Whales in the Mariana Islands Region*. Encinitas, CA: Bio-Waves, Inc.
- Yang, W.-C., L.-S. Chou, P. D. Jepson, R. L. Brownell, Jr., D. Cowan, P.-H. Chang, H.-I. Chiou, C.-J. Yao, T. K. Yamada, J.-T. Chiu, P.-J. Wang, and A. Fernández. (2008). Unusual cetacean mortality event in Taiwan, possibly linked to naval activities. *Veterinary Record*, 162, 184–186.
- Yazvenko, S. B., T. L. McDonald, S. A. Blokhin, S. R. Johnson, H. R. Melton, M. W. Newcomer, R. Nielson, and P. W. Wainwright. (2007). Feeding of western gray whales during a seismic survey near Sakhalin Island, Russia. *Environmental Monitoring and Assessment*, 134(1–3), 93–106.
- Yelverton, J. T., D. R. Richmond, E. R. Fletcher, and R. K. Jones. (1973). *Safe Distances From Underwater Explosions for Mammals and Birds*. Albuquerque, NM: Lovelace Foundation for Medical Education and Research.
- Ylitalo, G. M., J. E. Stein, T. Hom, L. L. Johnson, K. L. Tilbury, A. J. Hall, T. Rowles, D. Greig, L. J. Lowenstine, and F. M. D. Gulland. (2005). The role of organochlorines in cancer-associated mortality in California sea lions (*Zalophus californianus*). *Marine Pollution Bulletin*, 50, 30–39.
- Ylitalo, G. M., R. W. Baird, G. K. Yanagida, D. L. Webster, S. J. Chivers, J. L. Bolton, G. S. Schorr, and D. J. McSweeney. (2009). High levels of persistent organic pollutants measured in blubber of island-associated false killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands. *Marine Pollution Bulletin*, 58, 1922–1952.
- Yuen, M. M. L., P. E. Nachtigall, M. Breese, and A. Y. Supin. (2005). Behavioral and auditory evoked potential audiograms of a false killer whale (*Pseudorca crassidens*). *The Journal of the Acoustical Society of America*, 118(4), 2688–2695.
- Zellar, R., A. Pulkkinen, K. Moore, D. Reeb, E. Karakoylu, and O. Uritskaya. (2017). *Statistical Assessment of Cetacean Stranding Events in Cape Cod (Massachusetts, USA) Area OS21A-1345*. Greenbelt, MD: National Aeronautics and Space Administration.
- Zimmer, W. M. X., and P. L. Tyack. (2007). Repetitive shallow dives pose decompression risk in deep-diving beaked whales. *Marine Mammal Science*, 23(4), 888–925.
- Zoidis, A. M., M. A. Smultea, A. S. Frankel, J. L. Hopkins, A. Day, and A. S. McFarland. (2008). Vocalizations produced by humpback whale (*Megaptera novaeangliae*) calves recorded in Hawaii. *The Journal of the Acoustical Society of America*, 123(3), 1737–1746.

---

---

## 3.5 Sea Turtles



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

3.5 Sea Turtles..... 3.5-1

3.5.1 Affected Environment..... 3.5-1

3.5.1.1 Group Size..... 3.5-2

3.5.1.2 Habitat Use ..... 3.5-2

3.5.1.3 Dive Behavior..... 3.5-3

3.5.1.4 Hearing and Vocalization ..... 3.5-4

3.5.1.5 General Threats ..... 3.5-5

3.5.1.6 Green Sea Turtle (*Chelonia mydas*) ..... 3.5-7

3.5.1.7 Hawksbill Sea Turtle (*Eretmochelys imbricata*) ..... 3.5-8

3.5.1.8 Loggerhead Sea Turtle (*Caretta caretta*) ..... 3.5-9

3.5.1.9 Olive Ridley Sea Turtle (*Lepidochelys olivacea*) ..... 3.5-9

3.5.1.10 Leatherback Sea Turtle (*Dermochelys coriacea*) ..... 3.5-9

3.5.2 Environmental Consequences ..... 3.5-9

3.5.2.1 Acoustic Stressors..... 3.5-11

3.5.2.2 Explosive Stressors..... 3.5-23

3.5.2.3 Energy Stressors ..... 3.5-45

3.5.2.4 Physical Disturbance and Strike Stressors ..... 3.5-47

3.5.2.5 Entanglement Stressors..... 3.5-49

3.5.2.6 Ingestion Stressors..... 3.5-50

3.5.2.7 Secondary Stressors..... 3.5-51

3.5.3 Public Scoping Comments ..... 3.5-51

**List of Figures**

Figure 3.5-1: Dive Depth and Duration Summaries for Sea Turtle Species ..... 3.5-3

Figure 3.5-2: Generalized Dive Profiles and Activities Described for Sea Turtles..... 3.5-4

Figure 3.5-3: Composite Underwater Audiogram for Sea Turtles ..... 3.5-5

Figure 3.5-4: Auditory Weighting Function for Sea Turtles ..... 3.5-17

Figure 3.5-5: TTS and PTS Exposure Functions for Sonar and Other Transducers.....3.5-18  
Figure 3.5-6: Auditory Weighting Functions for Sea Turtles.....3.5-32  
Figure 3.5-7: TTS and PTS Exposure Functions for Impulsive Sounds.....3.5-33  
Figure 3.5-8: Green Sea Turtle Estimated Impacts per Year from the Maximum Number of Explosions  
During Training and Testing Under Alternative 1 .....3.5-41  
Figure 3.5-9: Green Sea Turtle Impacts Estimated per Year from Explosions During Training and Testing  
Under Alternative 2.....3.5-44

### List of Tables

Table 3.5-1: Endangered Species Act Status and Presence of Endangered Species Act Listed Sea Turtles in  
the Mariana Islands Training and Testing Study Area .....3.5-2  
Table 3.5-2: Criteria to Quantitatively Assess Non-Auditory Injury due to Underwater Explosions....3.5-31  
Table 3.5-3: TTS and PTS Peak Pressure Thresholds Derived for Sea Turtles Exposed to Impulsive Sounds  
.....3.5-33  
Table 3.5-4: Ranges to Non-Auditory Injury<sup>1</sup> (in meters) for Sea Turtles Exposed to Explosives as a  
Function of Animal Mass.....3.5-35  
Table 3.5-5: Ranges to Mortality for Sea Turtles Exposed to Explosives as a Function of Animal Mass<sup>1</sup>  
.....3.5-36  
Table 3.5-6: Peak Pressure Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives .....3.5-37  
Table 3.5-7: SEL Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives.....3.5-38

### 3.5 Sea Turtles

#### 3.5.1 Affected Environment

The purpose of this section is to supplement the analysis of impacts on sea turtles presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea. Information presented in the 2015 MITT Final EIS/OEIS that remains valid is noted as such and referenced to the appropriate sections. Any new or updated information describing the affected environment and analysis of impacts on sea turtles associated with the Proposed Action is provided in this section. Comments received from the public during scoping related to sea turtles are addressed in Section 3.5.3 (Public Scoping Comments).

Only at-sea and Farallon de Medinilla (FDM) training and testing activities are subject to this Supplemental EIS (SEIS)/OEIS. Therefore, only effects within the nearshore and pelagic habitats for sea turtles are analyzed. The analysis of sea turtle presence and nesting on land presented in the 2015 MITT Final EIS/OEIS remains valid and continues to support these activities conducted within the Marianas.

The five sea turtle species potentially found in the Study Area are the same as those presented in the 2015 MITT Final EIS/OEIS and all are listed under the Endangered Species Act (ESA) as endangered or threatened (green sea turtle [*Chelonia mydas*], hawksbill sea turtle [*Eretmochelys imbricata*], loggerhead sea turtle [*Caretta caretta*], olive ridley sea turtle [*Lepidochelys olivacea*], and leatherback sea turtle [*Dermochelys coriacea*]). There is no critical habitat designated for sea turtle species within the Study Area. Similar to the 2015 MITT Final EIS/OEIS, this section provides an overview of the species, distribution, and occurrence of sea turtles, as well as new information released since the publication of the 2015 document. The status, presence, and nesting occurrence of sea turtles in the Study Area are listed by region in Table 3.5-1. Since the publication of the 2015 MITT Final EIS/OEIS, the National Marine Fisheries Service (NMFS) classified the global distribution of green sea turtles into distinct population segments (DPS). Within the area analyzed in this SEIS/OEIS, the endangered green sea turtle in the Mariana Islands has been determined to be part of the Central West Pacific DPS.

The Navy also reviewed the status and distribution of other pelagic reptile species, such as sea snakes, to evaluate if these species should be included in this SEIS/OEIS. There are no verified records of sea snakes in nearshore waters of the Mariana Islands. Eldredge (2003) notes that the few anecdotal reports of sea snakes are probably the result of confusion between the sea krait *Laticauda colubrina* commonly found on Palau and the snake eel *Myrichthys colubrinus*, indigenous to Guam. In the early 1970s there was a newspaper report of a yellow-bellied sea snake (*Pelamis platurus*) found on a Saipan beach (Eldredge, 2003). Sea snake occurrence in both pelagic and nearshore waters of the Study Area is extremely rare; therefore, sea snakes are not included in this SEIS/OEIS.

The 2015 MITT Final EIS/OEIS provided a general overview of sea turtle dive behavior, group size, and general threats. New information since the publication of the 2015 MITT Final EIS/OEIS is included below to better understand potential stressors and impacts on sea turtles resulting from training and testing activities.

**Table 3.5-1: Endangered Species Act Status and Presence of Endangered Species Act Listed Sea Turtles in the Mariana Islands Training and Testing Study Area**

| Species Name and Regulatory Status                          |                                 |  | Presence in Study Area <sup>1</sup> |                  |
|---|---------------------------------|--|-------------------------------------|------------------|
| Common Name   | Scientific Name                 | Endangered Species Act Status          | Open Ocean/Transit Corridor         | Coastal/Ocean    |
| <b>Family Cheloniidae (hard-shelled sea turtles)</b>        |                                 |  |                                     |                  |
| Green sea turtle<br>(Central West Pacific DPS) <sup>2</sup> | <i>Chelonia mydas</i>           | Endangered                             | Yes                                 | Yes <sup>3</sup> |
| Hawksbill sea turtle  | <i>Eretmochelys imbricatata</i> | Endangered                             | Yes                                 | Yes <sup>4</sup> |
| Loggerhead sea turtle                                       | <i>Caretta caretta</i>          | Endangered/<br>Threatened <sup>4</sup> | Yes <sup>5</sup>                    | Yes <sup>5</sup> |
| Olive ridley sea turtle                                     | <i>Lepidochelys olivacea</i>    | Endangered/<br>Threatened <sup>6</sup> | Yes <sup>5</sup>                    | Yes <sup>5</sup> |
| <b>Family Dermochelyidae (leatherback sea turtle)</b>       |                                 |  |                                     |                  |
| Leatherback sea turtle                                      | <i>Dermochelys coriacea</i>     | Endangered                             | Yes <sup>5</sup>                    | Yes <sup>5</sup> |

<sup>1</sup> MITT Study Area = Mariana Islands Training and Testing Study Area

<sup>2</sup> In 2015, NMFS published a final rule that classifies green sea turtles within the Study Area as part of the Western Pacific Distinct Population Segment.

<sup>3</sup> Indicates nesting activity within the Study Area. Only green sea turtles and hawksbill sea turtles are known to nest in the Study Area.

<sup>4</sup> The Northeast Atlantic Ocean, Mediterranean Sea, North Indian Ocean, North Pacific Ocean, and South Pacific Ocean Distinct Population Segments are listed as Endangered; the Northwest Atlantic Ocean, South Atlantic Ocean, Southeast Indo-Pacific Ocean, and Southwest Indian Ocean Distinct Population Segments are listed as threatened.

<sup>5</sup> Species occurrence is only expected during migratory movements through the Study Area and therefore may be present, albeit at extremely low densities.

<sup>6</sup> Breeding populations of olive ridley sea turtles on the Pacific coast of Mexico are listed as endangered and all other populations are listed as threatened. Both threatened and endangered populations could occur in the Study Area.

### 3.5.1.1 Group Size

Sea turtles are generally solitary animals, but they tend to group during migrations and mating. Because they do not show territoriality, foraging areas often overlap. New hatchlings, which often emerge from nesting beaches in groups, are solitary until they reach sexual maturity (Bolten, 2003; Bowen et al., 2004; James et al., 2005; Schroeder et al., 2003).


### 3.5.1.2 Habitat Use

Sea turtles are dependent on beaches for nesting habitat, in locations that have sand deposits that are not inundated with tides or storm events prior to hatching. In the water, sea turtle habitat use is dependent on species and corresponds to dive behavior because of foraging and migration strategies, as well as behavior state (e.g., diving deep at night for resting purposes) (Rieth et al., 2011).

### 3.5.1.3 Dive Behavior

Sea turtle dive depth and duration varies by species, the age of the animal, the location of the animal, and the activity (foraging, resting, and migrating). In addition, their relative distribution through the water column is an important consideration when conducting acoustic exposure analyses. The following text briefly describes the dive behavior of each species. Dive durations are often a function of turtle size, with larger turtles being capable of diving to greater depths and for longer periods. Methods of collecting dive behavior data over the years have varied in study design, configuration of electronic tags, parameters collected in the field, and data analyses.


New information is available that improves the analysis for sea turtle dive behavior. Hochscheid (2014) has completed a species-specific summary for sea turtles within the Study Area that was not included in the 2015 MITT Final EIS/OEIS. Hochscheid (2014) collected data from 57 studies published between 1986 and 2013, which summarized depths and durations of dives of datasets including an overall total of 538 sea turtles. Figure 3.5-1 presents the ranges of maximum dive depths for each sea turtle species found in the Study Area. This summary is used to improve exposure analysis for stressors analyzed in Section 3.5.3 (Environmental Consequences).



Sources: Hochscheid (2014), Sakamoto et al. (1993), Rice and Balazs (2008), Gitschlag (1996), Salmon et al. (2004)

**Figure 3.5-1: Dive Depth and Duration Summaries for Sea Turtle Species**

Hochscheid (2014) also collected information on generalized dive profiles, with correlations to specific activities. Generalized dive profiles compiled from 11 different studies show 8 distinct profiles tied to specific activities. These profiles and activities are shown in Figure 3.5-2.



Sources: Hochscheid (2014); Rice and Balazs (2008), Sakamoto et al. (1993), Houghton et al. (2003), Fossette et al. (2007), Salmon et al. (2004), Hays et al. (2004); Southwood et al. (1999).


Notes: Profiles A-H, as reported in the literature and compiled by Hochscheid (2014). The depth and time arrows indicate the axis variables, but the figure does not represent true proportions of depths and durations for the various profiles. In other words, the depths can vary greatly, but behavioral activity seems to dictate the shape of the profile. Profiles G and H have only been described for shallow dives (less than 5 meters).

**Figure 3.5-2: Generalized Dive Profiles and Activities Described for Sea Turtles**

#### 3.5.1.4 Hearing and Vocalization

Sea turtle ears are adapted for hearing underwater and in air, with auditory structures that may receive sound via bone conduction (Lenhardt et al., 1985), via resonance of the middle ear cavity (Willis et al., 2013), or via standard tympanic middle ear path (Hetherington, 2008). Studies of hearing ability show that sea turtles' ranges of in-water hearing detection generally lie between 50 and 1,600 hertz (Hz), with maximum sensitivity between 100 and 400 Hz, and that hearing sensitivity drops off rapidly at higher frequencies. Sea turtles are also limited to low frequency hearing in air, with hearing detection in juveniles possible between 50 to 800 Hz, with a maximum hearing sensitivity around 300–400 Hz (Bartol & Ketten, 2006; Piniak et al., 2016). Hearing abilities have primarily been studied with sub-adult, juvenile, and hatchling subjects in four sea turtle species, including green (Bartol & Ketten, 2006; Ketten & Moein-Bartol, 2006; Piniak et al., 2016; Ridgway et al., 1969; Yudhana et al., 2010), olive ridley (Bartol & Ketten, 2006), loggerhead (Bartol et al., 1999; Lavender et al., 2014; Martin et al., 2012), and leatherback (Dow Piniak et al., 2012). Only one study examined the auditory capabilities of an adult sea turtle (Martin et al., 2012); the hearing range of the adult loggerhead sea turtle was similar to other measurements of juvenile and hatchling sea turtle hearing ranges.

Using existing data on sea turtle hearing sensitivity, the U.S. Department of the Navy (Navy) developed a composite sea turtle audiogram for underwater hearing (Figure 3.5-3), as described in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).



Source: U.S. Department of the Navy (2017a)

Notes: dB re 1 µPa = decibels referenced to 1 micropascal, kHz = kilohertz

**Figure 3.5-3: Composite Underwater Audiogram for Sea Turtles**

The role of underwater hearing in sea turtles is unclear. Sea turtles may use acoustic signals from their environment as guideposts during migration and as cues to identify their natal beaches (Lenhardt et al., 1983). However, they may rely more on other senses, such as vision and magnetic orientation, to interact with their environment (Avens, 2003; Narazaki et al., 2013).

Sea turtles are not known to vocalize underwater. Some sounds have been recorded during nesting activities ashore, including belch-like sounds and sighs (Mrosovsky, 1972), exhale/inhales, gular pumps, and grunts (Cook & Forrest, 2005) by nesting female leatherback sea turtles and low-frequency pulsed and harmonic sounds by leatherback embryos in eggs and hatchlings (Ferrara et al., 2014).

#### **3.5.1.5 General Threats**

The general threats to sea turtles are described in the 2015 MITT Final EIS/OEIS. New information is available that provides a more refined understanding of how marine debris, potential invasive species introductions, and climate change can potentially threaten sea turtle species within the Study Area. Since the publication of the 2015 MITT Final EIS/OEIS, NMFS has classified green sea turtles occurring within the Mariana Islands as the Central West Pacific DPS. By doing so, the NMFS further defined threats to green sea turtles within this DPS; these threats are described below under species-specific threats for the green sea turtle. Although the information summarized below is from more recent literature since the publication of the 2015 MITT Final EIS/OEIS, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid, including the general threats to sea turtles.

##### **3.5.1.5.1 Marine Debris**

Ingestion of marine debris can cause injury or mortality to sea turtles if the debris does not pass through the digestive track. The United Nations Environment Program estimates that approximately 6.4 million tons of anthropogenic debris enters the marine environment every year (United Nations Environmental Program, 2005). This estimate, however, does not account for cataclysmic events, such as the 2011 Japanese tsunami estimated to have generated 1.5 million tons of floating debris (Murray et al., 2015). Plastic is the primary type of debris found in marine and coastal environments, and plastics are the most common type of marine debris ingested by sea turtles (Schuyler et al., 2014). Sea turtles can mistake

debris for prey; one study found 37 percent of dead leatherback sea turtles ingested various types of plastic (Mrosofsky et al., 2009), and Narazaki et al. (2013) observed a loggerhead exhibiting hunting behavior on approach to a plastic bag, possibly mistaking the bag for a jelly fish. Ingesting even small amounts of plastic can cause an obstruction in a sea turtle's digestive track and mortality (Bjorndal et al., 1994; Bjorndal, 1997), and hatchlings are at risk for ingesting small plastic fragments. Plastics absorb toxins, such as bisphenol-A (commonly known as "BPA") and phthalates, as well as heavy metals from the ocean, and can be harmful to the tissues when ingested. (Fukuoka et al., 2016; Teuten et al., 2007). Life stage and feeding preference affects the likelihood of ingestion. Sea turtles living in oceanic or coastal environments and feeding in the open ocean or on the seafloor may encounter different types and densities of debris, and may therefore have different probabilities of ingesting debris. In 2014, Schuyler et al. (2014) reviewed 37 studies of debris ingestion by sea turtles, showing that young oceanic sea turtles are more likely to ingest debris (particularly plastic), and that green and loggerhead sea turtles were significantly more likely to ingest debris than other sea turtle species.

#### 3.5.1.5.2 Invasive Species

Impacts on sea turtles associated with invasive species primarily concern nest predation and prey base. Some of the invasive species introduced to the larger, more populated islands in the Mariana archipelago are known nest predators (e.g., rats, feral dogs and cats, pigs, ants). Nests on populated islands are also at risk for illegal poaching (Kolinski et al., 2006). In foraging grounds, sea turtles have been shown to adapt their foraging preferences for invasive seagrass and algae. Becking et al. (2014) showed green sea turtle foraging behavior shift to consumption of *Halophila stipulacea*, a rapidly spreading seagrass in the Caribbean. In Hawaii, green sea turtles in Kaneohe Bay have modified their diets over several decades to include seven non-native species (*Acanthophora spicifera*, *Hypnea musciformis*, *Gracilaria salicornia*, *Euclima denticulatum*, *Gracilaria tikvahiae*, *Kappaphycus striatum*, and *Kappaphycus alvarezii*), with non-native algae accounting for over 60 percent of sea turtle diet (Russell & Balazs, 2015).

Since the publication of the 2015 MITT Final EIS/OEIS, the Navy has funded the *Regional Biosecurity Plan for Micronesia and Hawaii*, completed in 2015. Volume I, Appendix K of the biosecurity plan addresses general biosecurity recommendations for Guam and the Commonwealth of the Northern Mariana Islands, and Appendix M includes recommendations for U.S. Department of Defense activities (U.S. Department of the Navy, 2015d). Volume III includes a risk assessment for marine environments (U.S. Department of the Navy, 2015c), and Volume IV includes a risk assessment for potential introductions on land in terrestrial environments (U.S. Department of the Navy, 2015b). The 2015 biosecurity plan describes ongoing measures that reduce the potential for transport and introduction of invasive species resulting from military training and testing activities. Some of these species have the potential to degrade sea turtle habitats, reduce prey availability, or directly harm sea turtles. Because of the Navy's active biosecurity program, it is unlikely that training and testing activities would result in invasive species' introductions that would impact sea turtles. Therefore, invasive species are not analyzed as a new stressor in this SEIS/OEIS.

#### 3.5.1.5.3 Climate Change

Since the publication of the MITT Final EIS/OEIS, the Navy has obtained and consolidated additional information to conceptualize the potential of climate change to threaten sea turtle species within the Study Area. Sea turtles are particularly susceptible to climate change effects because their life history, physiology, and behavior are extremely sensitive to environmental temperatures (Fuentes et al., 2013). Climate change models predict sea level rise and increased intensity of storms and hurricanes in tropical

sea turtle nesting areas (Patino-Martinez et al., 2008). These factors could significantly increase beach inundation and erosion, thus affecting water content of sea turtle nesting beaches and potentially inundating nests (Pike et al., 2015). Climate change may negatively impact turtles in multiple ways and at all life stages. These impacts may include the potential loss of nesting beaches due to sea level rise and increasingly intense storm surge (Patino-Martinez et al., 2008), feminization of turtle populations from elevated nest temperatures (and skewing populations from more males to females unless nesting shifts to northward cooler beaches) (Reneker & Kamel, 2016), decreased reproductive success (Clark & Gobler, 2016; Hawkes et al., 2006; Laloë et al., 2016; Pike, 2014), shifts in reproductive periodicity and latitudinal ranges (Birney et al., 2015; Pike, 2014), disruption of hatchling dispersal and migration, and indirect effects to food availability (Witt et al., 2010).

#### **3.5.1.6 Green Sea Turtle (*Chelonia mydas*)**

This section has been updated based on a change in the regulatory status of the green sea turtle and new information regarding trends and distributions of green sea turtles in nearshore waters of the Mariana Islands. As such, the life history and regulatory status descriptions for each sea turtle species differs in detail.

##### **3.5.1.6.1 Status and Management**

As presented in the 2015 MITT Final EIS/OEIS, green sea turtles are listed as threatened under the ESA throughout their Pacific range, except for the population that nests on the Pacific coast of Mexico (endangered). However, NMFS and United States Fish and Wildlife Service (USFWS) reclassified the species in 2016 into 11 DPSs, which maintains federal protections while providing a more tailored approach for managers to address specific threats facing different populations (see the NMFS and USFWS Final Rule published on April 6, 2016). Only the Central West Pacific DPS occurs within the Study Area. This DPS is listed as endangered under the ESA. Only this distinct population segment is discussed further in the document; however, it should be noted that minimal mixing (gene flow) may occur with other distinct population segments (Seminoff et al., 2015).

##### **3.5.1.6.2 Habitat and Geographic Range**

The habitat and geographic range of green sea turtles is described in the 2015 MITT Final EIS/OEIS. Following a review of recent literature, information on green sea turtles related to habitat and geographic range has not changed since the publication of the MITT Final EIS/OEIS. As such, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid. There is no critical habitat designated for this species in the Study Area. Seminoff et al. (2015) however, provides specific information for Central West Pacific DPS, and determined that this DPS is spatially bounded by the Asian continent to the west and north, the Solomon Islands to the south, the Marshall Islands in the east, and Palau in the west.

##### **3.5.1.6.2.1 Population and Abundance**

The population and abundance of green sea turtles is described in the 2015 MITT Final EIS/OEIS; however, new information is available for estimating abundance in waters within the Study Area. Martin et al. (2016) analyzed five decades of aerial surveys (from 1962 through 2012) to assess changes in marine megafauna on the insular coral reef ecosystem of Guam. Turtle observations increased and varied spatially around Guam, with the highest densities occurring along the south, east, and north coasts, particularly in areas having low human density, reefs with coral cover, and either seagrass beds or a marine protected area. Observed individuals per survey ranged from 1.1 to 44.6 across all years. Based on this information, Martin et al. (2016) calculated a population growth rate of approximately

90 percent over the past five decades. Based on studies of in-water capture rates (where swimmers would capture and tag individual sea turtles), Martin et al. (2016) estimated that 85 percent of the sea turtles in waters off of Guam are green sea turtles, while 15 percent are hawksbill sea turtles. The Navy is currently funding in-water tagging of sea turtles in waters off of Guam, Tinian, and Saipan. Since November 2015 when tagging began, Falcone et al. (2017) report that the majority of sea turtles observed or captured (65 of 68 total sea turtles observed, or 96 percent) have been green sea turtles.

#### **3.5.1.6.2.2 Predator-Prey Interactions**

The predator-prey interactions relevant to green sea turtles are described in the 2015 MITT Final EIS/OEIS. Following a review of recent literature, information on green sea turtles related to predator-prey interactions has not changed since the publication of the 2015 MITT Final EIS/OEIS. As such, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid. When NMFS classified green sea turtles within the Central West Pacific DPS, no information on predator-prey interactions were used that were not included in the 2015 MITT Final EIS/OEIS.

#### **3.5.1.6.2.3 Species-Specific Threats**

Since the publication of the 2015 MITT Final EIS/OEIS, the NMFS has further defined threats to green sea turtles included in the West Pacific DPS. Damage to seagrass beds and declines in seagrass distribution can reduce foraging habitat for green sea turtles (National Marine Fisheries Service & U.S. Fish and Wildlife Service, 1991; Seminoff et al., 2015; Williams, 1988). Green sea turtles are susceptible to the disease fibropapillomatosis, which causes tumor-like growths (fibropapillomas) resulting in reduced vision, disorientation, blindness, physical obstruction to swimming and feeding, increased susceptibility to parasites, and increased susceptibility to entanglement (Balazs, 1986; National Marine Fisheries Service & U.S. Fish and Wildlife Service, 1991; Patrício et al., 2016; Work & Balazs, 2013). The potential effects of disease and endoparasites also exist for green sea turtles found in the Central West Pacific Ocean. The loss of eggs to non-human predators is a severe problem in some areas. These predators include domestic animals, such as cats, dogs, and pigs, as well as wild species such as rats, mongoose, birds, monitor lizards, snakes, crabs, ants, and other invertebrates (Seminoff et al., 2015).

#### **3.5.1.7 Hawksbill Sea Turtle (*Eretmochelys imbricata*)**

The hawksbill sea turtle is listed as endangered under the ESA (35 Federal Register 8491). While the current listing as a single global population remains valid, data may support separating populations at least by ocean basin under the distinct population segment policy (National Marine Fisheries Service, 2013). The most recent status review was released in 2013 by the NMFS and USFWS (National Marine Fisheries Service, 2013). There is no critical habitat designated for this species in the Study Area. The regulatory status for the hawksbill sea turtle has remained unchanged since the publication of the 2015 MITT Final EIS/OEIS.

In addition, the life history information for hawksbill sea turtles occurring in nearshore and open ocean habitats within the Study Area has not changed since the publication of the 2015 MITT Final EIS/OEIS. New information is available for estimating hawksbill sea turtle population and abundance based off of five decades of aerial surveys in the nearshore waters of Guam. While Martin et al. (2016) estimated that approximately 15 percent of sea turtles observed in waters off of Guam are hawksbill sea turtles, tagging from November 2015 has revealed that only 4 percent of observed turtles are hawksbill sea turtles. Overall, the trend data over this time period suggests a dramatic increase in sea turtle populations in waters around Guam. The Navy is currently funding in-water tagging of sea turtles in waters off of Guam, Tinian, and Saipan.

### 3.5.1.8 Loggerhead Sea Turtle (*Caretta caretta*)

In 2009, a status review was conducted for the loggerhead identified nine distinct population segments within the global population (Conant et al., 2009). In 2011, NMFS and USFWS listed five of these distinct population segments as endangered and kept four as threatened under the ESA. Only the North Pacific Ocean distinct population segment occurs within the Study Area; however, mixing is known to occur between other populations in the Pacific and Indian Oceans, enabling a limited amount of gene flow with other distinct population segments (National Marine Fisheries Service & U.S. Fish and Wildlife Service, 2008). There is no critical habitat designated for this species in the Study Area. The regulatory status for the loggerhead sea turtle has remained unchanged since the publication of the 2015 MITT Final EIS/OEIS. In addition, the life history information for loggerhead sea turtles occurring in nearshore and open ocean habitats within the Study Area has not changed since the publication of the 2015 MITT Final EIS/OEIS. As such, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid.

### 3.5.1.9 Olive Ridley Sea Turtle (*Lepidochelys olivacea*)

Olive ridley sea turtles that nest along the Pacific coast of Mexico are listed as endangered under the ESA, while all other populations are listed under the ESA as threatened (43 Federal Register 32800). Based on genetic data, the worldwide olive ridley population is composed of four main lineages: east India, Indo-Western Pacific, Atlantic, and eastern Pacific Ocean (National Marine Fisheries Service & U.S. Fish and Wildlife Service, 2014; Shankar et al., 2004). Most olive ridley sea turtles found within the Study Area are of the Indo-Western Pacific lineage (National Marine Fisheries Service & U.S. Fish and Wildlife Service, 2014). There is no critical habitat designated for this species in the Study Area. The regulatory status for the olive ridley sea turtle has remained unchanged since the publication of the 2015 MITT Final EIS/OEIS.

In addition, the life history information for olive ridley sea turtles occurring in nearshore and open ocean habitats within the Study Area has not changed since the publication of the 2015 MITT Final EIS/OEIS. As such, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid.

### 3.5.1.10 Leatherback Sea Turtle (*Dermochelys coriacea*)

The leatherback sea turtle is listed as a single population and is classified as endangered under the ESA (35 Federal Register 8491). Recent information on population structure (through genetic studies) and distribution (through telemetry, tagging, and genetic studies) have led to an increased understanding and refinement of the global stock structure (Clark et al., 2010). There is no critical habitat designated for this species in the Study Area. The regulatory status for the leatherback sea turtle has remained unchanged since the publication of the 2015 MITT Final EIS/OEIS. In addition, the life history information for leatherback sea turtles occurring in nearshore and open ocean habitats within the Study Area has not changed since the publication of the 2015 MITT Final EIS/OEIS. As such, the information and analysis presented in the 2015 MITT Final EIS/OEIS remains valid.

## 3.5.2 Environmental Consequences

Under the Proposed Action for this SEIS/OEIS, there have been some modifications to the quantity and type of acoustic stressors under the two action alternatives. Additionally, one new substressor (high-energy lasers) is being analyzed because of its potential to affect marine species, as detailed in Section 3.0.4.3.2.2 (High-Energy Lasers).

In general, there have been no substantial changes to the activities analyzed as the Proposed Action in the 2015 MITT Final EIS/OEIS which would change the conclusions reached regarding populations of sea turtles in the Study Area. Acoustic stressors (sonar and other transducers) and explosives have occurred since the 2015 completion of the MITT Record of Decision and ESA Biological Opinion. There have been no known impacts on sea turtles that were not otherwise previously analyzed or accounted for in the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015a), the NMFS Biological Opinion pursuant to ESA (National Marine Fisheries Service, 2015a), or the USFWS Biological Opinion.

In this SEIS/OEIS, the Navy has reviewed the analysis of impacts from these ongoing activities and additionally analyzed the new or changing training and testing activities as projected into the reasonably foreseeable future. The projected future actions are based on evolving operational requirements, including those associated with any anticipated new platforms or systems not previously analyzed. The Navy has completed a literature review for information on sea turtles within the Study Area, which included a search for the best available science since the publication of the 2015 MITT Final EIS/OEIS. Where there has been no substantive or otherwise meaningful change in the action, science, or regulations, the Navy will rely on the previous 2015 MITT Final EIS/OEIS analysis. Where there has been substantive change in the action, science, or regulations, the information and analysis provided in in this SEIS/OEIS will supplement the 2015 MITT Final EIS/OEIS to support environmental compliance with applicable environmental statutes for sea turtles.

The stressors applicable to sea turtles in the Study Area for this SEIS/OEIS include the new stressor (high-energy lasers) and the same stressors considered in the 2015 MITT Final EIS/OEIS:

- **Acoustic** (sonar and other transducers, vessel noise, aircraft noise, and weapon noise)
- **Explosive** (in-air explosions and in-water explosions)
- **Energy** (in-water electromagnetic devices, high-energy lasers)
- **Physical disturbance and strike** (vessels and in-water devices, military expended materials, seafloor devices)
- **Entanglement** (wires and cables, decelerators/parachutes)
- **Ingestion** (military expended materials – munitions and military expended materials – other than munitions)
- **Secondary** (impacts on habitat, impacts on prey availability)

This section of this SEIS/OEIS evaluates how and to what degree potential impacts on sea turtles from stressors described in Section 3.0 (Introduction) may have changed since the analysis presented in the 2015 MITT Final EIS/OEIS was completed. Table 2.5-1 and Table 2.5-2 in Chapter 2 (Description of Proposed Action and Alternatives) list the proposed training and testing activities and include the number of times each activity would be conducted annually and the locations within the Study Area where the activity would typically occur under each alternative. The tables also present the same information for activities described in the 2015 MITT Final EIS/OEIS so that the proposed levels of training and testing under this SEIS/OEIS can be easily compared. The analysis in this SEIS/OEIS includes consideration of the Navy's standard operating procedures and mitigation that the Navy will implement to avoid or reduce potential impacts on sea turtles from acoustic, explosive, and physical disturbance and strike stressors. Mitigation for sea turtles will be coordinated with NMFS through the ESA consultation processes, and is detailed in Chapter 5 (Mitigation) and Appendix I (Geographic Mitigation Assessment) of this SEIS/OEIS.

In their biological opinion, NMFS determined that within the Study Area, only acoustic stressors and explosive stressors could potentially result in adverse effects on ESA-listed sea turtles from training and testing activities and that none of the other stressors would result in significant adverse impacts or jeopardize the continued existence of any ESA-listed sea turtle (National Oceanic and Atmospheric Administration, 2015).

The analysis presented in this section of this SEIS/OEIS also considers standard operating procedures that are described in Chapter 2 (Description of Proposed Action and Alternatives) and mitigation measures that are described in Chapter 5 (Mitigation). The Navy will implement these measures to avoid or reduce potential impacts on sea turtles from stressors associated with the proposed training and testing activities. Mitigation for sea turtles will be coordinated with NMFS through the ESA consultation process.

As presented in Section 3.0 (Introduction), since completion of the 2015 MITT Final EIS/OEIS there have been refinements made in the modeling of estimated impacts from sonar and other transducers and in-water explosives. These changes have been incorporated into the re-analysis of acoustic and explosive stressors presented in this SEIS/OEIS. In addition to the new effects criteria, weighting functions, and thresholds across multiple species, new information for sea turtles includes the integration of new sea turtle density data based on new survey data.

#### **3.5.2.1 Acoustic Stressors**

The analysis of effects to sea turtles follows the concepts outlined in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities). This section begins with a summary of relevant data regarding acoustic impacts on sea turtles in Section 3.5.2.1.1 (Background). This is followed by an analysis of estimated impacts on sea turtles due to specific Navy acoustic stressors (sonar and other transducers, vessel noise, aircraft noise, and weapon noise). Additional explanations of the acoustic terms and sound energy concepts used in this section are found in Appendix H (Acoustic and Explosive Concepts). Studies of the effects of sound on sea turtles are limited; therefore, where necessary, knowledge of impacts on other species from acoustic stressors is used to assess impacts on sea turtles.

The Navy will rely on the previous 2015 MITT Final EIS/OEIS for the analysis of vessel noise, aircraft noise, and weapon noise, and new applicable and emergent science in regard to these substressors is presented in the sections that follow. Due to new acoustic impact criteria, sea turtle densities, and acoustic effects model, the analysis provided in Section 3.5.2.1.2 (Impacts from Sonar and Other Transducers) of this SEIS/OEIS will supplant the 2015 MITT Final EIS/OEIS for sea turtles, and may result in changes to estimated impacts for some species since the 2015 MITT Final EIS/OEIS.

##### **3.5.2.1.1 Background**

The sections below include a survey and synthesis of best-available-science published in peer-reviewed journals, technical reports, and other scientific sources pertinent to impacts on sea turtles potentially resulting from Navy training and testing activities. Sea turtles could be exposed to a range of impacts depending on the sound source and context of the exposure. Exposures to sound-producing activities may result in auditory or non-auditory trauma, hearing loss resulting in temporary or permanent hearing threshold shift, auditory masking, physiological stress, or changes in behavior.

### 3.5.2.1.1.1 Injury

The high peak pressures close to some non-explosive impulsive underwater sound sources may be injurious, although there are no reported instances of injury to sea turtles caused by these sources. A Working Group organized under the American National Standards Institute-Accredited Standards Committee S3, Subcommittee 1, Animal Bioacoustics, developed sound exposure guidelines for fish and sea turtles (Popper et al., 2014), hereafter referred to as the *ANSI Sound Exposure Guidelines*. Lacking any data on non-auditory sea turtle injuries due to sonars, the working group estimated the risk to sea turtles from low-frequency sonar to be low and mid-frequency sonar to be non-existent.

As discussed in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities, specifically Section 3.0.4.7.1, Injury), mechanisms for non-auditory injury due to acoustic exposure have been hypothesized for diving breath-hold animals. Acoustically induced bubble formation, rectified diffusion, and acoustic resonance of air cavities are considered for their similarity to pathologies observed in marine mammals stranded coincident with sonar exposures but were found to not be likely causal mechanisms (Section 3.5.2.1.1.1, Injury), and findings are applicable to sea turtles.

Nitrogen decompression due to modifications to dive behavior has never been observed in sea turtles. Sea turtles are thought to deal with nitrogen loads in their blood and other tissues, caused by gas exchange from the lungs under conditions of high ambient pressure during diving, through anatomical, behavioral, and physiological adaptations (Lutcavage & Lutz, 1997). Although diving sea turtles experience gas supersaturation, gas embolism has only been observed in sea turtles bycaught in fisheries (Garcia-Parraga et al., 2014). Therefore, nitrogen decompression due to changes in diving behavior is not considered a potential consequence to diving reptiles.

### 3.5.2.1.1.2 Hearing Loss

Exposure to intense sound may result in hearing loss, typically quantified as threshold shift, which persists after cessation of the noise exposure. Threshold shift is a loss of hearing sensitivity at an affected frequency of hearing. This noise-induced hearing loss may manifest as temporary threshold shift (TTS), if hearing thresholds recover over time, or permanent threshold shift (PTS), if hearing thresholds do not recover to pre-exposure thresholds. Because studies on inducing threshold shift in reptiles are very limited (e.g., alligator lizards: Dew et al., 1993; Henry & Mulroy, 1995), are not sufficient to estimate TTS and PTS onset thresholds, and have not been conducted on any of the reptiles present in the Study Area, auditory threshold shift in sea turtles is considered to be consistent with general knowledge about noise-induced hearing loss described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

Because there are no data on auditory effects on sea turtles, the *ANSI Sound Exposure Guidelines* (Popper et al., 2014) do not include numeric sound exposure thresholds for auditory effects on sea turtles. Rather, the guidelines qualitatively estimate that sea turtles are less likely to incur TTS or PTS with increasing distance from various sound sources. The guidelines also suggest that data from fishes may be more relevant than data from marine mammals when estimating impacts on sea turtles, because, in general, fish hearing range is more similar to the limited hearing range of sea turtles. As shown in Section 3.5.1.4 (Hearing and Vocalization), sea turtle hearing is most sensitive around 100–400 Hz in-water, is limited over 1 kilohertz (kHz), and is much less sensitive than that of any marine mammal. Therefore, sound exposures from most mid-frequency and all high-frequency sound sources are not anticipated to affect sea turtle hearing, and sea turtles are likely only susceptible to auditory impacts when exposed to very high levels of sound within their limited hearing range.

### 3.5.2.1.1.3 Physiological Stress

A stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor. If the magnitude and duration of the stress response is too great or too long, then it can have negative consequences to the animal (e.g., decreased immune function, decreased reproduction). Physiological stress is typically analyzed by measuring stress hormones, other biochemical markers, or vital signs. Physiological stress has been measured for sea turtles during nesting (Flower et al., 2015; Valverde et al., 1999), capture and handling (Flower et al., 2015; Gregory & Schmid, 2001), and when caught in entanglement nets (Hoopes et al., 2000; Snoddy et al., 2009) and trawls (Stabenau et al., 1991). However, the stress caused by acoustic exposure has not been studied for reptiles. Therefore, the stress response in sea turtles in the Study Area due to acoustic exposures is considered to be consistent with general knowledge about physiological stress responses described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

Marine animals naturally experience stressors within their environment and as part of their life histories. Changing weather and ocean conditions, exposure to diseases and naturally occurring toxins, lack of prey availability, social interactions with members of the same species, nesting, and interactions with predators all contribute to stress. Anthropogenic sound-producing activities have the potential to provide additional stressors beyond those that naturally occur.

Due to the limited information about acoustically induced stress responses, the Navy conservatively assumes in its effects analysis that any physiological response (e.g., hearing loss or injury) or significant behavioral response is also associated with a stress response.

### 3.5.2.1.1.4 Masking

As described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), auditory masking occurs when one sound, distinguished as the “noise,” interferes with the detection or recognition of another sound or limits the distance over which other biologically relevant sounds, including those produced by prey, predators, or conspecifics, can be detected. Masking only occurs when the sound source is operating; therefore, direct masking effects stop immediately upon cessation of the sound-producing activity. Any sound above ambient noise and within an animal’s hearing range may potentially cause masking.

Compared to other marine animals, such as marine mammals that are highly adapted to use sound in the marine environment, marine reptile hearing is limited to lower frequencies and is less sensitive. Because marine reptiles likely use their hearing to detect broadband low-frequency sounds in their environment, the potential for masking would be limited to certain similar sound exposures. Only continuous human-generated sounds that have a significant low-frequency component, are not brief in duration, and are of sufficient received level, would create a meaningful masking situation (e.g., proximate vessel noise). Other intermittent, short-duration sound sources with low-frequency components (e.g., low-frequency sonars) would have more limited potential for masking depending on duty cycle.

There is evidence that reptiles may rely primarily on senses other than hearing for interacting with their environment, such as vision (Narazaki et al., 2013) and magnetic orientation (Avens, 2003; Putman et al., 2015). Any effect of masking may be mediated by reliance on other environmental inputs.

### 3.5.2.1.1.5 Behavioral Reactions

Behavioral responses fall into two major categories: Alterations in natural behavior patterns and avoidance. These types of reactions are not mutually exclusive and reactions may be combinations of behaviors or a sequence of behaviors. As described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), the response of a reptile to an anthropogenic sound would likely depend on the frequency, duration, temporal pattern, and amplitude of the sound as well as the animal's prior experience with the sound and the context in which the sound is encountered (i.e., what the animal is doing at the time of the exposure). Distance from the sound source and whether it is perceived as approaching or moving away may also affect the way a reptile responds to a sound.

Sea turtles may detect sources below 2 kHz but have limited hearing ability above 1 kHz. They likely detect most broadband sources (including vessel noise) and low-frequency sonars, so they may respond to these sources. Because auditory abilities are poor above 1 kHz, detection and consequent reaction to any mid-frequency source is unlikely.

In the *ANSI Sound Exposure Guidelines* (Popper et al., 2014), qualitative risk factors were developed to assess the potential for sea turtles to respond to various underwater sound sources. The guidelines state that there is a low likelihood that sea turtles would respond within tens of meters of low-frequency sonars, and that it is highly unlikely that sea turtles would respond to mid-frequency sources. The risk that sea turtles would respond to other broadband sources, such as shipping, is considered high within tens of meters of the sound source, but moderate to low at farther distances.

#### **Behavioral Reactions to Impulsive Sound Sources**

There are limited studies of reptile responses to sounds from impulsive sound sources, and all data come from sea turtles exposed to seismic air guns, although air guns are not used during MITT training or testing activities. These exposures consist of multiple air gun shots, either in close proximity or over long durations, so it is likely that observed responses may over-estimate responses to single or short-duration impulsive exposures. Studies of responses to air guns are used to inform reptile responses to other impulsive sounds (e.g., some weapon noise).

O'Hara and Wilcox (1990) attempted to create a sound barrier at the end of a canal using seismic air guns. They reported that loggerhead sea turtles kept in a 300-meter by 45-meter enclosure in a 10-meter deep canal maintained a minimum standoff range of 30 meters from air guns fired simultaneously at intervals of 15 seconds with strongest sound components within the 25–1,000 Hz frequency range. McCauley et al. (2000) estimated that the received sound pressure level (SPL) at which turtles avoided sound in the O'Hara and Wilcox (1990) experiment was 175–176 decibels referenced to 1 micropascal (dB re 1  $\mu$ Pa).

Moein Bartol et al. (1995) investigated the use of air guns to repel juvenile loggerhead sea turtles from hopper dredges. Sound frequencies of the air guns ranged from 100 to 1,000 Hz at three source SPLs: 175, 177, and 179 dB re 1  $\mu$ Pa at 1 m. The turtles avoided the air guns during the initial exposures (mean range of 24 m), but additional exposures on the same day and several days afterward did not elicit statistically significant avoidance behavior. They concluded that this was likely due to habituation.

McCauley et al. (2000) exposed a caged green and a caged loggerhead sea turtle to an approaching-departing single air gun to gauge behavioral responses. The trials showed that above a received SPL of 166 dB re 1  $\mu$ Pa, the turtles noticeably increased their swimming activity compared to nonoperational periods, with swimming time increasing as air gun SPLs increased during approach.

Above 175 dB re 1  $\mu$ Pa, behavior became more erratic, possibly indicating the turtles were in an agitated state. The authors noted that the point at which the turtles showed more erratic behavior and exhibited possible agitation would be expected to approximate the point at which active avoidance to air guns would occur for unrestrained turtles.

No obvious avoidance reactions by free-ranging sea turtles, such as swimming away, were observed during a multi-month seismic survey using air gun arrays, although fewer sea turtles were observed when the seismic air guns were active than when they were inactive (Weir, 2007). The author noted that sea state and the time of day affected both air gun operations and sea turtle surface basking behavior, making it difficult to draw conclusions from the data. However, DeRuiter and Doukara (2012) noted several possible startle or avoidance reactions to a seismic air gun array in the Mediterranean by loggerhead sea turtles that had been motionlessly basking at the water surface.

### **Behavioral Reactions to Sonar and Other Transducers**

Studies of sea turtle responses to non-impulsive sounds are very limited. Lenhardt (1994) used very low frequency vibrations (< 100 Hz) coupled to a shallow tank to elicit swimming behavior responses by two loggerhead sea turtles. Watwood et al. (2016) tagged green sea turtles with acoustic transponders and monitored them using acoustic telemetry arrays in Port Canaveral, FL. Sea turtles were monitored before, during, and after a routine pier-side submarine sonar test that utilized typical source levels, signals, and duty cycle. The sea turtles did not exhibit significant long-term displacement in this study. The authors note that Port Canaveral is an urban marine habitat and that resident sea turtles may be less likely to respond than naïve populations.

#### **3.5.2.1.1.6 Long-Term Consequences**

For the sea turtles present in the Study Area, long-term consequences to individuals and populations due to acoustic exposures have not been studied. Therefore, long-term consequences to sea turtles due to acoustic exposures are considered following the framework presented in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

The long-term consequences due to individual behavioral reactions and short-term (seconds to minutes) instances of physiological stress are especially difficult to predict because individual experience over time can create complex contingencies. It is more likely that any long-term consequences to an individual would be a result of costs accumulated over a season, year, or life stage due to multiple behavioral or stress responses resulting from exposures to multiple stressors over significant periods of time. Conversely, some sea turtles may habituate to or become tolerant of repeated acoustic exposures over time, learning to ignore a stimulus that in the past did not accompany any overt threat. For example, loggerhead sea turtles exposed to air guns with a source SPL of 179 dB re 1  $\mu$ Pa initially exhibited avoidance reactions. However, they may have habituated to the sound source after multiple exposures, since a habituation behavior was retained when exposures were separated by several days (Moein Bartol et al., 1995). Intermittent exposures are assumed to be less likely to have lasting consequences.

#### **3.5.2.1.2 Impacts from Sonar and Other Transducers**

The overall use of sonar and other transducers for training and testing would be similar to what is currently conducted (see Table 2.5-1 and Table 3.0-2 for details). Although individual activities may vary somewhat from those previously analyzed, the overall determinations presented in the 2015 MITT Final EIS/OEIS remain valid. In addition, some new systems using new technologies would be tested under Alternatives 1 and 2. The quantitative analysis has been updated since the 2015 MITT Final EIS/OEIS;

therefore, the new analysis is fully presented and described in further detail in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2017b).

Activities using sonars and other transducers would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). Sonar and other transducers proposed for use are transient in most locations because activities that involve sonar and other transducers take place at different locations and many platforms are generally moving throughout the Study Area. Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. General categories of these systems are described in Section 3.0.4.1 (Acoustic Stressors). The activities that use sonar and other transducers are described in Appendix A (Training and Testing Activities Descriptions).

Sonar-induced acoustic resonance and bubble formation phenomena are very unlikely to occur under realistic conditions, as discussed in Section 3.5.2.1.1.1 (Injury). Non-auditory injury (i.e., other than PTS) and mortality from sonar and other transducers is so unlikely as to be discountable under normal conditions and is therefore not considered further in this analysis.

Potential impacts considered from exposure to sonar and other transducers are hearing loss due to threshold shift (permanent or temporary), physiological stress, masking of other biologically relevant sounds, and changes in behaviors, as described in Sections 3.5.2.1.1.2 (Hearing Loss and Auditory Injury), Section 3.5.2.1.1.3 (Physiological Stress), Section 3.5.2.1.1.4 (Masking) and Section 3.5.2.1.1.5 (Behavioral Reactions).

#### **3.5.2.1.2.1 Methods for Analyzing Impacts from Sonar and Other Transducers**

The Navy performed a quantitative analysis to estimate the number of times that sea turtles could be affected by sonars and other transducers used during Navy training and testing activities. The Navy's quantitative analysis to determine impacts on sea turtles and marine mammals uses the Navy Acoustic Effects Model to produce initial estimates of the number of times these animals may experience these effects; these estimates are further refined by considering animal avoidance of sound-producing activities and implementation of mitigation. The steps of this quantitative analysis takes into account

- criteria and thresholds used to predict impacts from sonar and other transducers (see below);
- the density and spatial distribution of sea turtles; and
- the influence of environmental parameters (e.g., temperature, depth, salinity) on sound propagation when estimating the received sound level on the animals.


A further detailed explanation of this analysis is provided in the technical report titled *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

#### **Criteria and Thresholds Used to Predict Impacts from Sonar and Other Transducers**

##### **Auditory Weighting Functions**

Animals are not equally sensitive to noise at all frequencies. To capture the frequency-dependent nature of the effects of noise, auditory weighting functions are used. Auditory weighting functions are mathematical functions that adjust received sound levels to emphasize ranges of best hearing and de-emphasize ranges with less or no auditory sensitivity. The adjusted received sound level is referred to as a weighted received sound level.

The auditory weighting function for sea turtles is shown in Figure 3.5-4. The derivation of this weighting function is described in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a). The frequencies around the top portion of the function, where the amplitude is closest to zero, are emphasized, while the frequencies below and above this range (where amplitude declines) are de-emphasized, when summing acoustic energy received by a sea turtle.




Source: U.S. Department of the Navy (2017a)

Notes: dB = decibels, kHz = kilohertz, TU = sea turtle species group

**Figure 3.5-4: Auditory Weighting Function for Sea Turtles**

#### Hearing Loss from Sonar and Other Transducers

No studies of hearing loss have been conducted on sea turtles. Therefore, sea turtle susceptibility to hearing loss due to an acoustic exposure is evaluated using knowledge about sea turtle hearing abilities in combination with non-impulsive auditory effect data from other species (marine mammals and fish). This yields sea turtle exposure functions, shown in Figure 3.5-5, which are mathematical functions that relate the sound exposure levels (SELs) for onset of TTS or PTS to the frequency of the sonar sound exposure. The derivation of the sea turtle exposure functions are provided in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).



Source: U.S. Department of the Navy (2017a)

Notes: dB re 1 μPa<sup>2</sup>s: decibels referenced to 1 micropascal second squared, kHz = kilohertz. The solid black curve is the exposure function for TTS and the dashed black curve is the exposure function for PTS onset. Small dashed lines and asterisks indicate the SEL thresholds at the most sensitive frequency for TTS (200 dB) and PTS (220 dB).

**Figure 3.5-5: TTS and PTS Exposure Functions for Sonar and Other Transducers**

### Accounting for Mitigation

The Navy will implement mitigation measures to avoid or reduce potential impacts from active sonar on sea turtles, as described in Section 5.3.2.1 (Active Sonar). The benefits of mitigation are conservatively factored into the analysis for Alternative 1 and Alternative 2 of the Proposed Action for training and testing. The Navy’s mitigation measures are identical for both action alternatives.

Procedural mitigation measures include a power down or shut down (i.e., power off) of applicable active sonar sources when a sea turtle is observed in a mitigation zone. The mitigation zones for active sonar activities were designed to avoid or reduce the potential for sea turtles to be exposed to levels of sound that could result in auditory injury (i.e., PTS) from active sonar to the maximum extent practicable. The mitigation zones encompass the estimated ranges to injury (including PTS) for a given sonar exposure. Therefore, the impact analysis quantifies the potential for mitigation to reduce the risk of PTS. Two factors are considered when quantifying the effectiveness of mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., active sonar) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

In the quantitative analysis, consideration of mitigation measures means that, for activities where mitigation is feasible, some model-estimated PTS is considered mitigated to the level of TTS. The impact analysis does not analyze the potential for mitigation to reduce TTS or behavioral effects, even though mitigation could also reduce the likelihood of these effects. In practice, mitigation also protects all

unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

The ability to observe the range to PTS was estimated for each training or testing event. The ability of Navy Lookouts to detect sea turtles in or approaching the mitigation zone is dependent on the animal's presence at the surface and the characteristics of the animal that influence its sightability (such as size or surface active behavior). The behaviors and characteristics of some species may make them easier to detect. Environmental conditions under which the training or testing activity could take place are also considered such as the sea surface conditions, weather (e.g., fog or rain), and day versus night.

#### **3.5.2.1.2.2 Impact Ranges for Sonar and Other Transducers**

Because sea turtle hearing range is limited to a narrow range of frequencies and thresholds for auditory impacts are relatively high, there are few sonar sources that could result in exposures exceeding the sea turtle TTS and PTS thresholds. The representative bin of LF4 for PTS and TTS is zero meters. Ranges would be greater (i.e., up to tens of meters) for sonars and other transducers with higher source levels (within their hearing range); however, specific ranges cannot be provided in an unclassified document.

#### **3.5.2.1.2.3 Alternative 1**

Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. General categories and characteristics of sonar systems and the number of hours these sonars would be operated during training and testing activities under Alternative 1 are described in Section 3.0.1.2.4.1 (Sonar and Other Transducers). Activities using sonars and other transducers would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). Low-frequency sources are operated more frequently during testing activities than during training activities. Although the general impacts from sonar and other transducers during testing would be similar in severity to those described during training, there may be slightly more impacts during testing activities as sea turtles can detect low frequency sources.

Under Alternative 1, training and testing activities would fluctuate each year to account for the natural variation of training cycles and deployment schedules. Training and testing activities, including low-frequency sonars within sea turtle hearing range (<2 kHz), could take place throughout the Study Area.

The quantitative analysis, using the number of hours of sonar and other transducers for a maximum year of training activities under Alternative 1, predicts that no sea turtles of any species are likely to be exposed to the high received levels of sound from sonars or other transducers that could cause TTS or PTS. Exposures were only modeled for green, hawksbill, leatherback, and loggerhead sea turtles in the Study Area and transit corridor. Olive ridley sea turtle presence in the Study Area is limited, and density data does not exist due to low occurrence in this region. Only a limited number of sonars and other transducers with frequencies within the range of sea turtle hearing (<2 kHz) and high source levels have the potential to cause TTS and PTS.

The *ANSI Sound Exposure Guidelines* estimate that the risk of a sea turtle responding to a low-frequency sonar (less than 1 kHz) is low regardless of proximity to the source, and that there is no risk of a sea turtle responding to a mid-frequency sonar (1–10 kHz) (Popper et al., 2014). A sea turtle could respond to sounds detected within their limited hearing range if they are close enough to the source. The few studies of sea turtle reactions to sounds, discussed in Section 3.5.2.1.1.5 (Behavioral Reactions), suggest

that a behavioral response could consist of temporary avoidance, increased swim speed, or changes in depth, or that there may be no observable response. Use of sonar and other transducers would typically be transient and temporary and there is no evidence to suggest that any behavioral response would persist after a sound exposure. It is assumed that a stress response could accompany any behavioral response.

Implementation of mitigation may further reduce the already low risk of auditory impacts on sea turtles. Depending on the sonar source, mitigation includes powering down the sonar or ceasing active sonar transmission if a sea turtle is observed in the mitigation zone, as discussed in Section 3.5.2.1.2.1 (Methods for Analyzing Impacts from Sonar and Other Transducers – Accounting for Mitigation).

Although masking of biologically relevant sounds by the limited number of sonars and other transducers operated in sea turtle hearing range is possible, this may only occur in certain circumstances. Sea turtles most likely use sound to detect nearby broadband, continuous environmental sounds, such as the sounds of waves crashing on the beach. The use characteristics of low-frequency active sonars, including limited band width, beam directionality, limited beam width, relatively low source levels, low duty cycle, and limited duration of use, would both greatly limit the potential for a sea turtle to detect these sources and limit the potential for masking of broadband, continuous environmental sounds. In addition, broadband sources within sea turtle hearing range, such as countermeasures used during anti-submarine warfare, would typically be used in off-shore areas, not in near-shore areas where detection of beaches or concentrated vessel traffic is relevant.

Considering the above factors and the mitigation measures that would be implemented as described in Chapter 5 (Mitigation), long-term consequences to sea turtle individuals or populations would not be expected.

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 1 may affect ESA-listed green, hawksbill, leatherback, loggerhead, and olive ridley sea turtles.*

#### **3.5.2.1.2.4 Alternative 2**

Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. Use of sonar and other transducers would typically be transient and temporary. General categories and characteristics of sonar systems and the number of hours these sonars would be operated during training under Alternative 2 are described in Section 3.0.1.2.4.1 (Sonar and Other Transducers). Activities using sonars and other transducers would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). Low-frequency sources are operated more frequently during testing activities than during training activities. Although the general impacts from sonar and other transducers during testing would be similar in severity to those described during training, there may be slightly more impacts during testing activities as sea turtles can detect low frequency sources.

Under Alternative 2, the same type and tempo of training and testing activities could occur as Alternative 1, but would include five Joint multi-strike group exercises (i.e., Valiant Shield) over five years as compared to three under Alternative 1. Additionally, Alternative 2 contemplates three (vice two) small joint coordinated anti-submarine warfare exercises (Multi-Sail/Guam Exercises) per year with a 50 percent increase in associated unit-level events (e.g., missile exercise [surface-to-air]). This would result in an increase of sonar use compared to Alternative 1. There would also be an increase in the use

of active sonar during certain testing events. Alternative 2 reflects the maximum number of training and testing activities that could occur within a given year, and assumes that the maximum number of Fleet exercises would occur every year.

The quantitative analysis predicts that no sea turtles of any species are likely to be exposed to the high received levels of sound from sonars or other transducers that could cause TTS or PTS during a maximum year of training activities under Alternative 2. Although there would be an increase in sonar use compared to Alternative 1, potential for and type of impacts on sea turtles would be the similar. This is because sea turtles are capable of detecting only a limited number of sonars due to their limited hearing range. Olive ridley sea turtle presence in the Study Area is limited and density data does not exist due to low occurrence in this region. Therefore, exposures were only modeled for green, hawksbill, leatherback, and loggerhead sea turtles in the Study Area and transit corridor. The NMFS's 2015 Biological Opinion (National Oceanic and Atmospheric Administration, 2015) on training and testing activities analyzed in the 2015 MITT Final EIS/OEIS considered sonars and other transducers to result in take incidental to military activities for green and hawksbill sea turtles.

Considering the above factors and the mitigation measures that would be implemented as described in Chapter 5 (Mitigation), long-term consequences to sea turtle individuals or populations would not be expected.

*Pursuant to the ESA, the use of sonar and other transducers during training and testing activities as described under Alternative 2 may affect ESA-listed green, hawksbill, leatherback, loggerhead, and olive ridley sea turtles.*

#### **3.5.2.1.2.5 No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Sonar and other transducers as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for acoustics stressors on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

#### **3.5.2.1.3 Impacts from Vessel Noise**

Sea turtles may be exposed to noise from vessel movement. A detailed description of the acoustic characteristics and typical sound levels of vessel noise is in Section 3.0.4.1.2 (Vessel Noise). Vessel movements involve transits to and from ports to various locations within the Study Area, including commercial ship traffic and recreational vessels, in addition to U.S. Navy vessels. Many ongoing and proposed training and testing activities within the Study Area involve maneuvers by various types of surface ships, boats, and submarines (collectively referred to as vessels).

Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Tables 2.5-1 and 2.5-2 for proposed training and testing activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors (e.g., vessel noise) as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for vessel noise impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

*Pursuant to the ESA, vessel noise during training and testing activities, as described under Alternative 1 and Alternative 2, may affect ESA-listed green, hawksbill, leatherback, loggerhead and olive ridley sea turtles.*

#### **3.5.2.1.4 Impacts from Aircraft Noise**

Sea turtles may be exposed to aircraft-generated noise throughout the Study Area. Fixed- and rotary-wing aircraft are used during a variety of training and testing activities throughout the Study Area. Tilt-rotor impacts would be similar to fixed-wing or helicopter impacts, depending on the aircraft's mode. Most of these sounds would be concentrated around airbases and fixed ranges within the range complex. Aircraft produce extensive airborne noise from either turbofan or turbojet engines. An infrequent type of aircraft noise is the sonic boom, produced when the aircraft exceeds the speed of sound. Rotary-wing aircraft (helicopters) produce low-frequency sound and vibration (Pepper et al., 2003).

A detailed description of aircraft noise as a stressor is in Section 3.0.4.1.3 (Aircraft Noise). Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Tables 2.5-1 and 2.5-2 for proposed training and testing activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS.

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors (e.g., aircraft noise) as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for aircraft noise impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

*Pursuant to the ESA, aircraft noise during training and testing activities as described under Alternative 1 and Alternative 2 may affect ESA-listed green, hawksbill, leatherback, loggerhead and olive ridley sea turtles.*

### 3.5.2.1.5 Impacts from Weapon Noise

Sea turtles may be exposed to sounds caused by the firing of weapons, objects in flight, and impact of non-explosive munitions on the water's surface, which are described in Section 3.0.4.1.4 (Weapon Noise). In general, these are impulsive sounds generated in close vicinity to or at the water surface, with the exception of items that are launched underwater. The firing of a weapon may have several components of associated noise. Firing of guns could include sound generated in air by firing a gun (muzzle blast) and a crack sound due to a low-amplitude shock wave generated by a supersonic projectile flying through the air. Most in-air sound would be reflected at the air-water interface.

Underwater sounds would be strongest just below the surface and directly under the firing point. Any sound that enters the water only does so within a narrow cone below the firing point or path of the projectile. Vibration from the blast propagating through a ship's hull, the sound generated by the impact of an object with the water surface, and the sound generated by launching an object underwater are other sources of impulsive sound in the water. Sound due to missile and target launches is typically at a maximum at initiation of the booster rocket and rapidly fades as the missile or target travels downrange.

Activities may vary slightly from those previously analyzed in the 2015 MITT Final EIS/OEIS, but the overall determinations presented remain valid. Increases and decreases shown in Tables 2.5-1 and 2.5-2 for proposed training and testing activities under Alternative 1 and 2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. Implementation of mitigation may further reduce the already low risk of auditory impacts on sea turtles from weapon noise during large-caliber gunnery events, as discussed in Section 5.3.2.2 (Weapons Firing Noise).

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors (e.g., weapon noise) as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for weapon noise impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

*Pursuant to the ESA, weapon noise during training and testing activities as described under Alternative 1 and Alternative 2 may affect ESA-listed green, hawksbill, leatherback, loggerhead and olive ridley sea turtles.*

### 3.5.2.2 Explosive Stressors

Explosions in the water or near the water surface can introduce loud, impulsive, broadband sounds into the marine environment. Unlike other acoustic stressors, explosives release energy at a high rate producing a shock wave that can be injurious and even deadly. Therefore, explosive impacts on sea turtles are discussed separately from other acoustic stressors, even though the analysis of explosive impacts will rely on data for sea turtle impacts due to impulsive sound exposure where appropriate.

Explosives are usually described by their net explosive weight, which accounts for the weight and type of explosive material. Additional explanation of the acoustic and explosive terms and sound energy concepts used in this section is found in Appendix H (Acoustic and Explosive Concepts).

This section begins with a summary of relevant data regarding explosive impacts on sea turtles in Section 3.5.2.2.1 (Background). The ways in which an explosive exposure could result in immediate effects or lead to long-term consequences for an animal are explained in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), and this section follows that framework. Studies of the effects of sound and explosives on sea turtles are limited; therefore, where necessary, knowledge of impacts on other species from explosives is used to assess impacts on sea turtles.

Due to new acoustic impact criteria, sea turtle densities, and acoustics effects model, the analysis provided in Section 3.5.2.2.2 (Impacts from Explosives) of this SEIS/OEIS will supplant the 2015 MITT Final EIS/OEIS for sea turtles, and may result in changes to estimated impacts for some species since the 2015 MITT Final EIS/OEIS.

#### **3.5.2.2.1 Background**

The sections below include a survey and synthesis of best available science published in peer-reviewed journals, technical reports, and other scientific sources pertinent to impacts on sea turtles potentially resulting from Navy training and testing activities. Sea turtles could be exposed to a range of impacts depending on the explosive source and context of the exposure. In addition to acoustic impacts including temporary or permanent hearing loss, auditory masking, physiological stress, or changes in behavior; potential impacts from an explosive exposure can include non-lethal injury and mortality.

##### **3.5.2.2.1.1 Injury**

Because direct studies of explosive impacts on sea turtles have not been conducted, the below discussion of injurious effects is based on studies of other animals, generally mammals. The generalizations that can be made about in-water explosive injuries to other species should be applicable to reptiles, with consideration of the unique anatomy of sea turtles. For example, it is unknown if the sea turtle shell may afford it some protection from internal injury.

If an animal is exposed to an explosive blast underwater, the likelihood of injury depends on the charge size, the geometry of the exposure (distance to the charge, depth of the animal and the charge), and the size of the animal. In general, an animal would be less susceptible to injury near the water surface because the pressure wave reflected from the water surface would interfere with the direct path pressure wave, reducing positive pressure exposure. However, rapid under-pressure phase caused by the negative surface-reflected pressure wave above an underwater detonation may create a zone of cavitation that may contribute to potential injury. In general, blast injury susceptibility would increase with depth, until normal lung collapse (due to increasing hydrostatic pressure) and increasing ambient pressures again reduce susceptibility.

See Appendix H (Acoustic and Explosive Concepts) for an overview of explosive propagation and an explanation of explosive effects on gas cavities.

Primary blast injury is injury that results from the compression of a body exposed to a blast wave. This is usually observed as barotrauma of gas-containing structures (e.g., lung and gut) and structural damage to the auditory system (Greaves et al., 1943; Office of the Surgeon General, 1991; Richmond et al., 1973). The lungs are typically the first site to show any damage, while the solid organs (e.g., liver, spleen,

and kidney) are more resistant to blast injury (Clark & Ward, 1943). Recoverable injuries would include slight lung injury, such as capillary interstitial bleeding, and contusions to the gastrointestinal tract. More severe injuries would significantly reduce fitness and likely cause death in the wild. Rupture of the lung may also introduce air into the vascular system, producing air emboli that can cause a stroke or heart attack by restricting oxygen delivery to critical organs. In this discussion, primary blast injury to auditory tissues is considered gross structural tissue injury distinct from noise-induced hearing loss, which is considered below in Section 3.5.2.2.1.2 (Hearing Loss).

Data on observed injuries to sea turtles from explosives is generally limited to animals found following explosive removal of offshore structures (Viada et al., 2008), which can attract sea turtles for feeding opportunities or shelter. Klima et al. (1988) observed a turtle mortality subsequent to an oil platform removal blast, although sufficient information was not available to determine the animal's exposure. Klima et al. (1988) also placed small sea turtles (less than 7 kilograms) at varying distances from piling detonations. Some of the turtles were immediately knocked unconscious or exhibited vasodilation over the following weeks, but others at the same exposure distance exhibited no effects.

Incidental injuries to sea turtles due to a military explosions have been documented in a few instances. In one incident, a single 1,200-pound (lb.) trinitrotoluene (TNT) underwater charge was detonated off Panama City, FL in 1981. The charge was detonated at a mid-water depth of 120 feet (ft.). Although details are limited, the following were recorded: at a distance of 500–700 ft., a 400 lb. sea turtle was killed; at 1,200 ft., a 200–300 lb. sea turtle experienced “minor” injury; and at 2,000 ft. a 200–300 lb. sea turtle was not injured (O’Keeffe & Young, 1984). In another incident, two “immature” green sea turtles (size unspecified) were found dead about 100-150 ft. away from detonation of 20 lb. of C-4 in a shallow water environment.

Results from limited experimental data suggest two explosive metrics are predictive of explosive injury: peak pressure and impulse.

#### **Impulse as a Predictor of Explosive Injury**

Without measurements of the explosive exposures in the above incidents, it is difficult to draw conclusions about what amount of explosive exposure would be injurious to sea turtles. Studies of observed in-water explosive injuries showed that terrestrial mammals were more susceptible than comparably sized fish with swim bladders (Yelverton & Richmond, 1981), and that fish with swim bladders may have increased susceptibility to swim bladder oscillation injury depending on exposure geometry (Goertner, 1978; Wiley et al., 1981). Therefore, controlled tests with a variety of terrestrial mammals (mice, rats, dogs, pigs, sheep and other species) are the best available data sources on actual injury to similar-sized animals due to underwater exposure to explosions.

In the early 1970s, the Lovelace Foundation for Medical Education and Research conducted a series of tests in an artificial pond to determine the effects of underwater explosions on mammals, with the goal of determining safe ranges for human divers. The resulting data were summarized in two reports (Richmond et al., 1973; Yelverton et al., 1973). Specific physiological observations for each test animal are documented in Richmond et al. (1973). Gas-containing internal organs, such as lungs and intestines, were the principle damage sites in submerged terrestrial mammals, consistent with earlier studies of mammal exposures to underwater explosions (Clark & Ward, 1943; Greaves et al., 1943).

In the Lovelace studies, acoustic impulse was found to be the metric most related to degree of injury, and size of an animal's gas-containing cavities was thought to play a role in blast injury susceptibility. The proportion of lung volume to overall body size is similar between sea turtles and terrestrial

mammals, so the magnitude of lung damage in the tests may approximate the magnitude of injury to sea turtles when scaled for body size. Measurements of some shallower diving sea turtles (Hochscheid et al., 2007) show lung-to-body size ratios that are larger than terrestrial animals, whereas the lung-to-body mass ratio of the deeper diving leatherback sea turtle is smaller (Lutcavage et al., 1992). The use of test data with smaller lung-to-body ratios to set injury thresholds may result in a more conservative estimate of potential for damaging effects (i.e., lower thresholds) for animals with larger lung-to-body ratios.

For these shallow exposures of small terrestrial mammals (masses ranging from 3.4 to 50 kilograms) to underwater detonations, Richmond et al. (1973) reported that no blast injuries were observed when exposures were less than 6 lb. per square inch (in.) per millisecond (psi-ms) (40 pascal-seconds [Pa-s]), no instances of slight lung hemorrhage occurred below 20 psi-ms (140 Pa-s), and instances of no lung damage were observed in some exposures at higher levels up to 40 psi-ms (280 Pa-s). An impulse of 34 psi-ms (230 Pa-s) resulted in about 50 percent incidence of slight lung hemorrhage. About half of the animals had gastrointestinal tract contusions (with slight ulceration, i.e., some perforation of the mucosal layer) at exposures of 25–27 psi-ms (170–190 Pa-s). Lung injuries were found to be slightly more prevalent than gastrointestinal tract injuries for the same exposure.

The Lovelace subject animals were exposed near the water surface; therefore, depth effects were not discernible in this data set. In addition, this data set included only small terrestrial animals, whereas adult sea turtles may be substantially larger and have respiratory structures adapted for the high pressures experienced at depth. Goertner (1982) examined how lung cavity size would affect susceptibility to blast injury by considering both size and depth in a bubble oscillation model of the lung, which is assumed to be applicable to sea turtles as well for this analysis. Animal depth relates to injury susceptibility in two ways: injury is related to the relative increase in explosive pressure over hydrostatic pressure, and lung collapse with depth reduces the potential for air cavity oscillatory damage. The time period over which an impulse must be delivered to cause damage is assumed to be related to the natural oscillation period of an animal's lung, which depends on lung size. Based on a study of green sea turtles, Berkson (1967) predicted sea turtle lung collapse would be complete around 80–160 m depth.

#### **Peak Pressure as a Predictor of Explosive Trauma**

High instantaneous peak pressures can cause damaging tissue distortion. Goertner (1982) suggested a peak overpressure gastrointestinal tract injury criterion because the size of gas bubbles in the gastrointestinal tract are variable, and their oscillation period could be short relative to primary blast wave exposure duration. The potential for gastrointestinal tract injury, therefore, may not be adequately modeled by the single oscillation bubble methodology used to estimate lung injury due to impulse. Like impulse, however, high instantaneous pressures may damage many parts of the body, but damage to the gastrointestinal tract is used as an indicator of any peak pressure-induced injury due to its vulnerability.

Older military reports documenting exposure of human divers to blast exposure generally describe peak pressure exposures around 100 lb. psi (237 dB re 1  $\mu$ Pa peak) to feel like a slight pressure or stinging sensation on skin, with no enduring effects (Christian & Gaspin, 1974). Around 200 psi, the shock wave felt like a blow to the head and chest. Data from the Lovelace Foundation experiments show instances of gastrointestinal tract contusions after exposures up to 1,147 psi peak pressure, while exposures of up to 588 psi peak pressure resulted in many instances of no observed gastrointestinal tract effects. The lowest exposure for which slight contusions to the gastrointestinal tract were reported was 237 dB re 1  $\mu$ Pa peak. As a vulnerable gas-containing organ, the gastrointestinal tract is vulnerable to both high

peak pressure and high impulse, which may vary to differing extents due to blast exposure conditions (i.e., animal depth, distance from the charge). This likely explains the range of effects seen at similar peak pressure exposure levels and shows the utility of considering both peak pressure and impulse when analyzing the potential for injury due to explosives.

The *ANSI Sound Exposure Guidelines* (Popper et al., 2014) recommended peak pressure guidelines for sea turtle injury from explosives. Lacking any direct data for sea turtles, these recommendations were based on fish data. Of the fish data available, the working group conservatively chose the study with the lowest peak pressures associated with fish mortality to set guidelines (Hubbs & Rechnitzer, 1952), and did not consider the Lovelace studies discussed above.

### **Fragmentation**

Fragments produced by exploding munitions at or near the surface may present a high speed strike hazard for an animal at or near the surface. In water, however, fragmentation velocities decrease rapidly due to drag (Swisdak & Montanaro, 1992). Because blast waves propagate efficiently through water, the range to injury from the blast wave would likely extend beyond the range of fragmentation risk.

#### **3.5.2.2.1.2 Hearing Loss**

An underwater explosion produces broadband, impulsive sound that can cause noise-induced hearing loss, typically quantified as threshold shift, which persists after cessation of the noise exposure. This noise-induced hearing loss may manifest as TTS or PTS. Because studies on inducing threshold shift in reptiles are very limited (e.g., alligator lizards: Dew et al., 1993; Henry & Mulroy, 1995) and have not been conducted on any of the sea turtles present in the Study Area, auditory threshold shift in reptiles is considered to be consistent with general knowledge about noise-induced hearing loss described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

Little is known about how sea turtles use sound in their environment. The *ANSI Sound Exposure Guidelines* (Popper et al., 2014) do not suggest numeric sound exposure thresholds for auditory effects on sea turtles due to lack of data. Rather, the guidelines qualitatively advise that sea turtles are less likely to incur TTS or PTS with increasing distance from an explosive. The guidelines also suggest that data from fishes may be more relevant than data from marine mammals when estimating auditory impacts on sea turtles, because, in general, fish hearing range is more similar to the limited hearing range of sea turtles. As shown in Section 3.5.1.4 (Hearing and Vocalization), sea turtle hearing is most sensitive around 100–400 Hz in-water, is limited over 1 kHz, and is much less sensitive than that of any marine mammal.

#### **3.5.2.2.1.3 Physiological Stress**

A stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor. If the magnitude and duration of the stress response is too great or too long, it can have negative consequences to the animal (e.g. decreased immune function, decreased reproduction). Physiological stress is typically analyzed by measuring stress hormones, other biochemical markers, or vital signs. Physiological stress has been measured for sea turtles during nesting (Flower et al., 2015; Valverde et al., 1999) and capture and handling (Flower et al., 2015; Gregory & Schmid, 2001), but the stress caused by acoustic exposure has not been studied for reptiles. Therefore, the stress response in sea turtles in the Study Area due to acoustic exposures is considered to be consistent with general knowledge about physiological stress responses described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

Marine animals naturally experience stressors within their environment and as part of their life histories. Changing weather and ocean conditions, exposure to diseases and naturally occurring toxins, lack of prey availability, social interactions with members of the same species, nesting, and interactions with predators all contribute to stress. Anthropogenic sound-producing activities have the potential to provide additional stressors beyond those that naturally occur.

Due to the limited information about acoustically induced stress responses, the Navy conservatively assumes in its effect analysis that any physiological response (e.g., hearing loss or injury) or significant behavioral response is also associated with a stress response.

#### **3.5.2.2.1.4 Masking**

As described in Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities), auditory masking occurs when one sound, distinguished as the “noise,” interferes with the detection or recognition of another sound or limits the distance over which other biologically relevant sounds can be detected. Masking only occurs when the sound source is operating; therefore, direct masking effects stop immediately upon cessation of the sound-producing activity. Any unwanted sound above ambient noise and within an animal’s hearing range may potentially cause masking which can interfere with an animal’s ability to detect, understand, or recognize biologically relevant sounds of interest.

Masking occurs in all vertebrate groups and can effectively limit the distance over which an animal can communicate and detect biologically relevant sounds. The effect of masking has not been studied for marine reptiles. The potential for masking in reptiles would be limited to certain sound exposures due to their limited hearing range to broadband low-frequency sounds and lower sensitivity to noise in the marine environment. Only continuous human-generated sounds that have a significant low-frequency component, are not of brief duration, and are of sufficient received level could create a meaningful masking situation. While explosives produce intense, broadband sounds with significant low-frequency content, these sounds are very brief with limited potential to mask relevant sounds.

There is evidence that reptiles may rely primarily on senses other than hearing for interacting with their environment, such as vision (Narazaki et al., 2013) and magnetic orientation (Avens, 2003; Putman et al., 2015). Any effect of masking may be mediated by reliance on other environmental inputs.

#### **3.5.2.2.1.5 Behavioral Reactions**

There are no observations of behavioral reactions by sea turtles to exposure to explosive sounds. Impulsive signals, particularly at close range, have a rapid rise time and higher instantaneous peak pressure than other signal types, making them more likely to cause startle responses or avoidance responses. Although explosive sources are more energetic than air guns, the few studies of sea turtle responses to air guns, which are not used during MITT training or testing activities, may show the types of behavioral responses that sea turtles may have towards explosives. General research findings regarding behavioral reactions from sea turtles due to exposure to impulsive sounds, such as those associated with explosions, are discussed in detail in Behavioral Reactions to Impulsive Sound Sources under Section 3.5.2.1 (Acoustic Stressors).

#### **3.5.2.2.1.6 Long-Term Consequences**

For sea turtles present in the Study Area, long-term consequences to individuals and populations due to acoustic exposures have not been studied. Therefore, long term consequences to sea turtles due to

explosive exposures are considered following Section 3.0.4.7 (Conceptual Framework for Assessing Effects from Acoustic and Explosive Activities).

Long-term consequences to a population are determined by examining changes in the population growth rate. Physical effects that could lead to a reduction in the population growth rate include mortality or injury, which could remove animals from the reproductive pool, and permanent hearing impairment, which could impact navigation. The long-term consequences due to individual behavioral reactions and short-term instances of physiological stress are especially difficult to predict because individual experience over time can create complex contingencies. It is more likely that any long-term consequences to an individual would be a result of costs accumulated over a season, year, or life stage due to multiple behavioral or stress responses resulting from exposures to multiple stressors over significant periods of time. Conversely, some sea turtles may habituate to or become tolerant of repeated acoustic exposures over time, learning to ignore a stimulus that in the past did not accompany any overt threat. For example, loggerhead sea turtles exposed to air guns with a source SPL of 179 dB re 1  $\mu$ Pa initially exhibited avoidance reactions. However, they may have habituated to the sound source after multiple exposures since a habituation behavior was retained when exposures were separated by several days (Moein Bartol et al., 1995). More research is needed to better understand the long-term consequences of human-made noise on sea turtles, although intermittent exposures are assumed to be less likely to have lasting consequences.

#### **3.5.2.2.2 Impacts from Explosives**

Sea turtles could be exposed to energy, sound, and fragments from explosions in the water and near the water surface associated with the proposed activities. Energy and sound from an explosion is capable of causing mortality, injury, hearing loss, a behavioral response, masking, or physiological stress, depending on the level and duration of exposure. The death of an animal would eliminate future reproductive potential, which is considered in the analysis of potential long-term consequences to the population. Exposures that result in non-auditory injuries may limit an animal's ability interpret the surrounding environment. Impairment of these abilities can decrease an individual's chance of survival or affect its ability to reproduce. Temporary threshold shift can also impair an animal's abilities, although the individual may recover quickly with little significant effect. Overall, the locations, types, and severity of predicted impacts for the use of explosives during training and testing activities would be similar to what is currently conducted, with the addition of several new testing activities as described in Table 2.5-1. Although individual activities may vary in the number of events or ordnances some from those previously analyzed, the overall determinations presented in the 2015 MITT Final EIS/OEIS remain valid, and has been developed further under the current SEIS/EIS.

The quantitative analysis has been improved upon and updated since the 2015 MITT Final EIS/OEIS; therefore, the new analysis is fully presented and described in further detail in the technical report *Quantitative Analysis for Estimating Acoustic and Explosive Impacts to Marine Mammals and Sea Turtles* (U.S. Department of the Navy, 2018a).

#### **3.5.2.2.2.1 Methods for Analyzing Impacts from Explosives**

Potential impacts considered are mortality, injury, hearing loss due to threshold shift (permanent or temporary), masking of other biologically relevant sounds, physiological stress, and changes in behavior.

The Navy's quantitative analysis to determine impacts on sea turtles and marine mammals uses the Navy Acoustic Effects Model to produce initial estimates of the number of times these animals may experience these effects; these estimates are further refined by considering animal avoidance of sound-

producing activities and implementation of mitigation. The steps of this quantitative analysis are described in Section 3.0.1.2 (Navy's Quantitative Analysis to Determine Impacts on Sea Turtles and Marine Mammals), which takes into account

- criteria and thresholds used to predict impacts from explosives (see below),
- the density and spatial distribution of sea turtles, and
- the influence of environmental parameters (e.g., temperature, depth, salinity) on sound propagation and explosive energy when estimating the received sound level and pressure on the animals.

A further detailed explanation of this analysis is provided in the technical report titled *Quantitative Analysis for Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2018a).

### **Criteria and Thresholds used to Predict Impacts on Sea Turtles from Explosives**

#### **Mortality and Injury from Explosives**

As discussed above in Section 3.5.2.2.1.1 (Injury), two metrics have been identified as predictive of injury: impulse and peak pressure. Peak pressure contributes to the "crack" or "stinging" sensation of a blast wave, compared to the "thump" associated with received impulse. Older military reports documenting exposure of human divers to blast exposure generally describe peak pressure exposures around 100 psi (237 dB re 1  $\mu$ Pa SPL peak) to feel like slight pressure or stinging sensation on skin, with no enduring effects (Christian & Gaspin, 1974).

Two sets of thresholds are provided for use in non-auditory injury assessment. The exposure thresholds are used to estimate the number of animals that may be affected during Navy training and testing activities (Table 3.5-2). The thresholds for the farthest range to effect are based on the received level at which 1 percent risk is predicted and are useful for assessing potential effects to sea turtles and marine mammals, and the range at which mitigation could be effective. Increasing animal mass and increasing animal depth both increase the impulse thresholds (i.e., decrease susceptibility), whereas smaller mass and decreased animal depth reduce the impulse thresholds (i.e., increase susceptibility). For impact assessment, sea turtle populations are assumed to be 5 percent adult and 95 percent sub-adult. This adult to sub-adult population ratio is estimated from what is known about the population age structure for sea turtles. Sea turtles typically lay multiple clutches of 100 or more eggs with little parental investment and generally have low survival in early life. However, sea turtles that are able to survive past early life generally have high age-specific survival in later life.

The derivation of these injury criteria and the species mass estimates are provided in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).

**Table 3.5-2: Criteria to Quantitatively Assess Non-Auditory Injury due to Underwater Explosions**

| <i>Impact Category</i> | <i>Exposure Threshold</i>                                | <i>Threshold for Farthest Range to Effect</i>            |
|------------------------|--|--|
| Mortality <sup>1</sup> | $144M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s  | $103M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s  |
| Injury <sup>1</sup>    | $65.8M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s | $47.5M^{1/3} \left(1 + \frac{D}{10.1}\right)^{1/6}$ Pa-s |
|                        | 243 dB re 1 μPa SPL peak                                 | 237 dB re 1 μPa SPL peak                                 |

<sup>1</sup> Impulse delivered over 20 percent of the estimated lung resonance period. See U.S. Department of the Navy (2017a).

<sup>2</sup> Threshold for 1 percent risk used to assess mitigation effectiveness.


Note: dB re 1 μPa = decibels referenced to 1 micropascal, SPL = sound pressure level, M = animal mass (kg), D = animal depth (m), and Pa-s = Pascal-second

When explosive munitions (e.g., a bomb or missile) detonates, fragments of the weapon are thrown at high-velocity from the detonation point, which can injure or kill sea turtles if they are struck. Risk of fragment injury reduces exponentially with distance as the fragment density is reduced. Fragments underwater tend to be larger than fragments produced by in-air explosions (Swisdak & Montanaro, 1992). Underwater, the friction of the water would quickly slow these fragments to a point where they no longer pose a threat. On the other hand, the blast wave from an explosive detonation moves efficiently through the seawater. Because the ranges to mortality and injury due to exposure to the blast wave are likely to far exceed the zone where fragments could injure or kill an animal, the above thresholds are assumed to encompass risk due to fragmentation.

**Auditory Weighting Functions**

Animals are not equally sensitive to noise at all frequencies. To capture the frequency-dependent nature of the effects of noise, auditory weighting functions are used. Auditory weighting functions are mathematical functions that adjust received sound levels to emphasize ranges of best hearing and de-emphasize ranges with less or no auditory sensitivity. The adjusted received sound level is referred to as a weighted received sound level.

The auditory weighting function for sea turtles is shown in Figure 3.5-6. The derivation of this weighting function is described in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a). The frequencies around the top portion of the function, where the amplitude is closest to zero, are emphasized, while the frequencies below and above this range (where amplitude declines) are de-emphasized, when summing acoustic energy received by a sea turtle.



Source: U.S. Department of the Navy (2017a)


Notes: dB = decibels, kHz = kilohertz, TU = sea turtle hearing group

**Figure 3.5-6: Auditory Weighting Functions for Sea Turtles**

#### Hearing Loss from Explosives

No studies of hearing loss have been conducted on sea turtles. Therefore, sea turtle susceptibility to hearing loss due to an acoustic exposure is evaluated using knowledge about sea turtle hearing abilities in combination with non-impulsive auditory effect data from other species (marine mammals and fish). This yields sea turtle exposure functions, shown in Figure 3.5-7, which are mathematical functions that relate the SELs for onset of TTS or PTS to the frequency of the sonar sound exposure. The derivation of the sea turtle exposure functions are provided in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).

For impulsive sounds, hearing loss in other species has also been observed to be related to the unweighted peak pressure of a received sound. Because this data does not exist for sea turtles, unweighted peak pressure thresholds for TTS and PTS were developed by applying relationships observed between impulsive peak pressure TTS thresholds and auditory sensitivity in marine mammals to sea turtles. This results in dual-metric hearing loss criteria for sea turtles for impulsive sound exposure: the SEL-based exposure functions in Figure 3.5-7 and the peak pressure thresholds in Table 3.5-3. The derivation of the sea turtle impulsive peak pressure TTS and PTS thresholds are provided in the technical report *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)* (U.S. Department of the Navy, 2017a).



Notes: kHz = kilohertz, SEL = Sound Exposure Level, dB re 1  $\mu\text{Pa}^2\text{s}$  = decibels referenced to 1 micropascal squared second. The solid black curve is the exposure function for TTS onset and the dashed black curve is the exposure function for PTS onset. Small dashed lines and asterisks indicate the SEL thresholds and most sensitive frequency for TTS and PTS.

**Figure 3.5-7: TTS and PTS Exposure Functions for Impulsive Sounds**

**Table 3.5-3: TTS and PTS Peak Pressure Thresholds Derived for Sea Turtles Exposed to Impulsive Sounds**

| <i>Auditory Effect</i> | <i>Unweighted Peak Pressure Threshold</i> |
|------------------------|---|
| TTS                    | 226 dB re 1 $\mu\text{Pa}$ SPL peak       |
| PTS                    | 232 dB re 1 $\mu\text{Pa}$ SPL peak       |

Notes: dB re 1  $\mu\text{Pa}$  = decibels referenced to 1 micropascal, PTS = permanent threshold shift, SPL = sound pressure level, TTS = temporary threshold shift

**Accounting for Mitigation**

The Navy will implement mitigation measures to avoid or reduce potential impacts from explosives on sea turtles, as described in Section 5.3.3 (Explosive Stressors). The benefits of mitigation are conservatively factored into the analysis for Alternative 1 and Alternative 2 of the Proposed Action for training and testing. The Navy’s mitigation measures are identical for both action alternatives.

Procedural mitigation measures include delaying or ceasing applicable detonations when a sea turtle is observed in a mitigation zone. The mitigation zones for explosives extend beyond the respective average ranges to mortality. The mitigation zones encompass the estimated ranges to mortality for a given explosive. Therefore, the impact analysis quantifies the potential for mitigation to reduce the risk of

mortality due to exposure to explosives. Two factors are considered when quantifying the effectiveness of mitigation: (1) the extent to which the type of mitigation proposed for a sound-producing activity (e.g., active sonar) allows for observation of the mitigation zone prior to and during the activity; and (2) the sightability of each species that may be present in the mitigation zone, which is determined by species-specific characteristics and the viewing platform. A detailed explanation of the analysis is provided in the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (U.S. Department of the Navy, 2017a).

In the quantitative analysis, consideration of mitigation measures means that, for activities where mitigation is feasible, model-estimated mortality is considered mitigated to the level of injury. The impact analysis does not analyze the potential for mitigation to reduce TTS or behavioral effects, even though mitigation could also reduce the likelihood of these effects. In practice, mitigation also protects all unobserved (below the surface) animals in the vicinity, including other species, in addition to the observed animal. However, the analysis assumes that only animals sighted at the water surface would be protected by the applied mitigation. The analysis, therefore, does not capture the protection afforded to all marine species that may be near or within the mitigation zone.

#### **3.5.2.2.2.2 Impact Ranges for Explosives**

The following section provides the range (distance) over which specific physiological or behavioral effects are expected to occur based on the explosive criteria and the explosive propagation calculations from the Navy Acoustic Effects Model (Section 3.5.2.2.2.1, Methods for Analyzing Impacts from Explosives). The range to effects are shown for a range of explosive bins, from E1 (up to 0.25 lb. net explosive weight) to E12 (up to 1,000 lb. net explosive weight). Ranges are determined by modeling the distance that noise from an explosion would need to propagate to reach exposure level thresholds specific to a hearing group that would cause TTS, PTS, non-auditory injury, and mortality. Range to effects is important information in not only predicting impacts from explosives, but also in verifying the accuracy of model results against real-world situations and assessing the level of impact that will likely be mitigated within applicable mitigation zones.

Table 3.5-4 shows the minimum, average, and maximum ranges due to varying propagation conditions to non-auditory injury based on the larger of the range to slight lung injury or gastrointestinal tract injury for representative animal masses ranging from 10 to 1,000 kilograms and different explosive bins ranging from 0.25 to 1,000 lb. net explosive weight. Animals within these water volumes would be expected to receive minor injuries at the outer ranges, increasing to more substantial injuries, and finally mortality as an animal approaches the detonation point. Ranges to mortality, based on animal mass, are shown in Table 3.5-5.

The following tables (Table 3.5-6 and Table 3.5-7) show the minimum, average, and maximum ranges to onset of auditory and behavioral effects based on the thresholds described in Section 3.5.2.2.2.1 (Methods for Analyzing Impacts from Explosives). Ranges are provided for a representative source depth and cluster size (the number of rounds fired [or buoys dropped] within a very short duration) for each bin. For events with multiple explosions, sound from successive explosions can be expected to accumulate and increase the range to the onset of an impact based on SEL thresholds. Modeled ranges to TTS and PTS based on peak pressure for a single explosion generally exceed the modeled ranges based on SEL even when accumulated for multiple explosions. Peak pressure based ranges are estimated using the best available science; however, data on peak pressure at far distances from explosions are very limited. For additional information on how ranges to impacts from explosions were

estimated, see the technical report *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing Ranges* (U.S. Department of the Navy, 2017b).

**Table 3.5-4: Ranges to Non-Auditory Injury<sup>1</sup> (in meters) for Sea Turtles Exposed to Explosives as a Function of Animal Mass**

| Bin | Range to Non-Auditory Injury (meters) for Various Animal Mass Intervals (kg) <sup>1</sup> |                  |                  |
|-----|---|------------------|------------------|
|     | 10 kg   | 250 kg           | 1,000 kg         |
| E1  | 12<br>(11–13)   | 12<br>(11–13)    | 12<br>(11–13)    |
| E2  | 16<br>(15–16)   | 16<br>(15–16)    | 16<br>(15–16)    |
| E3  | 25<br>(25–25)   | 25<br>(25–25)    | 25<br>(25–25)    |
| E4  | 30<br>(30–35)   | 30<br>(30–35)    | 30<br>(30–35)    |
| E5  | 40<br>(40–65)   | 40<br>(40–50)    | 40<br>(40–50)    |
| E6  | 52<br>(50–60)   | 52<br>(50–55)    | 52<br>(50–55)    |
| E7  | 110<br>(110–110)  | 75<br>(75–75)    | 75<br>(75–75)    |
| E8  | 93<br>(90–150)  | 91<br>(90–95)    | 91<br>(90–95)    |
| E9  | 123<br>(120–270)  | 123<br>(120–140) | 123<br>(120–130) |
| E10 | 155<br>(150–420)  | 155<br>(150–240) | 155<br>(150–160) |
| E11 | 398<br>(380–420)  | 219<br>(170–260) | 172<br>(160–220) |
| E12 | 195<br>(190–650)  | 195<br>(190–380) | 195<br>(190–200) |

<sup>1</sup> Average distance (m) to non-auditory injury is depicted above the minimum and maximum distances which are in parentheses. The ranges depicted are the further of the ranges for gastrointestinal tract injury or slight lung injury for an explosive bin and animal mass interval combination.

**Table 3.5-5: Ranges to Mortality for Sea Turtles Exposed to Explosives as a Function of Animal Mass<sup>1</sup>**

| <i>Bin</i> | <i>Ranges to Mortality (meters) for Various Animal Mass Intervals (kg)<sup>1</sup></i> |                |                 |
|------------|--|----------------|-----------------|
|            | <i>10 kg</i>   | <i>250 kg</i>  | <i>1,000 kg</i> |
| E1         | 2<br>(2-3)   | 1<br>(0-1)     | 0<br>(0-0)      |
| E2         | 4<br>(3-4)   | 1<br>(1-2)     | 1<br>(1-1)      |
| E3         | 8<br>(6-9)   | 4<br>(3-6)     | 2<br>(2-2)      |
| E4         | 13<br>(11-15)  | 7<br>(5-9)     | 4<br>(4-5)      |
| E5         | 12<br>(11-30)  | 7<br>(5-18)    | 4<br>(4-7)      |
| E6         | 15<br>(14-25)  | 9<br>(7-17)    | 5<br>(5-9)      |
| E7         | 51<br>(50-55)  | 27<br>(23-30)  | 17<br>(17-17)   |
| E8         | 40<br>(24-65)  | 22<br>(12-40)  | 14<br>(9-21)    |
| E9         | 31<br>(30-35)  | 20<br>(16-24)  | 13<br>(12-13)   |
| E10        | 54<br>(40-170)   | 24<br>(20-25)  | 16<br>(15-17)   |
| E11        | 194<br>(180-210)   | 96<br>(70-130) | 53<br>(50-55)   |
| E12        | 83<br>(50-260)   | 31<br>(25-90)  | 20<br>(19-20)   |

<sup>1</sup> Average distance (m) to mortality is depicted above the minimum and maximum distances which are in parentheses.

**Table 3.5-6: Peak Pressure Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives**

| <i>Range to Effects for Explosives: Sea turtles<sup>1</sup></i> |                              |                     |                              |                              |
|---|------------------------------|---------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E1  | 0.1                          | 1                   | 35<br>(35-35)                | 68<br>(65-70)                |
|   |                              | 18                  | 35<br>(35-35)                | 68<br>(65-70)                |
| E2  | 0.1                          | 1                   | 48<br>(45-50)                | 87<br>(80-90)                |
|   |                              | 5                   | 48<br>(45-50)                | 87<br>(80-90)                |
| E3  | 0.1                          | 1                   | 81<br>(75-85)                | 145<br>(140-150)             |
|   |                              | 12                  | 81<br>(75-85)                | 145<br>(140-150)             |
|   | 18.25                        | 1                   | 80<br>(80-80)                | 150<br>(150-150)             |
|   |                              | 12                  | 80<br>(80-80)                | 150<br>(150-150)             |
| E4  | 10                           | 2                   | 100<br>(100-100)             | 192<br>(190-200)             |
|   | 60                           | 2                   | 101<br>(100-110)             | 194<br>(190-220)             |
| E5  | 0.1                          | 20                  | 125<br>(120-130)             | 235<br>(230-250)             |
|   | 30                           | 20                  | 138<br>(130-160)             | 257<br>(240-290)             |
| E6  | 0.1                          | 1                   | 163<br>(160-170)             | 292<br>(270-320)             |
|   | 30                           | 1                   | 160<br>(160-160)             | 300<br>(300-300)             |
| E7  | 28                           | 1                   | 240<br>(240-240)             | 442<br>(440-450)             |
| E8  | 0.1                          | 1                   | 273<br>(260-280)             | 451<br>(370-500)             |
|   | 45.75                        | 1                   | 281<br>(280-300)             | 527<br>(525-575)             |
| E9  | 0.1                          | 1                   | 355<br>(320-380)             | 566<br>(440-675)             |

**Table 3.5-6: Peak Pressure Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives (continued)**

| <i>Range to Effects for Explosives: Sea turtles<sup>1</sup></i> |                              |                     |                              |                              |
|---|------------------------------|---------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E10   | 0.1                          | 1                   | 432<br>(360–550)             | 690<br>(480–1,025)           |
| E11   | 45.75                        | 1                   | 540<br>(525–625)             | 977<br>(950–1,025)           |
|   | 91.4                         | 1                   | 558<br>(500–800)             | 1,053<br>(825–2,025)         |
| E12   | 0.1                          | 1                   | 509<br>(410–575)             | 784<br>(550–1,025)           |
|   |                              | 4                   | 509<br>(410–575)             | 784<br>(550–1,025)           |

<sup>1</sup>Average distance (m) to TTS and PTS are depicted above the minimum and maximum distances which are in parentheses. Values depict ranges to TTS and PTS based on the peak pressure metric.

**Table 3.5-7: SEL Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives**

| <i>Range to Effects for Explosives: Sea turtles<sup>1</sup></i> |                              |                     |                              |                              |
|---|------------------------------|---------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E1  | 0.1                          | 1                   | 0<br>(0–0)                   | 0<br>(0–0)                   |
|   |                              | 18                  | 0<br>(0–0)                   | 2<br>(2–2)                   |
| E2  | 0.1                          | 1                   | 0<br>(0–0)                   | 1<br>(1–1)                   |
|   |                              | 5                   | 0<br>(0–0)                   | 2<br>(2–2)                   |
| E3  | 0.1                          | 1                   | 0<br>(0–0)                   | 3<br>(2–3)                   |
|   |                              | 12                  | 2<br>(1–2)                   | 8<br>(8–18)                  |
|   | 18.25                        | 1                   | 3<br>(3–3)                   | 17<br>(16–17)                |
|   |                              | 12                  | 10<br>(10–10)                | 70<br>(70–70)                |

**Table 3.5-7: SEL Based Ranges to TTS and PTS for Sea Turtles Exposed to Explosives  
 (continued)**

| <i>Range to Effects for Explosives: Sea turtles<sup>1</sup></i> |                              |                     |                              |                              |
|---|------------------------------|---------------------|------------------------------|------------------------------|
| <i>Bin</i>  | <i>Source Depth (meters)</i> | <i>Cluster Size</i> | <i>Range to PTS (meters)</i> | <i>Range to TTS (meters)</i> |
| E4  | 10                           | 2                   | 7<br>(7-8)                   | 52<br>(50-55)                |
|   | 60                           | 2                   | 7<br>(7-7)                   | 35<br>(35-35)                |
| E5  | 0.1                          | 20                  | 5<br>(5-5)                   | 36<br>(25-270)               |
|   | 30                           | 20                  | 48<br>(40-65)                | 293<br>(240-400)             |
| E6  | 0.1                          | 1                   | 2<br>(2-2)                   | 10<br>(10-180)               |
|   | 30                           | 1                   | 14<br>(14-14)                | 95<br>(95-95)                |
| E7  | 28                           | 1                   | 30<br>(30-30)                | 200<br>(200-200)             |
| E8  | 0.1                          | 1                   | 5<br>(5-5)                   | 39<br>(25-290)               |
|   | 45.75                        | 1                   | 40<br>(40-40)                | 271<br>(270-280)             |
| E9  | 0.1                          | 1                   | 9<br>(9-9)                   | 87<br>(40-410)               |
| E10   | 0.1                          | 1                   | 13<br>(13-270)               | 164<br>(60-1,000)            |
| E11   | 45.75                        | 1                   | 170<br>(170-180)             | 832<br>(750-850)             |
|   | 91.4                         | 1                   | 150<br>(150-170)             | 794<br>(750-875)             |
| E12   | 0.1                          | 1                   | 31<br>(18-120)               | 200<br>(80-950)              |
|   |                              | 4                   | 59<br>(30-380)               | 377<br>(140-5,025)           |

<sup>1</sup>Average distance (m) to TTS and PTS are depicted above the minimum and maximum distances which are in parentheses. Values depict ranges to TTS and PTS based on the SEL metric.

### 3.5.2.2.2.3 Presentation of Estimated Impacts from the Quantitative Analysis

The results of the analysis of potential impacts to sea turtles from explosives as described in Section 3.5.2.2.2.1 (Methods for Analyzing Impacts from Explosives) are discussed below. Estimated numbers of potential impacts from the quantitative analysis for sea turtles are presented below. The most likely regions and activity categories from which the impacts could occur are displayed in the figures. There is a potential for impacts to occur anywhere within the Study Area where sound and energy from explosives and the species overlap, although only areas or categories where 0.5 percent of the impact, or greater, are estimated to occur are graphically represented on the species specific figures below. All (i.e., grand total) estimated impacts are included in the graphics, regardless of region or category.

The numbers of activities planned can vary slightly from year-to-year. Results are presented for a maximum explosive use year; however, during most years, explosive use would be less resulting in fewer potential impacts. The number of explosives used are described in Section 3.0.4.2 (Explosive Stressors).

Ranges to effect (see Table 3.5-4 through Table 3.5-7) were developed in the Navy Acoustic Effects Model based on the thresholds for TTS, PTS, injury, and mortality discussed above.

### 3.5.2.2.2.4 Alternative 1

Activities using explosives would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). General characteristics, quantities, and net explosive weights of in-water explosives used during training and testing activities under Alternative 1 are provided in Section 3.0.4.2 (Explosive Stressors). Quantities and locations of fragment-producing explosives during training and testing activities under Alternative 1 are shown in 3.0.4.4.4 (Military Expended Materials).


Under Alternative 1, there could be fluctuation in the number of explosions that could occur annually, although potential impacts would be similar from year to year. The number of impulsive sources in this SEIS/OEIS compared with the totals analyzed in the 2015 MITT Final EIS/OEIS are described in Tables 2-7 and 2-8.

The number of torpedo testing events (both explosive and non-explosive) planned under Alternative 1 testing can vary slightly from year-to-year however all other training and testing activities would remain consistent from year-to-year. Alternative 1 results are presented for a maximum explosive use year; however, during most years, explosive use would be less resulting in fewer potential impacts, as described in Section 3.0.4.2 (Explosive Stressors).

Detonations would typically occur in waters greater than 200 ft. in depth, and greater than 3 nautical miles from shore, with the exception of existing mine warfare areas, including Outer Apra Harbor, Piti, and Agat. Section 5.3.3 (Explosive Stressors) outlines the procedural mitigation measures for explosive stressors to reduce potential impacts on biological resources.

The quantitative analysis, using a maximum year of training and testing activities, estimates that no sea turtles would be killed, however, a small number of green sea turtles would be exposed to levels of explosive sound and energy in the outer Apra Harbor that could cause TTS or PTS (Figure 3.5-8). The quantitative analysis predicts that no hawksbill, leatherback, or loggerhead sea turtles are likely to be exposed to the levels of explosive sound and energy that could cause TTS or PTS during training and testing activities under Alternative 1 (for impact tables, see Appendix E, Estimated Marine Mammal and Sea Turtle Impacts from Exposure to Acoustic and Explosive Stressors Under Navy Training and Testing Activities). Olive ridley sea turtle presence in the Study Area is limited and density data does not exist

due to low occurrence in this region. Therefore, exposures were only modeled for green, hawksbill, leatherback, and loggerhead sea turtles in the Study Area and transit corridor. Fractional estimated impacts per region and activity area represent the probability that the number of estimated impacts by effect would occur in a certain region or be due to a certain activity category.



**Figure 3.5-8: Green Sea Turtle Estimated Impacts per Year from the Maximum Number of Explosions During Training and Testing Under Alternative 1**

Threshold shifts and injuries could reduce the fitness of an individual animal, causing a reduction in foraging success, reproduction, or increased susceptibility to predators. This reduction in fitness would be temporary for recoverable impacts, such as TTS, but there could be long-term consequences to some individuals. However, no population-level impact is expected due to the low number of estimated injuries for any sea turtle species relative to total population size. This can also be assumed for olive ridley turtles if exposed to explosions.

As discussed in Section 5.3.3 (Explosive Stressors), procedural mitigation includes ceasing explosive detonations (e.g., ceasing deployment of an explosive bomb) if a sea turtle is observed in the mitigation zone whenever and wherever applicable activities occur. In addition to this procedural mitigation, the Navy will implement mitigation to avoid or reduce impacts from explosions on seafloor resources in mitigation areas throughout the Study Area, as described in Section 5.4.1 (Mitigation Areas for Seafloor

Resources). This will further reduce the potential for impacts on sea turtles that shelter and feed on shallow-water coral reefs, live hard bottom, artificial reefs, and shipwrecks.

Sea turtle hearing is less sensitive than other marine animal hearing (i.e., marine mammals), and the role of their underwater hearing is unclear. Sea turtle's limited hearing range (<2 kHz) is most likely used to detect nearby broadband, continuous environmental sounds, such as the sounds of waves crashing on the beach, that may be important for identifying their habitat. Recovery from a hearing threshold shift begins almost immediately after the noise exposure ceases. A temporary threshold shift is expected to take a few minutes to a few days, depending on the severity of the initial shift, to fully recover (U.S. Department of the Navy, 2017a). If any hearing loss remains after recovery, that remaining hearing threshold shift is permanent. Because explosions produce broadband sounds with low-frequency content, hearing loss due to explosive sound could occur across a sea turtle's very limited hearing range, reducing the distance over which relevant sounds, such as beach sounds, may be detected for the duration of the threshold shift.

Some sea turtles may behaviorally respond to the sound of an explosive. A sea turtle's behavioral response to a single detonation or explosive cluster is expected to be limited to a short-term (seconds to minutes) startle response, as the duration of noise from these events is very brief. Limited research and observations from air gun studies (See Section 3.5.2.2.2.1, Methods for Analyzing Impacts from Explosives) suggest that if sea turtles are exposed to repetitive impulsive sounds in close proximity, they may react by increasing swim speed, avoiding the source, or changing their position in the water column. There is no evidence to suggest that any behavioral response would persist beyond the sound exposure. Because the duration of most explosive events is brief, the potential for masking is low. The *ANSI Sound Exposure Guidelines* (Popper et al., 2014) consider masking to not be a concern for sea turtles exposed to explosions.

A physiological stress response is assumed to accompany any injury, hearing loss, or behavioral reaction. A stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor. While the stress response is a normal function for an animal dealing with natural stressors in their environment, chronic stress responses could reduce an individual's fitness. Due to the low number of estimated impacts, it is not likely that any sea turtle would experience repeated stress responses due to explosive impacts.

*Pursuant to the ESA, use of explosives during training and testing activities as described under Alternative 1 may affect ESA-listed green, hawksbill, leatherback, loggerhead, and olive ridley sea turtles.*

#### **3.5.2.2.2.5 Alternative 2**

Activities using explosives would be conducted as described in Chapter 2 (Description of Proposed Action and Alternatives) and Appendix A (Training and Testing Activities Descriptions). General characteristics, quantities, and net explosive weights of in-water explosives used during training under Alternative 2 are provided in Section 3.0.4.2 (Explosive Stressors). Quantities and locations of fragment-producing explosives during training under Alternative 2 are shown in 3.0.4.4.4 (Military Expended Materials).

Under Alternative 2, there could be fluctuation in the amount of explosions that could occur annually, although potential impacts would be similar from year to year. The number of impulsive sources in this SEIS/OEIS compared with the totals analyzed in the 2015 MITT Final EIS/OEIS are described in Tables 2-7 and 2-8.


The numbers of activities planned under Alternative 2 are consistent from year-to-year and would increase slightly compared to activities planned under Alternative 1. The numbers of explosives used under each alternative are described in Section 3.0.4.2 (Explosive Stressors).

Under Alternative 2, it is possible that impacts would be slightly increased in some years, as explosive use would fluctuate. The quantitative analysis, using a maximum year of training and testing activities, estimates that no sea turtles would be killed, however, a small number of green sea turtles would be exposed to levels of explosive sound and energy in the outer Apra Harbor that could cause TTS or PTS (Figure 3.5-9). The quantitative analysis predicts that no hawksbill, leatherback, or loggerhead sea turtles are likely to be exposed to the levels of explosive sound and energy that could cause TTS, PTS, or injury during training and testing activities under Alternative 2. Olive ridley sea turtle presence in the Study Area is limited and density data does not exist due to low occurrence in this region. Therefore, exposures were only modeled for green, hawksbill, leatherback, and loggerhead sea turtles in the Study Area and transit corridor. Fractional estimated impacts per region and activity area represent the probability that the number of estimated impacts by effect would occur in a certain region or be due to a certain activity category.

Threshold shifts and injuries could reduce the fitness of an individual animal, causing a reduction in foraging success, reproduction, or increased susceptibility to predators. This reduction in fitness would be temporary for recoverable impacts, such as TTS, but there could be long-term consequences to some individuals. However, no population-level impact is expected due to the low number of estimated injuries for any sea turtle species relative to total population size. This can also be assumed for olive ridley turtles exposed to explosions.

As discussed in Section 5.3.3 (Explosive Stressors), procedural mitigation includes ceasing explosive detonations (e.g., ceasing deployment of an explosive bomb) if a sea turtle is observed in the mitigation zone whenever and wherever applicable activities occur. In addition to this procedural mitigation, the Navy will implement mitigation to avoid or reduce impacts from explosions on seafloor resources in mitigation areas throughout the Study Area, as described in Section 5.4.1 (Mitigation Areas for Seafloor Resources). This would further reduce the potential for impacts on sea turtles that shelter and feed on shallow-water coral reefs, live hard bottom, artificial reefs, and shipwrecks.

Sea turtle hearing is less sensitive than other marine animals (i.e., marine mammals), and the role of their underwater hearing is unclear. Sea turtle's limited hearing range (<2 kHz) is most likely used to detect nearby broadband, continuous environmental sounds, such as the sounds of waves crashing on the beach, that may be important for identifying their habitat. Recovery from a hearing threshold shift begins almost immediately after the noise exposure ceases. A temporary threshold shift is expected to take a few minutes to a few days, depending on the severity of the initial shift, to fully recover (U.S. Department of the Navy, 2017a). If any hearing loss remains after recovery, that remaining hearing threshold shift is permanent. Because explosions produce broadband sounds with low-frequency content, hearing loss due to explosive sound could occur across a sea turtle's very limited hearing range, reducing the distance over which relevant sounds, such as beach sounds, may be detected for the duration of the threshold shift.



**Figure 3.5-9: Green Sea Turtle Impacts Estimated per Year from Explosions During Training and Testing Under Alternative 2**

Some sea turtles may behaviorally respond to the sound of an explosive. A sea turtle’s behavioral response to a single detonation or explosive cluster is expected to be limited to a short-term (seconds to minutes) startle response, as the duration of noise from these events is very brief. Limited research and observations from air gun studies suggest that if sea turtles are exposed to repetitive impulsive sounds in close proximity, they may react by increasing swim speed, avoiding the source, or changing their position in the water column. There is no evidence to suggest that any behavioral response would persist beyond the sound exposure. Because the duration of most explosive events is brief, the potential for masking is low. The *ANSI Sound Exposure Guidelines* (Popper et al., 2014) consider masking to not be a concern for sea turtles exposed to explosions.

A physiological stress response is assumed to accompany any injury, hearing loss, or behavioral reaction. A stress response is a suite of physiological changes that are meant to help an organism mitigate the impact of a stressor. While the stress response is a normal function for an animal dealing with natural stressors in their environment, chronic stress responses could reduce an individual’s fitness. Due to the low number of estimated impacts, it is not likely that any sea turtle would experience repeated stress responses due to explosive impacts.

*Pursuant to the ESA, use of explosives during training and testing activities as described under Alternative 2 may affect ESA-listed green, hawksbill, leatherback, loggerhead, and olive ridley sea turtles.*

#### **3.5.2.2.2.6 No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosive stressors (e.g., explosive shock wave and sound, explosive fragments) as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer explosive stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for explosive impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

#### **3.5.2.3 Energy Stressors**

Energy stressors are discussed in Section 3.0.4.3. Energy stressors that may impact sea turtles include in-water electromagnetic devices and high-energy lasers. With the increased use of undersea power cables associated with offshore energy generation, there has been renewed scientific interest in electromagnetic fields possibly affecting migrating marine animals (Brothers & Lohmann, 2015; Endres et al., 2016; Gill et al., 2014; Kremers et al., 2014; Kremers et al., 2016; Putman et al., 2015; Zellar et al., 2017). There is no new information that changes the basis of the conclusion. These additional scientific findings do not change in any way the rationale for the dismissal of in-water electromagnetic devices as presented in the 2015 analyses. While the number of training and testing activities using in-water electromagnetic devices would change slightly under this SEIS/OEIS, the analysis presented in the 2015 MITT Final EIS/OEIS, Section 3.5.4.3 (Energy Stressors), and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Marine Fisheries Service, 2015b) remains valid for in-water electromagnetic devices.

High-energy laser use was not covered in the 2015 MITT Final EIS/OEIS and represents a new activity analyzed in this SEIS/OEIS. As discussed in Section 3.0.4.3.2.2 (High-Energy Lasers), high-energy lasers are designed to disable surface targets, rendering them immobile. The primary concern for high-energy weapons testing is the potential for a sea turtle to be struck by a high-energy laser beam at or near the water's surface, which could result in injury or death, resulting from traumatic burns from the beam. Sea turtles could be exposed to a laser only if the beam missed the target. Should the laser strike the sea surface, individual sea turtles at or near the surface could be exposed. The potential for exposure to a high-energy laser beam decreases as the water depth increases. Because laser platforms are typically helicopters and ships, sea turtles at sea would likely transit away or submerge in response to other stressors, such as ship or aircraft noise, although some sea turtles may not exhibit a response to an oncoming vessel or aircraft, increasing the risk of contact with the laser beam.

High-energy laser events typically have short ranges from the source to target, the aim point being a surface target, with an inherent precision of the weapon and its targeting system. The Navy conducted statistical modeling to estimate the probability of a sea turtle being struck by a high-energy laser during training and testing activities. As a basis for modeling the probability of high-energy laser strike, the Navy used green sea turtles as a proxy in the modeling because green sea turtles had the highest

average density estimates among sea turtle species (U.S. Department of the Navy, 2018b). Appendix J (Statistical Probability Analysis for Estimating Direct Strike Impact and Number of Potential Exposures from Military Expended Materials) shows the results of the modeling. As shown in Table J-2 of Appendix J, the modeling results suggest that estimate an extremely low probability of a direct strike by a high-energy laser on a sea turtle, with a reasonable assumption that no strike of sea turtles would occur (U.S. Department of the Navy, 2018b).

#### **3.5.2.3.1 Impacts from In-Water Electromagnetic Devices Under Alternative 1**

Under Alternative 1, the number of proposed training and testing activities involving the use of in-water electromagnetic devices would decrease in comparison to the 2015 MITT Final EIS/OEIS (Table 3.0-9). The activities would occur in the same locations and in a similar manner as were analyzed previously.

Therefore, impacts on sea turtles under Alternative 1 from energy stressors, including in-water electromagnetic devices, would be negligible.

*Pursuant to the ESA, the use of in-water electromagnetic devices during training and testing activities as described under Alternative 1 would have no effect on ESA-listed sea turtles.*

#### **3.5.2.3.2 Impacts from In-Water Electromagnetic Devices Under Alternative 2**

Under Alternative 2, the number of proposed training and testing activities involving the use of in-water electromagnetic devices would decrease in comparison to the 2015 MITT Final EIS/OEIS (Table 3.0-9). The activities would occur in the same locations and in a similar manner as were analyzed previously and above for Alternative 1.

Therefore, impacts on sea turtles under Alternative 2 from energy stressors, including in-water electromagnetic devices, would be negligible.

*Pursuant to the ESA, the use of in-water electromagnetic devices during training and testing activities, as described under Alternative 2 would have no effect on ESA-listed sea turtles.*

#### **3.5.2.3.3 Impacts from In-Water Electromagnetic Devices Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Energy stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer energy stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for energy impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

#### **3.5.2.3.4 Impacts from High-Energy Lasers Under Alternative 1**

Alternative 1 would introduce high-energy lasers into the Study Area, which is analyzed in this SEIS/OEIS as a new substressor not previously analyzed in the 2015 MITT Final EIS/OEIS. As stated previously, the Navy conducted statistical modeling to estimate the number of potential exposures of sea turtles to high-energy laser beams. Under Alternative 1, the modeling estimated 0.000025 annual exposures, an extremely low estimate (see Table J-2, in Appendix J, Statistical Probability Analysis for Estimating Direct

Strike Impact and Number of Potential Exposures from Military Expended Materials). Based on the very low number of annual exposures, the characteristics of activities that would use high-energy lasers (e.g., short range distance from source to target, high-precision targeting, short duration of the energized beam), and likely avoidance behavior of sea turtles to other stressors (e.g., vessel or aircraft noise), there is a reasonable assurance that there is no risk to sea turtles from high-energy laser use within the Study Area.

*Pursuant to the ESA, the use of high-energy lasers during training and testing activities as described under Alternative 1 would have no effect on ESA-listed sea turtles.*

#### **3.5.2.3.5 Impacts from High-Energy Lasers Under Alternative 2**

Under Alternative 2, the number of proposed activities involving the use of high-energy lasers would increase from Alternative 1 (Table 3.0-10) and the 2015 MITT Final EIS/OEIS. The increase in the number of events that use high-energy lasers is reflected in the Navy's statistical modeling of potential exposures of sea turtles with a slight increase in the model's estimates. As shown in Table J-2 in Appendix J (Statistical Probability Analysis for Estimating Direct Strike Impact and Number of Potential Exposures from Military Expended Materials) high-energy laser use under Alternative 2 would result in 0.000027 exposures every year. As with Alternative 1, based on the very low number of annual exposures, the characteristics of activities that would use high-energy lasers (e.g., short range distance from source to target, high-precision targeting, short duration of the energized beam), and likely avoidance behavior of sea turtles to other stressors (e.g., vessel or aircraft noise), there is a reasonable assurance that there is no risk to sea turtles from high-energy laser use within the Study Area.

*Pursuant to the ESA, the use of in-water electromagnetic devices and high-energy lasers during training and testing activities as described under Alternative 2 would have no effect on ESA-listed sea turtles.*

#### **3.5.2.3.6 Impacts from High-Energy Lasers Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Energy stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer energy stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for energy impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

#### **3.5.2.4 Physical Disturbance and Strike Stressors**

Physical disturbance and strike stressors are discussed in Section 3.0.4.4 (Physical Disturbance and Strike Stressors). Physical disturbance and strike stressors that may impact sea turtles include (1) vessels and in-water devices, (2) military expended materials, and (3) seafloor devices. The annual number of events including vessels and in-water devices, the annual number of military expended materials, and the annual number of events including seafloor devices are shown in Tables 3.0-12 through 3.0-18. The Navy will implement further mitigation measures to avoid or reduce potential impacts of towed in-water devices, non-explosive practice munitions, and vessel movements (see Sections 5.3.4.1 through 5.3.4.3).

There have been no known instances of physical disturbance or strike to any sea turtle in the Study Area as a result of Navy training and testing activities prior to or since the 2015 MITT Final EIS/OEIS.

#### **3.5.2.4.1 Impacts from Physical Disturbance and Strike Stressors Under Alternative 1**

Under Alternative 1, analysis of the individual substressors including the use of vessels and in-water devices, military expended materials, and seafloor devices presented in Section 3.0.4.4 (Physical Disturbance and Strike Stressors) indicates that those items having the most potential to affect sea turtles have decreased in comparison to the 2015 MITT Final EIS/OEIS (Tables 3.0-12 through 3.0-18). The number of small-caliber munitions would increase under Alternative 1. Small-caliber munitions are inert, are meant to be aimed at targets, and are not long-range weapons. As a result, sea turtles are extremely unlikely to be disturbed or struck by expended small-caliber munitions. Given the reduction in physical disturbance and strike stressors for this SEIS/OEIS, the findings presented in the 2015 MITT Final EIS/OEIS, Section 3.5.2.4 (Physical Disturbance and Strike Stressors), and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015) remain valid. There has been no new science applicable to a further understanding of physical disturbance and strike stressors in the Study Area. In areas outside the Study Area (e.g., Hawaii and Southern California), there have been recorded military vessel strikes of sea turtles. However, these are areas where the number of military vessels is much higher and training and testing activities occur more often than in the Study Area.

*Pursuant to the ESA, the use of vessels and in-water devices, military expended materials, and seafloor devices as described under Alternative 1 may affect ESA-listed sea turtles.*

#### **3.5.2.4.2 Impacts from Physical Disturbance and Strike Stressors Under Alternative 2**

Under Alternative 2, physical disturbance and strike stressors during training and testing activities would decrease compared to the 2015 MITT Final EIS/OEIS (Tables 3.0-12 through 3.0-18), assuming the dismissal of small-caliber munitions use for the reasons noted above. Under Alternative 2, there would be additional physical disturbance and strike stressors in comparison to Alternative 1, but the conclusions remain the same. Therefore, the potential for strikes of sea turtles from vessels and in-water devices, military expended materials, and seafloor devices are unlikely to occur.

*Pursuant to the ESA, the use of vessels and in-water devices, military expended materials, and seafloor devices as described under Alternative 2 may affect ESA-listed sea turtles.*

#### **3.5.2.4.3 Impacts from Physical Disturbance and Strike Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Physical disturbance and strike stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer physical disturbance and strike stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for physical disturbance and strike impacts on individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

### 3.5.2.5 Entanglement Stressors

Entanglement stressors are discussed in Section 3.0.4.5. Entanglement stressors considered for sea turtles include (1) fiber optic cables and guidance wires, and (2) decelerators/parachutes. The annual number of wires and cables and decelerators/parachutes proposed under the alternatives and in comparison to current ongoing activities are presented in Tables 3.0-20 through 3.0-22. There have been no known instances of any sea turtle being entangled in wires and cables, or decelerators/parachutes associated with training and testing activities prior to or since the 2015 MITT Final EIS/OEIS.

#### 3.5.2.5.1 Impacts from Entanglement Stressors Under Alternative 1

Under Alternative 1, the annual number of entanglement stressors would decrease compared to the 2015 MITT Final EIS/OEIS (Tables 3.0-20 through 3.0-22). Therefore, the analysis from the 2015 MITT Final EIS/OEIS remains valid. The analysis presented in the 2015 MITT Final EIS/OEIS (Section 3.5.2.5, Entanglement Stressors) and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015) determined that impacts on sea turtles from entanglement stressors are not anticipated.

*Pursuant to the ESA, the use of fiber optic cable and guidance wires and decelerators/parachutes as described under Alternative 1 may affect ESA-listed sea turtles.*

#### 3.5.2.5.2 Impacts from Entanglement Stressors Under Alternative 2

Under Alternative 2, the number of entanglement stressors would decrease in comparison to current ongoing activities for fiber optic cable and decelerators/parachutes but would increase for the annual number of expended guidance wire (Tables 3.0-20 through 3.0-22). In comparison to Alternative 1, there would be a slight increase under Alternative 2 for entanglement stressors; however, the combined number of annual entanglement stressors (fiber optic cable, guidance wire, and decelerators/parachutes) decreases when compared to the 2015 MITT Final EIS/OEIS. Therefore, the analysis and conclusions presented in the 2015 MITT Final EIS/OEIS (Section 3.5.2.5, Entanglement Stressors) and the NMFS Biological Opinion for the 2015 MITT Final EIS/OEIS (National Oceanic and Atmospheric Administration, 2015) remain valid. Impacts on sea turtles from entanglement stressors are not anticipated.

*Pursuant to the ESA, the use of fiber optic cable and guidance wires and decelerators/parachutes as described above under Alternative 2 may affect ESA-listed sea turtles.*

#### 3.5.2.5.3 Impacts from Entanglement Stressors Under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Entanglement stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer entanglement stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for entanglement of individual sea turtles, but would not measurably improve the status of sea turtle populations or subpopulations.

### 3.5.2.6 Ingestion Stressors

Ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) are discussed in Section 3.0.4.6. Types of materials that could become ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) during training and testing in the Study Area include non-explosive practice munitions (small- and medium-caliber), fragments from explosives, fragments from targets, chaff, flare casings (including plastic end caps and pistons), and decelerators/parachutes. The annual number of events including military expended materials are shown in Tables 3.0-14 through 3.0-17 and Tables 3.0-22 through 3.0-24. As discussed in Section 3.5.4.6.3 (Impacts from Munitions) of the 2015 MITT Final EIS/OEIS, the number of munitions and explosive munitions fragments that an individual sea turtle could encounter would generally be low, based on the patchy distribution of both the munitions and the habitats where sea turtles forage. For the more numerous small-caliber munitions, these expended material-type items are inert, small in size, do not resemble prey items, and end up as part of the seafloor, where they are unlikely to be encountered by most sea turtles. In addition, it is assumed for sea turtle species that may feed at the seafloor, that they would not ingest every munition or munition's fragment encountered; if a munition or munition's fragment were ingested, an animal may attempt to reject it when it realizes the item is not food.

#### 3.5.2.6.1 Impacts from Ingestion Stressors Under Alternative 1

Under Alternative 1, analysis of the individual substressors presented in Section 3.0.4.6 (Ingestion Stressors) indicates that those items considered ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) having the most potential to affect sea turtles have decreased (Tables 3.0-14 through 3.0-17 and Tables 3.0-22 through 3.0-24). For the reasons noted above, the Navy has determined that potential impacts from ingestion stressors would not be substantially different from the 2015 MITT Final EIS/OEIS. In the 2015 analysis of training and testing activities within the Study Area, NMFS determined that ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) would not result in harassment or harm of sea turtles or jeopardize the continued existence of any sea turtle species (National Oceanic and Atmospheric Administration, 2015). The activities expending munitions and other military expended materials analyzed in this SEIS/OEIS under Alternative 1 are not a significant change over what was analyzed in the 2015 MITT Final EIS/OEIS, and there has been no new science necessitating a revision of the 2015 conclusions in that regard. Impacts on sea turtles from ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) in the Study Area are not anticipated.

*Pursuant to the ESA, the use of munitions and other military expended materials as described under Alternative 1 may affect ESA-listed sea turtles.*

#### 3.5.2.6.2 Impacts from Ingestion Stressors Under Alternative 2

Under Alternative 2, of the number of military expended materials would decrease compared to the 2015 MITT Final EIS/OEIS, with the exception of increased use of small-caliber munitions (Tables 3.0-14 through 3.0-17 and Tables 3.0-22 through 3.0-24). Under Alternative 2, increases as compared to Alternative 1 do not change the impact conclusions for ingestion stressors (military expended materials – munitions and military expended materials – other than munitions) as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, impacts on sea turtles from ingestion of military expended materials under Alternative 2 are not expected.

*Pursuant to the ESA, the use of munitions and other military expended materials as described under Alternative 2 may affect ESA-listed sea turtles.*

### **3.5.2.6.3 Impacts from Ingestion Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Ingestion stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer ingestion stressors within the marine environment where Navy training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for individual sea turtles to ingest items expended during training and testing activities, but would not measurably improve the status of sea turtle populations or subpopulations.

### **3.5.2.7 Secondary Stressors**

As discussed in Section 3.5.3.6 (Secondary Stressors) of the 2015 MITT Final EIS/OEIS, secondary stressors from training and testing activities could pose indirect impacts on sea turtles via habitat degradation or an effect on prey availability. These stressors include (1) explosives, (2) explosive byproducts and unexploded ordnance, (3) metals, and (4) chemicals. Analyses of the potential impacts on sediments and water quality from the proposed training and testing activities are discussed in detail in Section 3.1 (Sediments and Water Quality) of the 2015 MITT Final EIS/OEIS. The analysis of explosives, explosive byproducts, metals, chemicals, and the transmission of diseases and parasites and their potential to indirectly impact sea turtles has not appreciably changed and is presented in detail in Section 3.5.4.7 (Secondary Stressors) of the 2015 MITT Final EIS/OEIS.

The analysis concluded that the relatively low solubility of most explosives and their degradation products means that concentrations of these contaminants in the marine environment, from either high-order or low-order detonations, are relatively low and readily diluted. Given that the concentration of unexploded ordnance, explosion byproducts, metals, and other chemicals would never exceed that of a World War II dump site where minimal concentrations were detected only within a few feet of the ordnance (Briggs et al., 2016; Edwards et al., 2016; Kelley et al., 2016; Smith & Marx, 2016), indirect impacts on sea turtles from the Proposed Action would be negligible and would have no long-term effect on habitat or prey.

### **3.5.3 Public Scoping Comments**

The public raised a number of issues during the scoping period in regard to Sea Turtles. The issues are summarized in the list below.

- **Public comments concerning a lack of sea turtle information in waters surrounding FDM** – One commenter noted a lack of studies documenting the condition of sea turtles in waters surrounding FDM. Multi-year dive studies conducted by Smith and Marx (2016) have reported roughly comparable numbers of sea turtles (only green sea turtles and hawksbill sea turtles have been observed in waters surrounding FDM) during every survey between 1999 and 2012. None of the specimens seen by the authors had any visible fibropapiloma tumors, barnacles, lesions, or other visible abnormalities. The number of sea turtle sightings during each dive session was

low, ranging from 0.13 to 0.36 per biologist per dive in each year. For comparative purposes, some study sites off Oahu, Hawaii which have been surveyed since 1999 by the authors have averaged more than 10 sea turtles sighted per dive during all seasons. This equates to 28 times higher than the FDM densities. The precipitous sea cliffs, lack of suitable haulout sites or beaches preclude nesting or basking at FDM. In addition, Smith and Marx (2016) noted that no sea turtle remains, such as carapace or bone fragments, have ever been sighted or reported at FDM (the authors have encountered such remains at various locations in the Bahamas, Cayman Islands, Hawaiian Islands and Malaysia, which support resident sea turtle populations). In summary, sea turtles around FDM probably represent transient individuals and not a resident population. Although waters surrounding FDM likely maintain healthy foraging grounds for transient turtles, they do not congregate in high concentrations in these waters.

- **Public comments on habitat, prey availability, and overall health of sea turtles** – The Navy received comments expressing concerns over impacts on the general marine environment from military training and testing activities. The Navy has included a detailed summary of recent published studies that describe multi-year dive studies conducted by Smith and Marx (2016), which provide an indication of habitat quality in waters surrounding a location of concentrated and intensive military activities. The results of these surveys are included in Section 3.1.1.1.3 (Farallon de Medinilla) of this SEIS/OEIS. Throughout all dive surveys, the coral fauna at FDM were observed to be healthy and robust, which suggests healthy foraging habitats for sea turtles. The nearshore physical environment and basic habitat types at FDM have remained unchanged over the 13 years of survey activity. These conclusions are based on (1) a limited amount of physical damage, (2) very low levels of partial mortality and disease (less than 1 percent of all species observed), (3) absence of excessive mucus production, (4) good coral recruitment, (5) complete recovery by 2012 of the 2007 bleaching event, and (6) a limited number of macrobioeroders and an absence of invasive crown of thorns starfish (*Acanthaster planci*). These factors suggest that potential impacts from training and testing activities are not sufficient as to adversely impact water quality, substantiated by repeated dive surveys discussed above (Smith & Marx, 2016), and thereby reduce habitat quality for sea turtle populations.

## REFERENCES

- Avens, L. (2003). Use of multiple orientation cues by juvenile loggerhead sea turtles *Caretta caretta*. *The Journal of Experimental Biology*, 206(23), 4317–4325.
- Balazs, G. H. (1986). Fibropapillomas in Hawaiian green turtles. *Marine Turtle Newsletter*, 39, 1–3.
- Bartol, S. M., J. A. Musick, and M. L. Lenhardt. (1999). Auditory evoked potentials of the loggerhead sea turtle (*Caretta caretta*). *Copeia*, 1999(3), 836–840.
- Bartol, S. M., and D. R. Ketten. (2006). *Turtle and Tuna Hearing* (NOAA Technical Memorandum NMFS-PIFSC-7). Honolulu, HI: Pacific Islands Fisheries Science Center.
- Becking, L. E., T. C. J. M. van Bussel, A. O. Debrot, and M. J. A. Christianen. (2014). First record of a Caribbean green turtle (*Chelonia mydas*) grazing on invasive seagrass (*Halophila stipulacea*). *Caribbean Journal of Science*, 48(2–3), 162–163.
- Berkson, H. (1967). Physiological adjustments to deep diving in the Pacific green turtle (*Chelonia mydas agassizii*). *Comparative Biochemistry and Physiology*, 21(3), 507–524.
- Birney, K., M. Byrd, S. Hastings, S. Herron, B. Shafritz, and R. Freedman. (2015). *Report on the 2014 Vessel Speed Reduction Incentive Trial in the Santa Barbara Channel* (Protecting Blue Whales and Blue Skies). Santa Barbara, CA: National Marine Sanctuaries Channel Islands, Santa Barbara County Air Pollution Control District, Environmental Defense Center, and the National Marine Sanctuary Foundation.
- Bjorndal, K. A., A. B. Bolten, and C. Lagueux. (1994). Ingestion of Marine Debris by Juvenile Sea Turtles in Coastal Florida Habitats. *Marine Pollution Bulletin*, 28(3), 154–158.
- Bjorndal, K. A. (1997). Foraging ecology and nutrition of sea turtles. In P. L. Lutz & J. A. Musick (Eds.), *The Biology of Sea Turtles* (pp. 199–231). Boca Raton, FL: CRC Press.
- Bolten, A. B. (2003). Active swimmers-passive drifters: The oceanic juvenile stage of loggerheads in the Atlantic system. In A. B. Bolten & B. E. Witherington (Eds.), *Loggerhead Sea Turtles* (pp. 63–78). Washington, DC: Smithsonian Books.
- Bowen, B. W., A. L. Bass, S.-M. Chow, M. Bostrom, K. A. Bjorndal, A. B. Bolten, T. Okuyama, B. M. Bolker, S. Epperly, E. Lacasella, D. Shaver, M. Dodd, S. R. Hopkins-Murphy, J. A. Musick, M. Swingle, K. Rankin-Baransky, W. Teas, W. N. Witzell, and P. H. Dutton. (2004). Natal homing in juvenile loggerhead turtles (*Caretta caretta*). *Molecular Ecology*, 13, 3797–3808.
- Briggs, C., S. M. Shjegstad, J. A. K. Silva, and M. H. Edwards. (2016). Distribution of chemical warfare agent, energetics, and metals in sediments at a deep-water discarded military munitions site. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 63–69.
- Brothers, J. R., and K. J. Lohmann. (2015). Evidence for geomagnetic imprinting and magnetic navigation in the natal homing of sea turtles. *Current Biology*, 25(3), 392–396.
- Christian, E. A., and J. B. Gaspin. (1974). *Swimmer Safe Standards from Underwater Explosions*. Navy Science Assistance Program Project No. PHP-11-73. White Oak, MD: Naval Ordnance Laboratory.
- Clark, C. W., M. W. Brown, and P. Corkeron. (2010). Visual and acoustic surveys for North Atlantic right whales, *Eubalaena glacialis*, in Cape Cod Bay, Massachusetts, 2001–2005: Management implications. *Marine Mammal Science*, 26(4), 837–843.

- Clark, H. R., and C. J. Gobler. (2016). Diurnal fluctuations in CO<sub>2</sub> and dissolved oxygen concentrations do not provide a refuge from hypoxia and acidification for early-life-stage bivalves. *Marine Ecology Progress Series*, 558, 1–14.
- Clark, S. L., and J. W. Ward. (1943). The effects of rapid compression waves on animals submerged in water. *Surgery, Gynecology & Obstetrics*, 77, 403–412.
- Conant, T. A., P. H. Dutton, T. Eguchi, S. P. Epperly, C. C. Fahy, M. H. Godfrey, S. L. MacPherson, E. E. Possardt, B. A. Schroeder, J. A. Seminoff, M. L. Snover, C. M. Upite, and B. E. Witherington. (2009). *Loggerhead sea turtle (Caretta caretta) 2009 status review under the U.S. Endangered Species Act* (Report of the loggerhead biological review team to the National Marine Fisheries Service, August 2009). Silver Spring, MD: Loggerhead Biological Review Team.
- Cook, S. L., and T. G. Forrest. (2005). Sounds produced by nesting leatherback sea turtles (*Dermochelys coriacea*). *Herpetological Review*, 36(4), 387–389.
- DeRuiter, S. L., and K. L. Doukara. (2012). Loggerhead turtles dive in response to airgun sound exposure. *Endangered Species Research*, 16(1), 55–63.
- Dew, L. A., R. G. Owen, and M. J. Mulroy. (1993). Changes in size and shape of auditory hair cells in vivo during noise-induced temporary threshold shift. *Hearing Research*, 66(1), 99–107.
- Dow Piniak, W. E., S. A. Eckert, C. A. Harms, and E. M. Stringer. (2012). *Underwater Hearing Sensitivity of the Leatherback Sea Turtle (Dermochelys coriacea): Assessing the Potential Effect of Anthropogenic Noise* (OCS Study BOEM 2012-01156). Herndon, VA: U.S. Department of the Interior, Bureau of Ocean Energy Management.
- Edwards, M. H., S. M. Shjegstad, R. Wilkens, J. C. King, G. Carton, D. Bala, B. Bingham, M. C. Bissonnette, C. Briggs, N. S. Bruso, R. Camilli, M. Cremer, R. B. Davis, E. H. DeCarlo, C. DuVal, D. J. Fornari, I. Kaneakua-Pia, C. D. Kelley, S. Koide, C. L. Mah, T. Kerby, G. J. Kurras, M. R. Rognstad, L. Sheild, J. Silva, B. Wellington, and M. V. Woerkom. (2016). The Hawaii undersea military munitions assessment. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 4–13.
- Eldredge, L. G. (2003). The marine reptiles and mammals of Guam. *Micronesica*, 35-36, 653–660.
- Endres, C. S., N. F. Putman, D. A. Ernst, J. A. Kurth, C. M. F. Lohmann, and K. J. Lohmann. (2016). Multi-modal homing in sea turtles: Modeling dual use of geomagnetic and chemical cues in island-finding. *Frontiers in Behavioral Neuroscience*, 10, 19.
- Falcone, E. A., G. S. Schorr, S. L. Watwood, S. L. DeRuiter, A. N. Zerbini, R. D. Andrews, R. P. Morrissey, and D. J. Moretti. (2017). Diving behaviour of Cuvier's beaked whales exposed to two types of military sonar. *Royal Society Open Science*, 4(170629), 1–21.
- Ferrara, C. R., R. C. Vogt, M. R. Harfush, R. S. Sousa-Lima, E. Albavera, and A. Tavera. (2014). First evidence of leatherback turtle (*Dermochelys coriacea*) embryos and hatchlings emitting sounds. *Chelonian Conservation and Biology*, 13(1), 110–114.
- Flower, J. E., T. M. Norton, K. M. Andrews, S. E. Nelson, Jr., C. E. Parker, L. M. Romero, and M. A. Mitchell. (2015). Baseline plasma corticosterone, haematological and biochemical results in nesting and rehabilitating loggerhead sea turtles (*Caretta caretta*). *Conservation Physiology*, 3(1), cov003.
- Fossette, S., S. Ferraroli, H. Tanaka, Y. Ropert-Coudert, N. Arai, K. Sato, Y. Naito, Y. Le Maho, and J. Georges. (2007). Dispersal and dive patterns in gravid leatherback turtles during the nesting season in French Guiana. *Marine Ecology Progress Series*, 338, 233–247.

- Fuentes, M. M. P. B., D. A. Pike, A. Dimatteo, and B. P. Wallace. (2013). Resilience of marine turtle regional management units to climate change. *Global Change Biology*, 19(5), 1399–1406.
- Fukuoka, T., M. Yamane, C. Kinoshita, T. Narazaki, G. J. Marshall, K. J. Abernathy, N. Miyazaki, and K. Sato. (2016). The feeding habit of sea turtles influences their reaction to artificial marine debris. *Scientific Reports*, 6, 28015.
- Garcia-Parraga, D., J. L. Crespo-Picazo, Y. B. de Quiros, V. Cervera, L. Marti-Bonmati, J. Diaz-Delgado, M. Arbelo, M. J. Moore, P. D. Jepson, and A. Fernandez. (2014). Decompression sickness ('the bends') in sea turtles. *Diseases of Aquatic Organisms*, 111(3), 191–205.
- Gill, A. B., I. Gloyne-Philips, J. Kimber, and P. Sigray. (2014). Marine Renewable Energy, Electromagnetic (EM) Fields and EM-Sensitive Animals. In M. Shields & A. Payne (Eds.), *Marine Renewable Energy Technology and Environmental Interactions. Humanity and the Sea* (pp. 61–79). Dordrecht, Netherlands: Springer.
- Gitschlag, G. R. (1996). Migration and diving behavior of Kemp's ridley (Garman) sea turtles along the U.S. southeastern Atlantic coast. *Journal of Experimental Marine Biology and Ecology*, 205, 115–135.
- Goertner, J. F. (1978). *Dynamical Model for Explosion Injury to Fish*. Dalgren, VA: U.S. Department of the Navy, Naval Surface Weapons Center.
- Goertner, J. F. (1982). *Prediction of Underwater Explosion Safe Ranges for Sea Mammals*. Dalgren, VA: Naval Surface Weapons Center.
- Greaves, F. C., R. H. Draeger, O. A. Brines, J. S. Shaver, and E. L. Corey. (1943). An experimental study of concussion. *United States Naval Medical Bulletin*, 41(1), 339–352.
- Gregory, L. F., and J. R. Schmid. (2001). Stress response and sexing of wild Kemp's ridley sea turtles (*Lepidochelys kempii*) in the Northeastern Gulf of Mexico. *General and Comparative Endocrinology*, 124, 66–74.
- Hawkes, L. A., A. C. Broderick, M. S. Coyne, M. H. Godfrey, L.-F. Lopez-Jurado, P. Lopez-Suarez, S. E. Merino, N. Varo-Cruz, and B. J. Godley. (2006). Phenotypically linked dichotomy in sea turtle foraging requires multiple conservation approaches. *Current Biology*, 16, 990–995.
- Hays, G. C., J. D. R. Houghton, C. Isaacs, R. S. King, C. Lloyd, and P. Lovell. (2004). First records of oceanic dive profiles for leatherback turtles, *Dermochelys coriacea*, indicate behavioural plasticity associated with long-distance migration. *Animal Behaviour*, 67, 733–743.
- Henry, W. R., and M. J. Mulroy. (1995). Afferent synaptic changes in auditory hair cells during noise-induced temporary threshold shift. *Hearing Research*, 84(1), 81–90.
- Hetherington, T. (2008). Comparative anatomy and function of hearing in aquatic amphibians, reptiles, and birds. In J. G. M. Thewissen & S. Nummela (Eds.), *Sensory Evolution on the Threshold* (pp. 182–209). Berkeley, CA: University of California Press.
- Hochscheid, S., C. R. McMahon, C. J. A. Bradshaw, F. Maffucci, F. Bentivegna, and G. C. Hays. (2007). Allometric scaling of lung volume and its consequences for marine turtle diving performance. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 148(2), 360–367.
- Hochscheid, S. (2014). Why we mind sea turtles' underwater business: A review on the study of diving behavior. *Journal of Experimental Marine Biology and Ecology*, 450, 118–136.

- Hoopes, L. A., A. M. Landry Jr., and E. K. Stabenau. (2000). Physiological effects of capturing Kemp's ridley sea turtles, *Lepidochelys kempii*, in entanglement nets. *Canadian Journal of Zoology*, 78(11), 1941–1947.
- Houghton, J. D. R., M. J. Callow, and G. C. Hays. (2003). Habitat utilization by juvenile hawksbill turtles (*Eretmochelys imbricata*, Linnaeus, 1766) around a shallow water coral reef. *Journal of Natural History*, 37, 1269–1280.
- Hubbs, C., and A. Rechnitzer. (1952). Report on experiments designed to determine effects of underwater explosions on fish life. *California Fish and Game*, 38, 333–366.
- James, M. C., S. A. Eckert, and R. A. Myers. (2005). Migratory and reproductive movements of male leatherback turtles (*Dermochelys coriacea*). *Marine Biology*, 147, 845–853.
- Kelley, C., G. Carton, M. Tomlinson, and A. Gleason. (2016). Analysis of towed camera images to determine the effects of disposed mustard-filled bombs on the deep water benthic community off south Oahu. *Deep Sea Research Part II: Topical Studies in Oceanography*, 128, 34–42.
- Ketten, D. R., and S. Moein-Bartol. (2006). *Functional Measures of Sea Turtle Hearing*. Woods Hole, MA: Woods Hole Oceanographic Institution.
- Klima, E. F., G. R. Gitschlag, and M. L. Renaud. (1988). Impacts of the explosive removal of offshore petroleum platforms on sea turtles and dolphins. *Marine Fisheries Review*, 50(3), 33–42.
- Kolinski, S. P., R. K. Hoeke, S. R. Holzwarth, L. I. Ilo, E. F. Cox, R. C. O'Conner, and P. S. Vroom. (2006). Nearshore distribution and an abundance estimate for green sea turtles, *Chelonia mydas*, at Rota Island, Commonwealth of the Northern Mariana Islands. *Pacific Science*, 60(4), 509–522.
- Kremers, D., J. Lopez Marulanda, M. Hausberger, and A. Lemasson. (2014). Behavioural evidence of magnetoreception in dolphins: Detection of experimental magnetic fields. *Die Naturwissenschaften*, 101(11), 907–911.
- Kremers, D., A. Celerier, B. Schaal, S. Campagna, M. Trabalon, M. Boye, M. Hausberger, and A. Lemasson. (2016). Sensory Perception in Cetaceans: Part II—Promising Experimental Approaches to Study Chemoreception in Dolphins. *Frontiers in Ecology and Evolution*, 4(50), 1–9.
- Laloë, J.-O., N. Esteban, J. Berkel, and G. C. Hays. (2016). Sand temperatures for nesting sea turtles in the Caribbean: Implications for hatchling sex ratios in the face of climate change. *Journal of Experimental Marine Biology and Ecology*, 474, 92–99.
- Lavender, A. L., S. M. Bartol, and I. K. Bartol. (2014). Ontogenetic investigation of underwater hearing capabilities in loggerhead sea turtles (*Caretta caretta*) using a dual testing approach. *The Journal of Experimental Biology*, 217(Pt 14), 2580–2589.
- Lenhardt, M. L., S. Bellmund, R. A. Byles, S. W. Harkins, and J. A. Musick. (1983). Marine turtle reception of bone-conducted sound. *The Journal of Auditory Research*, 23, 119–125.
- Lenhardt, M. L., R. C. Klinger, and J. A. Musick. (1985). Marine turtle middle-ear anatomy. *The Journal of Auditory Research*, 25, 66–72.
- Lenhardt, M. L. (1994). *Seismic and very low frequency sound induced behaviors in captive loggerhead marine turtles (Caretta caretta)*. Paper presented at the Proceedings of the Fourteenth Annual Symposium on Sea Turtle Biology and Conservation. Hilton Head, SC.
- Lutcavage, M. E., P. G. Bushnell, and D. R. Jones. (1992). Oxygen stores and aerobic metabolism in the leatherback sea turtle. *Canadian Journal of Zoology*, 70(2), 348–351.

- Lutcavage, M. E., and P. L. Lutz. (1997). Diving Physiology. In P. L. Lutz & J. A. Musick (Eds.), *The Biology of Sea Turtles* (pp. 277–296). Boca Raton, FL: CRC Press.
- Martin, K. J., S. C. Alessi, J. C. Gaspard, A. D. Tucker, G. B. Bauer, and D. A. Mann. (2012). Underwater hearing in the loggerhead turtle (*Caretta caretta*): A comparison of behavioral and auditory evoked potential audiograms. *The Journal of Experimental Biology*, 215(17), 3001–3009.
- Martin, S. L., K. S. Van Houtan, T. T. Jones, C. F. Aguon, J. T. Gutierrez, R. B. Tibbatts, S. B. Wusstig, and J. D. Bass. (2016). Five decades of marine megafauna surveys from Micronesia. *Frontiers in Marine Science*, 2(116), 1–13.
- McCauley, R. D., J. Fewtrell, A. J. Duncan, C. Jenner, M.-N. Jenner, J. D. Penrose, R. I. T. Prince, A. Adhitya, J. Murdoch, and K. A. McCabe. (2000). *Marine Seismic Surveys: Analysis and Propagation of Air-gun Signals; and Effects of Air-gun Exposure on Humpback Whales, Sea Turtles, Fishes and Squid*. Bentley, Australia: Centre for Marine Science and Technology.
- Moein Bartol, S. E., J. A. Musick, J. A. Keinath, D. E. Barnard, M. L. Lenhardt, and R. George. (1995). Evaluation of Seismic Sources for Repelling Sea Turtles from Hopper Dredges. In L. Z. Hales (Ed.), *Sea Turtle Research Program: Summary Report* (Vol. Technical Report CERC-95, pp. 90–93). Kings Bay, GA: U.S. Army Engineer Division, South Atlantic, Atlanta, GA and U.S. Naval Submarine Base.
- Mrosovsky, N. (1972). Spectrographs of the sounds of leatherback turtles. *Herpetologica*, 28(3), 256–258.
- Mrosovsky, N., G. D. Ryan, and M. C. James. (2009). Leatherback turtles: The menace of plastic. *Marine Pollution Bulletin*, 58(2), 287–289.
- Murray, C. C., A. Bychkov, T. Therriault, H. Maki, and N. Wallace. (2015). The impact of Japanese tsunami debris on North America. *PICES Press*, 23(1), 28.
- Narazaki, T., K. Sato, K. J. Abernathy, G. J. Marshall, and N. Miyazaki. (2013). Loggerhead turtles (*Caretta caretta*) use vision to forage on gelatinous prey in mid-water. *PLoS ONE*, 8(6), e66043.
- National Marine Fisheries Service, and U.S. Fish and Wildlife Service. (1991). *Recovery Plan for U.S. Populations of Atlantic Green Turtle (Chelonia mydas)*. Washington, DC: National Marine Fisheries Service.
- National Marine Fisheries Service, and U.S. Fish and Wildlife Service. (2008). *Recovery Plan for the Northwest Atlantic Population of the Loggerhead Sea Turtle (Caretta caretta), Second Revision*. Silver Spring, MD: National Marine Fisheries Service.
- National Marine Fisheries Service. (2013). *2013 Annual Report of a Comprehensive Assessment of Marine Mammal, Marine Turtle, and Seabird Abundance and Spatial Distribution in US Waters of the Western North Atlantic Ocean*. Woods Hole, MA and Miami, FL: Northeast Fisheries Science Center and Southeast Fisheries Science Center.
- National Marine Fisheries Service, and U.S. Fish and Wildlife Service. (2014). *Olive Ridley Sea Turtle (Lepidochelys olivacea) 5-Year Review: Summary and Evaluation*. Silver Spring, MD: National Marine Fisheries Service Office of Protected Resources and U.S. Fish and Wildlife Service Southeast Region.
- National Marine Fisheries Service. (2015a). *National Marine Fisheries Service Endangered Species Act Section 7 Consultation Biological Opinion and Conference Report; Biological Opinion and Conference Report on Mariana Islands Training and Testing and Issuance of an MMPA Rule and*

- LOA. Silver Spring, MD: Endangered Species Act Interagency Cooperation Division of the Office of Protected Resources, National Marine Fisheries Service.
- National Marine Fisheries Service. (2015b). *Endangered Species Act Section 7 Biological Opinion and Conference Report U.S. Navy's Surveillance Towed Array Sensor System Low Frequency Active Sonar Routine Training, Testing, and Military Operations and NOAA's National Marine Fisheries Service, Office of Protected Resources' issuance of four letters of authorization for the U.S. Navy to "take" marine mammals incidental to Surveillance Towed Array Sensor System Low Frequency Active Sonar Routine Training, Testing, and Military Operations in areas of the Pacific Ocean for the period of August 15, 2015 through August 14, 2016 pursuant to the fiveyear MMPA regulation.* (FPR-2015-9119). Washington, DC: U.S. Department of the Navy.
- National Oceanic and Atmospheric Administration. (2015). Takes of marine mammals incidental to specified activities; U.S. Navy training and testing activities in the Mariana Islands Training and Testing Study Area. *Federal Register*, 80(148), 46112–46171.
- O'Hara, J., and J. R. Wilcox. (1990). Avoidance responses of loggerhead turtles, *Caretta caretta*, to low frequency sound. *Copeia*, 1990(2), 564–567.
- O'Keefe, D. J., and G. A. Young. (1984). *Handbook on the Environmental Effects of Underwater Explosions*. Silver Spring, MD: U.S. Navy, Naval Surface Weapons Center (Code R14).
- Office of the Surgeon General. (1991). Conventional warfare ballistic, blast, and burn injuries. In R. Zajitchuk, Col. (Ed.), *U.S.A. Textbook of Military Medicine*. Washington, DC: Office of the Surgeon General.
- Patino-Martinez, J., A. Marco, L. Quinones, and B. Godley. (2008). Globally significant nesting of the leatherback turtle (*Dermochelys coriacea*) on the Caribbean coast of Colombia and Panama. *Biological Conservation*, 141(8), 1982–1988.
- Patrício, A. R., C. E. Diez, R. P. Van Dam, and B. J. Godley. (2016). Novel insights into the dynamics of green turtle fibropapillomatosis. *Marine Ecology Progress Series*, 547, 247–255.
- Pepper, C. B., M. A. Nascarella, and R. J. Kendall. (2003). A review of the effects of aircraft noise on wildlife and humans, current control mechanisms, and the need for further study. *Environmental Management*, 32(4), 418–432.
- Pike, D. A. (2014). Forecasting the viability of sea turtle eggs in a warming world. *Global Change Biology*, 20(1), 7–15.
- Pike, D. A., E. A. Roznik, and I. Bell. (2015). Nest inundation from sea-level rise threatens sea turtle population viability. *Royal Society Open Science*, 2(7), 150127.
- Piniak, W. E. D., D. A. Mann, C. A. Harms, T. T. Jones, and S. A. Eckert. (2016). Hearing in the juvenile green sea turtle (*Chelonia mydas*): A comparison of underwater and aerial hearing using auditory evoked potentials. *PLoS ONE*, 11(10), e0159711.
- Popper, A. N., A. D. Hawkins, R. R. Fay, D. A. Mann, S. M. Bartol, T. J. Carlson, S. Coombs, W. T. Ellison, R. L. Gentry, M. B. Halvorsen, S. Løkkeborg, P. H. Rogers, B. L. Southall, D. G. Zeddies, and W. N. Tavolga. (2014). *ASA S3/SC1.4 TR-2014 Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI*. New York, NY and London, United Kingdom: Acoustical Society of America Press and Springer Briefs in Oceanography.

- Putman, N. F., P. Verley, C. S. Endres, and K. J. Lohmann. (2015). Magnetic navigation behavior and the oceanic ecology of young loggerhead sea turtles. *Journal of Experimental Biology*, 218(7), 1044–1050.
- Reneker, J. L., and S. J. Kamel. (2016). Climate change increases the production of female hatchlings at a northern sea turtle rookery. *Ecology*, 97(12), 3257–3264.
- Rice, M. R., and G. H. Balazs. (2008). Diving behavior of the Hawaiian green turtle (*Chelonia mydas*) during oceanic migrations. *Journal of Experimental Marine Biology and Ecology*, 356(1–2), 121–127.
- Richmond, D. R., J. T. Yelverton, and E. R. Fletcher. (1973). *Far-Field Underwater-Blast Injuries Produced by Small Charges*. Washington, DC: Lovelace Foundation for Medical Education and Research, Defense Nuclear Agency.
- Ridgway, S. H., E. G. Wever, J. G. McCormick, J. Palin, and J. H. Anderson. (1969). Hearing in the giant sea turtle, *Chelonia mydas*. *Proceedings of the National Academy of Sciences U.S.A.*, 64(3), 884–890.
- Rieth, T. M., T. L. Hunt, C. Lipo, and J. M. Wilmshurst. (2011). The 13th century Polynesian colonization of Hawaii Island. *Journal of Archaeological Science*, 28, 2740–2749.
- Russell, D. J., and G. H. Balazs. (2015). Increased use of non-native algae species in the diet of the green turtle (*Chelonia mydas*) in a primary pasture ecosystem in Hawaii. *Aquatic Ecosystem Health & Management*, 18(3), 342–346.
- Sakamoto, W., K. Sato, H. Tanaka, and Y. Naito. (1993). Diving patterns and swimming environment of two loggerhead turtles during internesting. *Nippon Suisan Gakkaishi*, 59(7), 1129–1137.
- Salmon, M., T. T. Jones, and K. W. Horch. (2004). Ontogeny of diving and feeding behavior in juvenile seaturtles: Leatherback seaturtles (*Dermochelys coriacea* L) and green seaturtles (*Chelonia mydas* L) in the Florida current. *Journal of Herpetology*, 38(1), 36–43.
- Schroeder, B. A., A. M. Foley, and D. A. Bagley. (2003). Nesting patterns, reproductive migrations, and adult foraging areas of loggerhead turtles. In A. B. Bolten & B. E. Witherington (Eds.), *Loggerhead Sea Turtles* (pp. 114–124). Washington, DC: Smithsonian Institution Press.
- Schuyler, Q., B. D. Hardesty, C. Wilcox, and K. Townsend. (2014). Global analysis of anthropogenic debris ingestion by sea turtles. *Conservation Biology*, 28(1), 129–139.
- Seminoff, J. A., C. D. Allen, G. H. Balazs, P. H. Dutton, T. Eguchi, H. L. Haas, S. A. Hargrove, M. P. Jensen, D. L. Klemm, A. M. Lauritsen, S. L. MacPherson, P. Opay, E. E. Possardt, S. L. Pultz, E. E. Seney, K. S. Van Houtan, and R. S. Waples. (2015). *Status Review of the Green Turtle (Chelonia mydas) Under the U.S. Endangered Species Act*. (NOAA Technical Memorandum NMFS-SWFSC-592). La Jolla, CA: Southwest Fisheries Science Center.
- Shankar, K., J. Ramadevi, B. C. Choudhary, L. Singh, and R. K. Aggarwal. (2004). Phylogeography of olive ridley turtles (*Lepidochelys olivacea*) on the east coast of India: Implications for conservation theory. *Molecular Ecology*, 13, 1899–1909.
- Smith, S. H., and D. E. Marx, Jr. (2016). De-facto marine protection from a Navy bombing range: Farallon de Medinilla, Mariana Archipelago, 1997 to 2012. *Marine Pollution Bulletin*, 102(1), 187–198.
- Snoddy, J. E., M. Landon, G. Blanvillain, and A. Southwood. (2009). Blood biochemistry of sea turtles captured in gillnets in the lower Cape Fear River, North Carolina, USA. *Journal of Wildlife Management*, 73(8), 1394–1401.

- Southwood, A. L., R. D. Andrews, M. E. Lutcavage, F. V. Paladino, N. H. West, R. H. George, and D. R. Jones. (1999). Heart rates and diving behavior of leatherback sea turtles in the eastern Pacific Ocean. *The Journal of Experimental Biology*, 202, 1115–1125.
- Stabenau, E. K., T. A. Heming, and J. F. Mitchell. (1991). Respiratory, acid-base and ionic status of Kemp's ridley sea turtles (*Lepidochelys kempii*) subjected to trawling. *Comparative Biochemistry and Physiology Part A: Physiology*, 99(1), 107–111.
- Swisdak, M. M., Jr., and P. E. Montanaro. (1992). *Airblast and Fragmentation Hazards Produced by Underwater Explosions*. Silver Spring, MD: Naval Surface Warfare Center.
- Teuten, E. L., S. J. Rowland, T. S. Galloway, and R. C. Thompson. (2007). Potential for plastics to transport hydrophobic contaminants. *Environmental Science and Technology*, 41(22), 7759–7764.
- U.S. Department of the Navy. (2015a). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2015b). *Regional Biosecurity Plan for Micronesia and Hawaii, Volume IV*. Mangilao, GU: University of Guam and the Secretariat of the Pacific Community.
- U.S. Department of the Navy. (2015c). *Regional Biosecurity Plan for Micronesia and Hawaii, Volume III*. Mangilao, GU: University of Guam and the Secretariat of the Pacific Community.
- U.S. Department of the Navy. (2015d). *Regional Biosecurity Plan for Micronesia and Hawaii, Volume I*. Mangilao, GU: University of Guam and the Secretariat of the Pacific Community.
- U.S. Department of the Navy. (2017a). *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)*. San Diego, CA: Space and Naval Warfare System Command, Pacific.
- U.S. Department of the Navy. (2017b). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing* (Technical Report prepared by Space and Naval Warfare Systems Center Pacific). San Diego, CA: Naval Undersea Warfare Center.
- U.S. Department of the Navy. (2018a). *2017 Living Marine Resources (LMR) Program Annual Report*. Port Hueneme, CA: Naval Facilities Engineering.
- U.S. Department of the Navy. (2018b). *2017 U.S. Navy Annual Marine Species Monitoring Report for the Pacific: A Multi-Range-Complex Monitoring Report For Hawaii-Southern California Training and Testing (HSTT), Mariana Islands Training and Testing (MITT), Northwest Training and Testing (NWTT), and the Gulf of Alaska Temporary Maritime Activities Area (GOA TMAA)*. Silver Spring, MD: National Marine Fisheries Service.
- United Nations Environmental Program. (2005). *Marine Litter: An analytical overview*. Nairobi, Kenya: United Nations Environment Programme's Regional Seas Programme.
- Valverde, R. A., D. W. Owens, D. S. MacKenzie, and M. S. Amoss. (1999). Basal and stress-induced corticosterone levels in olive ridley sea turtles (*Lepidochelys olivacea*) in relation to their mass nesting behavior. *Journal of Experimental Zoology*, 284(6), 652–662.
- Viada, S. T., R. M. Hammer, R. Racca, D. Hannay, M. J. Thompson, B. J. Balcom, and N. W. Phillips. (2008). Review of potential impacts to sea turtles from underwater explosive removal of offshore structures. *Environmental Impact Assessment Review*, 28, 267–285.

- Watwood, S. L., J. D. Iafate, E. A. Reyier, and W. E. Redfoot. (2016). Behavioral Response of Reef Fish and Green Sea Turtles to Mid-Frequency Sonar. In A. N. Popper & A. Hawkins (Eds.), *The Effects of Noise on Aquatic Life II* (pp. 1213–1221). New York, NY: Springer.
- Weir, C. R. (2007). Observations of marine turtles in relation to seismic airgun sound off Angola. *Marine Turtle Newsletter*, 116, 17–20.
- Wiley, M. L., J. B. Gaspin, and J. F. Goertner. (1981). Effects of underwater explosions on fish with a dynamical model to predict fishkill. *Ocean Science and Engineering*, 6(2), 223–284.
- Williams, S. L. (1988). *Thalassia testudinum* productivity and grazing by green turtles in a highly disturbed seagrass bed. *Marine Biology*, 98, 447–455.
- Willis, K. L., J. Christensen-Dalsgaard, D. R. Ketten, and C. E. Carr. (2013). Middle ear cavity morphology is consistent with an aquatic origin for testudines. *PLoS ONE*, 8(1), e54086.
- Witt, M. J., L. A. Hawkes, M. H. Godfrey, B. J. Godley, and A. C. Broderick. (2010). Predicting the impacts of climate change on a globally distributed species: The case of the loggerhead turtle. *The Journal of Experimental Biology*, 213(6), 901–911.
- Work, T. M., and G. H. Balazs. (2013). Tumors in sea turtles: The insidious menace of fibropapillomatosis. *The Wildlife Professional*, Fall 2013, 44–47.
- Yelverton, J. T., D. R. Richmond, E. R. Fletcher, and R. K. Jones. (1973). *Safe Distances From Underwater Explosions for Mammals and Birds*. Albuquerque, NM: Lovelace Foundation for Medical Education and Research.
- Yelverton, J. T., and D. R. Richmond. (1981). *Underwater Explosion Damage Risk Criteria for Fish, Birds, and Mammals*. Paper presented at the 102nd Meeting of the Acoustical Society of America. Miami Beach, FL.
- Yudhana, A., J. Din, Sundari, S. Abdullah, and R. B. R. Hassan. (2010). Green turtle hearing identification based on frequency spectral analysis. *Applied Physics Research*, 2(1), 125–134.
- Zellar, R., A. Pulkkinen, K. Moore, D. Reeb, E. Karakoylu, and O. Uritskaya. (2017). *Statistical Assessment of Cetacean Stranding Events in Cape Cod (Massachusetts, USA) Area OS21A-1345*. Greenbelt, MD: National Aeronautics and Space Administration.

This page intentionally left blank.

---

---

## 3.6 Marine Birds



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|         |   |        |
|---------|---|--------|
| 3.6     | Marine Birds.....   | 3.6-1  |
| 3.6.1   | Affected Environment.....   | 3.6-1  |
| 3.6.1.1 | Group Size.....   | 3.6-2  |
| 3.6.1.2 | Diving Behavior.....  | 3.6-2  |
| 3.6.1.3 | Flight Altitudes.....   | 3.6-2  |
| 3.6.1.4 | Distance from Shore.....  | 3.6-3  |
| 3.6.1.5 | Hearing and Vocalization.....   | 3.6-3  |
| 3.6.1.6 | General Threats.....  | 3.6-5  |
| 3.6.1.7 | Rookery Locations and Breeding Activities on FDM.....                           | 3.6-5  |
| 3.6.2   | Environmental Consequences.....   | 3.6-6  |
| 3.6.2.1 | Acoustic Stressors.....   | 3.6-11 |
| 3.6.2.2 | Explosives Stressors (explosive shock wave and sound, explosive fragments)..... | 3.6-19 |
| 3.6.2.3 | Energy Stressors.....   | 3.6-22 |
| 3.6.2.4 | Physical Disturbance and Strike Stressors.....                                  | 3.6-25 |
| 3.6.2.5 | Ingestion Stressors.....  | 3.6-28 |
| 3.6.2.6 | Secondary Stressors (Impacts on Habitat; Impacts on Prey Availability)<br>..... | 3.6-29 |
| 3.6.2.7 | Endangered Species Act Determinations.....                                      | 3.6-30 |
| 3.6.2.8 | Migratory Bird Treaty Act.....  | 3.6-30 |
| 3.6.3   | Public Scoping Comments.....  | 3.6-30 |

## List of Figures

There are no figures in this section.

## List of Tables

|  |       |
|--|-------|
| Table 3.6-1: Number of Events by Activity Type on Farallon de Medinilla.....               | 3.6-8 |
| Table 3.6-2: Number of Munitions by Activity Type on Farallon de Medinilla.....            | 3.6-9 |
| Table 3.6-3: Munitions Use on Farallon de Medinilla, Net Explosive Weight Comparisons..... | 3.6-9 |

### 3.6 Marine Birds

The purpose of this section is to supplement the analysis of impacts on marine birds presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement /Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon de Medinilla (FDM). Information presented in the 2015 MITT Final EIS/OEIS that remains valid is noted as such and referenced in the appropriate sections. Any new or updated information describing the affected environment and analysis of impacts on marine birds associated with the Proposed Action is provided in this section. Comments received from the public during scoping related to marine birds are addressed in Section 3.6.3 (Public Scoping Comments).

#### 3.6.1 Affected Environment

As presented in the 2015 MITT Final EIS/OEIS, the habitat found within the MITT Study Area supports a wide diversity of resident and migratory marine birds, with regionally important rookeries for numerous species on FDM. Descriptions of the climate, productivity, and oceanographic conditions were presented in the 2015 MITT Final EIS/OEIS, as well as important rookery locations throughout the Mariana Islands. Because FDM is the only land area within the Study Area that would be impacted by the proposed changes in activities described in Chapter 2 (Description of Proposed Action and Alternatives), the other rookery locations analyzed in the 2015 MITT Final EIS/OEIS are not included in the Study Area for this Supplemental EIS (SEIS)/OEIS. The species assemblage in open ocean portions of the Study Area has not changed, nor has the status of rookeries on FDM changed since the publication of the 2015 MITT Final EIS/OEIS. As such, the general description in the 2015 MITT Final EIS/OEIS of the existing conditions within the Study Area remains valid.

#### Endangered Species Act Listed Marine Bird Species

Three marine birds present in the Study Area are listed under the Endangered Species Act (ESA) as threatened or endangered species. The short-tailed albatross (*Phoebastria albatrus*) and Hawaiian petrel (*Pterodroma sandwichensis*) are listed as endangered, and the Newell's shearwater (*Puffinus auricularis newelli*)<sup>1</sup> is listed as threatened (U.S. Fish and Wildlife Service, 2010, 2015). None of these species have been observed on FDM or within other rookery locations for other species within the Mariana Islands. The short-tailed albatross, Hawaiian petrel, and Newell's shearwater nest outside the Study Area and are thought to occur only rarely within the Study Area (U.S. Fish and Wildlife Service, 2010, 2015), and there would be little to no overlap with at-sea training and testing activities. The 2015 MITT Final EIS/OEIS relied in part on information collected in 2007 from the Navy-funded Mariana Islands Sea Turtle and Cetacean Survey (U.S. Department of the Navy, 2007). Because the short-tailed albatross, Hawaiian petrel, and Newell's shearwater were not expected to be impacted by activities analyzed in the 2015 MITT Final EIS/OEIS, the Section 7(a)(2) ESA consultation between the Navy and U.S. Fish and Wildlife Service (USFWS) did not include these species (U.S. Fish and Wildlife Service, 2015). No new survey information is available on at-sea observations of marine birds and shorebirds that would change

---

<sup>1</sup> The current taxonomic classification of this species holds that the Newell's shearwater is a subspecies of the Townsend's shearwater. In some instances, this subspecies is also named the Newell's Townsend's shearwater; however, both Newell's shearwater and Newell's Townsend's shearwater refer to the same subspecies (scientific name *Puffinus auricularis newelli*).

the analysis from the 2015 MITT Final EIS/OEIS. As such, the description regarding at-sea observations of marine birds presented in the 2015 MITT Final EIS/OEIS remains valid.

#### **3.6.1.1 Group Size**

Section 3.6.2.1 of the 2015 MITT Final EIS/OEIS included a description of marine bird group sizes and reasons why some marine birds congregate in groups. Within the Study Area, the largest grouping of marine birds is anticipated during large upwelling events for feeding, and on-land rookery locations (at FDM). There is no new information that changes the basis of the conclusion on the group size analysis from the 2015 MITT Final EIS/OEIS. As such, the additional description regarding group sizes of marine birds presented in the 2015 MITT Final EIS/OEIS remains valid.

#### **3.6.1.2 Diving Behavior**

Section 3.6.2.2 of the 2015 MITT Final EIS/OEIS describes dive behaviors exhibited by different types of marine birds. Marine birds will dive to various depths in pursuit of prey items, exhibiting plunge diving. Many of the marine bird species found in the Study Area will dive, skim, or grasp prey at the water's surface or within the upper portion (1–2 meters [m]) of the water column (Cook et al., 2011; Jiménez et al., 2012; Sibley, 2014), although some marine birds will dive to depths greater than 30 m in pursuit of prey, with dive durations lasting from a few seconds to several minutes for deep diving marine birds. Dive durations are correlated with depth and range from a few seconds in shallow divers to several minutes in alcids (Ponganis, 2015). No new information is available on dive behavior that would alter the analysis from the 2015 MITT Final EIS/OEIS. As such, the additional description regarding dive behavior presented in the 2015 MITT Final EIS/OEIS remains valid.

#### **3.6.1.3 Flight Altitudes**

While foraging birds will be present near the water surface, migrating birds may fly at various altitudes. Flight altitudes for birds have traditionally been estimated from on the ground (or boat) observations, or from planes; however, flight altitude information increasingly relies on radar studies and telemetry techniques, where the bird's measured altitude is subtracted from the ground elevation (Poessel et al., 2018). Jongbloed (2016) completed a literature review to determine flight height of marine birds to assess potential risks from wind turbine collisions. This review found that most seabird species fly beneath the rotor blade altitudes of offshore wind turbines, which reduces the risk for collision. Some species such as sea ducks and loons may be commonly seen flying just above the water's surface, but the same species can also be spotted flying high enough (5,800 feet [ft.]) that they are barely visible through binoculars (Lincoln et al., 1998). While there is considerable variation, the favored altitude for most small birds appears to be between 500 ft. (152 m) and 1,000 ft. (305 m). Radar studies have demonstrated that 95 percent of the migratory movements occur at less than 10,000 ft. (3,050 m), with the bulk of the movements occurring under 3,000 ft. (914 m) (Lincoln et al., 1998). Weather factors may also influence flight heights. Tarroux et al. (2016) examined the flying tactics of Antarctic petrels, (*Thalassoica Antarctica*), in Antarctica revealing the flexibility of flight strategies. Birds tend to fly higher with favorable wind conditions, and fly at near ground level during strong winds. Birds were found to adjust their speed and heading during stronger winds to limit drift, however, they were able to tolerate a limited amount of drift (Tarroux et al., 2016). This was also found by Stumpf et al. (2011) for marbled murrelets using radar to quantify flight heights off of the Olympic Peninsula and by Sanzenbacher et al. (2014) off of Northern California. In summary, most marine birds can be expected to fly relatively close to the surface, but may range upwards in altitude depending on a number of factors such as wind speed and direction, precipitation avoidance, time of day or night, foraging behaviors,

migration, and distance to coast. In general the flight altitude of low migrating birds is likely distinctly lower offshore than along the coast or inland of islands within the Mariana Archipelago.

#### **3.6.1.4 Distance from Shore**

Pelagic ranges, as a function of distance from shore, can range widely for different species. Much of the recent research regarding abundance and distribution as a function of distance from shore for marine birds was conducted to better understand potential impacts on marine birds from offshore energy development. Spiegel et al. (2017) tracked the movements of over 400 individuals of three species (northern gannets, red-footed loon, and surf-scooter) over the course of five years off of the mid-Atlantic coast. In winter, all three species exhibited a largely near-shore, coastal, or in-shore distribution. Habitat use was concentrated in or around large bays, with the most extensive use at bay mouths. Northern gannets ranged much farther offshore than the other two species, and covered a much larger area (including instances of individuals using both the Gulf of Mexico and the mid-Atlantic within a single season). Spiegel et al. (2017) determined that the differences among species distributions were likely due to differences in motility and distribution of their preferred prey. In summary, marine bird distance from shore can depend on a variety of factors, such as physiological abilities of a particular species to tolerate long distance and duration flights, mobility of prey, and seasonal variations in ranges.

Pelagic marine birds are widely distributed throughout the Marianas, but they tend to congregate in areas of high productivity and prey availability. The 2015 MITT Final EIS/OEIS relied on information collected in 2007 from the Navy-funded Mariana Islands Sea Turtle and Cetacean Survey (U.S. Department of the Navy, 2007). No new information is available on at-sea observations of marine birds and shorebirds that would change the analysis from the 2015 MITT Final EIS/OEIS. As such, the description regarding at-sea observations of marine birds presented in the 2015 MITT Final EIS/OEIS remains valid.

#### **3.6.1.5 Hearing and Vocalization**

Section 3.6.2.3 of the 2015 MITT Final EIS/OEIS includes a description of marine bird hearing in air, as well as under water. The Navy's literature review of updated information since the publication of the 2015 MITT Final EIS/OEIS has found new information regarding in-air and underwater hearing sensitivities of marine birds.

The Navy conducted a literature search for new information since the publication of the 2015 MITT Final EIS/OEIS on bird hearing and vocalizations that may change the analysis of potential impacts on birds. New information regarding hearing sensitivities of waterbirds, including various duck species and lesser scaups, is summarized below, along with recent publications that show differences in hearing sensitivities between freshwater divers and pelagic birds. This information is summarized below with an overview of the most current best available science regarding bird hearing and vocalization.

Although hearing range and sensitivity has been measured for many land birds, little is known of seabird hearing. The majority of the published literature on bird hearing focuses on terrestrial birds and their ability to hear in air. A review of 32 terrestrial and marine species indicates that birds generally have greatest hearing sensitivity between 1 and 4 kilohertz (kHz) (Beason, 2004; Dooling, 2002). Very few can hear below 20 hertz, most have an upper frequency hearing limit of 10 kHz, and none exhibit hearing at frequencies higher than 15 kHz (Dooling, 2002; Dooling & Popper, 2000). Hearing capabilities have been studied for only a few seabirds (Beason, 2004; Beuter et al., 1986; Crowell et al., 2015; Johansen et al., 2016; Thiessen, 1958; Wever et al., 1969); these studies show that seabird hearing ranges and sensitivity in air are consistent with what is known about bird hearing in general.

Auditory abilities have been measured in 10 diving bird species in-air using electrophysiological techniques (Crowell et al., 2015; Maxwell et al., 2017). All species tested had the best hearing sensitivity from 1 to 3 kHz. The red-throated loon (*Gavia stellata*) and northern gannet (*Morus bassanus*) (both non-duck species) had the highest thresholds while the lesser scaup (*Aythya affinis*) and ruddy duck (*Oxyura jamaicensis*) (both duck species) had the lowest thresholds (Crowell et al., 2015). Auditory sensitivity varied amongst the species tested, spanning over 30 decibels (dB) in the frequency range of best hearing. While electrophysiological techniques provide insight into hearing abilities, auditory sensitivity is more accurately obtained using behavioral techniques. Crowell et al. (2016) used behavioral methods to obtain an in-air audiogram of the lesser scaup. Hearing frequency range in air was similar to other birds, with best sensitivity at 2.86 kHz with a threshold of 14 dB referenced to (re) 20 micropascals ( $\mu\text{Pa}$ ).

Crowell et al. (2015) also compared the vocalizations of the same 10 diving bird species to the region of highest sensitivity of in-air hearing. Of the birds studied, vocalizations of only eight species were obtained due to the relatively silent nature of two of the species. The peak frequency of the vocalizations of seven of the eight species fell within the range of highest sensitivity of in-air hearing. Crowell et al. (2015) suggested that the colonial nesters tested had relatively reduced hearing sensitivity because they relied on individually distinctive vocalizations over short ranges. Additionally, Crowell et al. (2015) observed that the species with more sensitive hearing were those associated with freshwater habitats, which are relatively quieter compared to marine habitats with wind and wave noise.

Although important to seabirds in air, it is unknown if seabirds use hearing or vocalizations underwater for foraging, communication, predator avoidance or navigation (Crowell, 2016; Dooling & Therrien, 2012). Some scientists suggest that birds must rely on vision rather than hearing while underwater (Hetherington, 2008), while others suggest birds must rely on an alternative sense in order to coordinate cooperative foraging and foraging in low light conditions (e.g., night, depth) (Dooling & Therrien, 2012).

There is little known about the hearing abilities of birds underwater (Dooling & Therrien, 2012). In air, the size of the bird is usually correlated with the sensitivity to sound (Johansen et al., 2016); for example, songbirds tend to be more sensitive to higher frequencies and larger non-songbirds tend to be more sensitive to lower frequencies (Dooling & Popper, 2000). Two studies have tested the ability of a single diving bird, a great cormorant (*Phalacrocorax carbo sinensis*), to respond to underwater sounds (Hansen et al., 2017; Johansen et al., 2016). These studies suggest that the cormorant's hearing in air is less sensitive than birds of similar size; however, the hearing capabilities in water are better than what would be expected for a purely in-air adapted ear (Johansen et al., 2016). The frequency range of best hearing underwater was observed to be narrower than the frequency range of best hearing in air, with greatest sensitivity underwater observed around 2 kHz (about 71 dB re 1  $\mu\text{Pa}$  based on behavioral responses). Although results were not sufficient to be used to generate an audiogram, Therrien (2014) also examined underwater hearing sensitivity of long-tailed ducks (*Clangula hyemalis*) by examining behavioral responses. The research showed that auditory thresholds at frequencies within the expected range of best sensitivity (1, 2, and 2.86 kHz) are expected to be between 77 and 127 dB re 1  $\mu\text{Pa}$ .

Diving birds may not hear as well underwater, compared to other (non-avian) species, based on adaptations to protect their ears from pressure changes (Dooling & Therrien, 2012). Because reproduction and communication with conspecifics occurs in air, adaptations for diving may have evolved to protect in-air hearing ability and may contribute to reduced sensitivity underwater (Hetherington, 2008). There are many anatomical adaptations in diving birds that may reduce sensitivity both in air and underwater. Anatomical ear adaptations are not well investigated, but include cavernous

tissue in the meatus and middle ear that may fill with blood during dives to compensate for increased pressure on the tympanum, active muscular control of the meatus to prevent water entering the ear, and interlocking feathers to create a waterproof outer covering (Crowell et al., 2015; Rijke, 1970; Sade et al., 2008). The northern gannet, a plunge diver, has unique adaptations to hitting the water at high speeds, including additional air spaces in the head and neck to cushion the impact and a thicker tympanic membrane than similar-sized birds (Crowell et al., 2015). All of these adaptations could explain why best hearing frequencies are narrower under water than on the surface or in flight.

This new information increases the understanding of bird auditory abilities; however, no new information is available on bird hearing that would alter the analysis from the 2015 MITT Final EIS/OEIS. As such, the additional description regarding dive behavior presented in the 2015 MITT Final EIS/OEIS remains valid.

#### **3.6.1.6 General Threats**

Section 3.6.2.4 (General Threats) of the 2015 MITT Final EIS/OEIS described the general threats facing marine birds within the Study Area. No new information is available that would change the characterization of threats described in the 2015 document; therefore, the description regarding general threats presented in the 2015 MITT Final EIS/OEIS remains valid. Since the publication of the 2015 MITT Final EIS/OEIS, more complete information regarding potential climate change-related impacts on water quality, which in turn may impact prey base and rookery resiliency to storm events, has become available and been included in this SEIS/OEIS. Section 3.1 (Sediments and Water Quality) describes the updated information included in this SEIS/OEIS in regards to potential impacts on water quality from climate change. These changes (e.g., air and sea temperatures, precipitation, frequency and intensity of storms, pH level of sea water, sea level rise) may potentially impact marine birds by reducing overall marine productivity and biodiversity, which could affect the food resources, distribution, and reproductive success of marine birds (Duffy, 2011; Frost et al., 2017; Lorrain et al., 2017; Ostrom et al., 2017; Ramírez et al., 2017; Trainer, 2017). In the long term, climate change could be the largest threat to marine birds.

On FDM, the primary threats to marine bird rookeries include invasive species currently on the island (e.g., rodents that prey on marine bird eggs and chicks). Extensive biosecurity planning by range operators is in place for land-based training activities on FDM to prevent the accidental introduction of other invasive species, such as the brown treesnake (U.S. Fish and Wildlife Service, 2010, 2015). A more detailed description of stressors on the terrestrial environment of FDM is provided in Section 3.10 (Terrestrial Species and Habitats).

#### **3.6.1.7 Rookery Locations and Breeding Activities on FDM**

The 2015 MITT Final EIS/OEIS included a summary of statistical analyses conducted on marine bird counts collected since 1997 and the findings from a non-published technical report produced by the same authors as the published report. Since the publication of the 2015 MITT Final EIS/OEIS, Camp et al. (2016) published this information. During the 159 counts conducted between February 1997 and August 2014, the numbers detected during each count ranged from 0 to 447 for brown booby, 6 to 404 for masked booby, and 42 to 915 for red-footed booby. From 1997 to 2014, there is some evidence that masked and red-footed booby populations on FDM have declined, while brown booby populations have increased. However, the general conclusion is that all three species exhibited population fluctuations over time. Combined with the level of variability observed in the count data, this precluded any definite conclusions about long-term population trends (i.e., the data showed no statistically significant trends)

(Camp et al., 2016). Since the publication of the 2015 MITT Final EIS/OEIS, one additional aerial survey was completed in September 2016. Because of a lack of commercial helicopter transit services, surveys have not been conducted since 2016.

### 3.6.2 Environmental Consequences

In the 2015 MITT Final EIS/OEIS, the Navy considered all potential stressors associated with ongoing training and testing activities in the Mariana Islands and then analyzed their potential impacts on marine birds in the Study Area. In this SEIS/OEIS, the Navy has reviewed the analysis of impacts from these ongoing activities and additionally analyzed new or changing training and testing activities as projected into the reasonably foreseeable future. The Navy has completed a literature review for information on marine birds within the Study Area, which included a search for the best available science since the publication of the 2015 MITT Final EIS/OEIS. Where there has been no substantive or otherwise meaningful change in the action, science, or regulations, the Navy will rely on the previous 2015 MITT Final EIS/OEIS analysis. Where there has been substantive change in the action, science, or regulations, the information and analysis provided in this SEIS/OEIS will supplement the 2015 MITT Final EIS/OEIS to support environmental compliance with applicable environmental statutes for marine birds.

In the alternatives descriptions for this SEIS/OEIS, there have been some modifications to the quantity and type of acoustic stressors under the two action alternatives compared to the 2015 MITT Final EIS/OEIS. In addition, the analysis of potential impacts associated with sonar and other transducers has been improved by incorporating additional information regarding marine bird hearing abilities in water. There have been no updates to the assessment of potential impacts on marine birds from other acoustic sources (e.g., vessel noise, airguns, weapons firing noise, and aircraft noise).

The stressors applicable to birds include the new stressor (high-energy laser) and the same stressors considered in the 2015 MITT Final EIS/OEIS. High-energy laser is detailed in Section 3.0.4.3.2.2 (High-Energy Lasers) and analyzed under the energy stressor category for potential impacts on birds (see Section 3.6.2.3, Energy Stressors).

In general, there have been no substantial changes to the activities analyzed in the 2015 MITT Final EIS/OEIS, which would change the conclusions reached regarding populations of marine birds in the Study Area. Table 2.5.1 and Table 2.5-2 in Chapter 2 (Description of Proposed Action and Alternatives) list the proposed training and testing activities and include the number of times each activity would be conducted annually and the locations within the Study Area where the activity would typically occur under each alternative. The tables also present the same information for activities described in the 2015 MITT Final EIS/OEIS so that the proposed levels of training and testing under this SEIS/OEIS can be compared. The increased use of FDM for training activities proposed in this SEIS/OEIS necessitates FDM-focused analysis for some stressor categories.

Use of acoustic stressors (sonar and other active acoustic sources) and use of explosives have occurred since the 2015 completion of the MITT Final EIS/OEIS Record of Decision. There have been no known adverse effects to marine birds or population impacts that were not otherwise previously analyzed or accounted for in the 2015 MITT Final EIS/OEIS (U.S. Department of the Navy, 2015). The potential stressors associated with the training and testing activities in the Study Area included the following:

- Acoustic stressors (sonar and other transducers, vessel noise, aircraft noise, and weapons noise)
- Explosives (in-air explosions and in-water explosions)
- Energy (in-water and in-air electromagnetic devices, high-energy lasers)

- Physical disturbance and strike stressors (vessels and in-water devices, military expended materials, seafloor devices)
- Ingestion (military expended materials – munitions, military expended materials other than munitions)
- Secondary stressors (impacts on habitat; impacts on prey availability)

During the preparation of the 2015 MITT Final EIS/OEIS, the Navy assessed the potential for wires and cables, along with decelerators/parachutes, to entangle marine birds. The Navy determined at that time that these materials would not present entanglement risks for marine birds because these items would be expended outside of their range of foraging abilities. During the Navy's literature review, no new information regarding fiber optic cables and guidance wires and decelerators/parachutes was found that would alter this conclusion in the 2015 MITT Final EIS/OEIS; therefore, they are not analyzed further in this SEIS/OEIS.

#### **Analysis of Stressors on Farallon de Medinilla**

#### **Analysis of Proposed Increases in Number of Events, Munitions, and Net Explosive Weight on Farallon de Medinilla**

Under Alternative 1 and Alternative 2, there would be an overall increase in the number of training events and munitions used on FDM, which would increase the number of exposures to explosives noise, weapons firing noise, and aircraft overflights to deliver munitions to the impact zones on FDM. The types of explosive munitions used on FDM include explosive bombs (less than or equal to 2,000 pounds [lb.]), missiles, rockets, explosive grenades and mortars, medium-caliber projectiles, and large-caliber projectiles (see Table 3.0-19). The calculations for the increases in the number of events proposed on FDM are shown on Table 3.6-1. Table 3.6-2 shows the calculations for the proposed increases in the number of explosive and non-explosive munitions expended on FDM. These increases in events and munitions will result in an increase in net explosive weight (NEW) of explosives over the course of a training year. The calculations for NEW expended on NEW resulting from proposed training activities are shown on Table 3.6-3. The NEW for each ordnance type may vary within each class. Based on these NEW ranges within each explosives bin, the Navy calculated the range of total munitions' NEW under each alternative proposed in the SEIS/OEIS by multiplying the number of munitions used by the low and high NEW ranges for each ordnance type. Based on these calculations, the following assumptions are presented as additional analysis for the SEIS/OEIS:

- In terms of the number of events, there would be an increase of less than 2 percent over what was analyzed previously in the 2015 MITT Final EIS/OEIS. No new activity types are proposed in the SEIS/OEIS from what were previously analyzed in the 2015 MITT Final EIS/OEIS. Some activity types, however, would increase in the number of events per year and/or the number of ordnance items expended. Other activities would not change compared to what was analyzed previously in the 2015 MITT Final EIS/OEIS, and therefore would not contribute to an increase in NEW or the number of munitions expended on FDM. Examples of these training activities include Gunnery Exercise (Air-to-Ground) and Bombing Exercise (Air-to-Ground). Table 3.6-1 shows the number of events that would occur under each alternative compared to what was analyzed in the 2015 MITT Final EIS/OEIS.
- In terms of munitions item numbers, there would be an increase of approximately 9 percent over what was analyzed previously in the 2015 MITT Final EIS/OEIS in the total number of munitions

used on FDM. Most of these increases are associated with small-caliber rounds, which do not contribute to increases in NEW. Table 3.6-2 shows the number of munitions proposed under each alternative compared to what was analyzed in the 2015 MITT Final EIS/OEIS.

- In terms of NEW, explosives used on FDM would increase by less than 1 percent compared to what was analyzed in the 2015 MITT Final EIS/OEIS (see calculations in Table 3.6-3).

Taken together, the increase in the number of events per year or the amount of ordnance used during events would result in more ordnance use on FDM and an increase in the amount of NEW expended on FDM each year. Although the amount of increased NEW is negligible (less than 1 percent), the number of events per year and the number of ordnance items (most of which are small and medium projectiles) expended increases the potential exposure to stressors associated with ordnance use. Factors that limit the potential for additional adverse impacts, however, include maintaining the same ordnance type and targeting restrictions included as part of the 2015 MITT Final EIS/OEIS. All ordnance expended on FDM would target existing impact zones, with the same ordnance restrictions imposed on all FDM activities and with the same avoidance and minimization measures in place as with the 2015 MITT Final EIS/OEIS (see Section 5.5 Terrestrial Mitigation Measures to be Implemented).

**Table 3.6-1: Number of Events by Activity Type on Farallon de Medinilla**

| Activity   | Number of Events         |               |               | Percent Increase from 2015 MITT Final EIS/OEIS |               |
|--|--------------------------|---------------|---------------|--|---------------|
|  | 2015 MITT Final EIS/OEIS | Alternative 1 | Alternative 2 | Alternative 1                                  | Alternative 2 |
| Marine Air Ground Task Force Exercise (Amphibious) – Battalion | 4                        | 4             | 4             | 0.0%   | 0.0%          |
| Naval Surface Fire Support Exercise – Land-based target        | 10                       | 10            | 15            | 0.0%   | 40.0%         |
| Bombing Exercise (Air-to-Ground)                               | 2,300                    | 2,300         | 2,300         | 0.0%   | 0.0%          |
| Gunnery Exercise (Air-to-Ground)                               | 96                       | 96            | 96            | 0.0%   | 0.0%          |
| Missile Exercise   | 85                       | 115           | 115           | 30.0%  | 30.0%         |
| Direct Action (Tactical Control Party)                         | 18                       | 18            | 18            | 0.0%   | 0.0%          |
| <b>Total</b>   | <b>2,513</b>             | <b>2,543</b>  | <b>2,548</b>  | <b>1.2%</b>                                    | <b>1.4%</b>   |

Notes: EIS = Environmental Impact Statement, MITT = Mariana Islands Training and Testing, NEPM = Non-explosive practice munition, OEIS = Overseas EIS

**Table 3.6-2: Number of Munitions by Activity Type on Farallon de Medinilla**

| Activity  | Munitions Type                | Number of Munitions      |                |                | Percent Increase from 2015 MITT Final EIS/OEIS |               |
|---|-------------------------------|--------------------------|----------------|----------------|--|---------------|
|   |                               | 2015 MITT Final EIS/OEIS | Alternative 1  | Alternative 2  | Alternative 1                                  | Alternative 2 |
| Naval Surface Fire Support Exercise – Land-based target | NEPM Rounds                   | 1,800                    | -              | -              | NA   | NA            |
|   | Explosive large-cal rounds    | 1,000                    | 2,800          | 4,200          | 94.7%  | 123.1%        |
| Bombing Exercise (Air-to-Ground)                        | NEPM Rounds                   | 2,670                    | 2,670          | 2,670          | 0.0%   | 0.0%          |
|   | Explosive Rounds              | 6,242                    | 6,242          | 6,242          | 0.0%   | 0.0%          |
| Gunnery Exercise (Air-to-Ground)                        | Small-cal rounds              | 24,000                   | 24,000         | 24,000         | 0.0%   | 0.0%          |
|   | Med-cal rounds                | 94,150                   | 94,650         | 94,650         | 0.5%   | 0.5%          |
|   | Explosive med-cal rounds      | 17,350                   | 17,500         | 17,500         | 0.9%   | 0.9%          |
|   | Explosive large-cal rounds    | 200                      | 200            | 200            | 0.0%   | 0.0%          |
| Missile Exercise  | Explosive rockets             | 2,000                    | 2,000          | 2,000          | 0.0%   | 0.0%          |
|   | Explosive missiles            | 85                       | 115            | 115            | 30.0%  | 30.0%         |
| Direct Action (Tactical Control Party)                  | Small-cal rounds              | 18,000                   | 30,000         | 30,000         | 50.0%  | 50.0%         |
|   | Medium-cal explosives         | -                        | 1,000          | 1,000          | NA   | NA            |
|   | Explosives (grenades/mortars) | 600                      | 1,000          | 1,000          | 50.0%  | 50.0%         |
| <b>Total</b>  |                               | <b>166,297</b>           | <b>181,177</b> | <b>182,577</b> | <b>8.6%</b>                                    | <b>9.3%</b>   |

Notes: EIS = Environmental Impact Statement, MITT = Mariana Islands Training and Testing, NEPM = Non-explosive practice munition, OEIS = Overseas EIS

**Table 3.6-3: Munitions Use on Farallon de Medinilla, Net Explosive Weight Comparisons**

| Explosive Munitions used at FDM | Bin     | NEW Range <sup>1</sup> | NEW                         |                             |                             | Percent Increase from 2015 MITT Final EIS/OEIS |                     |
|---------------------------------|---------|------------------------|-----------------------------|-----------------------------|-----------------------------|--|---------------------|
|                                 |         |                        | 2015 MITT Final EIS/OEIS    | Alternative 1               | Alternative 2               | Alternative 1                                  | Alternative 2       |
| Explosive Bombs ≤ 2,000 lb.     | E10-E13 | 250-1,740              | 3,121,000-10,861,080        | 3,121,000-10,861,080        | 3,121,000-10,861,080        | 0.0%   | 0.0%                |
| Missiles                        | E6      | 10-20                  | 850-1,700                   | 1,150-2,300                 | 1,150-2,300                 | 30.0%  | 30.0%               |
| Large-caliber Projectiles       | E5      | 5-10                   | 6,000-12,000                | 15,000 – 30,000             | 22,000 – 44,000             | 85.7%  | 114.3%              |
| Medium-caliber Projectiles      | E4      | 2.5-5                  | 43,375-86,750               | 46,250 - 92,500             | 46,250 - 92,500             | 6.4%   | 6.4%                |
| Rockets                         | E4      | 2.5-5                  | 1,000-20,000                | 1,000-20,000                | 1,000-20,000                | 0.0%   | 0.0%                |
| Explosive Grenades and Mortars  | E2      | 0.25-0.5               | 150-300                     | 250-500                     | 250-500                     | 50.0%  | 50.0%               |
| <b>Total</b>                    |         |                        | <b>3,172,375-10,981,830</b> | <b>3,184,650-11,006,380</b> | <b>3,191,650-11,020,380</b> | <b>0.22 – 0.39%</b>                            | <b>0.35 – 0.61%</b> |

<sup>1</sup> NEW (Net Explosive Weight) measured in lb. (pounds)

Notes: EIS = Environmental Impact Statement, FDM = Farallon de Medinilla, lb. = pounds, MITT = Mariana Islands Training and Testing, NEW = Net Explosive Weight (lb.)

### Population-Level Impact Analysis

Under the Migratory Bird Treaty Act (MBTA) regulations applicable to military readiness activities (50 Code of Federal Regulation [CFR] part 21), the stressors introduced during training and testing activities would not result in a significant adverse effect on marine birds protected under the MBTA. While this determination is applicable to all marine birds that occur in the Study Area, the Navy carried out a focused analysis for marine birds known to breed within the Study Area, particularly for breeding marine birds on FDM. The Navy identified two birds in particular that have a heightened concern with regards to 50 CFR Part 21—the great frigatebird and the masked booby.

In the 2015 MITT Final EIS/OEIS, the Navy assessed the significance of injury and mortality of individual masked boobies and great frigatebirds relative to the viability of these species' populations. The populations of the masked booby and great frigatebird were defined based on (1) the distribution of subspecies *S. d. personata* and *F. m. palmerstoni*, (2) the colony locations within these distributions, and (3) the number of individual birds associated with these colonies. The Navy then compared the number of masked boobies and great frigatebirds that are found within the colonies within the Marianas (particularly FDM) to that of the regional population within the western and central Pacific.

Because the numbers of activities described in this SEIS/OEIS potentially affecting these birds and the amount of NEW used on FDM do not appreciably differ from what was analyzed previously, the conclusions within the 2015 MITT Final EIS/OEIS remain valid. These conclusions are summarized below:

- The great frigatebird may occasionally nest on FDM, which is one of only two small breeding colonies known to exist within the Mariana Islands (the other is located on Maug in the northern portion of the archipelago). FDM does not appear to be a temporally or spatially stable rookery location. Compared to the numbers of great frigatebirds estimated throughout central and western Pacific (10,000 pairs in the Hawaiian Islands, with other colonies on Howland, Baker, Jarvis, Johnston Atoll, and Christmas Island), and the apparent low numbers of great frigatebirds from historic times through the present within the Mariana archipelago, the direct and indirect effects of military activities on FDM would not represent a significant adverse impact on the population of the great frigatebird.
- For the masked booby, FDM is the largest breeding colony in Mariana Islands. The colony numbers recorded by the Navy appear to be stable, and the data do not suggest any significant declines of masked booby numbers. Although the masked booby may be subject to short- and long-term impacts of military use of FDM and individuals likely suffer injury and mortality from some activities, FDM continues to support a relatively stable rookery. In the central and western Pacific, 2,500 pairs are estimated within the Northwestern Hawaiian Islands, Jarvis (up to 1,200 pairs), Barker Island (over 1,500 pairs), and smaller colonies in American Samoa, Palmyra, Johnson Atoll, and northern islands in the Mariana archipelago (Maug, Uracas, Guguan, and FDM). Based on the long-term use and stability of the masked booby breeding population on FDM and the wide geographic range and abundance of the masked booby throughout the Pacific, the effects of military use of FDM would not represent a significant adverse impact on the population of the masked booby.
- Pursuant with the Department of Defense's obligations under 50 CFR Part 21, the Department of Defense will continue to implement training restrictions on FDM (see Section 5.5, Terrestrial Mitigation Measures to be Implemented) and monitoring of bird populations on FDM.

### 3.6.2.1 Acoustic Stressors

Section 3.6.3.1 (Acoustic Stressors) in the 2015 MITT Final EIS/OEIS provided an overview of marine bird hearing, including an explanation of how birds can suffer injury, hearing loss, and physiological stress, as well as various behavioral reactions exhibited by birds when a noise event induces a response. In addition, long-term consequences associated with noise-induced impacts are discussed in the 2015 MITT Final EIS/OEIS in Section 3.6.3.1 (Acoustic Stressors).

#### 3.6.2.1.1 Impacts from Sonar and Other Transducer Stressors Under Alternative 1

Under Alternative 1, the number of sonar hours used in the Study Area during training and testing activities compared to the number analyzed in the 2015 MITT Final EIS/OEIS (see Table 3.0-2 and Table 3.0-3 in this SEIS/OEIS) would decrease overall. Therefore, the analysis in the 2015 MITT Final EIS/OEIS remains valid. Decreases in sonar hours shown for activities proposed under Alternative 1 would have no appreciable change on the impact analysis or conclusions for acoustic stressors presented in the 2015 MITT Final EIS/OEIS, based on the analysis below.

Sonar and other transducers emit sound waves into the water to detect objects, safely navigate, and communicate. Use of sonar and other transducers would typically be transient and temporary. General categories of sonar systems are described in Section 3.0.4.1 (Acoustic Stressors).

Information regarding the impacts of sonar on birds is unavailable, and little is known about the ability for birds to hear underwater. The limited information and data from other species suggest the range of best hearing may shift to lower frequencies in water (Dooling & Therrien, 2012; Johansen et al., 2016; Therrien, 2014). Because few birds can hear above 10 kHz in air, it is likely that the only sonar sources they may be able to detect are low- and mid-frequency sources. Other than pursuit diving species, the exposure to birds by these sounds is likely to be negligible because they spend only a very short time underwater (plunge-diving or surface-dipping) or forage only at the water surface. Pursuit divers may remain underwater for minutes, increasing the chance of underwater sound exposure.

In addition to diving behavior, the likelihood of a bird being exposed to underwater sound depends on factors such as duty cycle (defined as the percentage of the time during which a sound is generated over a total operational period), whether the source is moving or stationary, and other activities that might be occurring in the area. When used, continuously active sonars transmit more frequently (greater than 80 percent duty cycle) than traditional sonars, but at a substantially lower source level. However, it should be noted that active sonar is rarely used continuously throughout the listed activities, and many sources are mobile. For moving sources such as hull-mounted sonar, the likelihood of an individual bird being repeatedly exposed to a sound source over a short period of time is low because the training and testing activities are transient, and sonar use and bird diving are intermittent. The potential for birds to be exposed to intense sound associated with stationary sonar sources would likely be limited for some training and testing activities because other activities occurring in conjunction may cause them to leave the immediate area. For example, birds would likely react to helicopter noise during dipping sonar exercises by flushing from the immediate area.

Injury due to acoustic resonance of air space in the lungs due to sonar and other transducers is unlikely in birds. Unlike mammals, birds have compact, rigid lungs with strong pulmonary capillaries that do not change much in diameter when exposed to extreme pressure changes (Baerwald et al., 2008), leading to resonant frequencies lower than the frequencies used for Navy sources.

A physiological impact, such as hearing loss, would likely only occur if a marine bird were close to an intense sound source. Hearing loss is typically quantified in terms of threshold shift—the amount (in dB) that hearing thresholds at one or more specified frequencies are elevated, compared to their pre-exposure values, at some specific time after the noise exposure. The amount of threshold shift measured usually decreases with increasing recovery time—the amount of time that has elapsed since a noise exposure. If the threshold shift eventually returns to zero (i.e., the hearing threshold returns to the pre-exposure value), the threshold shift is called a temporary threshold shift (TTS). If the threshold shift does not completely recover (the threshold remains elevated compared to the pre-exposure value), the remaining threshold shift is called a permanent threshold shift (PTS). By definition, TTS is a function of the recovery time, therefore comparing the severity of noise exposures based on the amount of induced TTS can only be done if the recovery times are also taken into account. For example, a 20 dB TTS measured 24 hours post-exposure indicates a more hazardous exposure than one producing 20 dB of TTS measured only two minutes after exposure; if the TTS is 20 dB after 24 hours, the TTS measured after two minutes would have likely been much higher. Conversely, if 20 dB of TTS was measured after two minutes, the TTS measured after 24 hours would likely have been much smaller.

In general, birds are less susceptible to both TTS and PTS than mammals (Saunders & Dooling, 1974). Diving birds have adaptations to protect the middle ear and tympanum from pressure changes during diving that may affect hearing (Dooling & Therrien, 2012). While some adaptations may exist to aid in underwater hearing, other adaptations to protect in-air hearing may limit aspects of underwater hearing (Hetherington, 2008). Because of these reasons, the likelihood of a diving bird experiencing an underwater exposure to sonar or other transducer that could result in an impact on hearing is considered low.

Because there is no new information since the 2015 MITT Final EIS/OEIS that would change the previous analysis for potential impacts on ESA-listed marine bird species, the conclusions in the 2015 MITT Final EIS/OEIS remain valid. The described training and testing activities would present no measurable chance for interaction with ESA-listed marine bird species (e.g., short-tailed albatross, Hawaiian petrel, Newell's shearwater). In the 2015 MITT Final EIS/OEIS and during consultation between the Navy and USFWS, the Navy determined that the use of sonar and other transducers would have no effect on ESA-listed marine birds.

Because of the small numbers of birds potentially exposed to stressors associated with sonar and other transducers, and the low potential of any injurious exposure to sonar and other transducers while birds are under water, marine bird population impacts would not occur.

*Pursuant to the ESA, acoustic stressors from the use of sonar and other transducers during training and testing activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters. This determination is consistent with the previous consultation between the Navy and USFWS for activities described in the 2015 MITT Final EIS/OEIS.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from sonar and other transducers during training and testing activities described under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

### 3.6.2.1.2 Impacts from Sonar and Other Transducer Stressors under Alternative 2

As with Alternative 1, the number of sonar hours used under Alternative 2 in the Study Area during training and testing activities compared to the number analyzed in the 2015 MITT Final EIS/OEIS (Table 3.0-2 and Table 3.0-3) would decrease overall. Therefore, the analysis in the 2015 MITT Final EIS/OEIS remains valid. Decreases in the number of training and testing activities would potentially decrease the level of acoustic stressors in the Study Area. The conclusions for ESA-listed species presented in the 2015 MITT Final EIS/OEIS is the same as for Alternative 1 in this SEIS/OEIS.

As with Alternative 1, taken together, the small numbers of birds potentially exposed to stressors associated with sonar and other transducers under Alternative 2, and the low potential of any injurious exposure to sonar and other transducers while birds are under water, there would be no impacts on marine bird populations.

*Pursuant to the ESA, acoustic stressors from the use of sonar and other transducers during training and testing activities, as described under Alternative 2, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters. This determination is consistent with the previous consultation between the Navy and USFWS for activities described in the 2015 MITT Final EIS/OEIS.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from sonar and other transducers during training and testing activities described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

### 3.6.2.1.2.1 Impacts from Sonar and Other Transducer Stressors under the No Action Alternative

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on marine birds from sonar and other transducers, but would not measurably improve the overall distribution or abundance of marine birds.

### 3.6.2.1.3 Impacts from Aircraft Noise

Section 3.6.3.1.3.1 (Fixed-Wing Aircraft) and Section 3.6.1.3.2 (Helicopters) of the 2015 MITT Final EIS/OEIS discuss the different types of aircraft and the noise they generate, along with a summary of potential responses marine birds may exhibit. Since the publication of the 2015 MITT Final EIS/OEIS, no new information was identified during the Navy's literature review that would substantially alter the assessment of potential impacts on marine birds from aircraft noise. Therefore, the information contained in Section 3.6.3.1.3.3 (Vessels) of the 2015 MITT Final EIS/OEIS remains valid.

Birds in areas that may experience repeated exposure often habituate and do not respond behaviorally (Larkin et al., 1996; National Park Service, 1994; Plumpton, 2006). Throughout the Study Area, repeated exposure of individual birds or groups of birds is unlikely based on the dispersed nature of the overflights and the capability of birds to avoid or rapidly vacate an area of disturbance. Therefore, the

general health of individual birds would not be compromised. Occasional startle or alert reactions to aircraft noise are not likely to disrupt major behavior patterns (such as migrating, breeding, feeding, and sheltering) or to result in serious injury to any birds.

Training and testing activities where aircraft are used typically occur further offshore; however, increased use of FDM may increase the potential for aircraft strike of birds. Therefore, for the purposes of this SEIS/OEIS, only the use of aircraft related to FDM training activities are discussed below under the alternatives analysis for birds for this stressor category.

#### **3.6.2.1.3.1 Impacts from Aircraft Noise Stressors Under Alternative 1**

Under Alternative 1, the number of proposed activities including aircraft would decrease overall throughout the study area (see Table 3.0-11). In the open ocean, marine birds would be exposed to additional aircraft noise sources, but these activities are spread out throughout the Study Area. Because of the increase in munitions use at FDM, however, aircraft overflights over FDM would increase, depending on the delivery platform. For example, some of the increases are associated with ship to surface, while others may involve helicopters and fixed-wing aircraft. Therefore, the analysis in this section focuses on FDM, where actual aircraft overflights would likely increase.

Increased training activities under Alternative 1 would increase the potential for noise exposures for birds on FDM because the increase in the number of training activities would require more aircraft to fly over the island (potentially at low altitude) and land on the island to deliver and pick up personnel. As shown in Table 2.5-1, activities that would increase aircraft overflights include Missile Exercise and Direct Action (Tactical Air Control Party) activities.

Aircraft overflights are expected to elicit short-term behavioral responses in nesting birds at FDM. Based on studies from other nesting bird areas (Barnas et al., 2018; Bowles, 1995; Larkin et al., 1996), any period away from the nest would last a few seconds to a few minutes, which is likely not long enough for opportunistic predation of a nest (e.g., by rats on FDM). The 2015 MITT Final EIS/OEIS analyzed other adverse effects, such as damage to eggs and startling of juveniles and adults.

Anecdotally, some birds typically take flight while roosting or nesting during quarterly helicopter-based marine bird surveys over FDM; birds that are stationary and not on the wing are counted (U.S. Department of the Navy, 2013a). Although no studies are available specific to marine bird responses to low-level overflights over FDM, other studies of shorebird responses to military aircraft overflights are helpful. Black (2005), studied the effects of low-altitude (less than 500 ft. [152 m] above ground level) military training flights with sound levels from 55 to 100 A-weighted decibels on wading bird colonies (i.e., great egret, snowy egret, tricolored heron, and little blue heron). The training flights involved three or four aircraft and occurred once or twice per day. This study concluded that the reproductive activity—including nest success, nestling survival, and nestling chronology—was independent of F-16 overflights. Dependent variables were more strongly related to ecological factors, including location and physical characteristics of the colony and climatology. Another study on the effects of circling fixed-wing aircraft and helicopter overflights on wading bird colonies found that at altitudes of 195 ft. (59 m) to 390 ft. (119 m), there was no reaction in nearly 75 percent of the 220 observations. Ninety percent displayed no reaction or merely looked toward the direction of the noise source. Another 6 percent stood up, 3 percent walked from the nest, and 2 percent flushed (but were without active nests) and returned within five minutes (Kushlan, 1978).

These studies, coupled with anecdotal observations on FDM during quarterly marine bird monitoring surveys, suggest that aircraft overflights do not have harmful effects on nesting and roosting marine birds on FDM, and that the behavioral responses are short term (Camp et al., 2016).

Although some degree of disturbance is expected from the increase in aircraft noise over FDM, the island will likely continue to serve as an important rookery for regional species without long-term significant impacts on marine bird populations. As discussed in Section 3.6.1.7 (Rookery Locations and Breeding Activities on FDM), Camp et al. (2016) published results of multi-year population monitoring of three species of boobies on FDM, showing that is some evidence that masked and red-footed booby populations on FDM have declined, while brown booby populations have increased. However, the general conclusion is that all three species exhibited population fluctuations over time. Combined with the level of variability observed in the count data, this precluded any definite conclusions about long-term population trends (i.e., the data were non-significant) (Camp et al., 2016).

Because of the dispersed nature of overflights in open ocean training areas, birds or groups of birds in pelagic environments would not likely be exposed to repeated overflights. Because any exposures would be infrequent, and these exposures would not cause injury, population impacts would not occur for bird species in the open ocean.

Aircraft activity described in this SEIS/OEIS would present no measurable chance for interaction with ESA-listed marine bird species (e.g., short-tailed albatross, Hawaiian petrel, Newell's shearwater). In the 2015 MITT Final EIS/OEIS and during consultation between the Navy and USFWS, the Navy determined that aircraft activity would have no effect on ESA-listed marine birds. Although the amount of training and testing activities using aircraft would increase compared to the 2015 MITT Final EIS/OEIS, the potential for geographic and temporal overlap would remain negligible; therefore, the conclusions for ESA-listed species presented in the 2015 MITT Final EIS/OEIS is the same as for Alternative 1 in this SEIS/OEIS.

*Pursuant to the ESA, aircraft noise during training activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from aircraft noise during training activities described under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.1.3.2 Impacts from Aircraft Noise Stressors Under Alternative 2**

Under Alternative 2, the number of activities including aircraft would decrease compared to levels analyzed under the 2015 MITT Final EIS/OEIS, but would be more than proposed under Alternative 1 (see Table 3.0-11).

As with Alternative 1, the dispersed nature of overflights in open ocean training areas under Alternative 2 would not likely expose birds or groups of birds in pelagic environments to repeated overflights. Because any exposures would be infrequent, and these exposures would not cause injury, population impacts would not occur for bird species in the open ocean.

No additional targets would be used, and this activity would be constrained by confining targeting to specific sites within designated impact zones. Because the same locations would be used for targeting activities, the impacts of Alternative 2 are the same as Alternative 1. In the open ocean, marine birds would be exposed to additional aircraft noise sources, but these activities are spread out throughout the Study Area, and the potential impacts of at-sea training and testing activities would not be discernable

from Alternative 1. Therefore, impacts on marine birds under Alternative 2 from aircraft noise would be negligible.

*Pursuant to the ESA, aircraft noise during training activities, as described under Alternative 2, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from aircraft noise during training activities described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.1.3.3 Impacts from Aircraft Noise Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on marine birds from aircraft noise, but would not measurably improve the overall distribution or abundance of marine birds.

#### **3.6.2.1.4 Impacts from Weapons Noise**

Sounds produced by weapons firing (muzzle blast), launch boosters, and projectile travel are potential stressors to birds and are discussed as impulsive noise under Section 3.6.3.1.2.2 (Explosions on Land and In-Air) in the 2015 MITT Final EIS/OEIS. Since the publication of the 2015 MITT Final EIS/OEIS, no new information was identified during the Navy's literature review that would substantially alter the assessment of potential impacts on marine birds from weapons noise.

##### **3.6.2.1.4.1 Impacts from Weapons Noise Stressors Under Alternative 1**

Under Alternative 1, the number of training and testing activities that would expose marine birds to weapons noise would decrease throughout the Study Area, compared to levels analyzed in the 2015 MITT Final EIS/OEIS (Table 3.0-14 and Table 3.0-16). A bird in the open ocean could be exposed to weapons noise if not already displaced by the visual or noise disturbance of a vessel supporting weapons-firing exercises. The firing of a weapon may have several components of associated noise. Firing of guns could include sound generated in air by firing a gun (muzzle blast) and a crack sound due to a low-amplitude shock wave generated by a supersonic projectile flying through the air. Most in-air sound would be reflected at the air-water interface. Underwater sounds would be strongest just below the surface and directly under the firing point. Any sound that enters the water only does so within a narrow cone below the firing point or path of the projectile. Vibration from the blast propagating through a ship's hull, the sound generated by the impact of an object with the water surface, and the sound generated by launching an object underwater are other sources of impulsive sound in the water.

Supersonic projectiles, which would be similar in size to shells fired from 5-inch/54 guns, would travel at approximately 2,600 ft./second, creating a bow shock wave. Pater and Shea (1981) measured the characteristics of a bow shock wave from a 5 inch projectile and found that the shock wave ranged from 40 to 147 dB re 20  $\mu$ Pa sound pressure level peak taken at the ground surface at 1,100 m from the firing

location and 190 m perpendicular from the trajectory (for safety reasons). Shells fired from a kinetic energy weapon are considered hypersonic, and would travel at about 6,500 ft./second, and peak pressures would be expected to be several dB higher than for shell velocities described by (Pater & Shea, 1981). By definition, bow shock waves, regardless of shell velocity, would travel at the speed of sound in air. Marine birds would be exposed to this type of noise for a very brief period of time (a few seconds), and would likely cause brief and temporary behavioral reactions described previously for other in-air noise disturbances.

Sound due to missile and target launches is typically at a maximum at initiation of the booster rocket and rapidly fades as the missile or target travels downrange. Birds foraging or migrating through a training area in the open ocean may respond by avoiding areas where weapons-firing exercises occur. Exposures of most marine birds would be infrequent, based on the brief duration and dispersed nature of the vessels, and the brief duration of the weapons-firing noise. If a bird responds to weapons noise, only short-term behavioral responses such as startle responses, head turning, or avoidance responses would be expected. Weapons noise near rookery locations (only at FDM) may induce startle responses, inducing birds to temporarily leave nests. Because impacts on individual birds, if any, are expected to be minor and limited, no long-term consequences to individuals are expected. Accordingly, there would be no consequences to any bird populations, and weapons noise would not have a significant adverse effect on populations of migratory bird species.

On FDM, however, marine birds would likely be exposed to increased weapons noise because of the increase in the number of explosive and non-explosive munitions (see Table 3.6-1, Table 3.6-2, and Table 3.6-3). Sources of weapons generating noise at the time of weapons firing on or near FDM include small-caliber rounds, rockets, medium-caliber projectiles, and large-caliber projectiles. Other munitions types used on FDM (e.g., non-explosive practice munition and explosive bombs and missiles) are launched far from the target (impact areas on FDM) or released from aircraft.

The potential impacts of explosives noise and weapons firing noise on FDM's wildlife are discussed in Section 3.10.3.1.1 (Impacts from Explosives and Weapons Firing Noise) in Section 3.10 (Terrestrial Species and Habitats) of this SEIS/OEIS and provides a summary of the different types of sounds, frequency ranges, and intensity of sounds generated from munitions use on FDM. Sources of noise from weapons firing that may be heard by marine birds on FDM include close-in weapons firing from vessels, helicopters, close-combat surface firing from fixed-wing aircraft, and surface firing, with the largest increase in munitions use resulting from small arms, medium-caliber explosives, and mortar and grenade use during Direct Action training activities. As shown in Table 3.6-1, the number of training events for this activity type would stay the same compared to what was previously analyzed in the 2015 MITT Final EIS/OEIS; however, the number of munitions used would increase during each training event (see Table 3.6-2). These training events would occur within the Northern Special Use Area and fire into the impact areas towards the south; therefore, more birds would be exposed to more weapons firing noise under Alternative 1 because of the increased number of small-caliber rounds, medium-caliber explosives, and grenades and mortars fired into impact areas from the Northern Special Use Area. The weapons-firing noise would likely be masked somewhat by natural sounds on FDM, such as waves and winds. The impulsive sound caused by weapon firings would have limited potential to mask any important biological sound simply because the duration of the impulse is brief, even when multiple shots are fired in series.

Although some degree of disturbance is expected from the increase in weapons noise on FDM, the island will likely continue to serve as an important rookery for regional species without long-term

significant impacts on marine bird populations. As discussed in Section 3.6.1.7 (Rookery Locations and Breeding Activities on FDM), Camp et al. (2016) published results of multi-year population monitoring of three species of boobies on FDM, showing that is some evidence that masked and red-footed booby populations on FDM have declined, while brown booby populations have increased. However, the general conclusion is that all three species exhibited population fluctuations over time. Combined with the level of variability observed in the count data, this precluded any definite conclusions about long-term population trends (i.e., the data were non-significant) (Camp et al., 2016).

Weapons noise would present no measurable chance for interaction with ESA-listed marine bird species (short-tailed albatross, Hawaiian petrel, Newell's shearwater). As discussed previously, ESA-listed marine bird species do not occur on FDM (or any other island within the Marianas) and have little to no overlap with the Study Area. In the 2015 MITT Final EIS/OEIS and during consultation between the Navy and USFWS, the Navy determined that weapons noise would have no effect on ESA-listed marine birds. Although the amount of training and testing activities would increase compared to the 2015 MITT Final EIS/OEIS, the potential for geographic and temporal overlap would remain negligible; therefore, the conclusions for ESA-listed species presented in the 2015 MITT Final EIS/OEIS is the same as for Alternative 1 in this SEIS/OEIS.

*Pursuant to the ESA, weapons noise during training and testing activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from weapons noise during training and testing activities described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.1.4.2 Impacts from Weapons Noise Stressors Under Alternative 2**

As with Alternative 1, under Alternative 2 the number of training and testing activities that would expose marine birds to weapons noise would decrease throughout the Study Area, compared to levels analyzed in the 2015 MITT Final EIS/OEIS (see Table 3.0-14 and Table 3.0-16). Compared to Alternative 1, there would be small increases in the number of activities using large-caliber and medium-caliber projectiles and missiles under Alternative 2 for at-sea training and testing activities. Therefore, Alternative 2 would introduce fewer weapons firing events than activities analyzed in the 2015 MITT Final EIS/OEIS for at-sea activities. Because activities would occur within the same locations as with Alternative 1, at-sea weapons firing activities would be widely dispersed, and marine birds would also be widely dispersed, the impacts of Alternative 2 are the same as for Alternative 1.

On FDM, the only training activity that would introduce weapons firing noise is Direct Action (tactical control party). As shown in Table 3.6-1, the number of training events for this activity type would stay the same compared to what was previously analyzed in the 2015 MITT Final EIS/OEIS and compared to Alternative 1; however, the number of munitions used would increase (see Table 3.6-2). These training events would occur within the Northern Special Use Area and fire into the impact areas towards the south; therefore, more birds would be exposed to more weapons firing noise under Alternative 2 because of the increased number of small-caliber rounds, medium-caliber explosives, and grenades and mortars fired into impact areas from the Northern Special Use Area. The weapons-firing noise would likely be masked somewhat by natural sounds on FDM, such as waves and winds. The impulsive sound

caused by weapon firings would have limited potential to mask any important biological sound simply because the duration of the impulse is brief, even when multiple shots are fired in series.

As with Alternative 1, some degree of disturbance is expected from the increase in weapons noise on FDM; however, the island will likely continue to serve as an important rookery for regional species without long-term significant impacts on marine bird populations. As discussed in Section 3.6.1.7 (Rookery Locations and Breeding Activities on FDM), Camp et al. (2016) published results of multi-year population monitoring of three species of boobies on FDM, showing that is some evidence that masked and red-footed booby populations on FDM have declined, while brown booby populations have increased. However, the general conclusion is that all three species exhibited population fluctuations over time. Combined with the level of variability observed in the count data, this precluded any definite conclusions about long-term population trends (i.e., the data were non-significant) (Camp et al., 2016). The same conclusions for Alternative 1 for MBTA-protected marine bird species at sea and on FDM, and ESA-listed marine bird species at sea, are applicable to Alternative 2.

*Pursuant to the ESA, weapons noise during training and testing activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, and Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from weapons noise during training and testing activities described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.1.4.3 Impacts from Weapons Noise Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Acoustic stressors would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer acoustic stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on marine birds from weapons noise, but would not measurably improve the overall distribution or abundance of marine birds.

#### **3.6.2.2 Explosives Stressors (explosive shock wave and sound, explosive fragments)**

Section 3.6.3.1.2 (Impacts from Explosives and Swimmer Defense Airguns) in the 2015 MITT Final EIS/OEIS discusses the sources and potential impacts of explosives noise on marine birds (e.g., injury, hearing loss, physiological stress, masking, and long-term consequences of exposures). Explosions in the water, near the water surface, on land (FDM), and in the air can introduce loud, impulsive, broadband sounds into the marine environment. However, unlike other acoustic stressors, explosives release energy at a high rate, producing a shock wave that can be injurious and even deadly. The information regarding training and testing activities in open ocean training environments that generate explosives noise has not changed since the publication of the 2015 MITT Final EIS/OEIS. Therefore, this section focuses on the potential for increased training activities to impact birds on FDM proposed under the alternatives described in this SEIS/OEIS.

Noise can result from direct munitions impacts (one object striking another), blasts (explosions that result in shock waves), bow shock waves (pressure waves from projectiles flying through the air), and substrate vibrations (combinations of explosion, recoil, or vehicle motion with the ground). Noise may be continuous (i.e., lasting for a long time without interruption) or impulse (i.e., short duration). Continuous impulses (e.g., helicopter rotor noise, bursts from rapid-fire weapons) represent an intermediate type of sound and, when repeated rapidly, may resemble continuous noise. These types of sound are distinguished here as they differ in their effects. Continuous sounds can result in hearing damage, while impulses typically elicit physiological or behavioral responses. Some birds may be killed or injured during these activities, or expend energy stores needed for migration to avoid perturbations generated by explosions.

Because the military will continue to implement mitigation measures designed to avoid or reduce impacts on terrestrial biological resources, all additional ordnance use would still be targeted at existing impact areas.

FDM has three impact areas, a special use area on the northern portion of the island, and a special use area on the land bridge. Targeting of areas inside of the special use areas and other areas outside of impact areas are prohibited. In other words, all areas outside of the impact areas are considered “no-fire areas.” Any ordnance that inadvertently lands outside of impact areas, including special use areas and in water, must be reported to Mariana Islands Range Complex Operations, in accordance with Commander, U.S. Naval Forces Marianas Instruction 3500.4A (U.S. Department of the Navy, 2011). The impact areas and special use areas are described below:

- **Northern Special Use Area.** Reserved for direct action (tactical air control party) type exercises and personnel recovery. This area is about 41 acres (ac.) (17 hectares [ha]) and includes a landing zone. Weapons may be fired from the special use area into impact areas, such as small-caliber rounds, grenades, and mortars.
- **Impact Area 1.** This area contains high-fidelity target structures and is comprised of vehicle shells and cargo containers. This area is authorized for inert ordnance only, and operators are required to report any live ordnance inadvertently dropped into Impact Area 1 to Mariana Islands Range Complex Operations. Impact Area 1 contains 10 targets of varying shapes and sizes, including four vehicles and six targets comprised of shipping containers.
- **Impact Area 2.** Impact Area 2 may be used for both live and inert ordnance. Strafing is permitted in this area. Impact Area 2 is about 22 ac. (9 ha).
- **Land Bridge.** The land bridge is designated as a “no target zone.” Operators are required to report ordnance observed impacting the land bridge.

**Impact Area 3.** This area is south of the land bridge and authorized for inert ordnance, although live ordnance may be used only with prior approval from Joint Region Marianas. Strafing is permitted in this area. Impact Area 3 is about 11 ac. (4.5 ha).

#### 3.6.2.2.1 Impacts from Explosive Stressors Under Alternative 1

Under Alternative 1, there would be an overall decrease throughout the Study Area in the number of explosive munitions used during training and testing activities compared to the number analyzed in the 2015 MITT Final EIS/OEIS (Table 3.0-16).

As shown in Table 3.6-1, there would be an increase in the number of events using FDM as a training location or target, with an increase in the number of munitions items expended on FDM (see Table 3.6-2).

Taken together, the increase in the number of training events per year or the amount of ordnance used during training events would result in an increase in the amount of NEW expended on FDM each year (see Table 3.6-1, Table 3.6-2, and Table 3.6-3). Although the amount of increased NEW is negligible, the potential exposure to stressors associated with ordnance use would increase under Alternative 1 compared to what was analyzed previously in the MITT Final EIS/OEIS. Factors that limit the potential for additional adverse impacts, however, include maintaining the same ordnance type and targeting restrictions included as part of the 2015 MITT Final EIS/OEIS. All ordnance expended on FDM would target existing impact zones, with the same ordnance restrictions imposed on all FDM activities and with the same avoidance and minimization measures in place (see Section 5.5, Terrestrial Mitigation Measures to be Implemented, and Table 5.5-1). Therefore, the increases in ordnance use on FDM do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The conclusions for ESA-listed marine bird species and other marine bird species protected by the MBTA included in the 2015 MITT Final EIS/OEIS remain valid.

Explosives would present no measurable chance for interaction with ESA-listed marine bird species (short-tailed albatross, Hawaiian petrel, Newell's shearwater). In the 2015 MITT Final EIS/OEIS and during consultation between the Navy and USFWS, the Navy determined that training and testing activities using explosives would have no effect on ESA-listed marine birds. Despite the continued at-sea use of explosives compared to the 2015 MITT Final EIS/OEIS, the potential for geographic and temporal overlap would remain negligible; therefore, the conclusions for ESA-listed species presented in the 2015 MITT Final EIS/OEIS is the same as for Alternative 1 in this SEIS/OEIS.

*Pursuant to the ESA, explosives used during training and testing activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from explosives stressors during training and testing activities using explosives described under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.2.2 Impacts from Explosive Stressors Under Alternative 2**

As with Alternative 1, there would be an overall decrease throughout the Study Area in the number of explosive munitions used during at-sea training and testing activities compared to the number analyzed in the 2015 MITT Final EIS/OEIS. The number of explosive stressors under Alternative 2 would increase slightly as compared to Alternative 1 (see Table 3.0-16), but the conclusions for at-sea activities remains the same. Under Alternative 2, there would be an increase in the number of training events using FDM as a training location or target (see Table 3.6-1), with an increase in the number of munitions items expended on FDM (see Table 3.6-2) compared to what was analyzed previously in the MITT Final EIS/OEIS and under Alternative 1.

Taken together, the increase in the number of training events per year or the amount of ordnance used during events would result in an increase in the amount of NEW expended on FDM each year (see Table 3.6-3). Although the amount of increased NEW is negligible, the potential exposure to stressors

associated with ordnance use would increase under Alternative 2 compared to what was analyzed previously in the MITT Final EIS/OEIS. Under Alternative 2, Naval Surface Firing Exercise events would expend more large-caliber projectiles, thereby slightly increasing the NEW expended under Alternative 2 compared to Alternative 1. Factors that limit the potential for additional adverse impacts, however, include maintaining the same ordnance type and targeting restrictions included as part of the 2015 MITT Final EIS/OEIS. All ordnance expended on FDM would target existing impact zones, with the same ordnance restrictions imposed on all FDM activities and with the same avoidance and minimization measures in place (see Section 5.5, Terrestrial Mitigation Measures to be Implemented, and Table 5.5-1). Therefore, the increases in ordnance use on FDM shown in Tables 2.5-1 and 2.5-2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The conclusions for ESA-listed marine bird species and other marine bird species protected by the MBTA included in the 2015 MITT Final EIS/OEIS remain valid.

Explosives would present no measurable chance for interaction with ESA-listed marine bird species (short-tailed albatross, Hawaiian petrel, Newell's shearwater). In the 2015 MITT Final EIS/OEIS and during consultation between the Navy and USFWS, the Navy determined that training and testing activities using explosives would have no effect on ESA-listed marine birds. Although the amount of training and testing activities using explosives would increase compared to the 2015 MITT Final EIS/OEIS, the potential for geographic and temporal overlap would remain negligible; therefore, the conclusions for ESA-listed species presented in the 2015 MITT Final EIS/OEIS is the same as for Alternative 1 in this SEIS/OEIS.

*Pursuant to the ESA, explosives used during training and testing activities, as described under Alternative 2, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from explosives stressors during training and testing activities using explosives described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

### **3.6.2.2.3 Impacts from Explosive Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosive stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer explosive stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on marine birds from explosive stressors, but would not measurably improve the overall distribution or abundance of marine birds.

### **3.6.2.3 Energy Stressors**

The energy stressors that may impact marine birds include (1) in-air electromagnetic devices and (2) high-energy lasers. However, as discussed in Section 3.0.4.3 (Energy Stressors), in-air electromagnetic energy would be widely dispersed throughout the Study Area, but more concentrated

in portions of the Study Area near ports, naval installations, and range complexes. Because these stressors are operated at power levels, altitudes, and distances from people and animals to ensure that energy received is well below levels that could disrupt behavior or cause injury and because most in-air electromagnetic energy is reflected by water, in-air electromagnetic energy would not impact birds and is not analyzed further in this section.

Since the publication of the 2015 MITT Final EIS/OEIS, additional information has become available that improves understanding of how in-air electromagnetic devices (such as radar) may impact marine birds. This new information is included in this SEIS/OEIS. Studies conducted on in-air electromagnetic sensitivity in birds have typically been associated with land, and little information exists specifically on seabird response to in-air electromagnetic changes at sea. Based on these studies, in-air electromagnetic effects can be categorized as thermal (i.e., capable of causing damage by heating tissue) or non-thermal. Thermal effects are most likely to occur when near high-power systems. Should such effects occur, they would likely cause birds to temporarily avoid the area receiving the electromagnetic radiation until the stressor ceases (Manville, 2016). Currently, questions exist about far-field, non-thermal effects from low-power, in-air electromagnetic devices. Manville (2016) performed a literature review of this topic. Although findings are not always consistent, Manville (2016) reported that several peer-reviewed studies have shown non-thermal effects can include (1) affecting behavior by preventing birds from using their magnetic compass, which may in turn affect migration; (2) fragmenting the DNA of reproductive cells, decreasing the reproductive capacity of living organisms; (3) increasing the permeability of the blood-brain barrier; (4) other behavioral effects; (5) other molecular, cellular, and metabolic changes; and (6) increasing cancer risk.

Many bird species return to the same stopover, wintering, and breeding areas every year and often follow the exact same or very similar migration routes (Akesson & Hedenstrom, 2007). However, ample evidence exists that displaced birds can successfully reorient and find their way when one or more cues are removed. For example, Haftorn et al. (1988) found that after removal from their nests and release into a different area, snow petrels (*Pagodroma nivea*) were able to successfully navigate back to their nests even when their ability to smell was removed. Furthermore, Wiltschko et al. (2011) and Wiltschko and Wiltschko (2005) report that electromagnetic pulses administered to birds during an experimental study on orientation do not deactivate the magnetite-based receptor mechanism in the upper beak altogether but instead cause the receptors to provide altered information, which in turn causes birds to orient in different directions. However, these impacts were temporary, and the ability of the birds to correctly orient themselves eventually returned.

Given (1) the information provided above; (2) the dispersed nature of Navy training and testing activities at sea; (3) the relatively small area around an emitting source that experiences high power electromagnetic pulses; and (4) the relatively low-level and dispersed use of these systems at sea, the following conclusions are reached:

- The chance that in-air electromagnetic devices would cause thermal damage to an individual marine bird is extremely low.
- It is possible, although unlikely, that some marine bird individuals would be exposed to levels of electromagnetic radiation that would cause discomfort, in which case they would likely avoid the immediate vicinity of testing and training activities.
- The strength of any avoidance response would decrease with increasing distance from the in-air electromagnetic device.

- No long-term or population-level impacts would occur.

There is only one new activity involving an energy stressor (i.e., high-energy lasers) that differs from activities with energy stressors that were previously analyzed in the 2015 MITT Final EIS/OEIS. Use of low-energy lasers was covered in the 2015 MITT Final EIS/OEIS in Section 3.0.5.2.2.3 (Lasers), but high-energy laser weapons were not part of the proposed action in the 2015 MITT Final EIS/OEIS. The use of high-energy lasers represents a new substressor used in an existing activity in this SEIS/OEIS. As discussed in this SEIS/OEIS, Section 3.0.4.3.2.2 (High-Energy Lasers), high-energy lasers are designed to disable surface targets, rendering them immobile. The primary concern is the potential for a marine bird to be struck with the laser beam at or near the water's surface, where extended exposure could result in injury or death due to traumatic burns from the beam.

Marine birds could be exposed to a laser only if they flew between the source and the target, or if the beam missed the target and a bird happened to be in the line of fire. Should the laser strike the sea surface, individual sea birds at or near the surface could be exposed. Because laser platforms are typically helicopters and ships, marine birds at sea would likely transit away or submerge in response to other stressors, such as ship or aircraft noise, although some marine birds may not exhibit a response to an oncoming vessel or aircraft, increasing the risk of contact with the laser beam. High-energy laser activities would only occur in open ocean locations (not close to land areas).

#### **3.6.2.3.1 Impacts from High-Energy Lasers Under Alternative 1**

Under Alternative 1, the number of proposed activities involving the use of high-energy lasers is shown in Table 3.0-10. High-energy lasers is a new substressor that was not analyzed in the 2015 MITT Final EIS/OEIS. As discussed above, impacts on marine birds from energy stressors should not be expected to occur.

A direct strike of a marine bird at the water's surface or within the beam path is extremely unlikely, and potential impacts on ESA-listed marine bird species are negligible. Therefore, the conclusions for ESA-listed marine bird species and other marine bird species protected by the MBTA included in the 2015 MITT Final EIS/OEIS remain valid.

During Section 7 ESA consultation between the Navy and USFWS, the Navy determined that the activities described in the 2015 MITT Final EIS/OEIS would have no effect on the ESA-listed Hawaiian petrel, short-tailed albatross, or Newell's shearwaters.

*Pursuant to the ESA, the use of high-energy lasers during training and testing activities, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's Townsend's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from energy stressors during training and testing activities using high-energy lasers described under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.3.2 Impacts from High-Energy Lasers Under Alternative 2**

Under Alternative 2, the use of high-energy lasers would increase as compared to Alternative 1 (Table 3.0-10), but there would be no change regarding the impact conclusions for energy stressors as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore,

impacts on marine birds under Alternative 2 from energy stressors, including high-energy lasers, should not be expected to occur.

*Pursuant to the ESA, the use of high energy lasers during training and testing activities, as described under Alternative 2, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from energy stressors during training and testing activities using high-energy lasers described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

### **3.6.2.3.3 Impacts from High-Energy Lasers Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Energy stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities. Discontinuing the training and testing activities would result in fewer energy stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for impacts on marine birds from energy stressors, but would not measurably improve the overall distribution or abundance of marine birds.

### **3.6.2.4 Physical Disturbance and Strike Stressors**

The physical disturbance and strike stressors that may impact marine birds include (1) aircraft and aerial targets, (2) vessels and in-water devices, (3) military expended materials, and (4) wildfires on FDM. For activities occurring at sea, the use of aircraft and aerial targets, vessels and in-water devices, and military expended materials would decrease under this SEIS/OEIS (Tables 3.0-11 through 3.0-17, and Table 3.0-19), with the exception of increased small caliber munitions use. Small-caliber munitions are inert, are meant to be aimed at targets, and are not long-range weapons. As a result, marine birds are extremely unlikely to be struck by expended small caliber munitions. Military expended materials would increase for training activities occurring on FDM. For the purposes of this SEIS/OEIS, only activities that occur on or over FDM and activities that would occur in the open ocean environment that have changed since the publication of the 2015 MITT Final EIS/OEIS are discussed in this section.

Physical disturbances may elicit short-term behavioral or physiological responses such as alert response, startle response, cessation of feeding, fleeing the immediate area, and a temporary increase in heart rate. These disturbances can also result in abnormal behavioral, growth, or reproductive impacts in nesting birds and can cause foraging and nesting birds to flush from or abandon their habitats or nests. Aircraft strikes often result in bird mortalities or injuries.

Physical disturbance on land may induce erosion, either from loosening of rock and soil from direct impacts (which facilitates transport of material by wind and rain) or from wildfires ignited by explosions (see Section 3.6.2.4.3, Physical Disturbance and Strike Stressors [Impacts from Wildfires]). Military use of FDM may contribute to ongoing soil disturbance and erosion from natural causes. FDM is comprised of highly weathered limestone overlain by a thin layer of clay soil (U.S. Department of the Navy, 2013b). Ordnance use, particularly within Impact Areas 2 and 3 (where explosive munitions use is permitted), would dislodge sediments that may potentially wash into nearshore waters of FDM. In addition to

natural wind and water erosion (including high-energy typhoon events), erosion caused by ordnance use would contribute to increased turbidity and siltation of habitats used by marine bird prey species.

Section 3.6.3.3.1 (Impacts from Aircraft and Aerial Targets) in the 2015 MITT Final EIS/OEIS discusses the potential impacts on birds from collisions with fixed-wing aircraft, helicopters, and aerial targets. Aircraft and aerial target strikes could occur during training and testing activities that use aircraft, particularly in nearshore areas, where birds are more concentrated in the Study Area. Training and testing activities where aircraft are used typically occur further offshore; however, increased use of FDM may increase the potential for aircraft strike of birds. Therefore, for the purposes of this SEIS/OEIS, only the use of aircraft related to FDM training activities are discussed below under the alternatives analysis for birds for this stressor category.

#### **3.6.2.4.1 Impacts from Physical Disturbance and Strike Stressors Under Alternative 1**

Under Alternative 1, physical disturbance and strike stressors associated with training and testing activities would decrease in comparison to the 2015 MITT Final EIS/OEIS (Tables 3.0-12 through 3.0-18), assuming the dismissal of small-caliber munitions use for the reasons noted above. Under Alternative 1, there would be increases in the numbers of large-caliber non-explosive practice munitions (Table 3.0-14) and the number of targets expended at sea (Table 3.0-17), but overall there would be a decrease in the number of combined physical disturbance and strike stressors on marine birds. Consistent with the conclusions provided in the 2015 MITT EIS/OEIS, impacts on marine birds from physical disturbance and strike stressors are not expected to occur.

On FDM, marine birds that nest and roost on the island would be exposed to military expended materials resulting from explosive munitions and non-explosive practice munitions. Explosive munitions increase the potential for a marine bird (or nest) to be struck because of fragments dispersed throughout the blast zone.

As shown in Table 3.6-1, there would be an increase in the number of training events using FDM as a training location or target, with an increase in the number of munitions items expended on FDM (see Table 3.6-2).

Taken together, the increase in the number of training events per year or the amount of ordnance used during events would result in an increase in the amount of NEW expended on FDM each year (see Table 3.6-3). Although the amount of increased NEW is negligible (0.22 to 0.39 percent, depending on the NEW range of various munition types), the potential exposure to physical disturbance and strike stressors associated with ordnance use would increase under Alternative 1 compared to what was analyzed previously in the MITT Final EIS/OEIS. Factors that limit the potential for additional adverse impacts from physical disturbance and strike, however, include maintaining the same ordnance type and targeting restrictions included as part of the 2015 MITT Final EIS/OEIS. All ordnance expended on FDM would target existing impact zones, with the same ordnance restrictions imposed on all FDM activities and with the same avoidance and minimization measures in place (see Section 5.5, Terrestrial Mitigation Measures to be Implemented, and Table 5.5-1). Therefore, the increases in ordnance use on FDM shown in Tables 2.5-1 and 2.5-2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The conclusions for ESA-listed marine bird species and other marine bird species protected by the MBTA included in the 2015 MITT Final EIS/OEIS remain valid.

During Section 7 ESA consultation between the Navy and USFWS, the Navy determined that the activities described in the 2015 MITT Final EIS/OEIS would have no effect on the ESA-listed Hawaiian petrel, short-tailed albatross, or Newell's shearwater.

*Pursuant to the ESA, training and testing activities that use aircraft and aerial targets, as described under Alternative 1, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from physical disturbance and strike stressors during training and testing activities described under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.4.2 Impacts from Physical Disturbance and Strike Stressors Under Alternative 2**

Under Alternative 2, physical disturbance and strike stressors associated with training and testing activities would decrease in comparison to the 2015 MITT Final EIS/OEIS (Tables 3.0-12 through 3.0-18) assuming the dismissal of small-caliber munitions use for the reasons noted above. Under Alternative 2, there would be increases in the numbers of large-caliber non-explosive practice munitions (Table 3.0-14) and the number of targets expended at sea (Table 3.0-17), but overall there would be a decrease in the number of combined physical disturbance and strike stressors on marine birds. Consistent with the conclusions provided in the 2015 MITT EIS/OEIS, impacts on marine birds from physical disturbance and strike stressors are not expected to occur.

On FDM under Alternative 2, there would be an increase in the number of training events using FDM as a training location or target (see Table 3.6-1), with an increase in the number of munitions items expended on FDM (see Table 3.6-2) compared to what was analyzed previously in the MITT Final EIS/OEIS and under Alternative 1.

Taken together, the increase in the number of training events per year or the amount of ordnance used during events would result in an increase in the amount of NEW expended on FDM each year (see Table 3.6-3). Although the amount of increased NEW is negligible (0.35 to 0.6 percent, depending on the NEW range of various munition types), the potential exposure to stressors associated with ordnance use would increase under Alternative 2 compared to what was analyzed previously in the MITT Final EIS/OEIS. Under Alternative 2, Naval Surface Firing Exercise events would expend more large-caliber projectiles, thereby slightly increasing the NEW expended under Alternative 2 compared to Alternative 1. Factors that limit the potential for additional adverse impacts, however, include maintaining the same ordnance type and targeting restrictions included as part of the 2015 MITT Final EIS/OEIS. All ordnance expended on FDM would target existing impact zones, with the same ordnance restrictions imposed on all FDM activities and with the same avoidance and minimization measures in place (see Section 5.5, Terrestrial Mitigation Measures to be Implemented, and Table 5.5-1). Therefore, the increases in ordnance use on FDM shown in Tables 2.5-1 and 2.5-2 do not appreciably change the impact conclusions presented in the 2015 MITT Final EIS/OEIS. The conclusions for ESA-listed marine bird species and other marine bird species protected by the MBTA included in the 2015 MITT Final EIS/OEIS remain valid.

*Pursuant to the ESA, training and testing activities that use aircraft and aerial targets, as described under Alternative 2, would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), the impacts from physical disturbance and strike stressors during training and testing activities described under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.4.3 Impacts from Physical Disturbance and Strike Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Physical disturbance and strike stressors would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer physical disturbance and strike stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for physical disturbance and strike impacts on marine birds, but would not measurably improve the overall distribution or abundance of marine birds.

#### **3.6.2.5 Ingestion Stressors**

As discussed in Section 3.6.3.4 (Ingestion Stressors) of the 2015 MITT Final EIS/OEIS, a variety of ingestible materials may be released into the marine environment by training and testing activities. Types of materials that could become ingestion stressors (military expended materials – munitions and military expended materials other than munitions) for marine birds during training and testing activities in the Study Area include non-explosive practice munitions (small and medium caliber), fragments from high-explosive munitions, fragments from targets, chaff, plastic end caps from chaff cartridges, the plastic compression pads, end caps from pistons and flares, and small decelerators/parachutes. Ingestion stressors would decrease with the exception of increased small-caliber munitions use (Table 3.0-14, Table 3.0-15, Tables 3.0-22 through 3.0-24). However, small-caliber munitions are inert, small in size, do not resemble prey items, and end up as part of the seafloor where they are unlikely to be encountered by marine birds. The number of munitions and explosive munitions fragments that an individual marine bird could encounter would generally be low, based on the patchy distribution of both the munitions and open water feeding habitats of marine birds. In addition, it is assumed an animal would not ingest every munition or munition fragment it encountered, and if a munition or munition fragment were ingested, an animal may attempt to reject it when it realizes the item is not food.

##### **3.6.2.5.1 Impacts from Ingestion Stressors Under the Alternative 1**

Under Alternative 1, ingestion stressors would decrease, with the exception of increased small-caliber munitions use (Table 3.0-14, Table 3.0-15, Tables 3.0-22 through 3.0-24). For the reasons noted above, the Navy has determined that potential impacts from ingestion stressors would not be substantially different from the 2015 MITT Final EIS/OEIS. Military expended materials would be only a minute portion of the floating debris that marine birds could encounter and accidentally ingest. While military expended materials may be a contributing factor to the harmful effects of manmade debris on some marine birds, an individual military expended material would not negatively impact a marine bird. The overall likelihood that individual birds would be negatively impacted by ingestion of military expended materials in the Study Area under Alternative 1 is considered low, but not discountable. Population-level effects would be very unlikely given the relatively small quantities and limited persistence of military expended materials in habitats where birds are most likely to forage. Because of the extreme low likelihood of geographic or temporal overlap with training and testing activities with ESA-listed marine birds, potential ingestion of expended materials by ESA-listed marine birds is considered negligible. Therefore, the analysis from the 2015 MITT Final EIS/OEIS remains valid.

*Pursuant to the ESA, potential ingestion stressors introduced by training and testing activities under Alternative 1 would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), potential ingestion stressors introduced by training and testing activities under Alternative 1 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.5.2 Impacts from Ingestion Stressors Under Alternative 2**

Under Alternative 2, ingestion stressors (military expended materials – munitions and military expended materials other than munitions) would decrease under this SEIS/OEIS in comparison to the ongoing activities, with the exception of increased small-caliber munitions use (Table 3.0-14, Table 3.0-15, Tables 3.0-22 through 3.0-24). Under Alternative 2, increases as compared to Alternative 1 do not change the impact conclusions for ingestion stressors as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS. Therefore, impacts on marine birds from ingestion of military expended materials under Alternative 2 would be negligible.

*Pursuant to the ESA, potential ingestion stressors introduced by training and testing activities under Alternative 2 would have no effect on ESA-listed Hawaiian petrels, short-tailed albatrosses, Newell's shearwaters.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), potential ingestion stressors introduced by training and testing activities under Alternative 2 would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

#### **3.6.2.5.3 Impacts from Ingestion Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Ingestion stressors would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer ingestion stressors within the marine environment where Navy activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for ingestion impacts on marine birds, but would not measurably improve the overall distribution or abundance of marine birds.

#### **3.6.2.6 Secondary Stressors (Impacts on Habitat; Impacts on Prey Availability)**

The potential for secondary stressors, defined as potential impacts on habitat and prey availability, to impact marine bird species was analyzed in Section 3.6.3.5 (Secondary Stressors) of the 2015 MITT Final EIS/OEIS. Training and testing activities analyzed in this SEIS/OEIS would not introduce additional secondary stressors or change the impacts of secondary stressors on marine bird species from what was analyzed previously in the 2015 MITT Final EIS/OEIS.

*Pursuant to the ESA, secondary stressors associated with training or testing activities proposed in this SEIS/OEIS would have no effect on the ESA-listed Hawaiian petrel, Newell's shearwater, or short-tailed albatross.*

*Under the MBTA regulations applicable to military readiness activities (50 CFR Part 21), secondary stressors associated with training or testing activities would not result in a significant adverse effect on populations of the great frigatebird, masked booby, or other marine bird populations.*

### **3.6.2.7 Endangered Species Act Determinations**

Since the publication of the 2015 MITT Final EIS/OEIS, there have been no updates to the regulatory status, life history information, or species-specific threats that would alter the analysis from the 2015 MITT Final EIS/OEIS for the short-tailed albatross, Hawaiian petrel, or Newell's shearwater. As such, the description regarding these marine bird species presented in the 2015 MITT Final EIS/OEIS remains valid. Because of the limited period of time that ESA-listed marine bird species would be within the Study Area and the extreme unlikelihood that these birds would be subject to stressors generated by training and testing activities within the Study Area, the Navy and the USFWS have not included these species in past at-sea training and testing consultations within the Study Area (U.S. Fish and Wildlife Service, 2010, 2015). Similar to these past consultations, the activities proposed in Chapter 2 (Description of Proposed Action and Alternatives) of this SEIS/OEIS would have no effect on ESA-listed marine birds.

### **3.6.2.8 Migratory Bird Treaty Act**

The Navy has conducted an analysis of the potential impacts of increasing the number of events, munitions, and NEW expended on FDM. Taken together with the statistical analysis of bird trends on FDM described above in Section 3.6.2 (Environmental Consequences) and determinations that no significant population impacts would occur, the small increases in events, munitions numbers, and expended NEW on FDM proposed in this SEIS/OEIS would not significantly impact bird populations, as defined in the MBTA regulations applicable to military readiness activities (50 CFR Part 21). While this determination is applicable to all marine birds that occur in the Study Area, the Navy carried out a focused analysis for marine birds known to breed on FDM (see the discussion for population-level analysis in Section 3.6.2, Environmental Consequences). Pursuant with the Department of Defense's obligations under 50 CFR Part 21, the Department of Defense will continue to implement training restrictions on FDM (see Section 5.5, Terrestrial Mitigation Measures to be Implemented) and monitoring of bird populations on FDM.

### **3.6.3 Public Scoping Comments**

The public raised a number of issues during the scoping period in regard to marine birds. The issues are summarized in the list below.

- **Public comments regarding potential impacts of marine bird species on FDM** – This SEIS/OEIS includes an analysis of potential impacts from the additional training activities proposed to occur at FDM. For acoustic stressors, Section 3.6.2.1.3 (Impacts from Aircraft Noise) and Section 3.6.2.1.4 (Impacts from Weapon Noise) include an analysis of how these stressor types may impact marine bird rookeries on FDM. For explosives stressors, Section 3.6.2.2 (Explosives Stressors) includes an analysis of how the proposed increase in munitions for missile exercises and direct action training activities could impact marine birds on the island. Stressor categories within physical disturbance and strike stressors, such as potential strike impacts from aircraft and impacts from wildfires, also include an FDM-focused analysis. While assessing these

potential impacts of activities proposed in this SEIS/OEIS, it is important to note that all of the activities would continue under the same targeting constraints as described in the 2015 MITT Final EIS/OEIS. Mitigation measures designed in cooperation with U.S. Fish and Wildlife Service personnel provide a level of protection for the northern end of the island (where booby colonies have persisted), while ordnance use is only allowed in designated impact zones (see Section 5.5, Terrestrial Mitigation Measures to be Implemented).

- **Public comments regarding the status of nesting birds on FDM** – This SEIS/OEIS has been updated to include recent published work that provides a statistical review of repeated marine bird surveys on FDM. Camp et al. (2016) analyzed marine bird survey data collected from aerial surveys from 1997 to 2014. As discussed in Section 3.6.1.7 (Rookery Locations and Breeding Activities within the Mariana Islands), there is some evidence that masked and red-footed booby populations on FDM have declined, while brown booby populations have increased, though none of these trends were statistically significant. The general conclusion is that all three species exhibited population fluctuations over time. These fluctuations, combined with the level of variability observed in the count data, precluded any definitive conclusions about long-term population trends (i.e., the data were non-significant) (Camp et al., 2016). This SEIS/OEIS also includes historical observations and more recent surveys, such as Lusk et al. (2000), to provide context for the trend data and statistical analyses of FDM marine bird populations. Aerial surveys are conducted more frequently over FDM than on-the-ground surveys, with the primary focus to monitor marine bird rookeries (namely, brown boobies, masked boobies, and red-footed boobies). These surveys are described in more detail, along with quantitative trend analysis of populations in Section 3.6.2.6.3 (Farallon de Medinilla) of the 2015 MITT Final EIS/OEIS. All of these studies are summarized and included the Joint Region Marianas Integrated Natural Resources Management Plan (U.S. Department of the Navy, 2018), which is shared with cooperating agencies (e.g., Guam Department of Agriculture Division of Aquatic and Wildlife Resources, CNMI Department of Land and Natural Resources Division of Fish and Wildlife, and U.S. Fish and Wildlife Service Pacific Islands Fish and Wildlife Office).

This page intentionally left blank.

## REFERENCES

- Akesson, S., and A. Hedenstrom. (2007). How migrants get there: Migratory performance and orientation. *BioScience*, 57(2), 123–133.
- Baerwald, E. F., G. H. D'Amours, B. J. Klug, and R. M. Barclay. (2008). Barotrauma is a significant cause of bat fatalities at wind turbines. *Current Biology*, 18(16), R695–R696.
- Barnas, A., R. Newman, C. J. Felege, M. P. Corcoran, S. D. Hervey, T. J. Stechmann, R. F. Rockwell, and S. N. Ellis-Felege. (2018). Evaluating behavioral responses of nesting lesser snow geese to unmanned aircraft surveys. *Ecology and Evolution*, 8, 1328–1338.
- Beason, R. (2004). *What Can Birds Hear?* Lincoln, NE: University of Nebraska.
- Beuter, K. J., R. Weiss, and B. Frankfurt. (1986). *Properties of the auditory system in birds and the effectiveness of acoustic scaring signals*. Paper presented at the Bird Strike Committee Europe, 18th Meeting Part I, 26–30 May 1986. Copenhagen, Denmark.
- Black, A. (2005). Short note: Light induced seabird mortality on vessels operating in the Southern Ocean: Incidents and mitigation measures. *Antarctic Science*, 17(1), 67–68.
- Bowles, A. E. (1995). Chapter 8: Responses of Wildlife to Noise. In R. L. Knight & K. J. Gutzwiller (Eds.), *Wildlife and Recreationists: Coexistence Through Management and Research*. Washington, DC: Island Press.
- Camp, R., C. Leopold, K. Brinck, and F. Juola. (2016). *Farallon de Medinilla seabird and Tinian moorhen analyses*. Hilo, HI: Hawaii Cooperative Studies Unit, University of Hawaii at Hilo.
- Cook, T. R., M. Hamann, L. Pichegru, F. Bonadonna, D. Grémillet, and P. G. Ryan. (2011). GPS and time-depth loggers reveal underwater foraging plasticity in a flying diver, the Cape Cormorant. *Marine Biology*, 159(2), 373–387.
- Crowell, S. C. (2016). Measuring in-air and underwater hearing in seabirds. *Advances in Experimental Medicine and Biology*, 875, 1155–1160.
- Crowell, S. E., A. M. Wells-Berlin, C. E. Carr, G. H. Olsen, R. E. Therrien, S. E. Ynuzzzi, and D. R. Ketten. (2015). A comparison of auditory brainstem responses across diving bird species. *Journal of Comparative Physiology A*, 201(8), 803–815.
- Crowell, S. E., A. M. Wells-Berlin, R. E. Therrien, S. E. Ynuzzzi, and C. E. Carr. (2016). In-air hearing of a diving duck: A comparison of psychoacoustic and auditory brainstem response thresholds. *The Journal of the Acoustical Society of America*, 139(5), 3001.
- Dooling, R. (2002). *Avian Hearing and the Avoidance of Wind Turbines*. College Park, MD: University of Maryland.
- Dooling, R. J., and A. N. Popper. (2000). Hearing in birds and reptiles. In R. J. Dooling, R. R. Fay, & A. N. Popper (Eds.), *Comparative Hearing in Birds and Reptiles* (Vol. 13, pp. 308–359). New York, NY: Springer-Verlag.
- Dooling, R. J., and S. C. Therrien. (2012). Hearing in birds: What changes from air to water. *Advances in Experimental Medicine and Biology*, 730, 77–82.
- Duffy, D. C. (2011). No room in the ark? Climate change and biodiversity in the Pacific Islands of Oceania. *Pacific Conservation Biology*, 17, 192–200.

- Frost, M. T., S. Dye, B. Stoker, P. Buckley, and J. Baxter. (2017). *Marine Climate Change Impacts: 10 Years' Experience of Science to Policy Reporting. Summary Report*. London, United Kingdom: Marine Climate Change Impacts Partnership.
- Haftorn, S., F. Mehlum, and C. Bech. (1988). Navigation to nest site in the snow petrel (*Pagodroma nivea*). *The Condor*, 90(2), 484–486.
- Hansen, K. A., A. Maxwell, U. Siebert, O. N. Larsen, and M. Wahlberg. (2017). Great cormorants (*Phalacrocorax carbo*) can detect auditory cues while diving. *The Science of Nature*, 104(5–6), 45.
- Hetherington, T. (2008). Comparative anatomy and function of hearing in aquatic amphibians, reptiles, and birds. In J. G. M. Thewissen & S. Nummela (Eds.), *Sensory Evolution on the Threshold* (pp. 182–209). Berkeley, CA: University of California Press.
- Jiménez, S., A. Domingo, M. Abreu, and A. Brazeiro. (2012). Bycatch susceptibility in pelagic longline fisheries: Are albatrosses affected by the diving behaviour of medium-sized petrels? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22(4), 436–445.
- Johansen, S., O. N. Larsen, J. Christensen-Dalsgaard, L. Seidelin, T. Huulvej, K. Jensen, S. G. Lunneryd, M. Bostrom, and M. Wahlberg. (2016). In-air and underwater hearing in the great cormorant (*Phalacrocorax carbo sinensis*). *Advances in Experimental Medicine Biology*, 875, 505–512.
- Jongbloed, R. H. (2016). *Flight height of seabirds. A literature study*. Ijmuiden, Netherlands: Institute for Marine Resources & Ecosystem Studies.
- Kushlan, J. A. (1978). Non-rigorous foraging by robbing egrets. *Ecology*, 59, 649–653.
- Larkin, R. P., L. L. Pater, and D. J. Tazlk. (1996). *Effects of Military Noise on Wildlife: A Literature Review* (USACERL Technical Report 96/21). Champaign, IL: Department of the Army, Construction Engineering Research Lab.
- Lincoln, F. C., S. R. Perterson, and J. L. Zimmerman. (1998). *Migration of Birds* (Migration of Birds Circular 16). Manhattan, KS: U.S. Department of the Interior, U.S. Fish & Wildlife Service.
- Lorrain, A., F. Houlbrèque, F. Benzoni, L. Barjon, L. Tremblay-Boyer, C. Menkes, D. P. Gillikin, C. Payri, H. Jourdan, and G. Boussarie. (2017). Seabirds supply nitrogen to reef-building corals on remote Pacific islets. *Scientific Reports*, 7(1), 3721.
- Lusk, M. R., P. Bruner, and C. Kessler. (2000). The Avifauna of Farallon De Medinilla, Mariana Islands. *Journal of Field Ornithology*, 71(1), 22–33.
- Manville, A. (2016). *A Briefing Memorandum: What We Know, Can Infer, and Don't Yet Know about Impacts from Thermal and Non-thermal Non-ionizing Radiation to Birds and Other Wildlife—for Public Release*. Washington, DC: U.S. Fish and Wildlife Service.
- Maxwell, A., K. A. Hansen, S. T. Ortiz, O. N. Larsen, U. Siebert, and M. Wahlberg. (2017). In-air hearing of the great cormorant (*Phalacrocorax carbo*). *Biology Open*, 6(4), 496–502.
- National Park Service. (1994). *Report on Effects of Aircraft Overflights on the National Park System* (Report to Congress prepared pursuant to Public Law 100-191, the National Parks Overflights Act of 1987). Washington, DC: National Park Service.
- Ostrom, P. H., A. E. Wiley, H. F. James, S. Rossman, W. A. Walker, E. F. Zipkin, and Y. Chikaraishi. (2017). Broad-scale trophic shift in the pelagic North Pacific revealed by an oceanic seabird. *Proceedings of the Royal Society B*, 284(1851), 20162436.

- Pater, L. L., and J. W. Shea. (1981). *Techniques for Reducing Gun Blast Noise Levels: An Experimental Study*. Dahlgren, VA: Naval Surface Weapons System.
- Plumpton, D. (2006). *Review of Studies Related to Aircraft Noise Disturbance of Waterfowl: A Technical Report in Support of the Supplemental Environmental Impact Statement for Introduction of F/A-18 E/F (Super Hornet) Aircraft to the East Coast of the United States*. Norfolk, VA: U.S. Department of the Navy.
- Poessel, S. A., A. E. Duerr, J. C. Hall, M. A. Braham, and T. E. Katzner. (2018). Improving estimation of flight altitude in wildlife telemetry studies. *Journal of Applied Ecology*, 55(4), 2064–2070.
- Ponganis, P. (2015). *Diving Physiology of Marine Mammals and Seabirds*. Cambridge, United Kingdom: Cambridge University Press.
- Ramírez, F., I. Afán, L. S. Davis, and A. Chiaradia. (2017). Climate impacts on global hot spots of marine biodiversity. *Science Advances*, 3(2), e1601198.
- Rijke, A. M. (1970). Wettability and phylogenetic development of feather structure in water birds. *The Journal of Experimental Biology*, 52(2), 469–479.
- Sade, J., Y. Handrich, J. Bernheim, and D. Cohen. (2008). Pressure equilibration in the penguin middle ear. *Acta Oto-Laryngologica*, 128(1), 18–21.
- Sanzenbacher, P. M., B. A. Cooper, J. H. Plissner, and J. Bond. (2014). Intra-annual patterns in passage rates and flight altitudes of Marbled Murrelets *Brachyramphus marmoratus* at inland sites in Northern California. *Marine Ornithology*, 42, 169–174.
- Saunders, J. C., and R. Dooling. (1974). Noise-induced threshold shift in the parakeet (*Melopsittacus undulatus*). *Proceedings of the National Academy of Sciences*, 71(5), 1962–1965.
- Sibley, D. (2014). *The Sibley Guide to Birds* (Second ed.). New York, NY: Alfred A. Knopf.
- Spiegel, C., A. Berlin, A. Gilbert, C. E. Gray, W. Montevecchi, I. Stenhouse, S. Ford, G. H. Olsen, J. Fiely, and L. Savoy. (2017). *Determining Fine-scale Use and Movement Patterns of Diving Bird Species in Federal Waters of the Mid-Atlantic United States Using Satellite Telemetry*. Sterling, VA: US Department of the Interior, Bureau of Ocean Energy Management.
- Stumpf, J. P., N. Denis, T. E. Hamer, G. Johnson, and J. Verschuyl. (2011). Flight height distribution and collision risk of the marbled murrelet *Brachyramphus marmoratus*: Methodology and preliminary results. *Marine Ornithology*, 39, 123–128.
- Tarroux, A., H. Weimerskirch, S.-H. Wang, D. H. Bromwich, Y. Cherel, A. Kato, Y. Ropert-Coudert, Ø. Varpe, N. G. Yoccoz, and S. Descamps. (2016). Flexible flight response to challenging wind conditions in a commuting Antarctic seabird: Do you catch the drift? *Animal Behaviour*, 113, 99–112.
- Therrien, S. C. (2014). *In-air and underwater hearing of diving birds*. (Unpublished doctoral dissertation). University of Maryland, College Park, MD. Retrieved from <http://hdl.handle.net/1903/2>.
- Thiessen, G. J. (1958). Threshold of hearing of a ring-billed gull. *The Journal of the Acoustical Society of America*, 30(11), 1047.
- Trainer, V. L. (2017). *Conditions Promoting Extreme Pseudo-nitzschia Events in the Eastern Pacific but not the Western Pacific* (PICES Scientific Report No. 53, 2017). Sidney, Canada: North Pacific Marine Science Organization Science Board, Marine Environmental Quality Committee.

- U.S. Department of the Navy. (2007). *Marine Mammal and Sea Turtle Survey and Density Estimates for Guam and the Commonwealth of the Northern Mariana Islands - Final Report*. Newport Beach, CA: Naval Facilities Engineering Command Pacific and Commander, U.S. Pacific Fleet.
- U.S. Department of the Navy. (2011). *Marianas Training Manual*. Naval Base Guam, Guam: Commander Joint Region Marianas.
- U.S. Department of the Navy. (2013a). *Annual Report: Wildlife Surveys on Tinian and FDM*. Naval Base Guam, Guam: Joint Region Marianas.
- U.S. Department of the Navy. (2013b). *2013 Joint Region Marianas Integrated Natural Resources Management Plan (Final)*. Naval Base Guam, Guam: Joint Region Marianas.
- U.S. Department of the Navy. (2015). *Final Mariana Islands Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, HI: Naval Facilities Engineering Command, Pacific.
- U.S. Department of the Navy. (2018). *Joint Region Mariana Islands Integrated Natural Resources Management Plan*. Saipan, Mariana Islands: Joint Region Mariana Islands.
- U.S. Fish and Wildlife Service. (2010). *Biological Opinion for the Mariana Islands Range Complex, Guam and the Commonwealth of the Northern Mariana Islands 2010–2015*. Honolulu, HI: U.S. Fish & Wildlife Service.
- U.S. Fish and Wildlife Service. (2015). *Biological Opinion for the Mariana Islands Training and Testing Program*. Honolulu, HI: U.S. Fish and Wildlife Service Pacific Islands Fish and Wildlife Office.
- Wever, E. G., P. N. Herman, J. A. Simmons, and D. R. Hertzler. (1969). Hearing in the blackfooted penguin (*Spheniscus demersus*), as represented by the cochlear potentials. *Proceedings of the National Academy of Sciences*, 63, 676–680.
- Wiltschko, R., S. Denzau, D. Gehring, P. Thalau, and W. Wiltschko. (2011). Magnetic orientation of migratory robins, *Erithacus rubecula*, under long-wavelength light. *The Journal of Experimental Biology*, 214(18), 3096–3101.
- Wiltschko, W., and R. Wiltschko. (2005). Magnetic orientation and magnetoreception in birds and other animals. *Journal of Comparative Physiology A* 191(8), 675–693.

---

---

## 3.7 Marine Vegetation



**Supplemental Environmental Impact Statement/  
Overseas Environmental Impact Statement  
Mariana Islands Training and Testing**

**TABLE OF CONTENTS**

|         |   |       |
|---------|---|-------|
| 3.7     | Marine Vegetation .....                         | 3.7-1 |
| 3.7.1   | Affected Environment.....                       | 3.7-1 |
| 3.7.1.1 | General Threats .....                           | 3.7-1 |
| 3.7.1.2 | Marine Vegetation Groups .....                  | 3.7-1 |
| 3.7.1.3 | Seagrasses.....                                 | 3.7-1 |
| 3.7.1.4 | Mangroves .....                                 | 3.7-1 |
| 3.7.2   | Environmental Consequences .....                | 3.7-1 |
| 3.7.2.1 | Explosive Stressors.....                        | 3.7-2 |
| 3.7.2.2 | Physical Disturbance and Strike Stressors ..... | 3.7-3 |
| 3.7.2.3 | Secondary Stressors.....                        | 3.7-5 |
| 3.7.3   | Public Scoping Comments .....                   | 3.7-5 |

**List of Figures**

There are no figures in this section.

**List of Tables**

There are no tables in this section.

This page intentionally left blank.

### **3.7 Marine Vegetation**

#### **3.7.1 Affected Environment**

The purpose of this section is to supplement the analysis of impacts on Marine Vegetation presented in the 2015 Mariana Islands Training and Testing (MITT) Final Environmental Impact Statement/Overseas Environmental Impact Statement (EIS/OEIS) with new information relevant to proposed changes in training and testing activities conducted at sea and on Farallon de Medinilla (FDM). New information made available since the publication of the 2015 MITT Final EIS/OEIS is included below to better understand potential stressors and impacts on Marine Vegetation resulting from training and testing activities. Comments received from the public during scoping related to Marine Vegetation are addressed in Section 3.7.3 (Public Scoping Comments).

##### **3.7.1.1 General Threats**

There is no new information on threats to marine vegetation in the MITT Study Area that would change the conclusions from the 2015 MITT Final EIS/OEIS.

##### **3.7.1.2 Marine Vegetation Groups**

There is no new information on marine vegetation groups (phylum Cyanobacteria [blue-green algae], phylum Dinophyta [dinoflagellates], phylum Chlorophyta [green algae], phylum Heterokontophyta [brown algae], phylum Rhodophyta [red algae], and phylum Spermatophyta [flowering plants]) that would change the basis of the conclusions from the 2015 MITT Final EIS/OEIS.

##### **3.7.1.3 Seagrasses**

There is no new information on seagrasses that would change the basis of the conclusions from the 2015 MITT Final EIS/OEIS.

##### **3.7.1.4 Mangroves**

There is no new information on mangroves that would change the basis of the conclusions from the 2015 MITT Final EIS/OEIS.

#### **3.7.2 Environmental Consequences**

The 2015 MITT Final EIS/OEIS considered training and testing activities that currently occur in the Study Area and considered potential stressors related to marine vegetation. With the exception of explosives, stressors analyzed are the same as those analyzed in the 2015 MITT Final EIS/OEIS. In the 2015 MITT Final EIS/OEIS, explosives were addressed under acoustic stressors; however, for purposes of this analysis, explosives are analyzed as a separate stressor. The following are stressors analyzed for marine vegetation from the 2015 MITT Final EIS/OEIS:

- Explosive (in-air explosions and in-water explosions)
- Physical disturbance and strike (vessels, in-water devices, military expended materials, and seafloor devices)
- Secondary stressors (impacts associated with sediments and water quality)

This section evaluates how and to what degree potential impacts on marine vegetation from stressors described in Section 3.0 (General Approach to Analysis) may have changed since the analysis presented in the 2015 MITT Final EIS/OEIS was completed. Proposed training and testing activities, the number of times each activity would be conducted annually, and the locations within the Study Area where the

activity would typically occur under each alternative are presented in Tables 2.4-1 and 2.4-2 in Chapter 2 (Description of Proposed Action and Alternatives). The tables also present the same information for activities described in the 2015 MITT Final EIS/OEIS so that the proposed levels of training and testing under this Supplemental EIS (SEIS)/OEIS can be easily compared.

The Navy conducted a review of federal and state regulations and standards relevant to marine vegetation and reviewed literature published since 2015 for new information that could inform the analysis presented in the 2015 MITT Final EIS/OEIS. The analysis presented in this section also considers standard operating procedures, which are discussed in Section 2.3.3 (Standard Operating Procedures) of this Draft SEIS/OEIS, and mitigation measures that are described in Chapter 5 (Mitigation). The Navy implements these measures to avoid or reduce potential impacts on marine vegetation from stressors associated with training and testing activities.

### **3.7.2.1 Explosive Stressors**

As stated in the 2015 MITT Final EIS/OEIS, the potential for an explosion to injure or destroy marine vegetation would depend on the amount of vegetation present, the number of munitions used, and their net explosive weight. In areas where marine vegetation and locations for explosions overlap, marine vegetation on the surface of the water, in the water column, or rooted in the seafloor may be impacted. Seafloor macroalgae and single-celled algae may overlap with underwater and sea surface explosion locations. If these vegetation types are near an explosion, only a small number of them are likely to be impacted. Much of the attached macroalgae grows on live hard bottom areas that would be mostly protected in accordance with Navy mitigation measures (see Chapter 5, Mitigation). Also, some seafloor macroalgae are resilient to high levels of wave action (Mach et al., 2007), which may aid in their ability to withstand underwater explosions that occur near them. Underwater explosions also may temporarily increase the turbidity (sediment suspended in the water) in nearby waters, incrementally reducing the amount of light available to marine vegetation. Reducing light availability decreases, albeit temporarily, the photosynthetic ability of marine vegetation.

Seagrasses may potentially be uprooted or damaged by sea surface or underwater explosions. Regrowth of seagrasses after uprooting can take up to 10 years (Dawes et al., 1997). Explosions may also temporarily increase the turbidity (sediment suspended in the water) in nearby waters, but the sediment would settle to pre-explosion conditions within a few hours to days. Sustained high levels of turbidity may reduce the amount of light that reaches vegetation, which it needs to survive. Seagrasses typically grow in waters that are sheltered from wave action, such as estuaries, lagoons, and bays (Phillips & Meñez, 1988), where most activities are not conducted. Detonations are unlikely to occur in areas with mangroves or sea grasses and would continue to occur in disturbed areas over the unvegetated seafloor such as the Agat Bay site, Piti, and Outer Apra Harbor sites.

#### **3.7.2.1.1 Impacts from Explosive Stressors Under Alternative 1**

Under Alternative 1, there would be an overall decrease in the number of explosives used in the Study Area during training and testing activities events compared to the number analyzed in the 2015 MITT Final EIS/OEIS (Table 3.0-7). Under Alternative 1, underwater detonations would increase for underwater demolition qualification/certification (Table 2.4-1). However, these activities would continue to occur in the same areas and would have no appreciable change in the impact analysis or conclusions for explosive stressors as presented in the 2015 MITT Final EIS/OEIS. Therefore, the analysis in the 2015 MITT Final EIS/OEIS remains valid.

As described in the 2015 MITT Final EIS/OEIS, underwater explosions conducted for training and testing activities may destroy or remove marine vegetation. However, exposure to these detonations would be limited to the vicinity of the explosions. For example, the offshore underwater mine neutralization sites are located in areas with water depths that are unlikely for marine vegetation to occur. Underwater and surface explosions conducted for training and testing activities are not expected to pose a risk to seagrass because (1) the impact area of underwater explosions is very small relative to seagrass distribution and (2) the low number of charges reduces the potential for impacts.

Therefore, the use of explosives is not expected to impact the long-term survival, annual reproductive success, and lifetime reproductive success of marine vegetation.

#### **3.7.2.1.2 Impacts from Explosive Stressors Under Alternative 2**

Under Alternative 2, the number of explosives used during training and testing activities would decrease compared to the numbers analyzed in the 2015 MITT Final EIS/OEIS and increase compared to Alternative 1 (Table 3.0-7). Under Alternative 2, increases in the number of underwater explosives would have no appreciable change on the impact conclusions for explosive stressors as summarized above under Alternative 1 and as presented in the 2015 MITT Final EIS/OEIS.

Therefore, explosive impacts on marine vegetation under Alternative 2 would be negligible.

#### **3.7.2.1.3 Impacts from Explosive Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Explosive stressors as listed above would not be introduced into the marine environment. Therefore, existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer explosive stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for explosive impacts on marine vegetation, but would not measurably improve the overall distribution or abundance of marine vegetation.

#### **3.7.2.2 Physical Disturbance and Strike Stressors**

This section analyzes the potential impacts on marine vegetation of the various types of physical disturbance and strike stressors during training and testing activities within the Study Area. Three types of physical disturbance and strike stressors are evaluated for their impacts on marine vegetation, including (1) vessels and in-water devices, (2) military expended materials, and (3) seafloor devices.

The evaluation of the impacts of physical disturbance stressors on marine vegetation focuses on proposed activities that may cause vegetation to be damaged by an object that is moving through the water (e.g., vessels and in-water devices), or dropped to the seafloor (e.g., military expended materials, anchors). Not all activities are proposed throughout the Study Area. Wherever appropriate, specific geographic areas of potential impact are identified.

As described in the 2015 MITT Final EIS/OEIS, vessel disturbance of marine vegetation would be limited to floating marine algae. Vessel movements may disperse or injure algal mats. Because algal distribution is patchy and mats may re-form following a disturbance, training and testing activities involving vessel movement would not impact the general health of marine algae.

### **3.7.2.2.1 Impacts from Physical Disturbance and Strike Stressors Under Alternative 1**

Under Alternative 1, the number of proposed training and testing events involving vessel movements would increase from those presented in the 2015 MITT Final EIS/OEIS (Table 3.0-12). In contrast, the use of towed in-water devices (Table 3.0-13) would decrease. The decrease in the number of in-water devices is unlikely to change the impact conclusion presented in the 2015 MITT Final EIS/OEIS. As stated in the 2015 MITT Final EIS/OEIS, the impact of vessels and in-water devices on marine vegetation would remain inconsequential because of (1) the quick recovery of most vegetation types; (2) the short-term nature of most vessel movements and local disturbances of the surface water, with some temporary increase in suspended sediment in shallow areas; and (3) the deployment of in-water devices at depths where they would not likely come in contact with marine vegetation.

Under Alternative 1, the number of military expended materials used for training and testing activities that has the potential to impact marine vegetation would generally increase (see Tables 3.0-14 through 3.0-17). However, these increases are not expected to pose a risk to marine algae or seagrasses because (1) the relative coverage of marine algae in the Study Area is low, (2) new growth may result from marine algae exposure to military expended materials, (3) the impact area of military expended materials is very small relative to marine algae distribution, and (4) seagrass overlap with areas where the stressor occurs is very limited. Based on these factors, potential impacts on marine algae and seagrass from military expended materials are not expected to result in detectable changes in their growth, survival, or propagation, and are not expected to result in population-level impacts.

Under Alternative 1, the number of seafloor devices used in shallow-water habitats during training and testing activities would decrease slightly from the number presented in the 2015 MITT Final EIS/OEIS (Table 3.0-18). Seafloor devices would pose a negligible risk to marine vegetation for the same reasons described above for military expended materials and no impacts on the long-term survival, reproductive success, and lifetime reproductive success would occur.

Therefore, physical disturbance and strike impacts on marine vegetation under Alternative 1 would be negligible.

### **3.7.2.2.2 Impacts from Physical Disturbance and Strike Stressors Under Alternative 2**

Under Alternative 2, the combined number of proposed training and testing events involving vessels and in-water devices (Table 3.0-12 and Table 3.0-13) would decrease slightly from those presented in the 2015 MITT Final EIS/OEIS. Military expended materials (Table 3.0-14, Table 3.0-15, and Table 3.0-16) combined would increase, and seafloor devices (Table 3.0-18) would decrease slightly from the number in the 2015 MITT Final EIS/OEIS. Increases in some physical disturbance and strike stressors such as military expended materials could increase the impact risk on marine vegetation but does not appreciably change the analysis or impact conclusions presented in the 2015 MITT Final EIS/OEIS and those summarized above under Alternative 1.

Therefore, physical disturbance and strike impacts on marine vegetation under Alternative 2 would be negligible.

### **3.7.2.2.3 Impacts from Physical Disturbance and Strike Stressors Under the No Action Alternative**

Under the No Action Alternative, proposed training and testing activities would not occur. Other military activities not associated with this Proposed Action would continue to occur. Physical disturbance and strike stressors as listed above would not be introduced into the marine environment. Therefore,

existing environmental conditions would either remain unchanged or would improve slightly after cessation of ongoing training and testing activities.

Discontinuing the training and testing activities would result in fewer physical disturbance and strike stressors within the marine environment where training and testing activities have historically been conducted. Therefore, discontinuing training and testing activities under the No Action Alternative would lessen the potential for physical disturbance and strike impacts on marine vegetation, but would not measurably improve the overall distribution or abundance of marine vegetation.

### 3.7.2.3 Secondary Stressors

Stressors from Navy training and testing activities could pose secondary or indirect impacts on marine vegetation via habitat, sediment, or water quality. Potential impacts on marine vegetation exposed to secondary stressors could occur indirectly through sediments and water quality. Components of these stressors that could pose indirect impacts include (1) explosives and byproducts; (2) metals; (3) chemicals; and (4) other materials such as targets, chaff, and plastics.

Section 3.1 (Sediments and Water Quality) considered the impacts on marine sediments and water quality from explosives and explosive byproducts, metals, chemicals other than explosives, and other materials (marine markers, flares, chaff, targets, and miscellaneous components of other materials). As stated in the 2015 MITT Final EIS/OEIS, one example of a localized impact on marine vegetation associated with water quality impacts could be the increase of cyanobacteria associated with munitions deposits in marine sediments. Cyanobacteria may proliferate when the iron is introduced to the marine environment, and this proliferation can affect surrounding habitats by releasing toxins or stimulating the growth of nuisance species (Schils, 2012). Introducing iron into the marine environment from munitions or infrastructure is not associated with red tide events; rather, these harmful events are more associated with natural causes (e.g., upwellings) and the effects of human activities (e.g., agricultural runoff and other coastal pollution) (Hayes et al., 2007; Whitton & Potts, 2008).

Sediments entering the nearshore environment from FDM as a result of natural processes or explosives associated with strike warfare could cause temporary water quality impacts, some of which may be in foraging areas used by marine organisms. By limiting the location and extent of target areas, along with the types of ordnance allowed within specific impact areas, the military minimizes the potential for soil transport and, thus, water quality impacts. Erosion as a result of training activities at FDM may contribute to deposition of soils into the nearshore areas of FDM, causing increased turbidity. Turbidity can impact vegetation communities by reducing the amount of light that reaches these organisms. The impacts of explosive byproducts on sediment and water quality would be indirect, short term, and local. Explosive ordnance could loosen the soil on FDM and runoff from surface drainage areas containing soil, and explosive byproducts could contaminate sediments and the surrounding ocean water.

### 3.7.3 Public Scoping Comments

The public raised a number of issues during the scoping period in regard to marine vegetation. The issues are summarized in the list below.

- **Direct impacts on seagrass from sedimentation around FDM and military expended materials as marine debris** – Direct impacts on seagrass from sedimentation around FDM occur due to explosive stressors. Explosives may temporarily increase the turbidity (sediment suspended in the water) of nearby waters, but the sediment would settle to pre-explosion conditions within a short amount of time (e.g., a few hours to days). Sustained high levels of turbidity may reduce

the amount of light that reaches vegetation, which it needs to survive. This scenario is not likely given the low number of explosions planned in areas with seagrass. Potential impacts on seagrass from military expended materials are not expected to result in detectable changes in their growth, survival, or propagation, and are not expected to result in population-level impacts. See Section 3.7.2.1 (Explosive Stressors) for further analysis of increased turbidity or sedimentation on marine vegetation including seagrasses in the Study Area including FDM. Military expended materials are discussed in Section 3.7.2.2 (Physical Disturbance and Strike Stressors) as a cause of physical disturbance and strike to marine vegetation.

- **Request survey of all seagrass beds in the Study Area and monitoring of the seagrass beds –** The analysis of impacts on marine vegetation, including seagrasses, concluded that increased turbidity may be caused by items used in training and testing activities; under the standard operating procedures, the Navy avoids the seafloor to the greatest extent practicable. Additionally, activities that have a greater potential to impact the seafloor, such as amphibious assaults, are conducted at high tide to limit such interactions. Anchorages are also scheduled to occur in specific locations, mainly areas that lack vegetation and that have been previously disturbed. Therefore, serious damage is not anticipated, and survey or mitigation measures are not warranted. In addition, the 2015 MITT Final EIS/OEIS includes maps showing areas of marine vegetation in Section 3.7 (Marine Vegetation).
- **Impact of unexploded ordnance on marine species –** Potential impacts on marine vegetation from unexploded ordnance are not expected to result in detectable changes in their growth, survival, or propagation, and are not expected to result in population-level impacts. The impact of unexploded ordnance to marine species, specifically to marine vegetation, is discussed in Section 3.7.2.2 (Physical Disturbance and Strike Stressors) as a cause of physical disturbance and strike to marine vegetation.
- **Impacts on marine species from chemical pollution and destruction of habitat –** The analysis concluded that neither state nor federal standards or guidelines for sediments or water quality would be violated as a result of the implementation of the proposed training and testing activities. Therefore, because these standards and guidelines are structured to protect human health and the environment, and the proposed activities do not violate them, no indirect impacts are anticipated on marine vegetation from the training and testing activities proposed in this SEIS/OEIS. Destruction of habitat is not anticipated to result from the implementation of training and testing activities proposed in this SEIS/OEIS. Impacts on marine species, specifically to marine vegetation from chemical pollution, is discussed in Section 3.7.2.5 (Secondary Stressors).
- **Impacts on marine species from the metals in the water (copper and lead) –** The analysis concluded that neither state nor federal standards or guidelines for sediments or water quality would be violated as a result of the implementation of the proposed training and testing activities. Therefore, because these standards and guidelines are structured to protect human health and the environment, and the proposed activities do not violate them, no indirect impacts are anticipated on marine vegetation from the training and testing activities proposed in this SEIS/OEIS. Impacts on marine species, specifically on marine vegetation, from metals in the water (such as copper and lead) are discussed in Section 3.7.2.3 (Secondary Stressors).

- **Deposition and resuspension of sediments to EFH from training activities** – The 2015 MITT EFHA concluded that any impacts from explosives or physical disturbance and strike stressors that could cause deposition or resuspension of sediments would be short term and minimal.
- **Erosion and sedimentation impacting EFH** – The 2015 MITT EFHA concluded that any impacts from explosives or physical disturbance and strike stressors that could cause erosion and sedimentation would be short term and minimal.
- **Unexploded ordnance being triggered after use and directly impacting EFH** – Unexploded ordnance that explodes due to being triggered post training and testing would be considered an explosive stressor and the 2015 MITT EFHA concluded that the impacts on attached macroalgae from explosives used during training and testing would be minimal and temporary to short term throughout the Study Area. This analysis remains valid. Given the available information, the impact of explosives used during training and testing on submerged rooted vegetation beds would be minimal.

## REFERENCES

- Dawes, C. J., J. Andorfer, C. Rose, C. Uranowski, and N. Ehringer. (1997). Regrowth of the seagrass, *Thalassia testudinum*, into propeller scars. *Aquatic Botany*, 59(1–2), 139–155.
- Hayes, P. K., N. A. El Semary, and P. Sanchez-Baracaldo. (2007). The taxonomy of cyanobacteria: Molecular insights into a difficult problem. In J. Brodie & J. Lewis (Eds.), *Unravelling the Algae: The Past, Present, and Future of Algal Systematics* (pp. 93–102). Boca Raton, FL: CRC Press.
- Mach, K. J., B. B. Hale, M. W. Denny, and D. V. Nelson. (2007). Death by small forces: A fracture and fatigue analysis of wave-swept macroalgae. *The Journal of Experimental Biology*, 210(13), 2231–2243.
- Phillips, R. C., and E. G. Meñez. (1988). Seagrasses. *Smithsonian Contributions to the Marine Sciences*, 34, 104.
- Schils, T. (2012). Episodic eruptions of volcanic ash trigger a reversible cascade of nuisance species outbreaks in pristine coral habitats. *PLoS ONE*, 7: e46639.
- Whitton, B. A., and M. Potts. (2008). *The Ecology of Cyanobacteria: Their Diversity in Time and Space*. The Netherlands: Kluwer Academic Publishers.